



Journal de l'École polytechnique

Mathématiques

Giovanni FORNI & Adam KANIGOWSKI

Multiple mixing and disjointness for time changes of bounded-type Heisenberg nilflows

Tome 7 (2020), p. 63-91.

<http://jep.centre-mersenne.org/item/JEP_2020__7__63_0>

© Les auteurs, 2020.

Certains droits réservés.



Cet article est mis à disposition selon les termes de la licence
LICENCE INTERNATIONALE D'ATTRIBUTION CREATIVE COMMONS BY 4.0.
<https://creativecommons.org/licenses/by/4.0/>

L'accès aux articles de la revue « Journal de l'École polytechnique — Mathématiques » (<http://jep.centre-mersenne.org/>), implique l'accord avec les conditions générales d'utilisation (<http://jep.centre-mersenne.org/legal/>).

Publié avec le soutien
du Centre National de la Recherche Scientifique



Publication membre du
Centre Mersenne pour l'édition scientifique ouverte
www.centre-mersenne.org

MULTIPLE MIXING AND DISJOINTNESS FOR TIME CHANGES OF BOUNDED-TYPE HEISENBERG NILFLOWS

BY GIOVANNI FORNI & ADAM KANIGOWSKI

ABSTRACT. — We study time changes of bounded type Heisenberg nilflows (ϕ_t) acting on the Heisenberg nilmanifold M . We show that for every positive $\tau \in W^s(M)$, $s > 7/2$, every non-trivial time change (ϕ_t^τ) enjoys the *Ratner property*. As a consequence, every mixing time change is mixing of all orders. Moreover, we show that for every $\tau \in W^s(M)$, $s > 9/2$ and every $p, q \in \mathbb{N}$, $p \neq q$, (ϕ_{pt}^τ) and (ϕ_{qt}^τ) are disjoint. As a consequence, *Sarnak conjecture* on Möbius disjointness holds for all such time changes.

RÉSUMÉ (Mélange multiple et disjonction pour les reparamétrisations des flots nilpotents de type borné)

Nous étudions les reparamétrisations (ϕ_t^τ) des flots nilpotents de Heisenberg de type borné sur une variété nilpotente de Heisenberg M . Nous montrons que, pour des fonctions positives $\tau \in W^s(M)$ (espace de Sobolev) avec $s > 7/2$, toute reparamétrisation non triviale (ϕ_t^τ) a la propriété de Ratner. En conséquence, toute reparamétrisation mélangeante est mélangeante de tous les ordres. De plus, nous montrons que pour toutes les fonctions $\tau \in W^s(M)$, avec $s > 9/2$ et pour tous $p, q \in \mathbb{N}$, $p \neq q$, les flots (ϕ_{pt}^τ) et (ϕ_{qt}^τ) sont disjoints. Il s'ensuit, en particulier, que la conjecture de Sarnak sur la disjonction de la fonction de Möbius est valable pour toutes ces reparamétrisations.

CONTENTS

1. Introduction.....	64
2. Definitions.....	68
3. Ratner's property.....	70
4. Disjointness criterion.....	71
5. Birkhoff sums over toral skew-shifts.....	76
6. Ratner's property: proof of Theorem 3.....	84
7. Disjointness: proof of Theorem 4.....	86
References.....	90

2010 MATHEMATICS SUBJECT CLASSIFICATION. — 37C40, 28D10.

KEYWORDS. — Nilflows, time-changes, Ratner property, multiple mixing, disjointness.

1. INTRODUCTION

In this paper we study ergodic properties of time changes of *Heisenberg nilflows*. Nilflows on (non-abelian) nilmanifolds are classical examples of systems which share some features from both the *elliptic* and the *parabolic* world. They always have a non-trivial Kronecker factor which is responsible for the elliptic behavior (in particular they are never weakly mixing). On the other hand, orthogonally to the elliptic factor they exhibit polynomial speed of divergence of nearby trajectories and are polynomially mixing, which are properties characteristic of parabolic systems.

We are interested in the lowest dimensional (non-abelian) situation, i.e., nilflows on 3-dimensional Heisenberg nilmanifolds. In [3] it was shown that, for every (ergodic) Heisenberg nilflow, there exists a dense set of smooth time changes which are mixing. This result was strengthened in [13], where it was shown that for a full measure set of Heisenberg nilflows a *generic* time change is mixing, and moreover one has a “stretched-polynomial” decay of correlations for any pair of sufficiently smooth observables. For Heisenberg nilflows of *bounded type* the decay of correlations is estimated in [13] to be polynomial, as expected according to the “parabolic paradigm” (see [18, §8.2.f]). The mixing result of [3] was generalized in [27] to a class of nilflows on higher step nilmanifolds, called quasi-Abelian, which includes suspension flows over toral skew-shifts, and then recently to all non-Abelian nilflows in [4]. These general results reach no conclusion about the speed of mixing.

For time changes of horocycle flows, polynomial decay of correlations, as well as the property that the maximal spectral type is Lebesgue, were proved in [14]. A different proof of the absolute continuity of the spectrum was given simultaneously and independently by R. Tiedra de Aldecoa [29] under the so-called Kushnirenko condition (a positivity condition which holds for small time changes) and later in general [30]. More recently, B. Fayad and the authors [6] have proved that the spectrum is Lebesgue of countable multiplicity, also for a class of Kochergin flows on the two-torus. Nothing is known about the spectrum of smooth-time changes of nilflows beyond the mixing results of [3] and [13]. In particular, even for Heisenberg nilflows of bounded type, it is unclear whether the spectrum has an absolutely continuous component.

It follows from [3] and [13] that by a time change one can alter the dynamical features of Heisenberg nilflows, i.e., the elliptic factor becomes trivial for the time-changed flow and the mixing property holds (with polynomial decay of correlations for bounded type nilflows). It is therefore natural to ask to what extent the time-changed flow can behave, roughly speaking, as a “prototypical” parabolic flow (there is no widely accepted formal definition of a parabolic system). One of the characteristic features of parabolic systems is the *Ratner property* which quantifies the polynomial speed of divergence of nearby trajectories. It was first established by M. Ratner in [25] in the class of horocycle flows and was applied to prove Ratner’s rigidity phenomena in this class. Moreover, in [26], M. Ratner showed that the Ratner property survives under C^1 smooth time changes of horocycle flows, hence similar rigidity phenomena hold for time changes. One of the most important consequences of this property is

that a mixing system with the Ratner property is mixing of all orders. Indeed, Ratner showed that the Ratner property implies so-called PID property of the set of joinings which in turn implies higher order mixing, as observed by del Junco and Rudolph.

Recently Ratner’s property (or its variants) was observed in a new class of (non-homogeneous) systems, that of smooth flows on surfaces with finitely many (saddle-like) singularities. In [7], the authors studied the case of smooth mixing flows on the two-torus and established the SWR-property.⁽¹⁾ This property allows to establish the Ratner-type divergence of orbits, either in the future or in the past (depending on points), and moreover it has the same dynamical consequences as the original Ratner property. Then the authors showed in particular that the SWR-property holds for a full measure set of mixing flows with logarithmic singularities (Arnol’d flows) thereby proving higher order mixing in this class. The result in [7] was strengthened in [20], where the authors showed that the SWR-property holds for a full measure set of Arnol’d flows on surfaces of higher genus.

It is therefore natural to ask whether a Ratner property holds in the class of Heisenberg nilflows. In [21] it is shown that Ratner’s property implies in particular that the Kronecker factor is trivial and hence no nilflow can enjoy it. The situation is very different for *non-trivial time changes* of Heisenberg nilflows.

Let \mathfrak{H} denote the 3-dimensional Heisenberg group and let \mathfrak{h} denote its Lie algebra. Let $M := \Gamma \backslash \mathfrak{H}$ denote a Heisenberg nilmanifold, that is, the quotient of \mathfrak{H} over a (co-compact) lattice $\Gamma < \mathfrak{H}$. For any $W \in \mathfrak{h}$, the flow (ϕ_t^W) generated by the projection to M of the left-invariant vector field W on \mathfrak{H} , is called a Heisenberg nilflow (see Section 2.4 for the definition).

For any $W \in \mathfrak{h}$ and any positive function $\tau \in C^1(M)$, let $(\phi_t^{W,\tau})$ denote the time change of the nilflow (ϕ_t^W) , that is, the flow generated by the vector field τW on M . The flow $(\phi_t^{W,\tau})$ is a reparametrization of the nilflow (ϕ_t^W) in the sense that there exists a function $\mathcal{F} : M \times \mathbb{R} \rightarrow \mathbb{R}$ such that

$$\phi_t^{W,\tau}(x) = \phi_{\mathcal{F}(x,t)}^W, \quad \text{for all } (x, t) \in M \times \mathbb{R}.$$

One can verify that the group property of the flow $(\phi_t^{W,\tau})$ implies that the function \mathcal{F} is a cocycle over $(\phi_t^{W,\tau})$, in the sense that

$$\mathcal{F}(x, t + s) = \mathcal{F}(\phi_t^{W,\tau}(x), s) + \mathcal{F}(x, t),$$

and it is uniquely determined by the above cocycle condition and by the identity

$$\frac{\partial}{\partial t} \mathcal{F}(x, t)|_{t=0} = \tau(x), \quad \text{for all } x \in M.$$

The time-change $(\phi_t^{W,\tau})$ is called *trivial* if the cocycle \mathcal{F} is cohomologous to a constant, that is, there exist a constant $c \in \mathbb{R}$ and a measurable function $u : M \rightarrow \mathbb{R}$ such that

$$\mathcal{F}(x, t) - c = (u \circ \phi_t^{W,\tau})(x) - u(x), \quad \text{for all } (x, t) \in M \times \mathbb{R}.$$

⁽¹⁾Acronym for switchable weak Ratner. It was also shown that the original Ratner property does not hold.

All trivial time changes $(\phi_t^{W,\tau})$ are measurably conjugate to the nilflow (ϕ_t^W) and the regularity of the conjugacy is equal to the regularity of the so-called *transfer function* u on M .

A vector field $W \in \mathfrak{h}$ and the corresponding flow (ϕ_t^W) are called of *bounded type* if their projections on the Kronecker factor (which are respectively a constant coefficients vector field and the corresponding linear flow on a 2-dimensional torus) are of bounded type.

Our first main result establishes the Ratner property (see Section 3 for the definition) for time changes of bounded type Heisenberg nilflows in a very strong sense. In fact our first main result is the following. For every $s > 0$, let $W^s(M)$ denote the standard Sobolev space. By the Sobolev embedding theorem we have that $W^s(M) \subset C^k(M)$, for every $s > 3/2 + k$.

THEOREM 1. — *Let $W \in \mathfrak{h}$ be of bounded type. For any positive function $\tau \in W^s(M)$ with $s > 7/2$, either the time change is trivial or the time-changed flow $(\phi_t^{W,\tau})$ enjoys the Ratner property.*

Recall that the famous Rokhlin problem asks whether mixing implies mixing of all orders. The above result implies that the answer to the Rokhlin problem is positive for smooth time changes of bounded type Heisenberg nilflows:

COROLLARY 1.1. — *Let $W \in \mathfrak{h}$ be of bounded type and let*

$$\mathcal{D}^s(W) = \{(\phi_t^{W,\tau}) : \tau \in W^s(M), \tau > 0\}.$$

Then, for any $s \geq 7/2$, every element of $\mathcal{D}^s(W)$ is mixing if and only if it is mixing of all orders. As proved in [3] and [13], for $s \geq 7/2$ and W of bounded type, mixing is generic in the set $\mathcal{D}^s(W)$.

Moreover, by [21], we have the following strong dichotomy for time changes of Heisenberg nilflows:

COROLLARY 1.2. — *Let $W \in \mathfrak{h}$ be of bounded type. Then for every positive function $\tau \in W^s(M)$ with $s > 7/2$, either the time change is trivial or $(\phi_t^{W,\tau})$ is mildly mixing (no non-trivial rigid factors).*

It turns out that Heisenberg nilflows of bounded type (as in Theorem 1) are the only known examples, beyond horocycle flows and their time changes, for which the original Ratner property holds.

Our second main result deals with disjointness properties of time changes of Heisenberg nilflows. It is based on a variant of a parabolic disjointness criterion from [22]. We have:

THEOREM 2. — *Let $W \in \mathfrak{h}$ be of bounded type. For any positive function $\tau \in W^s(M)$ with $s > 9/2$, if the time change is non-trivial, then the flows $(\phi_{pt}^{W,\tau})$ and $(\phi_{qt}^{W,\tau})$ are disjoint for all $p, q \in \mathbb{N}$, $p \neq q$.*

As mentioned above, if the time-change τ is trivial, then (ϕ_{pt}^W) and (ϕ_{qt}^W) are not disjoint (since (ϕ_t^W) has a non-trivial Kronecker factor). However, L. Flaminio, K. Frączek, M. Lemańczyk and J. Kułaga-Przymus [11] have recently proved that any ergodic nilsystem satisfies the AOP-property: as $p, q \rightarrow +\infty$, the set of joinings between (ϕ_{pt}^W) and (ϕ_{qt}^W) weakly converges to the product measure.

The above theorem should be compared with analogous disjointness results for other flows with Ratner’s property. It follows from the renormalization equation for the horocycle flow, which states that $g_s h_t = h_{e^{-2st}} g_s$ for all $s, t \in \mathbb{R}$, that h_{pt} and h_{qt} are isomorphic (and hence not disjoint) for any $p, q \in \mathbb{R} \setminus \{0\}$. In [26], joinings of time changes of horocycle flows were completely characterized by M. Ratner. From Ratner’s work one can derive that distinct powers of the same time change are disjoint unless the time change is trivial [10]. A different proof of this result, based on a new disjointness criterion for parabolic flows, was given recently in [22]. Moreover, in [22] it is proved that for almost every Arnol’d flow on \mathbb{T}^2 the same assertion as in Theorem 2 holds. Therefore, among known flows with Ratner’s property, the horocycle flow is the only one for which the conclusion of Theorem 2 does not hold. The heuristic reason for that is that the Ratner property for the horocycle flow depends only on the distance between points, and not on their position in space (since the space is homogeneous). In all other examples (for flows as in Theorem 2 in particular) the divergence depends also on position which allows to get stronger consequences (see Section 4).

Let us now briefly discuss the connection between Theorem 2 and Sarnak’s conjecture on Möbius disjointness [28], which is recently under extensive study, see e.g. [8]. We say that a continuous flow (T_t) on a compact metric space (X, d) is Möbius disjoint, if for every $F \in C(X)$ and every $x \in X$ and every $t \in \mathbb{R}$ we have

$$(1) \quad \lim_{N \rightarrow +\infty} \frac{1}{N} \sum_{n \leq N} F(T_{nt}x)\mu(n) = 0,$$

here μ denotes the classical Möbius function.⁽²⁾

Möbius disjointness for horocycle flows was proved by J. Bourgain, P. Sarnak and T. Ziegler [5]. Moreover, as explained in [10], it follows from Ratner’s work [26] that Möbius disjointness also holds for non-trivial time changes of horocycle flows. Moreover, in view of a criterion due to I. Kátai [23] and to Bourgain, Sarnak and Ziegler [5], for non-trivial time changes of horocycle flows the convergence in (1) is uniform in $x \in X$. It is an open question whether for horocycle flows the convergence in (1) (for fixed F and $t \in \mathbb{R} \setminus \{0\}$) is uniform with respect to $x \in X$. A corollary of Theorem 2, again by the Kátai-Bourgain-Sarnak-Ziegler (KBSZ) criterion [23], [5], is the following:

COROLLARY 1.3. — *Let $W \in \mathfrak{h}$ be of bounded type. For any positive function $\tau \in W^s(M)$ with $s > 9/2$, if the time change is non-trivial, then $(\phi_t^{W,\tau})$ is Möbius disjoint. Moreover, the convergence in formula (1) is uniform with respect to $x \in M$.*

⁽²⁾Sarnak’s conjecture states that every system of zero topological entropy is Möbius disjoint.

It follows from the work of B. Green and T. Tao [17] that, if the time change is trivial, then $(\phi_t^{W,\tau})$ is Möbius disjoint and the convergence in formula (1) is also uniform (the uniform convergence in (1) follows also from the AOP property, [11]).

The structure of the paper is as follows. In Section 2 we recall some basic definitions of the theory of joinings and recall the definition Heisenberg nilflows and their special flow representations over skew-shifts of the 2-torus. In Section 3 we recall the Ratner property and then formulate a version of it for special flows. In Section 4 we state a disjointness criterion (Proposition 4.1) and then formulate a version of it for special flows (Lemma 4.3). In Section 5 we derive from results of [9] (see also [12]) estimates on Birkhoff sums for smooth functions over skew-shifts of the 2-torus. Finally, in Sections 6 and 7 we prove our main theorems by applying the estimates from Section 5.

2. DEFINITIONS

2.1. JOININGS AND DISJOINTNESS. — We refer the reader to [16] for basic theory of joinings. Let $(\phi_t) : (X, \mathcal{B}, \mu) \rightarrow (X, \mathcal{B}, \mu)$ and $(\psi_t) : (Y, \mathcal{C}, \nu) \rightarrow (Y, \mathcal{C}, \nu)$ be two ergodic flows. A *joining* of (ϕ_t) and (ψ_t) is any $(\phi_t \times \psi_t)$ invariant measure such that $\rho(X \times B) = \mu(X)\nu(B)$ and $\rho(C \times Y) = \mu(C)\nu(Y)$. The set of joinings of (ϕ_t) and (ψ_t) is denoted by $J((\phi_t), (\psi_t))$. Notice that $\mu \otimes \nu \in J((\phi_t), (\psi_t))$. We say that (ϕ_t) and (ψ_t) are *disjoint* (denoting $(\phi_t) \perp (\psi_t)$) if $J((\phi_t), (\psi_t)) = \{\mu \otimes \nu\}$.

2.2. HEISENBERG NILFLOWS. — The (three-dimensional) Heisenberg group \mathfrak{H} is given by

$$\mathfrak{H} := \left\{ \begin{pmatrix} 1 & x & z \\ 0 & 1 & y \\ 0 & 0 & 1 \end{pmatrix} : x, y, z \in \mathbb{R} \right\}.$$

Let Γ be a lattice in \mathfrak{H} . A *Heisenberg manifold* M is a quotient $\Gamma \backslash \mathfrak{H}$. It is known that up to an automorphism of \mathfrak{H}

$$\Gamma = \Gamma_K = \left\{ \begin{pmatrix} 1 & m & p/K \\ 0 & 1 & n \\ 0 & 0 & 1 \end{pmatrix} : m, n, p \in \mathbb{Z} \right\},$$

where K is a positive integer. Notice that M has a (normalized) volume element vol given by the projection of the (bi-invariant) Haar measure on \mathfrak{H} .

Since the Abelianized Lie group $\overline{\mathfrak{H}} := \mathfrak{H}/[\mathfrak{H}, \mathfrak{H}]$ is isomorphic to \mathbb{R}^2 (as a Lie group) and $\overline{\Gamma}_K := \Gamma_K/[\Gamma_K, \Gamma_K]$ is a lattice in $\overline{\mathfrak{H}}$, the Heisenberg nilmanifold M fibers over a 2-dimensional torus $\overline{M} = \overline{\mathfrak{H}}/\overline{\Gamma}_K$, with fibers isomorphic to a circle.

Let W be any element of the Lie algebra \mathfrak{h} of \mathfrak{H} . The *Heisenberg nilflow* for W is given by

$$\phi_t^W(x) = x \exp(tW), \quad \text{for all } (x, t) \in M \times \mathbb{R}.$$

Every Heisenberg nilflow (ϕ_t^W) on M preserves the volume element vol on M . The classical ergodic theory of nilflows (see [2]) implies that a Heisenberg nilflow (ϕ_t^W) is uniquely ergodic iff it is ergodic iff it is minimal iff the projected flow on \overline{M} (which is isomorphic to its Kronecker factor) has rationally independent frequencies. More

generally, the Diophantine properties of a vector $W \in \mathfrak{h}$, and of the corresponding nilflow (ϕ_t^X) , under the renormalization dynamics introduced in [9] can be entirely read from the Diophantine properties of the projection \overline{W} of W onto the Abelianized Lie algebra $\overline{\mathfrak{h}} := \mathfrak{h}/[\mathfrak{h}, \mathfrak{h}]$, which is also isomorphic to \mathbb{R}^2 (as a Lie algebra). In particular, a vector $W \in \mathfrak{h}$ is called of *bounded type* if and only if its projection $\overline{W} \in \mathbb{R}^2$ is of bounded type.

Let $\tau \in L^1(M)$, $\tau > 0$. The flow $(\phi_t^{W,\tau})$ is called a *time change* (or a reparametrization) of the flow (ϕ_t^W) if

$$\phi_t^{W,\tau}(x) = \phi_{\mathcal{I}(x,t)}^W(x), \quad \text{for all } (x, t) \in M \times \mathbb{R},$$

where the $(\phi_t^{W,\tau})$ -cocycle $\mathcal{I}(x, t)$ is uniquely defined by the condition that

$$\frac{\partial}{\partial t} \mathcal{I}(x, t)|_{t=0} = \tau(x), \quad \text{for all } x \in M.$$

The above condition is equivalent to either of the following integral conditions

$$\int_0^t \tau(\phi_s^{W,\tau}(x)) ds = \mathcal{I}(x, t) \quad \text{and} \quad \int_0^{\mathcal{I}(x,t)} 1/\tau(\phi_s^W(x)) ds = t.$$

By definition the time change is trivial if the $(\phi_t^{W,\tau})$ -cocycle \mathcal{I} is cohomologous to a constant. It follows from the above formula that the time change is trivial if and only if the function τ is $(\phi_t^{W,\tau})$ -cohomologous to constant, in the sense that there exist a constant $c \in \mathbb{R}$ and a function $u : M \rightarrow \mathbb{R}$ such that

$$u \circ \phi_t^{W,\tau}(x) - u(x) = \tau(\phi_t^{W,\tau}(x)) - c, \quad \text{for all } (x, t) \in M \times \mathbb{R},$$

or, infinitesimally, if and only if $\tau W u = \tau - c$. Finally we see that this condition is equivalent to the property that the function $1/\tau$ is (ϕ_t^W) -cohomologous to constant, that is, $-W u/c = 1/\tau - 1/c$ or, equivalently,

$$(-u/c) \circ \phi_t^W - (-u/c) = 1/\tau(\phi_t^W(x)) - (1/c).$$

We have thus reduced the condition that a time change of a nilflow be trivial to the existence of solutions of cohomological equation for the nilflow itself, a question that can be analyzed by Fourier analysis [24], [9], [3], [12].

2.3. SPECIAL FLOWS. — Let $\Phi : (X, \mathcal{B}, \mu, d) \rightarrow (X, \mathcal{B}, \mu, d)$ be an ergodic automorphism of a compact metric probability space and let $f : X \rightarrow \mathbb{R}$ be strictly positive.

We recall that the special flow $(\Phi_t) := (\Phi_t^f)$ constructed above Φ and under f acts on $X^f := \{(x, s) : x \in X, 0 \leq s < f(x)\}$ by

$$\Phi_t(x, s) = (\Phi^{N(x,s,t)}(x), s + t - S_{N(x,s,t)}(f)(x)),$$

where $N(x, s, t)$ is the unique integer such that

$$0 \leq s + t - S_{N(x,s,t)}(f)(x) \leq f(\Phi^{N(x,s,t)}(x)),$$

and

$$S_N(f)(x) = \begin{cases} f(x) + \cdots + f(\Phi^{n-1}x) & \text{if } n > 0 \\ 0 & \text{if } n = 0 \\ -(f(\Phi^n x) + \cdots + f(\Phi^{-1}x)) & \text{if } n < 0. \end{cases}$$

Notice that the flow (Φ_t) preserves the measure $\mu^f = \mu \otimes \lambda_{\mathbb{R}}$ restricted to X^f , where $\lambda_{\mathbb{R}}$ denotes the Lebesgue measure on \mathbb{R} . Let moreover d^f denote the product metric (of d and the absolute value metric on \mathbb{R}).

2.4. SPECIAL FLOW REPRESENTATION OF NILFLOWS. — It is classical that an *ergodic* nilflow (ϕ_t^W) can be represented as a *special flow*, where the base automorphism $\Phi_{\alpha,\beta} : \mathbb{T}^2 \rightarrow \mathbb{T}^2$ is given by $\Phi_{\alpha,\beta}(x, y) = (x + \alpha, y + x + \beta)$ for $\alpha \in [0, 1) \setminus \mathbb{Q}$, $\beta \in \mathbb{R}$ and under a constant roof function $f(x, y) = C_W > 0$.

Let $(q_n)_{n=1}^{+\infty}$ denote the sequence of denominators of $\alpha \in [0, 1) \setminus \mathbb{Q}$. The vector field $W \in \mathfrak{h}$ is of *bounded type* if and only if α is of bounded type, i.e., there exists $C_\alpha > 0$ such that $q_{n+1} \leq C_\alpha q_n$ for every $n \in \mathbb{N}$.

For any function $f \in L^1(\mathbb{T}^2)$, $f > 0$, let $(\Phi_t^{f,\alpha,\beta})$ denote the special flow over $\Phi_{\alpha,\beta}$ and under f . Then every time change $\phi_t^{W,\tau}$ is isomorphic to a special flow $(\Phi_t^{f_\tau,\alpha,\beta})$, where the roof function f_τ is as smooth as τ . In view of the above representation, Theorems 1 and 2 are respectively equivalent to the following two theorems:

THEOREM 3. — *Let $\alpha \in \mathbb{R} \setminus \mathbb{Q}$ be of bounded type and let $f \in W^s(\mathbb{T}^2)$, with $s > 7/2$, be a positive function. Then the flow $(\Phi_t^{f,\alpha,\beta})$ has the Ratner property.*

THEOREM 4. — *Let $\alpha \in [0, 1) \setminus \mathbb{Q}$ be of bounded type and let $f \in W^s(\mathbb{T}^2)$, with $s > 9/2$, be a positive function. Then the flows $(\Phi_{pt}^{f,\alpha,\beta})$ and $(\Phi_{qt}^{f,\alpha,\beta})$ are disjoint for all $p, q \in \mathbb{N}$ with $p \neq q$.*

REMARK 2.1. — It seems to the authors that a necessary condition for the Ratner property (or any of its variants) to hold in

$$\mathcal{D}^s(W) = \{(\phi_t^{W,\tau}) : \tau \in W^s(M), \tau > 0, (\phi_t^{W,\tau}) \text{ is a non-trivial time-change}\}$$

is that W is of bounded type.

3. RATNER'S PROPERTY

Let $(\phi_t) : (X, \mathcal{B}, \mu, d) \rightarrow (X, \mathcal{B}, \mu, d)$ be an ergodic flow on a σ -compact metric probability space.

DEFINITION 3.1. — Let $P = \{-1, 1\}$ and let $t_0 \in \mathbb{R}$. The flow (ϕ_t) has the $R(t_0, P)$ -*property* if for every $\varepsilon > 0$ and $N \in \mathbb{N}$, there exist $\kappa = \kappa(\varepsilon)$, $\delta = \delta(\varepsilon, N)$ and a set $Z = Z(\varepsilon, N)$ with $\mu(Z) > 1 - \varepsilon$, such that: for every $x, y \in Z$ with $d(x, y) < \delta$ and x not in the orbit of y , there exist $p = p(x, y) \in P$ and $M = M(x, y)$, $L = L(x, y) \geq N$, $L/M \geq \kappa$, for which

$$\#\{n \in [M, M + L] \cap \mathbb{Z} : d(\phi_{nt_0}(x), \phi_{nt_0+p}(y)) < \varepsilon\} \geq (1 - \varepsilon)L.$$

The flow (ϕ_t) is said to have *Ratner's property* if

$$\{s \in \mathbb{R} : (\phi_t) \text{ has property } R(s, P)\}$$

is uncountable.

3.1. RATNER'S PROPERTY FOR SPECIAL FLOWS. — In what follows $\Phi : (X, \mathcal{B}, \mu, d) \rightarrow (X, \mathcal{B}, \mu, d)$ is an ergodic automorphism of a σ -compact metric probability space and $f \in L_1^+(X)$. We have the following proposition (see [19, Prop. 4.1]):

PROPOSITION 3.2. — *Let $P = \{-1, 1\}$. If for every $\varepsilon > 0$ and $N \in \mathbb{N}$ there exist $\kappa = \kappa(\varepsilon) > 0$, $\delta = \delta(\varepsilon, N) > 0$ and a set $Z = Z(\varepsilon, N) \subset X$ with $\mu(Z) \geq 1 - \varepsilon$, such that, for every $x, y \in Z$ with $d(x, y) \leq \delta$, there exist $M, L \geq N$, $L/M \geq \kappa$ and $p \in P$ such that for every $n \in [M, M + L] \cap \mathbb{Z}$*

$$d(\Phi^n x, \Phi^n y) < \varepsilon \quad \text{and} \quad |S_n(f)(x) - S_n(f)(y) - p| < \varepsilon,$$

then the special flow (Φ_t^f) satisfies the Ratner property.

In [19], Proposition 3.2 was proved for the SR-property, which is a modification of Ratner's property in which one allows for divergence either in the future or in the past. However the proof in [19] immediately extends to a proof of Proposition 3.2.

We will use Proposition 3.2 to prove Theorem 3.

4. DISJOINTNESS CRITERION

Let $(\phi_t) : (X, \mathcal{B}, \mu, d_1) \rightarrow (X, \mathcal{B}, \mu, d_1)$ and $(\psi_t) : (Y, \mathcal{C}, \nu, d_2) \rightarrow (Y, \mathcal{C}, \nu, d_2)$ be two weakly mixing flows (and X, Y are σ -compact). In this section we prove the following proposition:

PROPOSITION 4.1 (Disjointness criterion). — *Let $P' \subset \mathbb{R}$ be a compact set and let $v' \neq 0$. Fix $1 \geq c > 0$. Assume there exists $(A_k) \subset \text{Aut}(X_k, \mathcal{B}|_{X_k}, \mu|_{X_k})$, such that $\mu(X_k) \rightarrow \mu(X)$, $A_k \rightarrow \text{Id}$ uniformly. Assume moreover that for every $\varepsilon > 0$ and $N \in \mathbb{N}$ there exist $(E_k = E_k(\varepsilon)) \subset \mathcal{B}$ with $\mu(E_k) \geq c\mu(X)$, $0 < \kappa = \kappa(\varepsilon) < \varepsilon$, $\delta = \delta(\varepsilon, N) > 0$, a set $Z = Z(\varepsilon, N) \subset Y$ with $\nu(Z) \geq (1 - \varepsilon)\nu(Y)$, such that for all $y, y' \in Z$ satisfying $d_2(y, y') < \delta$, every k such that $d_1(A_k, \text{Id}) < \delta$ and every $x \in E_k$, $x' := A_k x$, there are $M \geq N$, $L \geq 1$, $L/M \geq \kappa$ and $V \in P'$, $v \in \{-v', v'\}$, for which the following holds:*

$$(2) \quad \max(d_1(\phi_t x, \phi_{t+V+v} x'), d_2(\psi_t y, \psi_{t+V} y')) < \varepsilon$$

$$\text{for } t \in U \subset [M, M + L] \text{ with } \lambda_{\mathbb{R}}(U) \geq (1 - \varepsilon)L.$$

Then (ϕ_t) and (ψ_t) are disjoint.

The proof of the above proposition follows similar lines to the proof of [22, Th. 3]. We provide a proof for completeness.

Proof. — Let $\rho \in J((\phi_t)_{t \in \mathbb{R}}, (\psi_t)_{t \in \mathbb{R}})$ be an ergodic joining with $\rho \neq \mu \times \nu$. Since (ϕ_t) is weakly mixing it follows that, for $v \in \{-v', v'\}$, the map ϕ_v is ergodic and hence disjoint from Id. Therefore, there exist $B_v \in \mathcal{B}$ and $C_v \in \mathcal{C}$ such that

$$(3) \quad |\rho(\phi_{-v}(B_v) \times C_v) - \rho(B_v \times C_v)| > \eta$$

for some $0 < \eta < 1$. Let $V_\varepsilon^1(B_v) := \{x \in X : d_1(x, B_v) < \varepsilon\}$ and similarly $V_\varepsilon^2(C_v) := \{y \in Y : d_2(y, C_v) < \varepsilon\}$. There exists $\varepsilon \in (0, c\eta/1000)$ such that

$$\max(|\mu(V_\varepsilon^1(B_v)) - \mu(B_v)|, |\nu(V_\varepsilon^2(C_v)) - \nu(C_v)|) < \eta/32.$$

Since ρ is a joining, by the triangle inequality, for each $t \in \mathbb{R}$, we have

$$(4) \quad |\rho(\phi_{-t}V_\varepsilon^1(B_v) \times V_\varepsilon^2(C_v)) - \rho(\phi_{-t}B_v \times C_v)| < \frac{\eta}{16}.$$

By applying Birkhoff point-wise ergodic theorem to the joining flow $(\phi_t \times \psi_t, \rho)$ and to the characteristic functions of the sets $\phi_{-v}B_v \times C_v$ and $\phi_{-v}V_\varepsilon^1(B_v) \times V_\varepsilon^2(C_v)$ for $v \in \{-v', v'\}$, it follows that there exist $N_0 \in \mathbb{N}$, $\kappa > 0$ and a set $U_1 \in \mathcal{B} \otimes \mathcal{C}$, with $\rho(U_1) > (1 - c/100)\rho(X \times Y)$, such that, for every $L, M \geq N_0$ with $L/M \geq \kappa$ and $v \in \{-v', v'\} \cup \{0\}$, and for all $(x, y) \in U_1$, we have

$$(5) \quad \left| \frac{1}{L} \int_M^{M+L} \chi_{\phi_{-v}B_v \times C_v}(\phi_t x, \psi_t y) dt - \rho(\phi_{-v}B_v \times C_v) \right| < \frac{\eta}{16},$$

$$(6) \quad \left| \frac{1}{L} \int_M^{M+L} \chi_{\phi_{-v}V_\varepsilon^1(B_v) \times V_\varepsilon^2(C_v)}(\phi_t x, \psi_t y) dt - \rho(\phi_{-v}V_\varepsilon^1(B_v) \times V_\varepsilon^2(C_v)) \right| < \frac{\eta}{16}.$$

Let $U_2 := U_1 \cap (X \times Z)$, where $Z = Z(\varepsilon, N_0)$ comes from our assumptions. Then $\rho(U_2) > (1 - c/50)\rho(X \times Y)$. Note also that since $X \times Y$ is σ -compact, the measure ρ is regular, and hence we can additionally assume that U_2 is compact. Define $\text{proj} : X \times Y \rightarrow X$, $\text{proj}(x, y) = x$. Then the fibers of proj are σ -compact, and since U_2 is compact, the fibers of the map $\text{proj}|_{U_2} : U_2 \rightarrow \text{proj}(U_2) \subset X$ are also σ -compact and $\text{proj}(U_2)$ is also compact. Thus, by Kunugui's selection theorem (see e.g. [15, Th. 4.1]), it follows that there exists a measurable (selection) $s_Y : \text{proj}(U_2) \rightarrow X \times Y$ such that $(x, s_Y(x)) \in U_2$. Note that $\mu(\text{proj}(U_2)) \geq \rho(U_2) > (1 - c/50)\mu(X)$. By Luzin's theorem there exists $X_{\text{cont}} \subset \text{proj}(U_2)$, with $\mu(X_{\text{cont}}) \geq (1 - c/50)\mu(X)$, such that s_Y is uniformly continuous on X_{cont} . Finally, we set

$$\tilde{U} := U_2 \cap (X_{\text{cont}} \times Y).$$

We have $\rho(\tilde{U}) > (1 - c/10)\rho(X \times Y)$. Moreover, for the set $U_X := \text{proj}(\tilde{U})$ we have $\mu(U_X) \geq \rho(\tilde{U}) > (1 - c/10)\rho(X \times Y)$. Hence, by the definitions of sequences (A_k) and $(E_k) = (E_k(\varepsilon))$, it follows that there exists $k_0 = k_0(\varepsilon)$ such that for $k \geq k_0$,

$$(7) \quad \mu(A_{-k}(U_X \cap X_k) \cap (U_X \cap X_k) \cap E_k) > 0.$$

Let $\delta = \delta(\varepsilon, N_0)$ come from the assumptions of our theorem. By the uniform continuity of $s_Y : X_{\text{cont}} \rightarrow Y$ it follows that there exists $0 < \delta' < \delta$ such that $d_1(x_1, x_2) < \delta'$ implies $d_2(s_Y(x_1), s_Y(x_2)) < \delta$ for each $x_1, x_2 \in X_{\text{cont}}$. Since $A_k \rightarrow \text{Id}$ uniformly and $\tilde{U} \subset X_{\text{cont}} \times Y$, there exists $k_1 = k_1(\varepsilon)$ such that for $k \geq k_1$, $d_2(s_Y(x), s_Y(A_k x)) < \delta$ for $x \in X_k \cap X_{\text{cont}}$. Fix $k \geq \max(k_0, k_1 + 1)$ (so that $d_1(A_k, \text{Id}) < \delta'$). Let $x \in$

$A_{-k}(U_X \cap X_k) \cap (U_X \cap X_k) \cap E_k$. Such a point does exist in view of (7). Set $x' = A_k x$, $y = s_Y(x)$, $y' = s_Y(x')$. By definition, $(x, y), (x', y') \in \tilde{U}$ and $d_2(y, y') < \delta$ and all other assumptions of our theorem are satisfied for $(x, y), (x', y')$ (so that we obtain M, L, V, v depending on (x, y) and (x', y') satisfying (2)).

We claim that

$$(8) \quad \rho(\phi_{-v}(B_v) \times C_v) > \rho(B_v \times C_v) - \frac{\eta}{2}.$$

Indeed, in view of (4), the estimate (8) follows if we can prove that

$$\rho(\phi_{-v}(V_\varepsilon^1(B_v)) \times V_\varepsilon^2(C_v)) > \rho(B_v \times C_v) - \frac{\eta}{4}.$$

Using (2) (for $v = v'$), (5) (for $v = 0$) and $(x, y) \in \tilde{U} \subset U_1$, we obtain that

$$\begin{aligned} \frac{1}{L} \int_M^{M+L} \chi_{V_\varepsilon^1(B_v) \times V_\varepsilon^2(C_v)}(\phi_{t+V+v}x', \psi_{t+V}y') dt \\ > \frac{1}{L} \int_M^{M+L} \chi_{B_v \times C_v}(\phi_t x, \psi_t y) dt - \varepsilon > \rho(B_v \times C_v) - \varepsilon - \frac{\eta}{16}. \end{aligned}$$

Hence to complete the proof of Claim (8), it is enough to show that

$$(9) \quad \frac{1}{L} \int_M^{M+L} \chi_{V_\varepsilon^1(B_v) \times V_\varepsilon^2(C_v)}(\phi_{t+V+v}x, \psi_{t+V}y) dt < \rho(\phi_{-v}V_\varepsilon^1(B_v) \times V_\varepsilon^2(C_v)) + \frac{\eta}{8}.$$

Notice however that

$$\begin{aligned} \frac{1}{L} \int_M^{M+L} \chi_{V_\varepsilon^1(B_v) \times V_\varepsilon^2(C_v)}(\phi_{t+V+v}x, \psi_{t+V}y) dt \\ = \frac{1}{L} \int_{M+V}^{M+V+L} \chi_{\phi_{-v}(V_\varepsilon^1(B_v)) \times V_\varepsilon^2(C_v)}(\phi_t x, \psi_t y) dt, \end{aligned}$$

hence the estimate in (9) follows from (6) with $M = M + V$ and $L = L$.

By a similar reasoning, we get

$$(10) \quad \rho(\phi_{-v}(B_v) \times C_v) < \rho(B_v \times C_v) + \frac{\eta}{2},$$

so putting together (8) and (10) we derive the estimate

$$|\rho(\phi_{-v}(B_v) \times C_v) - \rho(B_v \times C_v)| < \frac{\eta}{2}.$$

This however contradicts (3), hence the argument is complete. □

4.1. DISJOINTNESS CRITERION FOR SPECIAL FLOWS. — In this section we assume that $(\Phi_t) = (\Phi_t^f)$ and $(\Psi_t) = (\Psi_t^g)$ are special flows over ergodic $\Phi \in \text{Aut}(X, \mathcal{B}, \mu, d_1)$, $\Psi \in \text{Aut}(Y, \mathcal{C}, \nu, d_2)$ respectively and under $f \in L_+^1(X, \mathcal{B}, \mu)$, $g \in L_+^1(Y, \mathcal{C}, \nu)$. Let (Φ_t^f) act on X^f with metric d_1^f and (Ψ_t^g) act on X^g with metric d_2^g . For $(x, s) \in X^f$ and $t \in \mathbb{R}$ we denote by $n(x, s, t) \in \mathbb{Z}$ the unique number for which

$$S_{n(x,s,t)}(f)(x) \leq t + s < S_{n(x,s,t)+1}(f)(x),$$

i.e.,

$$\Phi_t^f(x, s) = (\Phi^{n(x,s,t)}x, s + t - S_{n(x,s,t)}(f)(x)).$$

We define $m(y, r, t)$ analogously for $(y, r) \in Y^g$. We tacitly assume that

f and g are bounded away from zero.

Before we state a disjointness criterion for special flows, we need the following general lemma:

LEMMA 4.2. — Fix ε such that $\inf_X f/10 > \varepsilon > 0$. Let $t \in \mathbb{R}$ and $(x, s), (x', s') \in X^f$ be such that $|s - s'| < \varepsilon^2$, $d_1(\Phi^{n(x,s,t)}x, \Phi^{n(x,s,t)}x') < \varepsilon^2$ and

$$\Phi_t^f(x, s) \in \{(x, s) : \varepsilon < s < f(x) - \varepsilon\}.$$

Let $V(t) = V(x, s, x', s', t) := S_{n(x,s,t)}(f)(x) - S_{n(x,s,t)}(f)(x')$. Then

$$d_1^f(\Phi_t^f(x, s), \Phi_{t-V(t)}^f(x', s')) \leq 2\varepsilon^2.$$

Proof. — Notice that since $\Phi_t^f(x, s) \in \{(x, s) : \varepsilon < s < f(x) - \varepsilon\}$ and $|s - s'| < \varepsilon^2$, we have

$$S_{n(x,s,t)}(f)(x') \leq t - V(t) + s' \leq S_{n(x,s,t)+1}(f)(x').$$

Therefore, $\Phi_{t-V(t)}^f(x', s') = (\Phi^{n(x,s,t)}x', t - V(t) + s' - S_{n(x,s,t)}(f)(x'))$. By definition $\Phi_t^f(x, s) = (\Phi^{n(x,s,t)}x, t + s - S_{n(x,s,t)}(f)(x))$. The statement follows by the definition of $V(t)$ since d_1^f is the product metric and we have $d_1(\Phi^{n(x,s,t)}x, \Phi^{n(x,s,t)}x') < \varepsilon^2$ and $|s - s'| \leq \varepsilon^2$. \square

LEMMA 4.3. — Let $V \in \mathbb{R}$ and $P = \{-p', p'\}$ for $p' \neq 0$ and $\zeta := \int_X f d\mu / \int_Y g d\nu$. Let $A_k \in \text{Aut}(X, \mathcal{B}, \mu)$, $A_k \rightarrow \text{Id}$ uniformly. Assume moreover that for every $\varepsilon' > 0$ and $N' \in \mathbb{N}$ there exist $0 < \kappa' = \kappa'(\varepsilon') < \varepsilon'$, $\delta' = \delta'(\varepsilon', N') > 0$, such that for all $y, y' \in Y$ satisfying $d_2(y, y') < \delta'$, every k such that $d_1(A_k, \text{Id}) < \delta'$ and every $x, x' := A_k x \in X$ there are $M' \geq N'$, $L \geq 1$, $L'/M' \geq \kappa'$ and $p \in P$ satisfying:

$$(11) \quad |S_{M'}(f)(x) - S_{M'}(f)(x')| < V,$$

$$(12) \quad |(S_{M'}(f)(x) - S_{M'}(f)(x')) - (S_{[\zeta M']}(g)(y) - S_{[\zeta M']}(g)(y')) - p| < \varepsilon',$$

$$(13) \quad d_1(\Phi^w x, \Phi^w x'), d_2(\Psi^w y, \Psi^w y') < \kappa'$$

for $w \in [0, \max(1, \zeta)(M' + L')] \cap \mathbb{Z}$; and for $h = \{f, g\}$,

$$(14) \quad |(S_w(h)(x) - S_w(h)(x')) - (S_u(h)(x) - S_u(h)(x'))| < \varepsilon',$$

for every $w, u \in [0, 2 \max(1, \zeta)(M' + L')]$, $|w - u| \leq L'$. Then (Φ_t^f) and (Ψ_t^g) are disjoint.

The proof of the above proposition follows similar lines (although is simpler) than the proof of [22, Prop. 4.1]. We provide a proof here for completeness.

Proof of Lemma 4.3. — We will show that the assumptions of Proposition 4.1 are satisfied. Let $P' := [-2V - |p'|, 2V + |p'|]$ and $v' = p'$. Let $c = 1$, and $A_k^f(x, s) = (A_k x, s)$ on X^f (then $A_k^f \rightarrow \text{Id}$ uniformly). Fix $\varepsilon > 0$ and $N \in \mathbb{N}$. Let $\varepsilon' = \varepsilon^4$, $N' =$ and let κ', δ' be as in Lemma 4.3. Define $\kappa := \kappa'^2$ and $\delta := \min(\varepsilon^{10}, \delta'^2)$.

By the ergodic theorem for Φ_t^f there exist $N'' \in \mathbb{R}$ and a set $E = E(\varepsilon) \subset X^f$, $\mu^f(E) > 1 - \varepsilon$, such that for every $M, L \geq N''$, $L/M \geq \kappa$ and every $(x, s) \in E$, we have

$$(15) \quad \lambda_{\mathbb{R}}(\{t \in [M, M + L] : \Phi_t^f(x, s) \in \{(x, s) : \varepsilon^{3/2} < s < f(x) - \varepsilon^{3/2}\}\}) \geq (1 - \varepsilon^{4/3})L$$

and for $t \in \mathbb{R}$ such that $n(x, s, t) \in [M, M + L]$, we have

$$(16) \quad |t - n(x, s, t) \int_X f d\mu| < \kappa'^2 t.$$

Similarly, by the ergodic theorem for Ψ_t^g there exist $N'' \in \mathbb{R}$ and $Z = Z(\varepsilon) \subset Y^g$, $\nu^g(Z) > 1 - \varepsilon$ such that for every $M, L \geq N''$, $L/M \geq \kappa$ and every $(y, r) \in Z$, we have

$$(17) \quad \lambda_{\mathbb{R}}(\{t \in [M, M + L] : \Psi_t^g(y, r) \in \{(y, r) : \varepsilon^{3/2} < r < g(y) - \varepsilon^{3/2}\}\}) \geq (1 - \varepsilon^{4/3})L,$$

and for $t \in \mathbb{R}$ such that $m(y, r, t) \in [M, M + L]$, we have

$$(18) \quad |t - m(y, r, t) \int_Y g d\nu| < \kappa'^2 t.$$

For $k \in \mathbb{N}$ let $E_k = E_k(\varepsilon) := E$. Fix $E \ni (x, s), (x', s) = A_k(x, s)$ with $d_1^f((x, s), (x', s)) < \delta$ and $(y, r), (y', r') \in Z$ with $d_2^g((y, r), (y', r')) < \delta$. Let M', L', p come from Lemma 4.3 for x, x' and y, y' . Define M, L by $n(x, s, M) \int_X f d\mu = M'$ and $n(x, s, M + L) \int_X f d\mu = M' + L'$. It follows by (16) that $L/M \geq \kappa$ and $M \geq N$.

Let $U \subset [M, M + L]$ be such that for $t \in U$, we have

$$\Phi_t^f(x, s) \in \{(x, s) : \varepsilon^{3/2} < s < f(x) - \varepsilon^{3/2}\}, \quad \Psi_t^g(y, r) \in \{(y, r) : \varepsilon^{3/2} < r < g(y) - \varepsilon^{3/2}\}$$

and

$$d_1(\Phi^{n(x,s,t)}x, \Phi^{n(x,s,t)}x'), d_2(\Psi^{m(y,r,t)}y, \Psi^{m(y,r,t)}y') < \varepsilon^2.$$

By (15), (17) and (13) it follows that $|U| \geq (1 - \varepsilon)L$. Let us then set

$$V(t) = S_{n(x,s,t)}(f)(x) - S_{n(x,s,t)}(f)(x'), \\ W(t) = S_{m(y,r,t)}(g)(y) - S_{m(y,r,t)}(g)(y').$$

By Lemma 4.2 (for (Φ_t^f) and (Ψ_t^g)) it follows that

$$d_1^f(\Phi_t^f(x, s), \Phi_{t-W(t)+(W(t)-V(t))}^f(x', s)) \leq 2\varepsilon^2$$

and

$$d_2^g(\Psi_t^g(y, r), \Psi_{t-W(t)}^g(y', r')) \leq 2\varepsilon^2.$$

Notice that for $t \in [M, M + L]$, by (14) (and (16),(18)), we have

$$|V(t) - V(M)|, |W(t) - W(M)| \leq \varepsilon^2$$

Moreover, by (12) and (14), we have

$$|V(M) - W(M) - p| < \varepsilon^2.$$

Hence

$$d_1^f(\Phi_t^f(x, s), \Phi_{t-W(M)+p}^f(x', s)) \leq 4\varepsilon^2.$$

and

$$d_2^g(\Psi_t^g(y, r), \Psi_{t-W(M)}^g(y', r')) \leq 4\varepsilon^2.$$

Finally, by (11) and (14) it follows that $|W(M)| \leq |V(M)| + p + 2\varepsilon^2 \leq 2V + p$ and hence $W(M) \in P$. This finishes the proof. \square

We will use Lemma 4.3 to prove Theorem 4.

5. BIRKHOFF SUMS OVER TORAL SKEW-SHIFTS

In what follows $\Phi_{\alpha, \beta}(x, y) = (x + \alpha, y + x + \beta)$ is a (linear) skew-shift on \mathbb{T}^2 , with α of bounded type and $g \in W^s(\mathbb{T}^2)$, with $s > 7/2$ and $\int_{\mathbb{T}^2} g d\lambda_{\mathbb{T}^2} = 0$, where $\lambda_{\mathbb{T}^2}$ denotes the normalized (Haar) Lebesgue measure on \mathbb{T}^2 . We also assume that g is not a coboundary (although some lemmas below are true also for coboundaries).

5.1. COHOMOLOGICAL EQUATION FOR SKEW-SHIFTS. — The cohomological equation for (linear) skew-shifts on \mathbb{T}^2 can be completely solved by Fourier series (see [24], [3]). Let $\Phi_{\alpha, \beta} : \mathbb{T}^2 \rightarrow \mathbb{T}^2$ be given for $\alpha \in [0, 1) \setminus \mathbb{Q}$ by the formula

$$\Phi_{\alpha, \beta}(x, y) = (x + \alpha, y + x + \beta).$$

It follows (see [1]) that

$$L^2(\mathbb{T}^2) = \bigoplus_{(m, n) \in \mathbb{Z}_{|n|} \times \mathbb{Z} \setminus \{0\}} H_{m, n},$$

where the spaces $H_{m, n}$ are $\Phi_{\alpha, \beta}$ -invariant and

$$H_{m, n} = \bigoplus_{j \in \mathbb{Z}} \mathbb{C} e_{m+jn, n} \subset L^2(\mathbb{T}^2)$$

with $e_{a, b}(x, y) = \exp(2\pi i(ax + by))$, for all $(a, b) \in \mathbb{Z}^2$.

THEOREM 5 ([24, Th. 11.25], [3, Th. 10]). — *For every $(m, n) \in \mathbb{Z}^2$, there exists a unique distributional obstruction to the existence of a smooth solution $u \in C^\infty(H_{m, n})$ of the cohomological equation*

$$u \circ \Phi_{\alpha, \beta} - u = g$$

with right hand side $g \in C^\infty(H_{m, n})$. Such an obstruction is the invariant distribution $D_{m, n} \in W^{-s}(\mathbb{T}^2)$ for all $s > 1/2$ defined as follows:

$$D_{m, n}(e_{a, b}) := \begin{cases} e^{-2\pi i[(\alpha m + \beta n)j + \alpha n \binom{j}{2}]} & \text{if } (a, b) = (m + jn, n); \\ 0 & \text{otherwise.} \end{cases}$$

The solution of the cohomological equation, for any function $g \in C^\infty(H_{m, n})$ such that $D_{m, n}(g) = 0$, is given by the following formula. If $g = \sum_{j \in \mathbb{Z}} g_j e_{m+jn, n}$, the solution

$u = \sum_{j \in \mathbb{Z}} u_j e_{m+jn, n}$ has Fourier coefficients:

$$\begin{aligned} u_j &= -e^{2\pi i[(\alpha m + \beta n)j + \alpha n \binom{j}{2}]} \sum_{k=-\infty}^j g_k e^{-2\pi i[(\alpha m + \beta n)k + \alpha n \binom{k}{2}]} \\ &= e^{2\pi i[(\alpha m + \beta n)j + \alpha n \binom{j}{2}]} \sum_{k=j+1}^{\infty} g_k e^{-2\pi i[(\alpha m + \beta n)k + \alpha n \binom{k}{2}]} \end{aligned}$$

If $g \in W^s(H_{m,n})$ for any $s > 1$ and $D_{m,n}(g) = 0$, then the above solution u belongs to $W^t(H_{m,n})$ for all $t < s - 1$ and there exists a constant $C_{s,t} > 0$ such that

$$\|u\|_t \leq C_{s,t} \|g\|_s.$$

The results below establish the quantitative behavior of the square mean of ergodic averages for smooth functions under the skew-shift.

LEMMA 5.1 ([3, Lem. 15] or [13, Lem. 8.1]). — Let $(m, n) \in \mathbb{Z}_{|n|} \times \mathbb{Z} \setminus \{0\}$ and let $s > 1/2$. There exists a constant $C_s > 0$ such that, for any $g \in W^s(H_{m,n})$,

$$\begin{aligned} C_s^{-1} |D_{m,n}(g)| &\leq \liminf_{N \rightarrow +\infty} \frac{1}{N^{1/2}} \left\| \sum_{k=0}^{N-1} g \circ \Phi_{\alpha, \beta}^k \right\|_{L^2(\mathbb{T}^2)} \\ &\leq \limsup_{N \rightarrow +\infty} \frac{1}{N^{1/2}} \left\| \sum_{k=0}^{N-1} g \circ \Phi_{\alpha, \beta}^k \right\|_{L^2(\mathbb{T}^2)} \leq C_s |D_{(m,n)}(g)|. \end{aligned}$$

5.2. GENERAL ESTIMATES

LEMMA 5.2. — There exists a constant $C_{\alpha, g} > 0$ such that, for every $N \in \mathbb{N}$ and every $(x, y) \in \mathbb{T}^2$, we have

$$|S_N(g)(x, y)| \leq C_{\alpha, g} N^{1/2}.$$

Proof. — Since $\alpha \in \mathbb{R} \setminus \mathbb{Q}$ is of bounded type, the statement follows from [12, Lem. 1.4.9] or from [13, Lem. 6.1 & Th. 6.2]. In fact, since any constant roof suspension of $\Phi_{\alpha, \beta}$ is smoothly isomorphic to a Heisenberg nilflow (ϕ_t^W) on a nilmanifold M , generated by a bounded type vector field $W \in \mathfrak{h}$, for any $g \in W^s(\mathbb{T}^2)$ and every $(x, y) \in \mathbb{T}^2$, there exist a function $G \in W^s(M)$ and $p \in M$ such that

$$S_N(g)(x, y) = \int_0^N G \circ \phi_t^W(p) dt.$$

By [12, Lem. 1.4.9], for any Heisenberg triple $\mathcal{F} := (X, Y, Z)$ and for any $\sigma > 2$, there exists a function $B_\sigma(\mathcal{F}, T)$ (defined [12, Eq. (1.71)]) such that, for any function $f \in W^\sigma(M)$ and for all $(p, T) \in M \times \mathbb{R}$,

$$\left| \frac{1}{T} \int_0^T f \circ \phi_t^X(p) dt \right| \leq \frac{B_\sigma(\mathcal{F}, T)}{T} \|f\|_\sigma.$$

For $X = W$ of bounded type, by definition there exists a constant $C > 0$ such that $B_\sigma(\mathcal{F}, T) \leq C_\sigma T^{1/2}$, hence we derive

$$|S_N(g)(x, y)| \leq \left| \int_0^N G \circ \phi_t^W(p) dt \right| \leq C_\sigma N^{1/2} \|G\|_s$$

(see also the comments after the proof of [12, Lem. 1.4.9]). Alternatively, from [13, Th. 6.2] we derive that for $a = (X, Y, Z)$ satisfying an explicit Diophantine condition (depending only on Y) and for any $f \in W^s(M)$, there exists a Hölder cocycle $\beta^f(a, p, T)$ such that

$$\left| \int_0^T f \circ \phi_t^X(p) dt - \beta^f(a, p, T) \right| \leq C_s(a) \|f\|_s,$$

and [13, Lem. 6.1] implies that whenever $X = W$ is of bounded type, the cocycle $\beta^f(a, p, T)$ satisfies the upper bound

$$|\beta^f(a, p, T)| \leq C_s(a) T^{1/2} \|f\|_s,$$

which again implies our statement. \square

LEMMA 5.3. — *There exists a constant $c_g'' > 0$ such that for every $N \in \mathbb{N}$,*

$$\|S_N(g)\|_{C^0(\mathbb{T}^2)} \geq c_g'' N^{1/2}.$$

Proof. — Since g has zero average, but it is not a coboundary, and $\alpha \in \mathbb{R} \setminus \mathbb{Q}$ has bounded type, we can assume that $g \in W^s(H_{m,n})$, for some (m, n) with $n \neq 0$. In fact, otherwise g is the pull back of a function on the circle \mathbb{T} , which belongs to $W^s(\mathbb{T})$ with $s > 7/2$, and since α is of constant type, it follows by Fourier series that g is a coboundary with transfer function $u \in W^t(\mathbb{T})$ for all $t < s - 1$ (in particular $u \in C^2(\mathbb{T})$).

By orthogonality of the decomposition $W^s(\mathbb{T}^2)$ as a direct sum of components $W^s(H_{m,n})$, for all $s \in \mathbb{R}$, we can assume that $g \in W^s(H_{m,n})$, for some (m, n) with $n \neq 0$, hence by Lemma 5.1 there exists $c_g''' > 0$ such that, for all $N \in \mathbb{N}$,

$$\|S_N(g)\|_{L^2(\mathbb{T}^2)} \geq \|S_N(g_{m,n})\|_{L^2(\mathbb{T}^2)} \geq c_g''' N^{1/2}.$$

This finishes the proof. \square

LEMMA 5.4. — *For every $\chi > 0$, $\zeta > 1$ there exists $V_{\chi, \zeta} > 1$ such that for every $T > 0$ there exists $n_\chi, K_\chi \in [T, V_{\chi, \zeta} T]$ for which*

$$\|S_{n_\chi}(g)\|_{C^0(\mathbb{T}^2)} \geq (1 - \chi) \zeta^{-1/2} \|S_{[\zeta n_\chi]}(g)\|_{C^0(\mathbb{T}^2)}$$

and, for $q = \zeta p$,

$$\|S_{qK_\chi}(g)\|_{C^0(\mathbb{T}^2)} \geq (1 - \chi) \zeta^{1/2} \|S_{pK_\chi}(g)\|_{C^0(\mathbb{T}^2)}.$$

Proof. — For $\chi > 0$ let $k \in \mathbb{N}$ be such that $(1 - \chi)^k < c''/C_{\alpha,g}$ and let $V_{\chi,\zeta} := \zeta^k$. By contradiction, if the statement is not true, then

$$\begin{aligned} c''T^{1/2} &\leq \|S_T(g)\|_{C^0(\mathbb{T}^2)} < (1 - \chi)\zeta^{-1/2}\|S_{[\zeta T]}(g)\|_{C^0(\mathbb{T}^2)} \leq \dots \\ &\leq (1 - \chi)^k \zeta^{-k/2}\|S_{[\zeta^k T]}(g)\|_{C^0(\mathbb{T}^2)} \leq C_{\alpha,g}(1 - \chi)^k T^{1/2}. \end{aligned}$$

This contradicts the choice of k . The proof of the second inequality follows the same lines. \square

For $(a, b), (c, d) \in \mathbb{T}^2$ let $d_1((a, b), (c, d)) = \|a - c\|$ and $d_2((a, b), (c, d)) = \|b - d\|$.

LEMMA 5.5. — *There exists $C' = C'_{\alpha,g} > 0$ such that, for every $(x, y), (x', y') \in \mathbb{T}^2$ and for every $n \in \mathbb{N}$, we have*

$$|S_n(g)(x, y) - S_n(g)(x', y')| \leq C'(n^{3/2}d_1((x, y), (x', y')) + n^{1/2}d_2((x, y), (x', y'))).$$

Proof. — By the mean value theorem for $S_n(g)$, we have for some $\theta_n \in \mathbb{T}^2$

$$(19) \quad |S_n(g)(x, y) - S_n(g)(x', y')| = \left| \frac{\partial S_n(g)}{\partial x}(\theta_n)(x - x') + \frac{\partial S_n(g)}{\partial y}(\theta_n)(y - y') \right|.$$

By the chain rule and Lemma 5.2

$$\left| \frac{\partial S_n(g)}{\partial y}(\theta_n) \right| = |S_n(\partial g/\partial y)(\theta_n)| \leq C_{\alpha,g}n^{1/2}.$$

Moreover, by the chain rule, we have

$$\left| \frac{\partial S_n(g)}{\partial x}(\theta_n) \right| \leq |S_n(\partial g/\partial x)(\theta_n)| + \left| \sum_{i=0}^{n-1} i \frac{\partial g}{\partial x}(\Phi_{\alpha,\beta}^i \theta_n) \right|$$

and by summation by parts

$$\left| \sum_{i=0}^{n-1} i \frac{\partial g}{\partial x}(\Phi_{\alpha,\beta}^i \theta_n) \right| \leq |nS_n(\partial g/\partial x)(\theta_n)| + \left| \sum_{r=0}^{n-1} S_r(\partial g/\partial x)(\theta_n) \right|.$$

By Lemma 5.2, for some $C > 0$,

$$|nS_n(\partial g/\partial x)(\theta_n)| < Cn^{3/2} \quad \text{and} \quad \left| \sum_{r=0}^{n-1} S_r(\partial g/\partial x)(\theta_n) \right| < Cn^{3/2}.$$

Using the above estimates in (19) finishes the proof. \square

LEMMA 5.6. — *Fix $q \in \mathbb{N}$. For every $\eta > 0$ there exists $D_\eta > 1$ such that for every $n, m \in \mathbb{N}$ and every $(x, y) \in \mathbb{T}^2$, we have*

$$\max_{i \in \{m, \dots, m + D_\eta n\}} |S_n(g)(\Phi_{\alpha,\beta}^{iq}(x, y))| \geq (1 - \eta)\|S_n(g)\|_{C^0(\mathbb{T}^2)}.$$

Proof. — Since α is of bounded type, there exists $D_\eta > 1$ such that for every $(x, y) \in \mathbb{T}^2$ and every $n \in \mathbb{N}$, the orbit $\{\Phi_{\alpha,\beta}^{iq}(x, y)\}_{i=0}^{D_\eta n}$ is $(\eta^2/n, \eta^2)$ -dense in \mathbb{T}^2 . Notice that, if $(a, b) \in \mathbb{T}^2$ is any point such that $|S_n(g)(a, b)| = \|S_n(g)\|_{C^0(\mathbb{T}^2)}$ and $\|a - c\| \leq \eta/n$ and $\|b - d\| \leq \eta^2$, then by Lemma 5.5, $|S_n(g)(a, b) - S_n(g)(c, d)| \leq 2C'\eta^2 n^{1/2}$, which together with Lemma 5.3 finishes the proof is $\eta > 0$ is small enough. \square

Recall that (q_n) denotes the sequence of denominators of α . The following simple lemma is a consequence of the pigeonhole principle:

LEMMA 5.7. — *Fix $p, q \in \mathbb{N}$. For every $\eta > 0$ there exists $L_\eta > 0$ such that for every $(x, y), (z, w) \in \mathbb{T}^2$ and for every $n \in \mathbb{N}$, there exists $\ell_{1,n}, \ell_{2,n} \in \{0, \dots, L_\eta\}$, $\ell_{1,n} \neq \ell_{2,n}$ such that*

$$d_2(\Phi_{\alpha,\beta}^{q\ell_{1,n}q_n}(x, y), (\Phi_{\alpha,\beta}^{q\ell_{2,n}q_n}(x, y))) < \eta \quad \text{and} \quad d_2(\Phi_{\alpha,\beta}^{p\ell_{1,n}q_n}(z, w), (\Phi_{\alpha,\beta}^{p\ell_{2,n}q_n}(z, w))) < \eta.$$

Proof. — Let $L_\eta := 2\eta^{-2}$. By the pigeonhole principle there exist positive integers $\ell_1, \dots, \ell_{[\eta^{-1}]+1} \in \{0, \dots, L_\eta\}$ such that $d_2(\Phi_{\alpha,\beta}^{q\ell_n q_n}(x, y), (\Phi_{\alpha,\beta}^{q\ell'_n q_n}(x, y))) < \eta$ for $\ell, \ell' \in \{\ell_1, \dots, \ell_{[\eta^{-1}]+1}\}$. Again by the pigeonhole principle, there exist positive integers $\ell_{1,n}, \ell_{2,n} \in \{\ell_1, \dots, \ell_{[\eta^{-1}]+1}\}$ such that $d_2(\Phi_{\alpha,\beta}^{p\ell_{1,n}q_n}(z, w), (\Phi_{\alpha,\beta}^{p\ell_{2,n}q_n}(z, w))) < \eta$. This finishes the proof. \square

LEMMA 5.8. — *There exists a constant $C''' > 0$ such that for any $(a, b), (c, d) \in \mathbb{T}^2$, for every $W, K \in \mathbb{N}$ with $W \leq \|a - c\|^{-1}$, we have*

$$\begin{aligned} & |S_K(g)(\Phi_{\alpha,\beta}^W(a, b)) - S_K(g)(\Phi_{\alpha,\beta}^W(c, d)) - WS_K\left(\frac{\partial g}{\partial y}\right)(\Phi_{\alpha,\beta}^W(a, b))(a - c)| \\ & \leq C'''(K^{1/2}W^2\|a - c\|^2 + K^{3/2}\|a - c\| + K^{1/2}\|b - d\|). \end{aligned}$$

Proof. — Let

$$\theta_W = (a + W\alpha, d + Wc + W(W - 1)\alpha/2)$$

and

$$\theta'_W = (a + W\alpha, b + Wc + W(W - 1)\alpha/2).$$

Then

$$\begin{aligned} S_K(g)(\Phi_{\alpha,\beta}^W(a, b)) - S_K(g)(\Phi_{\alpha,\beta}^W(c, d)) &= (S_K(g)(\Phi_{\alpha,\beta}^W(a, b)) - S_K(g)(\theta'_W)) \\ &+ (S_K(g)(\theta'_W) - S_K(g)(\theta_W)) + (S_K(g)(\theta_W) - S_K(g)(\Phi_{\alpha,\beta}^W(c, d))). \end{aligned}$$

By Lemma 5.5, we have

$$|S_K(g)(\theta'_W) - S_K(g)(\theta_W)| \leq C'K^{1/2}\|b - d\|$$

and

$$|S_K(g)(\theta_W) - S_K(g)(\Phi_{\alpha,\beta}^W(c, d))| \leq C'K^{3/2}\|a - c\|.$$

Finally, by Taylor formula and the chain rule, for some $\theta_K \in \mathbb{T}^2$,

$$\begin{aligned} S_K(g)(\Phi_{\alpha,\beta}^W(a, b)) - S_K(g)(\theta'_W) \\ = S_K(\partial g/\partial y)(\Phi_{\alpha,\beta}^W(a, b))W(a - c) + S_K(\partial^2 g/\partial^2 y)(\theta_K)(W(a - c))^2, \end{aligned}$$

and by Lemma 5.2, we get

$$S_K(\partial^2 g/\partial^2 y)(\theta_K)(W(a - c))^2 \leq C''K^{1/2}W^2\|a - c\|^2.$$

This finishes the proof. \square

5.3. ESTIMATES OF SECOND ORDER TERMS. — In what follows $p, q \in \mathbb{N}$, $\zeta = q/p$, and a zero-mean non-coboundary $h \in W^s(M)$, $s \geq 5/2$ is fixed. We assume WLOG that $q > p$, that is $\zeta > 1$.

The following lemma is important, it crucially uses the fact that $p \neq q$:

LEMMA 5.9. — *There exist $D', d' > 0$ such that, for any $(x, y), (z, w) \in \mathbb{T}^2$ and any $T > 0$, for some $n' \in [0, D'T]$, we have*

$$(20) \quad |S_{n'}(h)(z, w) - \zeta^{-1/2} S_{[\zeta n']}(h)(x, y)| \geq d'T^{1/2}.$$

Proof. — We will consider only numbers of the form $n' = pk$, for $k \in \mathbb{N}$. We will show that there exist $D', d' > 0$ such that

$$(21) \quad \left| \left(S_{pK}(h)(\Phi_{\alpha, \beta}^{pW}(z, w)) - S_{pK}(h)(\Phi_{\alpha, \beta}^{pW} \Phi_{\alpha, \beta}^{pQ}(z, w)) \right) - \zeta^{-1/2} \left(S_{qK}(h)(\Phi_{\alpha, \beta}^{qW} x, y) - S_{qK}(h)(\Phi_{\alpha, \beta}^{qW} \Phi_{\alpha, \beta}^{qQ}(x, y)) \right) \right| \geq 16d'T^{1/2},$$

for some $K, W, Q \leq D'T$. This will finish the proof since by cocycle identity

$$\begin{aligned} S_{pK}(h)(\Phi_{\alpha, \beta}^{pW}(z, w)) - S_{pK}(h)(\Phi_{\alpha, \beta}^{p(W+Q)}(z, w)) \\ = S_{p(W+Q)}(h)(z, w) - S_{p(K+W+Q)}(h)(z, w) + S_{p(K+W)}(h)(z, w) - S_{pW}(h)(z, w) \end{aligned}$$

and the same splitting for $S_{qK}(h)(\cdot)$. Hence (20) then holds for n' being one of $\{pW, p(K+W), p(W+Q), p(K+W+Q)\}$.

Let $\eta > 0$ be small. By Lemma 5.4, let $K \in [T, V_{\eta, \zeta} T]$ be such that

$$(22) \quad \|S_{qK}(\partial h / \partial y)\|_{C^0(\mathbb{T}^2)} \geq (1 - \eta) \zeta^{1/2} \|S_{pK}(\partial h / \partial y)\|_{C^0(\mathbb{T}^2)}.$$

Let now $n = n(K, \eta) > 0$ be the smallest number such that $\eta^{3/2} q_{n+1} / L_\eta \geq D_\eta q K$ (L_η from Lemma 5.7 and $D_\eta > 1$ from Lemma 5.6). Let $\ell_{1,n}, \ell_{2,n} \in \{0, \dots, L_\eta\}$ be as in Lemma 5.7 for n, x, y, z, w and let $\ell_n = \ell_{2,n} - \ell_{1,n}$. Denote $(\bar{x}, \bar{y}) = \Phi_{\alpha, \beta}^{q\ell_{1,n} q_n}(x, y)$ and $(\bar{z}, \bar{w}) = \Phi_{\alpha, \beta}^{p\ell_{1,n} q_n}(z, w)$.

Notice that by definition, $2(\eta^{1/2} q_{n+1} / \ell_n) - \eta^{1/2} q_{n+1} / \ell_n \geq D_\eta q K$, hence by Lemma 5.6 there exists $W \in [\eta^{1/2} q_{n+1} / \ell_n, 2\eta^{1/2} q_{n+1} / \ell_n]$ such that

$$(23) \quad |S_{qK}(\partial h / \partial y)(\Phi_{\alpha, \beta}^{qW}(\bar{x}, \bar{y}))| \geq (1 - \eta) \|S_{qK}(\partial h / \partial y)\|_{C^0(\mathbb{T}^2)} (\geq c' q^{1/2} K^{1/2}).$$

Therefore,

$$(24) \quad |S_{qK}(\partial h / \partial y)(\Phi_{\alpha, \beta}^{qW}(\bar{x}, \bar{y}))| q^2 W \ell_n \|q_n \alpha\| \geq c' q^{5/2} K^{1/2} W \ell_n \|q_n \alpha\|.$$

Since $W \ell_n \|q_n \alpha\| \in [\eta^{1/2} / 2, 2\eta^{1/2}]$, if $\eta \ll \min(p, q)$, we have for $b \in \{p, q\}$

$$\eta^{1/10} c' q^{5/2} K^{1/2} W \ell_n \|q_n \alpha\| \geq (bK)^{1/2} (W b^2 \ell_n \|q_n \alpha\|)^2,$$

similarly

$$\eta^{1/10} c' q^{5/2} K^{1/2} W \ell_n \|q_n \alpha\| \geq (bK)^{1/2} \eta,$$

and since $W \geq \eta^{1/2} q_{n+1} / 2\ell_n \geq \eta^{-1} D_\eta q K / 2$, for $\eta > 0$ sufficiently small, we have

$$\eta^{1/10} c' q^{5/2} K^{1/2} W \ell_n \|q_n \alpha\| \geq (bK)^{3/2} b \ell_n \|q_n \alpha\|.$$

Notice that by Lemma 5.8 for $(a, b) = (\bar{x}, \bar{y})$ and $(c, d) = \Phi_{\alpha, \beta}^{q\ell_n q_n}(\bar{x}, \bar{y})$, by formula (24) and the three above equations for $b = q$, we have

$$\begin{aligned} & |S_{qK}(h)(\Phi_{\alpha, \beta}^{qW}(\bar{x}, \bar{y})) - S_{qK}(h)(\Phi_{\alpha, \beta}^{qW}(\Phi_{\alpha, \beta}^{q\ell_n q_n}(\bar{x}, \bar{y})))| \\ & \geq |S_{qK}(\partial h / \partial y)(\Phi_{\alpha, \beta}^{qW}(\bar{x}, \bar{y}))| q^2 W \ell_n \|q_n \alpha\| \\ & \quad - C''' \left((qK)^{1/2} (W q^2 \ell_n \|q_n \alpha\|)^2 + (qK)^{1/2} \eta + (qK)^{3/2} q \ell_n \|q_n \alpha\| \right) \\ & \geq (1 - 3\eta^{1/10} C''') |S_{qK}(\partial h / \partial y)(\Phi_{\alpha, \beta}^{qW}(\bar{x}, \bar{y}))| q^2 W \ell_n \|q_n \alpha\|. \end{aligned}$$

Moreover, by (23) and (22), we have

$$|S_{qK}(\partial h / \partial y)(\Phi_{\alpha, \beta}^{qW}(\bar{x}, \bar{y}))| \geq (1 - \eta)^2 \zeta^{1/2} |S_{pK}(\partial h / \partial y)(\Phi_{\alpha, \beta}^{pW}(\bar{z}, \bar{w}))|.$$

Therefore, by Lemma 5.8, (24) and the three above equations for $b = p$, we have

$$\begin{aligned} & |S_{pK}(h)(\Phi_{\alpha, \beta}^{pW}(\bar{z}, \bar{w})) - S_{pK}(h)(\Phi_{\alpha, \beta}^{pW}(\Phi_{\alpha, \beta}^{p\ell_n q_n}(\bar{z}, \bar{w})))| \\ & \leq |S_{pK}(\partial h / \partial y)(\Phi_{\alpha, \beta}^{pW}(\bar{z}, \bar{w}))| p^2 W \ell_n \|q_n \alpha\| \\ & \quad + C''' \left((pK)^{1/2} (p^2 W \ell_n \|q_n \alpha\|)^2 + (pK)^{1/2} \eta + (pK)^{3/2} p \ell_n \|q_n \alpha\| \right) \\ & \leq (p^2 (1 - \eta)^{-2} \zeta^{-1/2} + 3\eta^{1/10} C'''' q^2) |S_{qK}(\partial h / \partial y)(\Phi_{\alpha, \beta}^{qW}(\bar{x}, \bar{y}))| W \ell_n \|q_n \alpha\|. \end{aligned}$$

Let $d_{p,q,\eta} := q^2 - p^2 (1 - \eta)^{-2} \zeta^{-1} - C'''' (3 + \zeta^{-1/2}) \eta^{1/10} q^2$; then $d_{p,q,\eta} > 0$ if $\eta > 0$ is small enough (since $q > p$).

We have by the above

$$\begin{aligned} & |S_{qK}(h)(\Phi_{\alpha, \beta}^{qW}(\bar{x}, \bar{y})) - S_{qK}(h)(\Phi_{\alpha, \beta}^{qW}(\Phi_{\alpha, \beta}^{q\ell_n q_n}(\bar{x}, \bar{y})))| \\ & \quad - \zeta^{-1/2} |S_{pK}(h)(\Phi_{\alpha, \beta}^{pW}(\bar{z}, \bar{w})) - S_{pK}(h)(\Phi_{\alpha, \beta}^{pW}(\Phi_{\alpha, \beta}^{p\ell_n q_n}(\bar{z}, \bar{w})))| \\ & \geq d_{p,q,\eta} |S_{qK}(\partial h / \partial y)(\Phi_{\alpha, \beta}^{qW}(\bar{x}, \bar{y}))| W \ell_n \|q_n \alpha\| \\ & \geq d'_{p,q,\eta} (qK)^{1/2} W \ell_n \|q_n \alpha\| \geq d''_{p,q,\eta} K^{1/2}. \end{aligned}$$

Since $K \geq T$ and by the definition of (\bar{x}, \bar{y}) , (\bar{z}, \bar{w}) , this finishes the proof of (21) with $K = K$, $W = W + \ell_{1,n} q_n$ and $Q = \ell_n q_n$, hence the proof of Lemma 5.9 is complete. \square

For $(x, y), (x, y') \in \mathbb{T}^2$, let $\delta_y = |y - y'|$. We also have the following lemma:

LEMMA 5.10. — *There exist $D'' = D''(\alpha, f, p, q) > 0$, $d'' = d''(\alpha, f, p, q) > 0$ such that for every $(x, y), (x, y'), (z, w), (z, w') \in \mathbb{T}^2$, if $T' := \min(\delta_y^{-2}, \delta_w^{-2}) \gg 1$, then there exists $s \in [0, D'' T']$ such that*

$$(25) \quad |(S_s(p^{-1}f)(z, w) - S_s(p^{-1}f)(z, w')) - (S_{[s]}(q^{-1}f)(x, y) - S_{[s]}(q^{-1}f)(x, y'))| \geq d''.$$

Proof. — Notice that by Taylor formula and the chain rule, for some $\theta \in \mathbb{T}^2$

$$\begin{aligned} & S_s(p^{-1}f)(z, w) - S_s(p^{-1}f)(z, w') \\ & = (w - w') p^{-1} S_s(\partial f / \partial y)(z, w) + (w - w')^2 p^{-1} S_s(\partial^2 f / \partial^2 y)(\theta), \end{aligned}$$

and by Lemma 5.2 for $\partial^2 f / \partial^2 y$

$$|(w - w')^2 p^{-1} S_s(\partial^2 f / \partial^2 y)(\theta)| \leq C_{\alpha, f} p^{-1} \delta_w^2 s^{1/2} < T'^{-1/3}$$

for $s \leq T'^{11/10}$. An analogous reasoning for $S_{[\zeta s]}(q^{-1} f)(x, y) - S_{[\zeta s]}(q^{-1} f)(x, y')$ shows that (25) follows by showing that there exist $D'', d'' > 0$ such that

$$(26) \quad |(w - w') p^{-1} S_s(\partial f / \partial y)(z, w) - (y - y') q^{-1} S_{[\zeta s]}(\partial f / \partial y)(x, y)| \geq 2d''$$

for some $s \in [0, D'' T']$.

To simplify notation let $h = \partial f / \partial y$. Let $\chi \in (0, 1)$ be a small number, $\chi \ll d' / D'$ (d', D' from Lemma 5.9). We will consider two cases.

$$(A) \quad | |(w - w') q^{1/2} / (y - y') p^{1/2} | - 1 | \geq \chi.$$

We will (WLOG) assume that $|(w - w') q^{1/2}| \geq (1 + \chi) |(y - y') p^{1/2}|$, the proof in the case $|(w - w') q^{1/2}| \leq (1 - \chi) |(y - y') p^{1/2}|$ is symmetric and follows the same lines. Notice that in this case

$$T' \leq \frac{2p}{q} \delta_w^{-2}.$$

Let $V_{\chi/3, \zeta} > 0$ come from Lemma 5.4 and let $n_0 \in [T', V_{\chi/3, \zeta} T']$ be as in the statement of Lemma 5.4 for $g = h$. Then

$$(27) \quad \|S_{n_0}(h)\|_{C_0(\mathbb{T}^2)} \geq (1 - \chi/3) \zeta^{-1/2} \|S_{[\zeta n_0]}(h)\|_{C_0(\mathbb{T}^2)}.$$

Let $D_{\chi/3} > 0$ come from Lemma 5.3 and let $u = u(n_0) \in \{0, \dots, D_{\chi/3} n_0\}$ be such that

$$(28) \quad |S_{n_0}(h)(\Phi_{\alpha, \beta}^u(z, w))| \geq (1 - \chi/3) \|S_{n_0}(h)\|_{C_0(\mathbb{T}^2)}.$$

Let $v = [\zeta(n_0 + u)] - [\zeta n_0]$. We will show that

$$(29) \quad p^{-1} |S_{n_0}(h)(\Phi_{\alpha, \beta}^u(z, w))| |w - w'| \geq (1 + \chi/100) q^{-1} |S_{[\zeta n_0]}(h)(\Phi_{\alpha, \beta}^v(x, y))| |y - y'|.$$

Then

$$\begin{aligned} & |(w - w') p^{-1} S_{n_0}(h)(\Phi_{\alpha, \beta}^u(z, w)) - (y - y') q^{-1} S_{[\zeta n_0]}(h)(\Phi_{\alpha, \beta}^v(x, y))| \\ & \geq (1 - (1 + \chi/100)^{-1}) |w - w'| p^{-1} |S_{n_0}(h)(\Phi_{\alpha, \beta}^u(z, w))| \geq d'' |w - w'| n_0^{1/2} \\ & \geq d''(\chi) \delta_w T'^{1/2} \geq d''(\chi), \end{aligned}$$

for some $d''(\chi) > 0$. But then by cocycle identity we know that (26) holds for $s = n_0$ or $s = n_0 + u$ and $d'(\chi) = d''(\chi)/2 > 0$.

Therefore, it only remains to show (29). By (28) and (27) and the assumptions of (A), we have

$$\begin{aligned} & p^{-1} |S_{n_0}(h)(\Phi_{\alpha, \beta}^u(z, w))| |w - w'| \\ & \geq p^{-1} (1 - \chi/3)^2 \zeta^{-1/2} \|S_{[\zeta n_0]}(h)\|_{C_0(\mathbb{T}^2)} (1 + \chi) \frac{p^{1/2}}{q^{1/2}} |y - y'| \\ & \geq q^{-1} (1 + \chi/100) |S_{[\zeta n_0]}(h)(\Phi_{\alpha, \beta}^v(x, y))| |y - y'|. \end{aligned}$$

This finishes the proof of (29) and hence also the proof of case (A).

$$\mathbf{(B)} \quad \left| |(w - w')q^{1/2}/(y - y')p^{1/2} - 1| \right| \leq \chi.$$

In this case the LHS in (26) is larger than

$$|w - w'|p^{-1}|S_s(h)(z, w) - \zeta^{-1/2}S_{[\zeta s]}(h)(x, y)| - \chi|y - y'|q^{-1}|S_{[\zeta s]}(h)(x, y)|.$$

By Lemma 5.9 for $T = T'$ and the definition of T' , there exists an $s_0 \in [0, D'T']$ such that

$$|w - w'|p^{-1}|S_{s_0}(h)(z, w) - \zeta^{-1/2}S_{[\zeta s_0]}(h)(x, y)| \geq p^{-1}d'.$$

Moreover, by Lemma 5.2 and the definition of T' for h it follows that

$$\chi|y - y'|q^{-1}|S_{[\zeta s_0]}(h)(x, y)| \leq C_{\alpha, h}(D')^{1/2}\zeta^{1/2}q^{-1}\chi < p^{-1}\frac{d'}{10},$$

if $\chi > 0$ is sufficiently small. This finishes the proof of Lemma 5.10. \square

6. RATNER'S PROPERTY: PROOF OF THEOREM 3

In this section we will use the estimates from Section 5 to prove Theorem 3. We will use Proposition 3.2. Before we do that, we will prove a crucial proposition: For $(x, y), (x', y') \in \mathbb{T}^2$ denote $|x - x'| = \delta_x$ and $|y - y'| = \delta_y$.

PROPOSITION 6.1. — *For any $\alpha \in \mathbb{R} \setminus \mathbb{Q}$ of bounded type and for any $f \in W^s(\mathbb{T}^2)$, $s > 7/2$, there exists a constant $D_{\alpha, f} > 0$ such that the following holds. For every $(x, y), (x', y') \in \mathbb{T}^2$, if $T := \min(\delta_x^{-2/3}, \delta_y^{-2})$, there exists $n_0 = n_0((x, y), (x', y')) \in [0, D_{\alpha, f}T] \cap \mathbb{Z}$ such that*

$$(30) \quad |S_{n_0}(f)(x, y) - S_{n_0}(f)(x', y')| > 1.$$

Proof. — Denote $a_n = S_n(f)(x, y) - S_n(f)(x', y')$, let $\eta < 1/100$ and let $D_\eta > 0$ come from Lemma 5.6 for $g = \partial f / \partial y$. Let moreover $c'' > 0$ be as in Lemma 5.3 for $g = \partial f / \partial y$ and $C' > c'' > 0$ be as in Lemma 5.5 for $g = f - \int_{\mathbb{T}^2} f d\lambda_{\mathbb{T}^2}$. Let $k := [100T/c''^2] + 1$. We will show that there exists $m \in [10C'k/c'', (10C'k/c'') + D_\eta k + 1] \cap \mathbb{Z}$ such that

$$(31) \quad |S_k(f)(\Phi_{\alpha, \beta}^m(x, y)) - S_k(f)(\Phi_{\alpha, \beta}^m(x', y'))| \geq 3.$$

This will finish the proof since then, by cocycle identity $|a_{m+k} - a_k| \geq 3$ and consequently (30) holds for $n_0 = m + k$ or $n_0 = k$. By Lemma 5.6 and 5.3 there exists $m_1 \in [10C'k/c'', (10C'k/c'') + D_\eta k + 1] \cap \mathbb{Z}$ and $m_2 \in [10m_1, 10m_1 + D_\eta k + 1]$ such that

$$(32) \quad |S_k(\partial f / \partial y)(\Phi_{\alpha, \beta}^m(x, y))| \geq (1 - \eta)c''k^{1/2}, \quad \text{for } m = m_1, m_2.$$

Moreover, since $m_2 \geq 10m_1$, by the triangle inequality we have

$$\max_{m \in \{m_1, m_2\}} |m(x - x') + (y - y')| \geq \max\left(\frac{m_1}{2}\delta_x, \frac{\delta_y}{2}\right).$$

Let now m denote the element above which attains the maximum. Since $m \geq 10C'k/c''$, we have

$$(33) \quad (1 - \eta)c''|m(x - x') + (y - y')|k^{1/2} \geq 2C'\delta_x k^{3/2}.$$

Let $z_m := (x + m\alpha, y' + mx' + m(m - 1)\alpha/2)$. Notice that

$$(34) \quad S_k(f)(\Phi_{\alpha,\beta}^m(x, y)) - S_k(f)(\Phi_{\alpha,\beta}^m(x', y')) = (S_k(f)(\Phi_{\alpha,\beta}^m(x, y)) - S_k(f)(z_m)) + (S_k(f)(z_m) - S_k(f)(\Phi_{\alpha,\beta}^m(x', y'))).$$

By Lemma 5.5 for $g = f - \int_{\mathbb{T}^2} f d\lambda_{\mathbb{T}^2}$, we have

$$|S_k(f)(\Phi_{\alpha,\beta}^m(x', y')) - S_k(f)(z_m)| \leq C' \delta_x k^{3/2}.$$

By Taylor formula (and chain rule), for some $\theta_k \in \mathbb{T}^2$,

$$S_k(f)(\Phi_{\alpha,\beta}^m(x, y)) - S_k(f)(z_m) = (m(x - x') + (y - y'))S_k(\partial f/\partial y)(\Phi_{\alpha,\beta}^m(x, y)) + (m(x - x') + (y - y'))^2 S_k(\partial^2 f/\partial y^2)(\theta_k).$$

Since $m \leq C'''T$ (for some $C''' > 0$), we have $|m(x - x') + (y - y')| \leq m\delta_x + \delta_y \leq T^{-1/3}$, for $T > 1$ large enough, and therefore by Lemma 5.2 for $g = \partial f/\partial y$, we have

$$|(m(x - x') + (y - y'))^2 S_k(\partial^2 f/\partial y^2)(\theta_k)| \leq 1/100.$$

Therefore, by (33), (32) and (34), we have

$$\begin{aligned} |S_k(f)(\Phi_{\alpha,\beta}^m(x, y)) - S_k(f)(\Phi_{\alpha,\beta}^m(x', y'))| &\geq (1 - \eta)c''|m\delta_x + \delta_y|k^{1/2} - 1/100 - C'\delta_x k^{3/2} \\ &\geq \frac{c''}{3}|m\delta_x + \delta_y|k^{1/2} - 1/100. \end{aligned}$$

However, since $m_1 \geq 10C'k/c'' > 10k$, we have

$$\frac{c''}{3}|m\delta_x + \delta_y|k^{1/2} \geq \frac{c''}{3} \max\left(\frac{m_1}{2}\delta_x, \frac{\delta_y}{2}\right)k^{1/2} \geq 2,$$

the last inequality since $k = [100T/c''^2] + 1$. This finishes the proof of (31) and hence also the proof of Proposition 6.1. \square

Now we can prove Theorem 3:

Proof of Theorem 3. — We will use Proposition 3.2. Fix $\varepsilon > 0$ and $N \in \mathbb{N}$. Let $Z = \mathbb{T}^2$ (see Definition 3.1), and take any $(x, y), (x', y') \in \mathbb{T}^2$ with $d((x, y), (x', y')) = \delta_x + \delta_y < \delta$ (we will specify δ and κ in the proof). Let

$$(35) \quad T := \min(\delta_x^{-2/3}, \delta_y^2).$$

Denote $a_k = S_k(f)(x, y) - S_k(f)(x', y')$ and let $D_{\alpha,f} > 0$ be as in Proposition 6.1. Notice that for every $k \in [0, 2D_{\alpha,f}T]$, we have

$$(36) \quad \begin{aligned} d_1((\Phi_{\alpha,\beta}^k(x, y)), (\Phi_{\alpha,\beta}^k(x', y'))) &= |x - x'| = \delta_x \\ \text{and} \quad d_2((\Phi_{\alpha,\beta}^k(x, y)), (\Phi_{\alpha,\beta}^k(x', y'))) &\leq k\delta_x + \delta_y \leq 2D_{\alpha,f}\delta_x^{1/3} + \delta_y, \end{aligned}$$

hence by Lemma 5.5 and by the cocycle identity, for any $\varepsilon > 0$ there exists $\delta_\varepsilon > 0$ such that for $\delta \in (0, \delta_\varepsilon)$, we have that $|a_{k+1} - a_k| < \varepsilon/4$ for every $k \in [0, D_{\alpha,f}T]$. Since by Proposition 6.1, there exists $n_0 \in [0, D_{\alpha,f}T]$ such that $|a_{n_0}| > 1$, it follows that there exists $M \in [0, D_{\alpha,f}T]$ such that

$$\min\{|a_M + 1|, |a_M - 1|\} \leq \varepsilon/3.$$

Notice that by Lemma 5.5, for every $N \in \mathbb{N}$ there exists $\delta_N > 0$ such that for $\delta \in (0, \delta_N)$, we have $M \geq N$. Let $L = \kappa M$. Notice that $[M, M + L] \subset [0, 2D_{\alpha, f}T]$ and therefore

$$(37) \quad d(\Phi_{\alpha, \beta}^n(x, y), \Phi_{\alpha, \beta}^n(x', y')) < \varepsilon, \quad \text{for every } n \in [M, M + L].$$

Moreover, by the cocycle identity, for every $n \in [M, M + L]$, we have

$$|a_n - a_M| = |S_{n-M}(f)(\Phi_{\alpha, \beta}^M(x, y)) - S_{n-M}(f)(\Phi_{\alpha, \beta}^M(x', y'))|.$$

By Lemma 5.5 (for $g = f - \int_{\mathbb{T}^2} f d\lambda_{\mathbb{T}^2}$), (36) for $k = M$ and (35), we have (since $n - M \leq L = \kappa M \leq D_{\alpha, f}\kappa T$), there exists $\kappa_\varepsilon > 0$ such that for $\kappa \in (0, \kappa_\varepsilon)$, we have

$$\begin{aligned} & |S_{n-M}(f)(\Phi_{\alpha, \beta}^M(x, y)) - S_{n-M}(f)(\Phi_{\alpha, \beta}^M(x', y'))| \\ & \leq C'(D_{\alpha, f}\kappa T)^{3/2}\delta_x + C'(D_{\alpha, f}\kappa T)^{1/2}(2D_{\alpha, f}\delta_x^{1/3} + \delta_y) \\ & \leq C'(D_{\alpha, f}\kappa)^{3/2} + C'(D_{\alpha, f}\kappa)^{1/2}(2D_{\alpha, f} + 1) \leq \varepsilon/3. \end{aligned}$$

Therefore, for every $n \in [M, M + L]$,

$$|a_n + 1| \leq |a_n - a_M| + |a_M + 1| \leq \varepsilon \quad \text{or} \quad |a_n - 1| \leq |a_n - a_M| + |a_M - 1| \leq \varepsilon.$$

This property, together with (37), finishes the proof of the hypotheses of Proposition 3.2 and hence also the proof of Theorem 3. \square

7. DISJOINTNESS: PROOF OF THEOREM 4

Notice that if (Φ_t^f) is a special flow over T and under f then (Φ_{rt}^f) is isomorphic to $(\Phi_t^{r^{-1}f})$. We will therefore use Lemma 4.3 for $(\Phi_t^{\alpha, p^{-1}f})$ and $(\Phi_t^{\alpha, q^{-1}f})$ (notice that by Corollary 1.2 we know that $(\Phi_t^{\alpha, f})$, and hence also $(\Phi_t^{\alpha, p^{-1}f})$ and $(\Phi_t^{\alpha, q^{-1}f})$, are weakly mixing).

We assume WLOG that $\zeta = q/p > 1$. Recall that for $(x, y), (x', y') \in \mathbb{T}^2$, $\delta_x = |x - x'|$ and $\delta_y = |y - y'|$. For the proof of Theorem 4 we need the following crucial proposition:

PROPOSITION 7.1. — *There exists $D''' = D'''(\alpha, f, p, q) > 0$ and $d''' = d'''(\alpha, f, p, q) > 0$ such that for every $(x, y), (x', y'), (z, w), (z, w') \in \mathbb{T}^2$ if*

$$T := \min(\delta_x^{-2/3}, \delta_y^{-2}, \delta_w^{-2}),$$

then for some $s \in [0, D'''T]$, we have

$$(38) \quad |(S_s(p^{-1}f)(z, w) - S_s(p^{-1}f)(z, w')) - (S_{[\zeta s]}(q^{-1}f)(x, y) - S_{[\zeta s]}(q^{-1}f)(x', y'))| \geq d'''.$$

Proof. — Let $C_{p, q} > 1$ be a large constant (specified at the end of Case 1). We will consider the following cases.

Case 1. $\delta_w \geq C_{p,q} \max(\delta_y, \delta_x^{1/3})$ or $\delta_w \leq C_{p,q}^{-1} \max(\delta_y, \delta_x^{1/3})$.

If $\delta_w \geq C_{p,q} \max(\delta_y, \delta_x^{1/3})$, then $T = \delta_w^{-2}$. By Proposition 6.1 for (z, w) and (z, w') we have

$$|S_s(p^{-1}f)(z, w) - S_s(p^{-1}f)(z, w')| \geq p^{-1}$$

for some $s \leq D_{\alpha,f}T$. Moreover, by Lemma 5.5, for (x, y) and (x', y') we have

$$\begin{aligned} & |S_{[\zeta s]}(q^{-1}f)(x, y) - S_{[\zeta s]}(q^{-1}f)(x', y')| \\ & \leq C'q^{-1}((\zeta D_{\alpha,f}T)^{3/2}\delta_x + (\zeta D_{\alpha,f}T)^{1/2}\delta_y) \leq \frac{C'q^{-1}\zeta^{3/2}D_{\alpha,f}^{3/2}}{C_{p,q}} \leq \frac{p^{-1}}{2}, \end{aligned}$$

if $C_{p,q} > 1$ is large enough.

Similarly, if $\delta_w \leq C_{p,q}^{-1} \max(\delta_y, \delta_x^{1/3})$, using Proposition 6.1 for (x, y) and (x', y') we have

$$|S_{[\zeta s]}(q^{-1}f)(x, y) - S_{[\zeta s]}(q^{-1}f)(x', y')| \geq q^{-1},$$

for some $s \leq \zeta^{-1}D_{\alpha,f}T$. Moreover, by Lemma 5.5, for (z, w) and (z, w') we have

$$|S_s(p^{-1}f)(z, w) - S_s(p^{-1}f)(z, w')| \leq p^{-1} \frac{C'D_{\alpha,f}^{1/2}\zeta^{-1/2}}{C_{p,q}} \leq \frac{q^{-1}}{2},$$

if $C_{p,q} > 1$ is large enough. This finishes the proof of Case 1.

Case 2. $C_{p,q} \max(\delta_y, \delta_x^{1/3}) \geq \delta_w \geq C_{p,q}^{-1} \max(\delta_y, \delta_x^{1/3})$.

Let $R' = R'_{p,q} > 1$ be a constant to be specified later (at the end of the proof of Subcase 1).

We consider two subcases.

Subcase 1. $R'\delta_x^{1/3} \leq \delta_y$. — Notice that, by Lemma 5.5, we have, for all $k \in \mathbb{N}$,

$$|S_{[\zeta k]}(q^{-1}f)(x', y') - S_{[\zeta k]}(q^{-1}f)(x, y')| \leq q^{-1}C'(\zeta k)^{3/2}\delta_x \leq \frac{q^{-1}C'(\zeta k)^{3/2}}{R'^3T^{3/2}}.$$

Moreover, in this case we have $T = \min(\delta_y^{-2}, \delta_w^{-2})$, hence by Lemma 5.10, there exists $s \in [0, D''T]$ such that

$$|(S_s(p^{-1}f)(z, w) - S_s(p^{-1}f)(z, w')) - (S_{[\zeta s]}(q^{-1}f)(x, y) - S_{[\zeta s]}(q^{-1}f)(x, y'))| \geq d''.$$

Hence if only $q^{-1}C'(\zeta D''T)^{3/2}/R'^3T^{3/2} \leq d''/2$ (which is true if R' is large enough), then (38) follows by triangle inequality.

Subcase 2. $R'\delta_x^{1/3} \geq \delta_y$. — Notice that by Lemma 5.5, for every $k, m \in \mathbb{N}$

$$(39) \quad |S_k(p^{-1}f)(\Phi_{\alpha,\beta}^m(z, w)) - S_k(p^{-1}f)(\Phi_{\alpha,\beta}^m(z, w'))| \leq C'p^{-1}k^{1/2}\delta_w.$$

Moreover, for every $m \in \mathbb{N}$ denote $h_m = (x' + m\alpha, y + mx + m(m-1)\alpha/2)$. Then

$$\begin{aligned} (40) \quad & S_{[\zeta k]}(q^{-1}f)(\Phi_{\alpha,\beta}^m(x, y)) - S_{[\zeta k]}(q^{-1}f)(\Phi_{\alpha,\beta}^m(x', y')) \\ & = (S_{[\zeta k]}(q^{-1}f)(\Phi_{\alpha,\beta}^m(x, y)) - S_{[\zeta k]}(q^{-1}f)(h_m)) \\ & \quad + (S_{[\zeta k]}(q^{-1}f)(h_m) - S_{[\zeta k]}(q^{-1}f)(\Phi_{\alpha,\beta}^m(x', y'))). \end{aligned}$$

By Lemma 5.5 for $g = f - \int_{\mathbb{T}^2} f d\lambda_{\mathbb{T}^2}$, we have

$$(41) \quad |S_{[\zeta k]}(q^{-1}f)(\Phi_{\alpha,\beta}^m(x, y)) - S_{[\zeta k]}(q^{-1}f)(h_m)| \leq q^{-1}\zeta^{3/2}C'\delta_x k^{3/2}.$$

By Taylor formula (and the chain rule), for some $\theta_k \in \mathbb{T}^2$,

$$(42) \quad \begin{aligned} & S_{[\zeta k]}(q^{-1}f)(\Phi_{\alpha,\beta}^m(x', y')) - S_{[\zeta k]}(q^{-1}f)(h_m) \\ &= (m(x - x') + (y - y'))S_{[\zeta k]}(q^{-1}\partial f/\partial y)(\Phi_{\alpha,\beta}^m(x', y')) \\ & \quad + (m(x - x') + (y - y'))^2 S_{[\zeta k]}(q^{-1}\partial^2 f/\partial y^2)(\theta_k). \end{aligned}$$

For $m \leq T^{1+1/10}$, we have $|m(x - x') + (y - y')| \leq m\delta_x + \delta_y \leq T^{-1/3}$ and therefore, by Lemma 5.2 for $g = \partial^2 f/\partial y^2$,

$$|(m(x - x') + (y - y'))^2 S_k(\partial^2 f/\partial y^2)(\theta_k)| \leq T^{-1/10}.$$

Let now $R'' \gg R'$ (to be specified later) and let

$$m \in [R'' \max(k, T), R'' \max(k, T) + D_\eta \zeta k]$$

(see Lemma 5.6) be such that

$$S_{[\zeta k]}(q^{-1}\partial f/\partial y)(\Phi_{\alpha,\beta}^m(x', y')) \geq c'q^{-1}\zeta^{1/2}k^{1/2}.$$

Then since $R'\delta_x^{1/3} \geq \delta_y$ and we are in Case 2, for $R'' > 1$ large enough we have the inequalities $m\delta_x \geq R''T\delta_x \geq 10\delta_y$, hence by (42) we have

$$(43) \quad |S_{[\zeta k]}(q^{-1}f)(\Phi_{\alpha,\beta}^m(x', y')) - S_{[\zeta k]}(q^{-1}f)(h_m)| \geq \frac{c'}{2}q^{-1}\zeta^{1/2}k^{1/2}m\delta_x.$$

Since $R'\delta_x^{1/3} \geq \delta_y$, $m \geq R'' \max(k, T)$ and we are in Case 2, for $R'' > 1$ large enough we have the following lower bound:

$$\frac{c'}{2}q^{-1}\zeta^{1/2}k^{1/2}m\delta_x \geq 10 \max(C'p^{-1}k^{1/2}\delta_w, q^{-1}\zeta^{3/2}C'k^{3/2}\delta_x)$$

and, for any $k \geq T$, we have $(c'/2)q^{-1}\zeta^{1/2}k^{1/2}m\delta_x \geq d''' > 0$. This, together with (40), (43), (41) and (39), implies that

$$\begin{aligned} & |(S_k(p^{-1}f)(\Phi_{\alpha,\beta}^m(z, w)) - S_k(p^{-1}f)(\Phi_{\alpha,\beta}^m(z, w')) - \\ & \quad (S_{[\zeta k]}(q^{-1}f)(\Phi_{\alpha,\beta}^m(x, y)) - S_{[\zeta k]}(q^{-1}f)(\Phi_{\alpha,\beta}^m(x', y')))| \geq d'''/2. \end{aligned}$$

Thus, by cocycle identity, (38) follows for $s = k$ or $s = k + m$. This finishes the proof. \square

We can now prove Theorem 4.

Proof of Theorem 4. — We will prove that the hypotheses of the disjointness criterion for special flows given in Lemma 4.3 are satisfied. Theorem 4 will then follow.

Let $A_k : (\mathbb{T}^2, \lambda_{\mathbb{T}^2}) \rightarrow (\mathbb{T}^2, \lambda_{\mathbb{T}^2})$ be given by $A_k(x, y) = (x, y + 1/k)$. Obviously $A_k \rightarrow \text{Id}$ uniformly. Fix $\varepsilon > 0$ and $N \in \mathbb{N}$. Let us consider any $(x, y), (x', y') \in \mathbb{T}^2$ with $d((x, y), (x', y')) < \delta$ and any $(z, w), (z, w') = A_k(z, w)$ with $k < \delta^{-1}$. Let

$$T := \min(\delta_x^{-2/3}, \delta_y^{-2}, \delta_w^{-2}).$$

Let us define $a_s := a_{1,s} - a_{2,s}$ with

$$\begin{aligned} a_{1,s} &:= S_s(p^{-1}f)(z, w) - S_s(p^{-1}f)(z, w'), \\ a_{2,s} &:= S_{[\zeta s]}(q^{-1}f)(x, y) - S_{[\zeta s]}(q^{-1}f)(x', y'). \end{aligned}$$

Let $D''' > 0$ be as in Proposition 7.1. For every $s \in [0, 2D'''T]$, we have

$$(44) \quad \begin{aligned} d_2((\Phi_{\alpha,\beta}^s(z, w)), (\Phi_{\alpha,\beta}^s(z, w'))) &= \delta_w, \\ d_1((\Phi_{\alpha,\beta}^s(x, y)), (\Phi_{\alpha,\beta}^s(x', y'))) &= \delta_x \\ \text{and} \quad d_2((\Phi_{\alpha,\beta}^s(x, y)), (\Phi_{\alpha,\beta}^s(x', y'))) &\leq s\delta_x + \delta_y \leq 2D''' \delta_x^{1/3} + \delta_y. \end{aligned}$$

Hence (since f is C^1), for every $\kappa > 0$ there exists $\delta_\kappa > 0$ such that, for $\delta \in (0, \delta_\kappa)$ and for every $s \in [0, 2D'''T]$, we have

$$|a_{s+1} - a_s| < \kappa.$$

Thus by Proposition 7.1 there exists $M \in [0, D'''T]$ such that

$$\min\{|a_M + d'''\}, |a_M - d'''\} \leq \varepsilon/3.$$

This gives (12). Notice that, for any given $N \in \mathbb{N}$, there exists $\delta_N > 0$ such that, for $\delta \in (0, \delta_N)$ we have $M \geq N$, and by Lemma 5.5 (for $a_{1,M}, a_{2,M}$), by the definition of $T > 0$, there exist $V > 0$, independent of $\varepsilon > 0$, $\delta > 0$ and $N \in \mathbb{N}$, such that

$$|a_M| \leq V,$$

for some $V > 0$. This gives (11).

Let $L = \kappa M$. Notice that $[0, (1 + \zeta)(M + L)] \subset [0, 2D'''T]$ and therefore by (44)

$$d(\Phi_{\alpha,\beta}^n(x, y), \Phi_{\alpha,\beta}^n(x', y')) < \varepsilon, \quad \text{for every } n \in [0, \max(1 + \zeta)M + L]$$

and analogously for $(z, w), (z, w')$. This gives (13).

By the cocycle identity and by Lemma 5.5, there exists $\kappa_\varepsilon''' > 0$ such that for $\kappa \in (0, \kappa_\varepsilon''')$, for all $u, w \in [0, 2\zeta(M + L)]$ with $|u - w| \leq \kappa M$, we have

$$\begin{aligned} |a_{1,w} - a_{1,u}| &= |S_{w-u}(p^{-1}f)(\Phi_{\alpha,\beta}^u(z, w)) - S_{w-u}(p^{-1}f)(\Phi_{\alpha,\beta}^u(z, w'))| \\ &\leq C''|u - w|^{1/2}\delta_w \leq C''\kappa^{1/2}D^{1/2}T^{1/2}\delta_w \leq \varepsilon. \end{aligned}$$

Analogously we show that $|a_{2,u} - a_{2,w}| \leq \varepsilon/10$:

$$|a_{2,w} - a_{2,u}| = |S_{\zeta w, u}(q^{-1}f)(\Phi_{\alpha,\beta}^{[\zeta u]}(x, y)) - S_{\zeta w, u}(q^{-1}f)(\Phi_{\alpha,\beta}^{[\zeta u]}(x', y'))|,$$

for some $\zeta_{w,u} \leq \zeta C_0|w - u|$. By Lemma 5.5 (for $g = f - \int_{\mathbb{T}^2} f d\lambda_{\mathbb{T}^2}$) and (44) for $s = [\zeta M]$, there exist $C''' > 0$ and $\kappa'_\varepsilon > 0$ such that for $\kappa \in (0, \kappa'_\varepsilon)$, we have (since $|w - u| \leq L = \kappa M \leq D''' \kappa T$) (for some $C'''' > 0$)

$$\begin{aligned} |S_{\zeta w, u}(q^{-1}f)(\Phi_{\alpha,\beta}^{[\zeta u]}(x, y)) - S_{\zeta w, u}(q^{-1}f)(\Phi_{\alpha,\beta}^{[\zeta u]}(x, y'))| \\ \leq C''''(D''' \kappa T)^{3/2}\delta_x + C''''(D''' \kappa T)^{1/2}(2D''' \delta_x^{1/3} + \delta_y) \\ \leq C''''(D''' \kappa)^{3/2} + \kappa^{1/2}C''''D''''(2D''' + 1) \leq \varepsilon/10. \end{aligned}$$

This finishes the proof of (14) and hence by Lemma 4.3 completes the proof of Theorem 4. \square

REFERENCES

- [1] H. ANZAI – “Ergodic skew product transformations on the torus”, *Osaka J. Math.* **3** (1951), p. 83–99.
- [2] L. AUSLANDER, L. GREEN & F. HAHN – *Flows on homogeneous spaces*, Annals of Math. Studies, vol. 53, Princeton University Press, Princeton, NJ, 1963.
- [3] A. AVILA, G. FORNI & C. ULCIGRAI – “Mixing for time-changes of Heisenberg nilflows”, *J. Differential Geom.* **89** (2011), no. 3, p. 369–410.
- [4] A. AVILA, D. RAVOTTI, G. FORNI & C. ULCIGRAI – “Mixing for smooth time-changes of general nilflows”, 2019, [arXiv:1905.11628](https://arxiv.org/abs/1905.11628).
- [5] J. BOURGAIN, P. SARNAK & T. ZIEGLER – “Disjointness of Moebius from horocycle flows”, in *From Fourier analysis and number theory to Radon transforms and geometry*, Dev. Math., vol. 28, Springer, New York, 2013, p. 67–83.
- [6] B. FAYAD, G. FORNI & A. KANIGOWSKI – “Lebesgue spectrum of countable multiplicity for conservative flows on the torus”, 2016, [arXiv:1609.03757v2](https://arxiv.org/abs/1609.03757v2).
- [7] B. FAYAD & A. KANIGOWSKI – “Multiple mixing for a class of conservative surface flows”, *Invent. Math.* **203** (2016), no. 2, p. 555–614.
- [8] S. FERENCZI, J. KULAGA-PRZYMUS & M. LEMAŃCZYK – “Sarnak’s conjecture: what’s new”, in *Ergodic theory and dynamical systems in their interactions with arithmetics and combinatorics (CIRM Jean-Morlet Chair, Fall 2016)*, Lect. Notes in Math., vol. 2213, Springer, Cham, 2018, p. 163–235.
- [9] L. FLAMINIO & G. FORNI – “Equidistribution of nilflows and applications to theta sums”, *Ergodic Theory Dynam. Systems* **26** (2006), no. 2, p. 409–433.
- [10] ———, “Orthogonal powers and Möbius conjecture for smooth time changes of horocycle flows”, *Electron. Res. Announc. Math. Sci.* **26** (2019), p. 16–23.
- [11] L. FLAMINIO, K. FRĄCZEK, J. KULAGA-PRZYMUS & M. LEMAŃCZYK – “Approximate orthogonality of powers for ergodic affine unipotent diffeomorphisms on nilmanifolds”, *Studia Math.* **244** (2019), no. 1, p. 43–97.
- [12] G. FORNI – “Effective equidistribution of nilflows and bounds on Weyl sums”, in *Dynamics and analytic number theory*, London Math. Soc. Lecture Note Ser., vol. 437, Cambridge Univ. Press, Cambridge, 2016, p. 136–188.
- [13] G. FORNI & A. KANIGOWSKI – “Time-changes of Heisenberg nilflows”, 2017, [arXiv:1711.05543v1](https://arxiv.org/abs/1711.05543v1).
- [14] G. FORNI & C. ULCIGRAI – “Time-changes of horocycle flows”, *J. Modern Dyn.* **6** (2012), no. 2, p. 251–273.
- [15] P. GABRIEL, M. LEMAŃCZYK & K. SCHMIDT – “Extensions of cocycles for hyperfinite actions and applications”, *Monatsh. Math.* **123** (1997), no. 3, p. 209–228.
- [16] E. GLASNER – *Ergodic theory via joinings*, Math. Surveys and Monographs, vol. 101, American Mathematical Society, Providence, RI, 2003.
- [17] B. GREEN & T. TAO – “The Möbius function is strongly orthogonal to nilsequences”, *Ann. of Math. (2)* **175** (2012), no. 2, p. 541–566.
- [18] B. HASSELBLATT & A. KATOK – “Principal structures”, in *Handbook of dynamical systems, Vol. 1A*, North-Holland, Amsterdam, 2002, p. 1–203.
- [19] A. KANIGOWSKI & J. KULAGA-PRZYMUS – “Ratner’s property and mild mixing for smooth flows on surfaces”, *Ergodic Theory Dynam. Systems* **36** (2016), no. 8, p. 2512–2537.
- [20] A. KANIGOWSKI, J. KULAGA-PRZYMUS & C. ULCIGRAI – “Multiple mixing and parabolic divergence in smooth area-preserving flows on higher genus surfaces”, *J. Eur. Math. Soc. (JEMS)* **21** (2019), no. 12, p. 3797–3855.
- [21] A. KANIGOWSKI & M. LEMAŃCZYK – “Flows with Ratner’s property have discrete essential centralizer”, *Studia Math.* **237** (2017), no. 2, p. 185–194.
- [22] A. KANIGOWSKI, M. LEMAŃCZYK & C. ULCIGRAI – “On disjointness properties of some parabolic flows”, 2018, [arXiv:1810.11576](https://arxiv.org/abs/1810.11576).
- [23] I. KÁTAI – “A remark on a theorem of H. Daboussi”, *Acta Math. Hungar.* **47** (1986), no. 1–2, p. 223–225.
- [24] A. KATOK – *Combinatorial constructions in ergodic theory and dynamics*, University Lect. Series, vol. 30, American Mathematical Society, Providence, RI, 2003.

- [25] M. RATNER – “Horocycle flows, joinings and rigidity of products”, *Ann. of Math. (2)* **118** (1983), no. 2, p. 277–313.
- [26] ———, “Rigid reparametrizations and cohomology for horocycle flows”, *Invent. Math.* **88** (1987), no. 2, p. 341–374.
- [27] D. RAVOTTI – “Mixing for suspension flows over skew-translations and time-changes of quasi-abelian filiform nilflows”, *Ergodic Theory Dynam. Systems* **39** (2018), no. 12, p. 3407–3436.
- [28] P. SARNAK – “Three lectures on the Möbius function randomness and dynamics”, 2011, <https://publications.ias.edu/sarnak/paper/512>.
- [29] R. TIEDRA DE ALDECOA – “Spectral analysis of time changes of horocycle flows”, *J. Modern Dyn.* **6** (2012), no. 2, p. 275–285.
- [30] ———, “Commutator methods for the spectral analysis of uniquely ergodic dynamical systems”, *Ergodic Theory Dynam. Systems* **35** (2015), no. 3, p. 944–967.

Manuscript received 17th February 2019
accepted 9th October 2019

GIOVANNI FORNI, Department of Mathematics, University of Maryland
4176 Campus Drive – William E. Kirwan Hall, College Park, MD 20742-4015, USA
E-mail : gforni@math.umd.edu

ADAM KANIGOWSKI, Department of Mathematics, University of Maryland
4176 Campus Drive – William E. Kirwan Hall, College Park, MD 20742-4015, USA
E-mail : adkanigowski@gmail.com