



Journal de l'École polytechnique

Mathématiques

Louis-Hadrien ROBERT & Emmanuel WAGNER

Symmetric Khovanov-Rozansky link homologies

Tome 7 (2020), p. 573-651.

http://jep.centre-mersenne.org/item/JEP_2020__7__573_0

© Les auteurs, 2020.

Certains droits réservés.



Cet article est mis à disposition selon les termes de la licence
LICENCE INTERNATIONALE D'ATTRIBUTION CREATIVE COMMONS BY 4.0.
<https://creativecommons.org/licenses/by/4.0/>

L'accès aux articles de la revue « Journal de l'École polytechnique — Mathématiques » (<http://jep.centre-mersenne.org/>), implique l'accord avec les conditions générales d'utilisation (<http://jep.centre-mersenne.org/legal/>).

Publié avec le soutien
du Centre National de la Recherche Scientifique



Publication membre du
Centre Mersenne pour l'édition scientifique ouverte
www.centre-mersenne.org

SYMMETRIC KHOVANOV-ROZANSKY LINK HOMOLOGIES

BY LOUIS-HADRIEN ROBERT & EMMANUEL WAGNER

ABSTRACT. — We provide a finite-dimensional categorification of the symmetric evaluation of sl_N -webs using foam technology. As an output we obtain a symmetric link homology theory categorifying the link invariant associated to symmetric powers of the standard representation of sl_N . The construction is made in an equivariant setting. We prove also that there is a spectral sequence from the Khovanov-Rozansky triply graded link homology to the symmetric one and provide along the way a foam interpretation of Soergel bimodules.

RÉSUMÉ (Homologies d'entrelacs de Khovanov–Rozansky symétriques). — On donne une catégorification de l'évaluation symétrique des toiles sl_N en utilisant les mousses. On en déduit des théories homologiques d'entrelacs qui catégorifient les invariants quantiques d'entrelacs associés aux puissances symétriques de la représentation standard de sl_N . Ces théories sont obtenues dans un cadre équivariant. On montre qu'il existe des suites spectrales de l'homologie triplement graduée de Khovanov-Rozansky vers ces homologies symétriques. On donne aussi une interprétation des bimodules de Soergel en terme de mousses.

CONTENTS

1. Introduction.....	574
2. MOY graphs.....	577
3. Foams.....	587
4. Soergel bimodules.....	603
5. One quotient and two approaches.....	610
6. Link homologies.....	625
Appendix A. Quantum link invariants and representations of $U_q(\mathfrak{gl}_N)$	639
Appendix B. Koszul resolutions of polynomial algebras.....	642
Appendix C. A pinch of algebraic geometry.....	648
References.....	649

2010 MATHEMATICS SUBJECT CLASSIFICATION. — 57R56, 57M27, 17B10, 17B37.

KEYWORDS. — Link homology, quantum invariant, foams, Soergel bimodules.

The authors thank the Newton Institute for Mathematical Sciences, the University of Burgundy and Marie Fillastre for their logistical support. L.-H. R. was supported by SwissMAP.

1. INTRODUCTION

In [RW17], we provided a combinatorial evaluation of the foams underlying the (exterior) colored Khovanov-Rozansky link homologies [CK08a, CK08b, MS09, Sus07, MSV09, Wu14, Yon11]. See [MW18] for an overview. This formula was the keystone to provide a down-to-earth treatment of these homologies, completely similar to the one in Khovanov's original paper [Kho05] or in his sl_3 paper [Kho04]. Immediate consequences of this formula were used by Ehrig-Tubbenhauer-Wedrich [ETW18] to prove functoriality of these homologies.

The present paper grew up as an attempt to provide a similar formula for foams underlying link homologies categorifying the Reshetikhin-Turaev invariants of links corresponding to symmetric powers of the standard representation of quantum sl_N . Providing manageable definitions of these link homologies is one of the keys of the program aiming at categorifying quantum invariants of 3-manifolds.

The first such link homologies were provided by Khovanov for the colored Jones polynomial [Kho05] (see as well [BW08]). There are nowadays many definitions of link homologies, e.g. using categorified projectors [SS14, FSS12, CK12, Roz14, Cau15, CH15] or using spectral sequences [Cau17] (see below for more details concerning this last one). They also fit in the higher representations techniques developed by Webster [Web17].

In addition, it has been conjectured in [GG18] that there exist symmetries between the categorification of Reshetikhin-Turaev invariants arising from exterior powers and symmetric powers of the standard representation of quantum sl_N . It has been proved by Tubbenhauer-Vaz-Wedrich at a decategorified level [TVW17]. Moreover the work of Rose-Tubbenhauer [RT16], Que elec-Rose [QR18] and Que elec-Rose-Sartori [QRS18] made clear that the planar graphical calculus underlying the description of the symmetric powers of the standard representation of quantum sl_N is for a large part similar to the one for the exterior powers. In the exterior case it was developed by Murakami-Ohtsuki-Yamada [MOY98] (called in this paper the *exterior* MOY calculus). Que elec, Rose and Sartori proved that the invariants differ by the initializations on colored circles [QR18, QRS18]. They work in an annular setting. We call it the *symmetric* MOY calculus.

An attentive reader may have noticed that we spoke about an attempt. Let us explain why one cannot provide such a formula in the symmetric case at the level of generality we had in [RW17] and therefore need to restrict the setup.

Foams are 2-dimensional CW-complexes which are naturally cobordisms between trivalent graphs (see below for an example). The closed formula of [RW17] provided a singular TQFT using the universal construction [BHMV95]. In this construction a facet of the foam is decorated with elements of a Frobenius algebra which is attached to the circle colored with the same color. Since the work of Bar-Natan and Khovanov the existence of the non-degenerate pairing on the algebra is rephrased in a topological type relation, known as the neck-cutting relation. Moreover, the TQFT feature also forces the evaluation of a planar graph times a circle to be the dimension of the vector

space, the universal construction associates to the planar graph. We emphasized in [RW17] that if such a construction works, not only circles are associated Frobenius algebras but all planar graphs which have a symmetry axis. In addition, the co-unit, for degree reasons, should be non-zero only on the maximal degree elements. All the previous properties would be forced if one could obtain a closed formula providing a categorification of the symmetric MOY calculus. Elementary computations show that a functor categorifying the symmetric MOY calculus cannot satisfy such properties essentially for degree reasons.

This is why we work in an annular setting. The drawback is that we cannot deal with general link diagrams. The benefit is that we can use part of the technology developed by Quelelec-Rose [QR18]. We obtain an evaluation in this restricted case and apply a restricted universal construction to obtain the following result:

THEOREM. — *There exists a finite-dimensional categorification of the symmetric MOY calculus yielding a categorification of the Reshetikhin-Turaev invariants of links corresponding to symmetric powers of the standard representation of quantum sl_N .*

We call these link homologies *symmetric* (colored) Khovanov-Rozansky link homologies. The construction applies in particular to the case where the representation is the standard representation of quantum sl_N . As shown on an example (Section 6.3), it provides in this case a different categorification of the (uncolored)- sl_N -link invariants than those of Khovanov and Khovanov-Rozansky. This observation is due to Quelelec-Rose-Sartori; actually they discuss in [QRS18] how the annular link homologies constructed by Quelelec and Rose in [QR18] can be specialized to be invariant under Reidemeister I and give link invariants in the 3-sphere.

Whereas the definition of the link homologies can be made only using the language of foams and symmetric polynomial, the proof of invariance at the moment requires a more algebraic treatment. This algebraic treatment uses Soergel bimodules and makes explicit the comparison with the work of Cautis [Cau17].⁽¹⁾ Cautis constructs a differential d_N on the Hochschild homology of Soergel bimodules compatible with the differential of the Rickard complex such that the total homology provides a categorification of Reshetikhin-Turaev invariants of links corresponding to symmetric powers of the standard representation of quantum sl_N . We provide here an explicit version of the additional differential in an equivariant setting. The non-equivariant case is studied by Cautis [Cau17] and investigated by Quelelec-Rose-Sartori [QRS18]. Hence, one consequence of our proof of invariance is the following.

THEOREM. — *There exists a spectral sequence whose first page is isomorphic to the (unreduced) colored triply graded link homology converging to the symmetric Khovanov-Rozansky link homologies.*

⁽¹⁾This strategy of proof was discussed with H. Quelelec and D. Rose.

We would like to stress that one can also see on the same picture the spectral sequences converging to the (exterior) colored Khovanov-Rozansky link homology [Ras15, Wed19]. Hence in some sense the only differential missing from the perspective of the work of Dunfield-Gukov-Rasmussen is d_0 which seems to be tackled by Dowlin [Dow17].

The definition of the link homologies in this paper, starts with the links presented as closures of braids, hence regarding functoriality questions it only makes sense to consider braid-like movie moves. It is an immediate consequence of our definitions and the work of Ehrig-Tubbenhauer-Wedrich concerning functoriality of the (exterior) colored Khovanov-Rozansky link homologies that the following holds.

THEOREM. — *The symmetric Khovanov-Rozansky link homologies are functorial with respect to braid-like movie moves.*

One very important feature is that our construction works in an equivariant setting and will allow with a little more work to define Rasmussen type invariants for braids. In another direction, the fact that we are restricted to braid closures naturally suggest that one will obtain Morton-Franks-Williams type inequalities in this case (see Wu [Wu13]). The interactions between the two previous directions seem to us worth pursuing.

To conclude, the construction is done over rationals and we think the following question deserves attention.

QUESTION. — *Can one make the content of this paper work over integers?*

The main obstruction so far is that in our proof of invariance we need to invert 2, just like for the stabilization in the triply graded homology [Rou17, WW17]. The strategy adopted by Krasner [Kra10b] might be a starting point.

Outline of the paper. — The paper is divided as follows. In the Section 2 we develop the symmetric MOY calculus. In the Section 3 we provide the needed definitions of the restricted class of foams we will be working with: disk-like and vinyl foams. We also give an overview on foams and the closed formula of [RW17]. In Section 4 we explain how to think of Soergel bimodules as spaces of disk-like foams and part of their homologies as vinyl foams. In Section 5 we define an evaluation of vinyl foams, providing a categorification of the symmetric MOY calculus. The rest of the section is devoted to rephrasing it in terms of an additional differential on the Hochschild homology of Soergel bimodules. The algebraic description is used in Section 6 to prove the invariance of the link homologies. The link homologies are constructed using the well-known Rickard complexes.

There are as well three appendices. The first one deals with the representation theory of quantum \mathfrak{gl}_N . The second is a reminder on Koszul resolutions and contains some technical homological lemmas. The third one present some inspiring algebraic geometry.

Acknowledgements. — The authors thank Christian Blanchet, François Costantino, Hoel Queelec, David Rose and Paul Wedrich for interesting discussions and Mikhail Khovanov and Daniel Tubbenhauer for their comments on a previous version of this work. We thank the referees for their numerous remarks which helped us to improve the paper exposition.

2. MOY GRAPHS

DEFINITION 2.1 ([MOY98]). — An *abstract MOY graph* is a finite oriented graph $\Gamma = (V; E)$ with a labeling of its edges $\ell : E \rightarrow \mathbb{N}_{>0}$ such that:

- the vertices are either univalent (we call this subset of vertices the *boundary* and denote it by $@$) or trivalent (these are the *internal vertices*),
- the flow given by labels and orientations is preserved along the trivalent vertices, meaning that every trivalent vertex follows one of the two models (*merge* and *split* vertices) drawn here.



The univalent vertices are either sinks or sources. We call the first *positive boundary points* and the later *negative boundary points*.


REMARK 2.2

(1) Sometimes it will be convenient to allow edges labeled by 0. However, these edges should be thought of as “irrelevant”. We simply delete them to recover the original definition.

(2) Each internal vertex has three adjacent edges. The label of one of these edges is strictly greater than the other two. This edge is called the *big* edge relative to this vertex. The two other edges are called the *small* edges relative to this vertex.

DEFINITION 2.3. — A *MOY graph* is the image of an abstract MOY graph Γ by a smooth embedding in the $[0; 1] \times [0; 1]$ such that:

- All the oriented tangent lines at vertices are equal.⁽²⁾



- The boundary of Γ is contained in $[0; 1] \times \{0; 1\}$.
- $[0; 1] \times \{0; 1\} \cap \Gamma = @$.
- the tangent lines at the boundary points of Γ are vertical, (that is, collinear with $\begin{pmatrix} 0 \\ 1 \end{pmatrix}$).

⁽²⁾In pictures which follows we may forget about this technical condition, since it is clear that we can always deform the embedding locally so that this condition is fulfilled.

In what follows it will be convenient to speak about *the tangent vector* at a point p of the graph Γ (or more precisely of its image in $[0;1] \times [0;1]$). By this we mean the only vector which is tangent to Γ , has norm 1 and whose orientation agrees with the one of Γ . Note that the condition on the embedding on vertices ensure that it is well-defined everywhere.

NOTATION 2.4. — In what follows, \underline{k} always denotes a finite sequence of integers (the empty sequence is allowed). If $\underline{k} = (k_1; \dots; k_r)$, r is the *length* of \underline{k} and $\sum_{i=1}^r k_i$ is the *level* of \underline{k} . If \underline{k} is a sequence of length 1 and level k , we abuse notation and write k instead of \underline{k} .

If we want to specify the boundary of a MOY graph Γ , we will speak about \underline{k}_1 -MOY graph- \underline{k}_0 (see the example in Figure 1 to understand the notations). If a MOY graph has an empty boundary we say that it is *closed*.

If Γ is an \underline{k}_1 -MOY graph- \underline{k}_0 , we denote by Γ^* the (\underline{k}_1) -MOY graph- (\underline{k}_0) , which is obtained from Γ by reversing all orientations.

REMARK 2.5. — MOY graphs are regarded up to ambient isotopy fixing the boundary. This fits into a category where objects are finite sequences of signed and labeled points in $]0;1[$, and morphisms are MOY graphs. The composition is then given by concatenation and rescaling (see Figure 1).




FIGURE 1. Examples of a MOY graphs: a $(4;3;1)$ -MOY graph- $(2;3;1;2)$, a $(2;2;5;3)$ -MOY graph- $(4;3;1)$ and their concatenation.

DEFINITION 2.6. — Let Γ be a closed MOY graph. The *rotational* of Γ is the sum of the rotational of the oriented circles appearing in the cabling of Γ . The *rotational* of a circle is $+1$ if it winds counterclockwisely and -1 if it winds clockwise. It is denoted by $\text{rot}(\Gamma)$. This definition is illustrated in Figure 2.

2.1. MOY CALCULI. — In their seminal paper, Murakami, Ohtsuki and Yamada [MOY98] gave a combinatorial definition of the colored $U_q(\mathfrak{sl}_N)$ framed link invariant. For clarity we refer to this construction as *exterior* MOY calculus since it calculates the Reshetikhin-Turaev invariant of a framed link labeled with exterior powers of V the standard representation of $U_q(\mathfrak{sl}_N)$. We denote by $h i_N$ this invariant (or simply by $h i$ when there is no ambiguity about N). This goes in two steps. We consider a framed link diagram labeled with integers between 0 and N (one



FIGURE 2. The rotational of the closed MOY graph depicted on the left is equal to $2 - 2 = 0$.

should think about an integer a as representing $\langle a \rangle_q V$ and we replace every crossing by a formal $\mathbb{Z}[q; q^{-1}]$ -linear combination of planar graphs following the formulas:

$$(2.1) \quad \left\langle \begin{array}{c} m \quad n \\ \diagdown \quad \diagup \\ \diagup \quad \diagdown \end{array} \right\rangle = \sum_{k=\max(0; m-n)}^m (-1)^m q^k \left\langle \begin{array}{c} m \quad n \\ \diagdown \quad \diagup \\ \diagup \quad \diagdown \\ \quad \quad k \end{array} \right\rangle$$

$$(2.2) \quad \left\langle \begin{array}{c} m \quad n \\ \diagup \quad \diagdown \\ \diagdown \quad \diagup \end{array} \right\rangle = \sum_{k=\max(0; m-n)}^m (-1)^m q^k \left\langle \begin{array}{c} m \quad n \\ \diagup \quad \diagdown \\ \diagdown \quad \diagup \\ \quad \quad k \end{array} \right\rangle$$

The first crossing in the formula is said to have *type* $(m; n; +)$, the second to have *type* $(m; n; -)$.

Finally we can evaluate these planar graphs (which are closed MOY graphs) using the following identities and their mirror images:

$$(2.3) \quad \left\langle \begin{array}{c} \bigcirc \\ \quad \quad k \end{array} \right\rangle = \begin{bmatrix} N \\ k \end{bmatrix}$$

$$(2.4) \quad \left\langle \begin{array}{c} i \quad j \quad k \\ \diagdown \quad \diagup \\ \diagup \quad \diagdown \\ \quad \quad j+k \\ i+j+k \end{array} \right\rangle = \left\langle \begin{array}{c} i \quad j \quad k \\ \diagup \quad \diagdown \\ \diagdown \quad \diagup \\ \quad \quad j+k \\ i+j+k \end{array} \right\rangle$$

$$(2.5) \quad \left\langle \begin{array}{c} m+n \\ \quad \quad \uparrow \\ m \quad \quad n \\ \quad \quad \downarrow \\ m+n \end{array} \right\rangle = \begin{bmatrix} m+n \\ m \end{bmatrix} \left\langle \begin{array}{c} \uparrow \\ m+n \end{array} \right\rangle$$

$$(2.6) \quad \left\langle \begin{array}{c} m+n \\ \quad \quad \downarrow \\ m+n \quad \quad n \\ \quad \quad \uparrow \\ m+n \end{array} \right\rangle = \begin{bmatrix} N & m \\ n & \end{bmatrix} \left\langle \begin{array}{c} \downarrow \\ m \end{array} \right\rangle$$

$$(2.7) \quad \left\langle \begin{array}{c} 1 \quad m \\ \uparrow \quad \downarrow \\ m \quad \quad 1 \\ \downarrow \quad \uparrow \\ m \quad \quad 1 \\ 1 \quad \quad m \end{array} \right\rangle = \left\langle \begin{array}{c} \uparrow \\ 1 \end{array} \right\rangle \left\langle \begin{array}{c} \downarrow \\ m \end{array} \right\rangle + [N \quad m \quad 1] \left\langle \begin{array}{c} 1 \quad m \\ \diagdown \quad \diagup \\ m \quad \quad 1 \\ \diagup \quad \diagdown \\ 1 \quad \quad m \end{array} \right\rangle$$

$$(2.8) \quad \left\langle \begin{array}{c} \begin{array}{ccc} \uparrow & & \uparrow \\ \leftarrow + n & \xrightarrow{n} & \leftarrow \\ \downarrow & & \downarrow \\ 1 & m+1 & 1 \end{array} \\ \end{array} \right\rangle = \begin{bmatrix} m & 1 \\ n & 1 \end{bmatrix} \left\langle \begin{array}{c} \begin{array}{ccc} \uparrow & & \uparrow \\ \leftarrow 1 & \xrightarrow{1} & \leftarrow \\ \downarrow & & \downarrow \\ 1 & m+1 & 1 \end{array} \\ \end{array} \right\rangle + \begin{bmatrix} m & 1 \\ n & 1 \end{bmatrix} \left\langle \begin{array}{c} \begin{array}{ccc} \uparrow & & \uparrow \\ \leftarrow & \xrightarrow{m} & \leftarrow \\ \downarrow & & \downarrow \\ 1 & m+1 & 1 \end{array} \\ \end{array} \right\rangle$$

$$(2.9) \quad \left\langle \begin{array}{c} \begin{array}{ccc} \uparrow & & \uparrow \\ \leftarrow n+k & \xrightarrow{n+k} & \leftarrow \\ \downarrow & & \downarrow \\ n & m+1 & n \end{array} \\ \end{array} \right\rangle = \sum_{j=\max(0, m-n)}^m \begin{bmatrix} \cdot & \cdot \\ k & j \end{bmatrix} \left\langle \begin{array}{c} \begin{array}{ccc} \uparrow & & \uparrow \\ \leftarrow j & \xrightarrow{j} & \leftarrow \\ \downarrow & & \downarrow \\ n & m+1 & n+j \end{array} \\ \end{array} \right\rangle$$

In the previous formulas, we used quantum integers and quantum binomials. These are symmetric Laurent polynomials in q defined by $[k] := \frac{q^k - q^{-k}}{q - q^{-1}}$ and

$$\begin{bmatrix} \cdot & \cdot \\ k & j \end{bmatrix} = \frac{\prod_{i=0}^{k-1} [\cdot - i]}{[k]!} \quad \text{where } [i]! = \prod_{j=1}^i [i]:$$

The first formal proof that these relations are enough to compute has been written by Wu [Wu14] and is based on a result of Kau man and Vogel [Kau13, App. 4]. In particular, this shows that there is a unique evaluation of MOY graphs which satisfies these relations. The coherence of these relations follows from the representation theoretic point of view. For more details we refer to [MOY98, MS09] and to Appendix A.

As pointed out by [TVW17], a similar story applies when one think about the integers labeling the strands of a link (which are now only required to be positive) as representing q -symmetric powers of V (i.e., $\text{Sym}_q V$). We denote by $\langle\langle \cdot \rangle\rangle_N$ this invariant (or simply by $\langle\langle \cdot \rangle\rangle_N$ when there is no ambiguity about N). This yields what we call the *symmetric* MOY calculus.

$$(2.10) \quad \left\langle\left\langle \begin{array}{c} \begin{array}{ccc} \uparrow & & \uparrow \\ \leftarrow m & \xrightarrow{n} & \leftarrow \\ \downarrow & & \downarrow \\ m & n & m \end{array} \\ \end{array} \right\rangle\right\rangle_N = \sum_{k=\max(0, m-n)}^m (-1)^m q^{k^2} \left\langle\left\langle \begin{array}{c} \begin{array}{ccc} \uparrow & & \uparrow \\ \leftarrow n+k & \xrightarrow{m} & \leftarrow \\ \downarrow & & \downarrow \\ n & m & k \end{array} \\ \end{array} \right\rangle\right\rangle_N$$

$$(2.11) \quad \left\langle\left\langle \begin{array}{c} \begin{array}{ccc} \uparrow & & \uparrow \\ \leftarrow m & \xrightarrow{n} & \leftarrow \\ \downarrow & & \downarrow \\ m & n & m \end{array} \\ \end{array} \right\rangle\right\rangle_N = \sum_{k=\max(0, m-n)}^m (-1)^m q^{m^2} \left\langle\left\langle \begin{array}{c} \begin{array}{ccc} \uparrow & & \uparrow \\ \leftarrow n+k & \xrightarrow{m} & \leftarrow \\ \downarrow & & \downarrow \\ n & m & k \end{array} \\ \end{array} \right\rangle\right\rangle_N$$

The formulas for evaluating MOY graphs become:

$$(2.12) \quad \left\langle\left\langle \begin{array}{c} \begin{array}{c} \circlearrowleft \\ \uparrow k \end{array} \\ \end{array} \right\rangle\right\rangle_N = \begin{bmatrix} N+k & 1 \\ k & 1 \end{bmatrix}$$

$$(2.13) \quad \left\langle\left\langle \begin{array}{c} i \quad j \quad k \\ \swarrow \quad \searrow \quad \nearrow \\ \downarrow \\ i+j+k \end{array} \right\rangle\right\rangle_N = \left\langle\left\langle \begin{array}{c} i \quad j \quad k \\ \swarrow \quad \searrow \quad \nearrow \\ \downarrow \\ i+j+k \end{array} \right\rangle\right\rangle_N$$

$$(2.14) \quad \left\langle\left\langle \begin{array}{c} m+n \\ \uparrow \quad \downarrow \\ m \quad n \\ \downarrow \quad \uparrow \\ m+n \end{array} \right\rangle\right\rangle_N = [m+n \atop m] \left\langle\left\langle \begin{array}{c} m+n \\ \uparrow \\ m+n \end{array} \right\rangle\right\rangle_N$$

$$(2.15) \quad \left\langle\left\langle \begin{array}{c} m+n \\ \uparrow \quad \downarrow \\ m \quad n \\ \downarrow \quad \uparrow \\ m+n \end{array} \right\rangle\right\rangle_N = [N+m+n \atop n \quad 1] \left\langle\left\langle \begin{array}{c} m \\ \uparrow \\ m \end{array} \right\rangle\right\rangle_N$$

$$(2.16) \quad \left\langle\left\langle \begin{array}{c} 1 \quad m \\ \uparrow \quad \downarrow \\ m+1 \quad 1 \\ \downarrow \quad \uparrow \\ 1 \quad m \end{array} \right\rangle\right\rangle_N = \left\langle\left\langle \begin{array}{c} 1 \\ \uparrow \\ 1 \\ \downarrow \\ m \end{array} \right\rangle\right\rangle_N + [N+m+1] \left\langle\left\langle \begin{array}{c} 1 \quad m \\ \swarrow \quad \searrow \\ \downarrow \\ 1 \quad m \end{array} \right\rangle\right\rangle_N$$

$$(2.17) \quad \left\langle\left\langle \begin{array}{c} n \\ \uparrow \quad \downarrow \\ m \quad n \\ \downarrow \quad \uparrow \\ 1 \quad m+1 \end{array} \right\rangle\right\rangle_N = [m \atop n \quad 1] \left\langle\left\langle \begin{array}{c} n \\ \uparrow \quad \downarrow \\ 1 \\ \downarrow \\ m+1 \end{array} \right\rangle\right\rangle_N + [m \atop n \quad 1] \left\langle\left\langle \begin{array}{c} n \\ \swarrow \quad \searrow \\ \downarrow \\ 1 \quad m+1 \end{array} \right\rangle\right\rangle_N$$

$$(2.18) \quad \left\langle\left\langle \begin{array}{c} m \quad n+k \\ \uparrow \quad \downarrow \\ n+k \quad m \\ \downarrow \quad \uparrow \\ n \quad m+k \end{array} \right\rangle\right\rangle_N = \sum_{j=\max(0, m-n)}^m [k \atop j] \left\langle\left\langle \begin{array}{c} m \quad n+k \\ \uparrow \quad \downarrow \\ n+k \quad m \\ \downarrow \quad \uparrow \\ n \quad m+k \end{array} \right\rangle\right\rangle_N$$

REMARK 2.7

(1) The proof of computability and uniqueness of Wu [Wu14] still works in the symmetric case. As before consistency follows from the representation theoretic point of view. We describe explicitly in Appendix A the $U_q(\mathfrak{gl}_N)$ -intertwiners between products of symmetric powers of the standard $U_q(\mathfrak{gl}_N)$ -module. One can check by brute force

computation that these morphisms satisfy the identities defining the symmetric MOY calculus.

(2) We choose a normalization making the polynomial associated by the symmetric MOY calculus with any planar graph a Laurent polynomial with positive coefficients. The skein formula we use for the crossings is not compatible with the choices made for the exterior MOY calculus. The braiding in the exterior MOY calculus is given by an R matrix of $U_q(\mathfrak{sl}_N)$, while for the symmetric MOY calculus it is given by its inverse. Hence the two calculi we present here cannot be merge into one bicolor calculus as it is done in [TVW17].

(3) In [TVW17], there is an overall sign which we choose to remove here. For recovering this sign one should multiply our symmetric evaluation of a MOY graph by $(-1)^{\text{rot}(\cdot)}$ (see Definition 2.6).

(4) Up to this sign the formulas are actually the same as the ones of the exterior MOY calculus applied to N . Since they do not involve N , the identities (2.4), (2.5), (2.8) and (2.9) of the exterior MOY calculus are the same as the identities (2.13), (2.14), (2.17) and (2.18) of the symmetric MOY calculus.

(5) In order to turn the framed invariants $h i_N$ and $\langle\langle \cdot \rangle\rangle_N$ into invariants of unframed links, one needs to renormalize them. For any link diagram D , we define:

$$RT_N(D) = (-1)^{e(D)} q^{k(D)} hDi_N$$

and

$$RT_N^S(D) = q^{k^S(D)} \langle\langle D \rangle\rangle_N;$$

where $e(D)$ (resp. $k(D)$, resp. $k^S(D)$) is the sum over all crossing x of D of e_x (resp. k_x , resp. k_x^S) defined by:

$$(k_x; k_x^S; e_x) = \begin{cases} (m(N+1-m); m(m+N-1); m) & \text{if } x \text{ is of type } (m; m; +), \\ (m(N+1-m); m(m+N-1); +m) & \text{if } x \text{ is of type } (m; m; -), \\ (0; 0; 0) & \text{else.} \end{cases}$$

While the case $N = 1$ is trivial in the exterior MOY calculus, it is not in the symmetric MOY calculus. However, the symmetric evaluation of a MOY graph for $N = 1$ is especially simple.

LEMMA 2.8. — Let Γ be a MOY graph. For every vertex v of Γ , let us denote by $W(v)$ the element of $\mathbb{N}[q; q^{-1}]$ given by the formulas:

$$W \left(\begin{array}{c} a+b \\ \uparrow \\ a \quad b \end{array} \right) := \begin{bmatrix} a+b \\ a \end{bmatrix} =: W \left(\begin{array}{c} a+b \\ \downarrow \\ a \quad b \end{array} \right):$$

The following identities hold in $\mathbb{Z}[q; q^{-1}]$:

$$\langle\langle \cdot \rangle\rangle_{N=1} = \prod_{v \text{ split}} W(v) = \prod_{v \text{ merge}} W(v) = \left(\prod_{v \text{ vertex of } \Gamma} W(v) \right)^{1=2}:$$

Sketch of the proof. — We only treat the “merge” part of the statement. Since there is a unique polynomial satisfying the symmetric MOY calculus for $N = 1$. It is enough to check that $W(\cdot) := \prod_{v \text{ merge}} W(v)$ satisfies the symmetric MOY calculus of $N = 1$. For instance, to prove that W satisfies identity (2.18), one shows the following q -binomial identity:

$$\begin{bmatrix} n+k \\ m \end{bmatrix} \begin{bmatrix} m+r \\ k \end{bmatrix} = \sum_{j=\max(0, m-n)}^n \begin{bmatrix} r \\ k-j \end{bmatrix} \begin{bmatrix} n \\ m-j \end{bmatrix} \begin{bmatrix} n+r+j \\ j \end{bmatrix}$$

which can be done by induction on $n + m$.

Another identity holds both in the exterior and the symmetric MOY calculi.

LEMMA 2.9. — *The following local identities and their mirror images hold:*

(2.19)

(2.20)

Sketch of proof. — For the exterior calculus: it is a consequence of identity (2.4), its mirror image and identity (2.5). For the symmetric calculus: it is a consequence of identity (2.13), its mirror image and identity (2.14).

2.2. BRAID-LIKE MOY GRAPHS. — In this section we introduce a special class of MOY graphs which contains in particular the ones appearing in the expansion of braids when using identities (2.1) and (2.2) to get rid of crossings. We call these graphs braid-like.


DEFINITION 2.10. — A MOY graph is *braid-like* if the scalar products of all its tangent vectors with (q) are strictly positive. In Figure 1, the leftmost MOY graph is braid-like, while the two others are not.

REMARK 2.11

(1) Braid-like MOY graphs are regarded up to ambient isotopy fixing the boundary and preserving the braid-like property. They fit into a category which is a non-full subcategory of the one described in Remark 2.5.

(2) The braid-likeness of a MOY graph implies that boundary points of $\text{on }]0;1[\text{ } f0g$ are negative, while the one on $]0;1[\text{ } f1g$ are positive.

(3) Every braid-like MOY graph can be obtained as vertical concatenation of MOY graphs of type:



(4) If Γ is a \underline{k}_1 -MOY graph- \underline{k}_0 , then the sum of the element of \underline{k}_1 is equal to the sum of the element of \underline{k}_0 , we call the number the *level* of Γ .

The following lemma, although quite elementary and completely combinatorial, is one of the keystones of this paper.

LEMMA 2.12. — *Let k be a positive integer and \underline{k} be a finite collection of positive integers of level k . We consider M the $\mathbb{Z}[q; q^{-1}]$ module generated by braid-like \underline{k} -MOY graphs- (k) and modded out by ambient isotopy and relations (2.4) and (2.5) (or (2.13) and (2.14)). The module M is generated by a braid-like tree. Moreover all braid-like trees are equal in M .*

Proof. — Let T be a braid-like \underline{k} -tree- k , that is, a braid-like \underline{k} -MOY graph- k which is a tree. Thanks to the relation (2.4), it is clear⁽³⁾ that all braid-like \underline{k} -tree- k are equal in M .

It is enough to show that any braid-like \underline{k} -MOY graph- k is equal in M to P (a Laurent polynomial in q) times a braid-like \underline{k} -tree- k . We show this simultaneously on all finite sequences of integers of level k by induction on the number of merge vertices. If there is no merge then Γ is a tree and there is nothing to show. If Γ contains a merge, we cut Γ horizontally into two parts, just below its highest merge. We obtain Γ_{top} and Γ_{bot} . The latter is a \underline{k}^0 -MOY graph- k and has one merge vertex less than Γ . Hence we can use the induction hypothesis to write $\Gamma_{\text{bot}} = P(q)T$ and choose the tree T to have a split vertex on its top part which is symmetric to the merge vertex below which we cut Γ . We now stack Γ_{top} onto T , and reduce the digon thanks to the relation (2.5), we obtain that Γ is equal in M to a Laurent polynomial times a braid-like \underline{k} -tree- k .

REMARK 2.13

(1) Note that with the representation theoretic interpretation of MOY-graph, this results should be interpreted as: the multiplicity of ${}^k_q V$ (resp. $\text{Sym}_q^k V$) in $\bigotimes_{i=1}^k {}^k_q V$ (resp. $\bigotimes_{i=1}^k \text{Sym}_q^k V$) is one (see Appendix A).

(2) This lemma says, that for any braid-like $(k_1; \dots; k_r)$ -MOY graph- (k) , there exists a Laurent polynomial $r(\cdot)$, such that $\Gamma = r(\cdot)T$ in the skein module, where T

⁽³⁾This is similar to saying that the associativity of a product allows to remove parentheses in arbitrary long product. Indeed the first relation can be seen as an associativity property and a tree as a (big) product.

is a braid-like tree. From the proof, one deduces that


$$r(\gamma) = \prod_{\substack{v \in V(\gamma) \\ v \text{ merge of type } (a; b; a+b)}} \begin{bmatrix} a+b \\ a \end{bmatrix} :$$

For a latter use it will be convenient to have a preferred tree.

DEFINITION 2.14. — Let \underline{k} be a finite sequence of positive integers which add up to k . We denote by $T_{\underline{k}}$ the braid like \underline{k} -tree- k which is obtained by this inductive definition:

- T_k is a single vertical strand,
- $T_{(k_1, \dots, k_r)}$ is obtained from $T_{(k_1, \dots, k_{r-2}, k_{r-1}+k_r)}$ by splitting its rightmost strand into two strands labeled by k_{r-1} and k_r .

This is probably better understood with the following figure:



2.3. VINYL GRAPHS. — We denote by A the annulus $\{x \in \mathbb{R}^2 \mid 1 < \|x\| < 2g\}$ and for all $x = \begin{pmatrix} x_1 \\ x_2 \end{pmatrix}$ in A , we denote by t_x the vector $\begin{pmatrix} x_2 \\ x_1 \end{pmatrix}$. A ray in \mathbb{R}^2 is a half-line which starts at O , the origin of \mathbb{R}^2 .

DEFINITION 2.15. — A vinyl graph is the image of an abstract closed MOY graph in A by a smooth⁽⁴⁾ embedding such that for every point x in the image of γ , the tangent vector at this point has a positive scalar product with t_x . The set of vinyl graphs is denoted by V . We define the level of a vinyl graph to be the rotational of the underlying MOY graph. If k is a non-negative integer, we denote by V_k the set of vinyl graph with rotational equal to k . Vinyl graphs are regarded up to ambient isotopy preserving A .




FIGURE 3. A vinyl record and a vinyl graph of level 7.

⁽⁴⁾The smoothness condition is the same as the one of Definition 2.3.

REMARK 2.16. — Let Γ be a vinyl graph with rotational k , and D be a ray which does not contain any vertices of Γ . Then the condition on the tangent vectors of Γ , implies that:

- the intersection points of the ray D with Γ are all transverse and positive,
- the sum of the labels of edges which intersects D is equal to k .

Informally, the level counts the numbers of tracks of a vinyl graph.

Of course, a natural way to obtain vinyl graphs is by closing braid-like MOY graphs.

NOTATION 2.17. — Let \underline{k} be a finite sequence of integers and Γ be a braid-like \underline{k} -MOY graph- \underline{k} . Then we denote by $\widehat{\Gamma}$ the vinyl graph obtained by closing up Γ . The level of $\widehat{\Gamma}$ equals the level of \underline{k} .




FIGURE 4. The vinyl graph $\widehat{\Gamma}$ is obtained by closing up the braid-like MOY graph Γ .

The following theorem from Que elec and Rose shows that the MOY relations (2.3), (2.4), (2.5), (2.9) and (2.19) (resp. (2.12), (2.13), (2.14), (2.18) and (2.20)) defines uniquely the exterior (resp. symmetric) MOY calculus for vinyl graphs.

THEOREM 2.18 ([QR18, Lem.5.2]). — *Let S_k be the $\mathbb{Z}[q; q^{-1}]$ -module generated by vinyl graphs of level k modded out by the relations (2.4), (2.5), (2.9) and (2.19). The module S_k is generated by vinyl graphs which are collections of circles of level k .*

The proof of this result is constructive. Que elec and Rose give an algorithm which reduces a vinyl graph using (2.4), (2.5), (2.9) and (2.19). Their algorithm produces a linear combination of collection of circles. Hence adding relation (2.3) (resp. relation (2.12)) we obtain the exterior (resp. symmetric) evaluation of any vinyl graph.

In what follows we categorify the symmetric MOY calculus of vinyl graphs. We construct a category TLF_N whose objects are vinyl graphs. Suppose that F is a functor from the category TLF_N to a category \mathcal{C} with a grading, such that the relations (2.4), (2.5), (2.9) and (2.19) are compatible with F (i.e., linear combinations translates into direct sums of objects with degree shifts). From the algorithm of Que elec and Rose, we can deduce the following proposition:

PROPOSITION 2.19 ([QR18, Proof of Prop. 5.1]). — For any vinyl graph Γ , there exist two $\mathbb{N}[q; q^{-1}]$ -linear combinations of collections of circles $\sum_i a_i C_i$ and $\sum_j b_j C_j^0$ such that:

$$F(\Gamma) = \bigoplus_i F(C_i) f a_i g + \bigoplus_j F(C_j^0) f b_j g;$$

In particular, if \mathcal{C} is a category of modules over an algebra A and collections of circles are mapped to finitely generated projective modules, then $F(\Gamma)$ is finitely generated and projective. Moreover, if A is a polynomial algebra, then $F(\Gamma)$ is free and relations (2.4), (2.5), (2.9) and (2.19) are satisfied by the graded rank of the modules.


3. FOAMS

Foams have been introduced in the realm of link homologies by Khovanov [Kho04]. They have been used by Blanchet [Bla10] to fix functoriality of link homologies. They are now widely used [QR16, LQR15, EST17].

3.1. DEFINITIONS. — In the first two subsections we summarize some of the results of [RW17]. However we think that familiarity with [RW17] is essential to fully understand the constructions done in Sections 3.3, 3.4 and 5.1.1. We fix a positive integer N .

DEFINITION 3.1. — An abstract foam F is a finite collection of facets $F(F) = \{f_i\}_{i \in I}$, that is, a finite set of oriented connected surfaces with boundary, together with the following data:

- A labeling $\ell : \{f_i\}_{i \in I} \rightarrow \{0, \dots, N\}$,
- A “gluing recipe” of the facets along their boundaries such that when glued together using the recipe a neighborhood of a point of the foam has three possible local models:




The letter appearing on a facet indicates the label of this facet. That is we have facets, bindings (which are compact oriented 1-manifolds) and singular points. Each binding carries:

- an orientation which agrees with the orientations of the facets with labels a and b and disagrees with the orientation of the facet with label $a + b$.
- a cyclic ordering of the three facets around it. When a foam is embedded in \mathbb{R}^3 , we require this cyclic ordering to agree with the left-hand rule⁽⁵⁾ with

⁽⁵⁾This agrees with Khovanov’s convention [Kho04].

respect to its orientation (the dotted circle in the middle indicates that the orientation of the binding points to the reader, a crossed circle indicates the other orientation, see Figure 6):



The cyclic orderings of the different bindings adjacent to a singular point should be compatible. This means that a neighborhood of the singular point is embeddable in \mathbb{R}^3 in a way that respects the left-hand rule for the four bindings adjacent to this singular point.

REMARK 3.2. — Let us explain shortly what is meant by “gluing recipe”. The boundaries of the facets forms a collection of circles. We denote it by S . The gluing recipe consists of:


- For a subset S^θ of S , a subdivision of each circle of S^θ into a finite number of closed intervals. This gives us a collection I of closed intervals.
- Partitions of $I \cap (S \cap S^\theta)$ into subsets of three elements. For every subset $(X_1; X_2; X_3)$ of this partition, three diffeomorphisms $\varphi_1 : X_2 \rightarrow X_3$, $\varphi_2 : X_3 \rightarrow X_1$, $\varphi_3 : X_1 \rightarrow X_2$ such that $\varphi_3 \circ \varphi_2 \circ \varphi_1 = \text{Id}_{X_2}$.

A foam is obtained by gluing the facets along the diffeomorphisms, provided that the conditions given in the previous definition are fulfilled.

DEFINITION 3.3. — A *decoration* of a foam F is a map $f \mapsto P_f$ which associates with any facet f of F an homogeneous symmetric polynomial P_f in $\dim(f)$ variables. A *decorated foam* is a foam together with a decoration.

From now on all foams are decorated.

DEFINITION 3.4. — A *closed foam* is a smoothly embedded abstract foam in \mathbb{R}^3 . Smoothness means that the facets are smoothly embedded and the different oriented tangent planes agree on bindings and singular points as depicted here



Just like for MOY graphs (see Definition 2.3), we will usually not care too much about the smoothness on bindings and singular points when drawing foams.




FIGURE 5. Example of a foam. The cyclic ordering on the central binding is (5;2;3).

The notion of foam extends naturally to the notion of foam with boundary. The boundary of a foam has a structure of a MOY graph. We require that the facets and bindings are locally orthogonal to the boundary to be able to glue them together. Probably the most local framework is given by the concept of *canopolis* of foams. We refer to [BN05, ETW18] for more details about this approach. In what follows we will consider:

- Foams in $\mathbb{R}^2 \times [0;1]$ where the boundary is contained in $\mathbb{R}^2 \times \{0;1\}$ (see Definition 3.5);
- Foams in the cube $[0;1]^3$, where the boundary is contained in $[0;1]^2 \times \{0;1\} \cup \{0;1\} \times [0;1]^2$ (see Section 3.3);
- Foams in the thickened annulus $A \times [0;1]$ where the boundary is contained in $A \times \{0;1\}$ (see Section 3.4).

DEFINITION 3.5. — The category Foam consists of the following data:

- Objects are closed MOY graphs,
- Morphisms from Γ_0 to Γ_1 are (ambient isotopy classes relatively to the boundary of) foams in $\mathbb{R}^2 \times [0;1]$ whose boundary is contained in $\mathbb{R}^2 \times \{0;1\}$. The part of the boundary in $\mathbb{R}^2 \times \{0\}$ (resp. $\mathbb{R}^2 \times \{1\}$) is required to be equal to Γ_0 (resp. Γ_1). Composition of morphisms is given by stacking foams and rescaling in the vertical direction.

DEFINITION 3.6. — If F is a foam (possibly with boundary), a *sub-surface* is a collection of oriented facets F' such that their union is a smooth oriented surface whose boundary is contained in the boundary of F .

3.2. REMINDER ON THE EXTERIOR EVALUATION OF FOAMS. — The combinatorial evaluation can be thought of as a higher dimensional state sum formula of the state sum of Murakami-Ohtsuki-Yamada [MOY98] for evaluating MOY graphs.

DEFINITION 3.7. — A *pigment* is an element of $P = \mathbb{N}^g$. The set P is endowed with the natural order.

A *coloring* of a foam F is a map $c: F \rightarrow P$, such that

- For each facet f , the number of elements $\#c(f)$ of $c(f)$ is equal to $\text{val}(f)$.

– For each binding joining a facet f_1 with label a , a facet f_2 with label b , and a facet f_3 with label $a + b$, we have $c(f_1) \upharpoonright c(f_2) = c(f_3)$. This condition is called the *flow condition*.

A *colored foam* is a foam together with a coloring. For a given foam F , the set of all its colorings is denoted $\text{col}_N(F)$.

A careful inspection of the local behavior of colorings in the neighborhood of bindings and singular points gives the following lemma:

LEMMA 3.8


(1) If $(F; c)$ is a colored foam and i is an element of \mathcal{P} , the union (with the identification coming from the gluing procedure) of all the facets which contain i in their colors is a surface. It is called the *monochrome surface* of $(F; c)$ associated with i and is denoted by $F_i(c)$. The restriction we imposed on the orientations of facets ensure that $F_i(c)$ is oriented.

(2) If $(F; c)$ is a colored foam and i and j are two distinct elements of \mathcal{P} , the union (with the identification coming from the gluing procedure) of all the facets which contain i or j but not both in their color set is a surface. It is called the *bichrome surface* of $(F; c)$ associated with $i; j$. This is the symmetric difference of $F_i(c)$ and $F_j(c)$ and is denoted by $F_{ij}(c)$. The restriction imposed on the orientations of facets ensures that $F_{ij}(c)$ can be oriented via taking the orientation of facets containing i and the reverse orientations on facets containing j .

(3) Let $i < j$ and consider a binding joining the facets f_1, f_2 and f_3 . Suppose that $i \in c(f_1), j \in c(f_2)$ and $i, j \in c(f_3)$. We say that the binding is *positive* with respect to $(i; j)$ if the cyclic order on the binding is $(f_1; f_2; f_3)$ and *negative* with respect to $(i; j)$ otherwise. The set $F_i(c) \setminus F_j(c) \setminus F_{ij}(c)$ is a collection of disjoint circles. Each of these circles is a union of bindings; for every circle the bindings are either all positive or all negative with respect to $(i; j)$.

Please note that the previous lemma contains the definition of *monochrome* and *bichrome surfaces*.

EXAMPLE 3.9 ([RW17, Ex. 2.7]). — Suppose $N = 4$ (and therefore $\mathcal{P} = \{1, 2, 3, 4\}$) and consider the colored foam $(F; c)$ given by the figure below



where the big digits represent labels. Note that the orientation of every facet can be deduced from the orientations of the bindings. Tables 1 and 2 describe the monochrome and bichrome surfaces as well as the values of $\overset{+}{ij}(c)$ for this colored foam.





$i \in \mathbb{P}$	Monochrome Surface $F_i(c)$	In words
1		Sphere (on the right)
2		Sphere (on the left)
3		Empty set
4		Sphere (on the left)

TABLE 1. The monochrome surfaces of Example 3.9.

$(i; j) \in \mathbb{P}$	1	2	3	4
1		Torus	Sphere (on the right)	Torus
2	2		Sphere (on the left)	Empty set
3	0	0		Sphere (on the left)
4	2	0	0	

TABLE 2. The bichrome surfaces (top right) and the $\overset{+}{ij}(c)$ (bottom left) of Example 3.9.

REMARK 3.10. — Monochrome surfaces of $(F; c)$ are sub-surfaces of F while, in general, bichrome surfaces are not in the sense of Definition 3.6.

DEFINITION 3.11. — Let $(F; c)$ be a colored foam and $i < j$ be two pigments. A circle in $F_i(c) \setminus F_j(c) \setminus F_{ij}(c)$ is *positive* (resp. *negative*) with respect to $(i; j)$ if it consists of positive (resp. negative) bindings. We denote by $\overset{+}{ij}(c)_F$ (resp. $\overset{-}{ij}(c)_F$) or simply

$\overset{+}{ij}(c)$ (resp. $\underset{-}{ij}(c)$) the number of positive (resp. negative) circles with respect to $(i; j)$. We set as well $ij(c) = \overset{+}{ij}(c) + \underset{-}{ij}(c)$. See Figure 6 for a pictorial definition.




FIGURE 6. A pictorial definition of the signs of the circle, we assume $i < j$. Recall that a dotted circle in the middle indicates that the orientation of the binding points to the reader and a crossed circle indicates the other orientation.

DEFINITION 3.12. — The *degree* deg_N of a foam F is the sum of the following contributions:

- For each facet f with label a , set $\text{deg}(f) = a(N - a) \chi(f)$, where χ stands for the Euler characteristic;
- For each interval binding e (i.e., not circle-like binding) surrounded by three facets with labels a, b and $a + b$, set $\text{deg}(e) = ab + (a + b)(N - a - b)$;
- For each singular point p surrounded with facets with labels $a, b, c, a + b, b + c, a + b + c$, set $\text{deg}(p) = ab + bc + cd + da + ac + bd$ with $d = N - a - b - c$;
- Thus,

$$\text{deg}_N(F) = \sum_f \text{deg}(f) + \sum_e \text{deg}(e) + \sum_p \text{deg}(p) + \sum_f \text{deg}(P_f);$$

where the variables of the polynomials P_f have degree 2.

REMARK 3.13

(1) The degree is additive with respect to the composition of foam. This is the same degree as in [QR16], but since we are not in a 2-categorical setting, the contributions of σ_0 and σ_1 to the degree are equal to 0.

(2) The degree can be thought of as an analogue of the Euler characteristic. The degree of the foam of Figure 5 is equal to -16 when $N = 6$.

DEFINITION 3.14. — If $(F; c)$ is a colored foam, define:

$$s(F; c) = \sum_{i=1}^N i \chi(F_i(c)) + \sum_{1 \leq i < j \leq N} \overset{+}{ij}(F; c);$$

$$P(F; c) = \prod_{f \text{ facet of } F} P_f(c(f)); \quad Q(F; c) = \prod_{1 \leq i < j \leq N} (x_i - x_j)^{\overset{+}{ij}(c)=2}$$

$$hF; c_i = (-1)^{s(F; c)} \frac{P(F; c)}{Q(F; c)};$$

In the definition of $P(F; c)$, $P_f(c(f))$ means the polynomial P evaluated on the variables $f x_i g_{i2c(f)}$. Since the polynomial P_f is symmetric, the order of the variables does not matter. A facet f is called *trivially decorated*, if $P_f = 1$. Define the *evaluation of the foam F* by:

$$hF i := \sum_{c \text{ coloring of } F} hF; c i :$$

REMARK 3.15. — Let F be a foam and denote by $\langle \cdot \rangle$ the product of all decorations of facets of label 0. Consider F^0 the foam obtained from F by removing the facets with label 0. There is a one-one correspondence between the colorings of F and the colorings of F^0 . For every coloring c of F and its corresponding coloring c^0 of F^0 , we have $hF; c i = hF^0; c^0 i$, and consequently $hF i = hF^0 i$.

PROPOSITION 3.16 ([RW17]). — Let F be a foam, then $hF i$ is an homogeneous element of $\mathbb{Q}[x_1; \dots; x_N]^{\mathbb{S}^N}$ of degree $\text{deg}_N(F)$.

PROPOSITION 3.17 ([RW17]). — The following local identities and their mirror images hold:

(3.1)

(3.2)

(3.3)


(3.4) $\langle \text{Diagram 1} \rangle = \sum_{2T(a;b)} (-1)^{bj} \langle \text{Diagram 2} \rangle;$

(3.5) $\langle \text{Diagram 3} \rangle = \sum_{2T(b;N \ a \ b)} (-1)^j \langle \text{Diagram 4} \rangle;$

(3.6) $\langle \text{Diagram 5} \rangle = \sum_{2T(r;s)} (-1)^{bj} \langle \text{Diagram 6} \rangle;$

(3.7) $\langle \text{Diagram 7} \rangle = \sum_{\substack{j=\max(0; m-n); \dots; m \\ 2T(k \ j; \ k+j)}} (-1)^{j+(k+j)(m-j)} \langle F_j \rangle;$

where



Moreover, in identities (3.1), (3.4), (3.5), (3.6) and (3.7), the terms on the right-hand sides are mutually orthogonal idempotents. In the previous formulas $T(a; b)$ denotes the set of all Young diagram contained in the rectangle of size $a \times b$, s_λ denotes the Schur polynomial associated with λ and c denote the Littlewood-Richardson constant. Further explanations of notations and conventions can be found in [RW17, App. 1].

In identity (3.7), h needs to be extended linearly to formal \mathbb{Q} -linear combinations of foams.

Using this evaluation and the universal construction idea (see [BHMV95]), we define a functor F_N from the category of foams to the category of $\mathbb{Q}[x_1, \dots, x_N]^{S_N}$ -module.

If \mathcal{G} is a MOY graph, consider the free graded $\mathbb{Q}[x_1, \dots, x_N]^{S_N}$ -module spanned by $\text{Hom}_{\text{Foam}}(\mathcal{G}; \cdot)$. We mod this space out by

$$\bigcap_{\mathcal{G} \in \text{Hom}_{\text{Foam}}(\mathcal{G}; \cdot)} \text{Ker} \left(\begin{array}{c} \text{Hom}_{\text{Foam}}(\mathcal{G}; \cdot) \rightarrow \mathbb{Q}[x_1, \dots, x_N]^{S_N} \\ F \mapsto h(G) \cdot F_i \end{array} \right);$$

We define $F_N(\mathcal{G})$ to be this quotient. The definition of F_N on morphisms follows. From Proposition 3.17, we deduce that the functor F_N categorifies the exterior MOY calculus.

COROLLARY 3.18 ([RW17]). — *Let \mathcal{G} be a closed MOY graph, then $F_N(\mathcal{G})$ is a free graded $\mathbb{Q}[x_1, \dots, x_N]^{S_N}$ -module of graded rank equal to $h(\mathcal{G})$.*

In Sections 3.3 and 3.4 we will work with elements of $F_N(\mathcal{G})$. Such elements are represented by $\mathbb{Q}[x_1, \dots, x_N]^{S_N}$ -linear combination of foams bounding \mathcal{G} . Since it is

more convenient to work with representatives of classes than with the classes themselves, we introduce the following terminology.

DEFINITION 3.19

(1) Let $\sum_i i F_i$ and $\sum_j j G_j$ be two elements of the free graded $\mathbb{Q}[x_1; \dots; x_N]^{S_N}$ -module spanned by $\text{Hom}_{\text{Foam}}(?; ?)$. We say that they are *N-equivalent* if they represent the same element in $F_N(\cdot)$.


(2) Let $\sum_i i F_i$ and $\sum_j j G_j$ two elements of the graded \mathbb{Q} -vector space generated by $\text{Hom}_{\text{Foam}}(?; ?)$. We say that they are *1-equivalent* if they are *N-equivalent* for all N in \mathbb{N} .

(3) Define $F_1(\cdot)$ the graded \mathbb{Q} -vector space generated by $\text{Hom}_{\text{Foam}}(?; ?)$ modded out by 1-equivalence.

REMARK 3.20. — The local identities (3.1), (3.3), (3.4), (3.6) and (3.7) can be translated into 1-equivalences, while the local identities (3.2) and (3.5) can only be translated into *N*-equivalences.

3.3. DISK-LIKE FOAMS (OR HOMPLYPT FOAMS). — For this section we fix a non-negative integer. We will work in \mathbb{R}^3 and we denote by P the plane spanned by $\begin{pmatrix} 1 \\ 0 \\ 0 \end{pmatrix}$ and $\begin{pmatrix} 0 \\ 0 \\ 1 \end{pmatrix}$:

We consider the cube $C = [0; 1]^3$ and will use the following parametrization of its boundary:



The symbols s denote the 6 squares of the boundary of C and the letters $f; h; l; r; b; t$ stand for **f**ront, **h**idden, **l**eft, **r**ight, **b**ottom and **t**op. The plan P is parallel to the square s_f and s_h .

DEFINITION 3.21. — Let F be a foam with boundary embedded in C . Suppose that the boundary of F is contained in $s \cdot [s_r [s_b [s_t$, and that the MOY-graphs $F \setminus s_f$, $F \setminus s_h$, $F \setminus s_b$ and $F \setminus s_t$ are all braid-like. We say that F is *disk-like* if for every point x of F , the normal line of the foam F at x is *not* parallel to P .

We say that F is a *rooted -foam of level k* if additionally:

- the restriction of F on s_b is a single strand labeled k ,
- the restriction of F on s_t is a braid-like MOY graph Γ ,
- the restriction of F on s_r and s_l are standard trees (see Definition 2.14).

REMARK 3.22. — The notion of *level* (see Remark 2.11) extends to disk-like foams.

The name disk-like comes from the following lemma.

LEMMA 3.23. — *Let F be a disk-like foam. Every non-empty connected subsurface of F is a disk whose boundary circle intersects each of the four squares $s_r; s_l; s_b$ and s_t non-trivially.*

Proof. — We consider a non-empty subsurface Σ of F . The condition on the normal vector of disk-like foams implies that the projection on the second (resp. the third) coordinate provides a Morse function with no critical points. This implies that Σ is diffeomorphic to its intersection with s_r (resp. s_b) times the interval. Since Σ is non-empty and connected $s_r \cap \Sigma$ is an interval. Finally, Σ is a disk which intersects non-trivially the four squares $s_r; s_l; s_b$ and s_t .

REMARK 3.24. — Let F be a disk-like foam. The condition on the normal vector implies that for all t in $[0; 1]$ the intersection of $ftg \subset [0; 1]^2$ (resp. $[0; 1] \times ftg$) with F is transverse. Moreover, if $ftg \subset [0; 1]^2$ (resp. $[0; 1] \times ftg$) does not contain any singular point of F , $ftg \subset [0; 1]^2 \setminus F$ (resp. $[0; 1] \times ftg \setminus F$) is a braid-like MOY graph. A very similar result (Corollary 3.37) is given a proper proof in the next subsection.

DEFINITION 3.25. — Let us fix a non-negative integer k . The 2-category DLF_k of disk-like foams of level k consists of the following data:

- Objects are finite sequences of positive integers of level k .
 - A 1-morphism from \underline{k}_0 to \underline{k}_1 is a braid-like \underline{k}_1 -MOY graph- \underline{k}_0 (it has level k).
- Composition is given by concatenation of braid-like MOY graphs.

- A 2-morphism from a braid-like \underline{k}_1 -MOY graph- \underline{k}_0 bot to a braid-like \underline{k}_1 -MOY graph- \underline{k}_0 top is an ambient isotopy class (relative to the boundary) of a disk-like foam F in the cube C such that:

- the intersection of F with s_b is equal to bot ,
- the intersection of F with s_t is equal to top ,
- the intersection of F with s_r is equal to $\underline{k}_0 \subset [0; 1]$,
- the intersection of F with s_l is equal to $\underline{k}_1 \subset [0; 1]$.

Compositions are given by stacking disk-like foams and re-scaling. This is illustrated in Figure 7.

The 2-category $\mathbb{D}LF_k$ is constructed as follows:

- Start from the 2-category DLF_k .
- Linearize the 2-hom-spaces over \mathbb{Q} .
- Mod out every 2-homspace by \sim -equivalence (disk-like foams are considered as foams from ? to the boundary vinyl graph).




FIGURE 7. Vertical (top) and horizontal (bottom) compositions of 2-morphisms in DLF_k .

DEFINITION 3.26. — Let \underline{k}_0 and \underline{k}_1 be two objects of the DLF_k and bot and top two 1-morphism from \underline{k}_0 to \underline{k}_1 . The *degree* of a 2-morphism $F : \text{bot} \rightarrow \text{top}$ is given by formula

$$\text{deg}^D(F) = \text{deg}_0(F) \frac{jjk_0j^2 + jjk_1j^2}{2};$$

where $\text{deg}_0(F)$ is the degree of F as an exterior 0-foam (see Definition 3.12) and if $\underline{k} := (k_1; \dots; k_n)$ is a finite sequence of non-negative integers, $jjk_j^2 := \sum_{i=1}^n k_i^2$.

Let top a 1-morphism from \underline{k}_0 to \underline{k}_1 . The *degree* of a rooted top -foam F is given by:

$$\text{deg}^r(F) = \text{deg}_0(F) \frac{jjk_0j^2 + jjk_1j^2 + 2k^2}{4};$$

REMARK 3.27. — One easily checks that with these definition the degree of 2-morphisms is additive with respect to vertical and horizontal compositions. In particular, the degrees of identity 2-morphisms are 0. Since the relations defining the \sim -equivalence are homogeneous, this degree induces a grading on the 2-homspaces of DLF_k . Moreover, the composition of a rooted bot -foam with an element of $\text{hom}(\text{bot}; \text{top})$ is a rooted top -foam, and the degree is additive with respect to this composition.

DEFINITION 3.28. — Let Γ be a braid-like MOY-graph, and F be a rooted top -foam. We say that F is *tree-like*, if $\text{ftg}_{[0;1]^2} \setminus F$ is a tree for all t in $[0;1]$. In particular, if $\text{ftg}_{[0;1]^2}$ does not contain any singular point of F , then $\text{ftg}_{[0;1]^2} \setminus F$ is a braid-like tree.

From Lemma 2.12 we derive the following lemma which tells us that disk-like foams are combinatorially very simple:

LEMMA 3.29. — Let k be a non-negative integer, Γ be the braid-like k -MOY graph consisting of one single strand labeled by k and F be a rooted top -foam (that is, a

disk-like foam which bound a circle with label k). Then F is 1-equivalent to a disk with label k decorated by $(-1)^{k(k+1)/2} \langle \widehat{F} \rangle_k$, where \widehat{F} is the foam obtained from F by capping it with a disk labeled by k .

Note that this makes sense since $\langle \widehat{F} \rangle_k$ is a symmetric polynomial in k variables.

Proof. — Let us denote by D the disk with label k decorated by $(-1)^{k(k+1)/2} \langle \widehat{F} \rangle_k$. It follows directly from the definition of the sl_k -evaluation of foams, that F is k -equivalent to D . The sign comes from the term $\sum_{i=1}^k i(F_i(c))^{-2}$ in the definition of $s(F; c)$ (see Definition 3.14). If $N < k$, then both D and F are N -equivalent to 0.

If $N > k$, we will see that the N -equivalence between F and D follows from their k -equivalence. We need to prove that for any foam G bounding a circle with label k , $hF \ G; ci_N = hD \ G; ci_N$. First note that thanks to identity (3.2), we can suppose that G is a decorated disk of label k . In this case, $D \ G$ is a decorated sphere of label k . Let c be a coloring of $D \ G$. The coloring c is given by the color $I(c) = (i_1(c); \dots; i_k(c))$ of this sphere. We have:

$$(3.8) \quad hD \ G; ci_N = \frac{hD \ G; i_k(x_{i_1(c)}; \dots; x_{i_k(c)})}{\prod_{j \in \text{Pr } I(c)} x_j^{i_j(c)}}$$

Note the foam $D \ G$ being a sphere, it admits only one sl_k -coloring. It can be obtained from c by replacing $i_a(c)$ by a in $I(c)$ for all $a \in \{1; \dots; k\}$.

Similarly, if c is a coloring of $F \ G$, it gives to the facet containing G a color $I(c) = (i_1(c); \dots; i_k(c))$. Let us denote by $f : I(c) \rightarrow \{1; \dots; k\}$ the one-to-one map given by $f(i_a) = a$ and $f(c)$ the sl_k -coloring of $F \ G$ induced from c by f . We have:

$$(3.9) \quad hF \ G; ci_N = \frac{hF \ G; f(c) i_k(x_{i_1(c)}; \dots; x_{i_k(c)})}{\prod_{j \in \text{Pr } I(c)} x_j^{i_j(c)}}$$

Combining (3.8) and (3.9) and keeping the same notations, we get:


$$\begin{aligned} hF \ G; ci_N &= \sum_{c \in \text{col}_N(F \ G)} hF \ G; ci_N = \sum_{\substack{I \in \mathcal{P} \\ \# I = k}} \sum_{c \in \text{col}_N(F \ G) \\ I(c) = I} hF \ G; ci_N \\ &= \sum_{\substack{I \in \mathcal{P} \\ \# I = k}} \sum_{c \in \text{col}_N(D \ G) \\ I(c) = I} hD \ G; ci_N = hD \ G; ci_N \end{aligned}$$

LEMMA 3.30. — *A rooted k -foam F is 1-equivalent to a \mathbb{Z} -linear combination of tree-like foams.*

Proof. — First, assume that F is a braid-like k -MOY graph- k . We will show that F is 1-equivalent to a \mathbb{Z} -linear combination of foams which are superposition of a tree-like k -foam on top of a disk-like foam which bounds a circle and conclude by Lemma 3.29

Proof. — This follows directly from the definition of the exterior evaluation of foams. The set of colorings of F and of F^θ are in one-to-one correspondence (because they are both in one-one correspondence with the set of colorings of their boundary). Let us denote c and c^θ two corresponding colorings of F and F^θ . The monochrome and the bichrome surfaces of $(F; c)$ and $(F^\theta; c^\theta)$ are diffeomorphic. The oriented arcs in $(F; c)$ and $(F^\theta; c)$ are in one-one correspondence preserving their orientation. Finally the condition on the decorations of F and F^θ ensures that their contributions to the evaluation are equal.

3.4. VINYL FOAMS (OR SYMMETRIC FOAMS). — In this part, we work in the thickened annulus $A = [0; 1]$. If $x := \begin{pmatrix} x_1 \\ x_2 \\ x_3 \end{pmatrix}$ is an element of $A = [0; 1]$, we denote by t_x the vector $\begin{pmatrix} x_2 \\ x_1 \\ 0 \end{pmatrix}$, by v the vector $\begin{pmatrix} 0 \\ 0 \\ 1 \end{pmatrix}$, and by P_x the affine plane containing x and spanned by t_x and v . If θ is an element of $[0; 2\pi[$, P is the half-plane $\left\{ \begin{pmatrix} \cos t \\ \sin t \\ t \end{pmatrix} \mid (t; t) \geq \mathbb{R}_+ \times \mathbb{R} \right\}$.



DEFINITION 3.33. — Let k be a non-negative integer and σ_0 and σ_1 two vinyl graphs of level k . Let F be a foam with boundary embedded in $A = [0; 1]$. Suppose that $F \setminus (A = f_0g) = \sigma_0$ and $F \setminus (A = f_1g) = \sigma_1$. We say F is a *vinyl $(\sigma_1$ -foam- σ_0 of level k* if for every point x of F , the normal line of F at x is *not* contained in P_x . See Figure 9 for an example.

REMARK 3.34. — Note that if we cut a vinyl foam F along a half plane P , we obtain a disk-like foam.

DEFINITION 3.35. — The category TLF_k of vinyl foams of level k consists of the following data:

- the objects are elements of V_k , i.e., vinyl graphs of level k ,
- morphisms from σ_0 to σ_1 are (ambient isotopy classes of) vinyl $(\sigma_1$ -foams- σ_0).

Composition is given by stacking vinyl foams together and rescaling. In the category TLF_k we have one distinguished object which consists of a single essential circle with label k denoted by S_k . The *degree* $\text{deg}^T(F)$ of a vinyl foam F is equal to $\text{deg}_0(F)$. Note that the degree is additive with respect to the composition in TLF_k .




FIGURE 9. An example of a vinyl foam

The name vinyl comes from the following lemma.

LEMMA 3.36. — *Let F be a vinyl $(-1, 0)$ -foam. Then any non-empty connected subsurface σ of F is an annulus. Moreover, for every t in $[0; 1]$, $A \cap \sigma_t$ is an essential circle in $A \cap \sigma_t$. Such annuli are called tubes.*

Proof. — The condition on the tangent plane of vinyl foams implies that the projection on the last coordinate is a Morse function for σ and that it has no critical points. The result follows.

COROLLARY 3.37. — *Let F be a vinyl $(-1, 0)$ -foam.*

- (1) *Let t be an element of $[0; 1]$ such that the intersection of $A \cap \sigma_t$ and F is generic (i.e., $A \cap \sigma_t$ does not contain any singular points of F and the intersection of $A \cap \sigma_t$ with the bindings of F is transverse). Then $F \setminus A \cap \sigma_t$ is a vinyl graph.*
- (2) *Let P be an element of $[0; 2[$ and assume that the intersection of F with P is generic. Then the intersection of F and P is braid-like.*

Before proving the statements, let us emphasize that there are only finitely many t 's (resp. s 's) for which the intersection of F and $A \cap \sigma_t$ (resp. P) is not generic.

Proof. — First note that every point of F is contained in a connected subsurface which intersects $A \cap \sigma_t$ non-trivially. Indeed we can cable foams just like we can cable MOY graphs (see Figure 2). This gives us a collection of sub-surfaces of F which covers it. Let us prove the first part. Let $x = \begin{pmatrix} x_1 \\ x_2 \\ t_0 \end{pmatrix}$ be a point in F and σ a connected subsurface of F containing x . Since F is vinyl, the scalar product of t_x with the tangent vector of $F \setminus A \cap \sigma_t$ is non-zero. Since σ is connected, this quantity is either always positive or always negative on σ . The graphs σ_0 and σ_1 being vinyl, it is positive. This proves that $F \setminus A \cap \sigma_t$ is vinyl.

The second part is similar but we consider the scalar product of $P \setminus F$ with $v = \begin{pmatrix} 0 \\ 0 \\ 1 \end{pmatrix}$.

The notion of tree-like foams developed in Section 3.3 extends mutatis mutandis to the concept of foams in the thickened annulus. The analogues of rooted \mathbb{Z} -foams are $\widehat{\mathbb{Z}}$ -foams- S_k . We have analogues of Lemmas 3.30 and 3.32:

LEMMA 3.38. — *Let \mathcal{G} be a vinyl graph of level k and F a vinyl \mathbb{Z} -foam- S_k . Then F is \mathbb{Z} -equivalent to a \mathbb{Z} -linear combination of tree-like foams.*

Proof. — First we choose a θ in $[0; 2\pi]$ such that the intersection of P with the foam F is generic. Thanks to Corollary 3.37, we know that this intersection is a braid-like \underline{k} -MOY graph- k B for some finite sequence \underline{k} of positive integers. We can suppose that F is locally diffeomorphic to $B \times [0; \pi]$. The algorithm described in Lemma 2.12 and the local identities (3.1) and (3.4) tells us that F is \mathbb{Z} -equivalent to a \mathbb{Z} -linear combination of vinyl foams such that the intersection with P is the canonical \underline{k} -tree- k . If we cut these foams along P , we can apply Lemma 3.30 on each of these foams. Gluing back the result along the canonical \underline{k} -tree- k gives us a \mathbb{Z} -linear combination of tree-like \mathbb{Z} -foams- S_k which is \mathbb{Z} -equivalent to F .

REMARK 3.39. — Just like for the disk-like context, thanks to the dot migration identity (3.3), we can move all non-trivial decorations of a tree-like foam on its leaves.

The proof of Lemma 3.32 can be easily adapted to the annular case. This gives the following lemma.

LEMMA 3.40. — *Let \mathcal{G} be a vinyl graph and F and F^0 be two tree-like \mathbb{Z} -foams- S_k with non-trivial decorations only on their leaves. Suppose furthermore that these decorations are the same for F and F^0 . Then F is \mathbb{Z} -equivalent to F^0 .*

4. SOERGEL BIMODULES

In this section we prove that for any braid-like MOY graph \mathcal{G} the space of rooted \mathbb{Z} -foams regarded up to \mathbb{Z} -equivalence is isomorphic to the Soergel bimodule associated with \mathcal{G} .

4.1. SOME POLYNOMIAL ALGEBRAS

NOTATION 4.1

(1) We denote the graded ring $\mathbb{Q}[T_1; \dots; T_N]^{S_N}$ by R_N , where the indeterminates T_i are homogeneous of degree 2. This is the q -degree.

(2) Denote by \mathcal{C} the category of \mathbb{Z} -graded finitely generated projective R_N -modules. If M is an object of \mathcal{C} , Mq^i denotes the same object where the degree has been shifted by i . This means that $(Mq^i)_j = M_{j-i}$. If $P(q) = \sum_i a_i q^i$ is a Laurent polynomial in q with positive integer coefficients, $MP(q)$ denotes the module

$$\bigoplus_i (Mq^i)^{a_i}.$$

(3) Let $\underline{k} = (k_1; \dots; k_\ell)$ be a finite sequence of positive integers of level k (if $k = 0$ the empty sequence is allowed). The group $\prod_{i=1}^{\ell} S_{k_i}$ is denoted by $S_{\underline{k}}$. We define the algebra $A_{\underline{k}}$:

$$A_{\underline{k}} := R_N[x_1; \dots; x_k]^{S_{\underline{k}}}.$$

The indeterminates x are homogeneous of degree 2. If $\underline{k} = (k)$ (that is, if \underline{k} has length 1), we write A_k instead of $A_{(k)}$.

(4) If Γ is a vinyl graph, denote by $F_{\Gamma}^T(\cdot)_{\mathbb{Q}}$ the graded \mathbb{Q} -vector space generated by vinyl Γ -foams- $S_{\underline{k}}$ modded out by Γ -equivalence (see Definition 3.19). Define $F_{\Gamma}^T(\cdot) := F_{\Gamma}^T(\cdot)_{\mathbb{Q}} \otimes R_N$. Since for all k , the exterior sl_k -evaluation of foams is homogeneous, the R_N -module $F_{\Gamma}^T(\cdot)$ is naturally graded.

Before dealing with Soergel bimodules, we state the following lemma which relates the algebra $A_{\underline{k}}$ with vinyl foams.

LEMMA 4.2. — *Let $\underline{k} := (k_1; \dots; k_\ell)$ be a finite sequence of positive integers and $S_{\underline{k}}$ be the vinyl graph which consists of ℓ oriented circles with labeling induced by \underline{k} . Then $F_{\Gamma}^T(S_{\underline{k}})$ is isomorphic to $A_{\underline{k}}$ as a graded R_N -module.*

Proof. — Let $k = \sum_i k_i$ and T be the canonical \underline{k} -tree- (k) . For i in $\{1; \dots; g\}$ and λ_i denotes a Young diagram with at most k_i lines. Denote $s_i^{(\lambda_i)}$ the Schur polynomial associated with λ_i in the variables $x_{1+r_i}; \dots; x_{k_i+r_i}$ where $r_i = \sum_{j=1}^{i-1} k_j$.

A R_N -base of $A_{\underline{k}}$ is given by $(s_1^{(\lambda_1)}; \dots; s_\ell^{(\lambda_\ell)}) =: \lambda$ where the λ_i 's take all possible shapes. Being given $\lambda = (\lambda_1; \dots; \lambda_\ell)$ a sequence of Young diagrams as described above, define F_{λ} to be the foam $T \times S^1$ where the i th leaf of F is decorated by the Schur polynomial $s_i^{(\lambda_i)}$.

Let us prove that the R_N -linear map sending $\lambda \in A_{\underline{k}}$ to $F_{\lambda} \in F_{\Gamma}^T(S_{\underline{k}})$ is bijective. It is surjective because of dots migration (3.3). Let $\sum_{j=1}^h \lambda_j = 0$ be a linear combination of monomials mapped to 0. Let ℓ_{\max} be the maximal length of all lines appearing in the Young diagrams of all the size of the Young diagram of $(\lambda_j)_{j=1; \dots; h}$. For $M = \ell_{\max}(k+1)$, the foams F_{λ_h} precomposed by a cup labeled by k are linearly independent in $F_M(S_{\underline{k}})$. Hence the coefficients λ_j must be all equal to 0.

4.2. SINGULAR SOERGEL BIMODULES. — We introduce singular Soergel bimodules. See for instance [Soe92, Str04, Kho07, Wil11] or [Wed19] for a pictorial description close to ours.

DEFINITION 4.3. — Let Γ be a braid-like \underline{k}_1 -MOY graph- \underline{k}_0 . If Γ has no trivalent vertices, we have $\underline{k}_0 = \underline{k}_1$ and we define $B(\Gamma)$ to be equal to $A_{\underline{k}_0}$ as a $A_{\underline{k}_0}$ -module- $A_{\underline{k}_0}$. If Γ has only one trivalent vertex (which is supposed to be of type $(a; b; a+b)$), then:

- if the length of \underline{k}_1 is equal to the length of \underline{k}_0 plus 1, we define $B(\Gamma)$ to be $A_{\underline{k}_1} q^{ab-2}$ as a $A_{\underline{k}_1}$ -module- $A_{\underline{k}_0}$;
- if the length of \underline{k}_0 is equal to the length of \underline{k}_1 plus 1, we define $B(\Gamma)$ to be $A_{\underline{k}_0} q^{ab-2}$ as a $A_{\underline{k}_1}$ -module- $A_{\underline{k}_0}$;

If Γ has more than one trivalent vertex, if necessary we perturb⁽⁷⁾ Γ to see it as a composition:


$$\Gamma = \Gamma_0 \circ \Gamma_1 \circ \dots \circ \Gamma_{t-1} \circ \Gamma_t$$

where Γ_i is a braid-like k^{i+1} -MOY graph- k^i with one trivalent vertex, for all i in $\{0, \dots, t\}$. The symbols k^i mean that Γ_i and Γ_{i-1} are glued along k^i . We have $k^0 = k_0$ and $k^{t+1} = k_1$. We define

$$B(\Gamma) := B(\Gamma_0) \otimes_{A_{k_0}} B(\Gamma_1) \otimes_{A_{k_1}} \dots \otimes_{A_{k_{t-1}}} B(\Gamma_t) \otimes_{A_{k_t}} B(\Gamma_{t+1})$$

The space $B(\Gamma)$ has a natural structure of A_{k_1} -module- A_{k_0} . It is called the *Soergel bimodule associated with Γ* . Note that the grading of $B(\Gamma)$ takes values either in \mathbb{Z} or in $\frac{1}{2} + \mathbb{Z}$.

EXAMPLE 4.4. — The singular Soergel bimodule associated with



is the following $A_{(2,1;2)}$ -module- $A_{(1,2;1;1)}$:

$$A_{(2,1;2)} \otimes_{A_{(2;3)}} A_{(2;3)} \otimes_{A_{(2;3)}} A_{(2;2;1)} \otimes_{A_{(4;1)}} A_{(4;1)} \otimes_{A_{(4;1)}} A_{(3;1;1)} \otimes_{A_{(3;1;1)}} A_{(1,2;1;1)} q^{13=2}$$

$$\otimes_{A_{(2;1;2)}} A_{(2;3)} \otimes_{A_{(2;2;1)}} A_{(4;1)} \otimes_{A_{(4;1)}} A_{(1,2;1;1)} q^{13=2}.$$

REMARK 4.5. — For Definition 4.3 to be valid, the isomorphism type of the bimodule $B(\Gamma)$ should not depend on the decomposition of Γ . This is clear since two such decompositions are related by the “commutation of faraway vertices” for which the isomorphism is clear (see as well Remark 4.13). The purpose of the grading shift introduced in the previous definition is to ensure compatibility of gradings in Proposition 4.15. In order to keep track of this overall shift, define

$$s(\Gamma) = \sum_{\substack{v \text{ vertex of} \\ \text{of type } (a; b; a + b)}} \frac{ab}{2}.$$

⁽⁷⁾In other words, we choose an ambient isotopy of the square such that the images of the trivalent vertices of Γ have distinct y -coordinate.

4.3. A 2-FUNCTOR. — The relationship between Soergel bimodules and foams has already been investigated, see for instance [RW16, Wed19]. However, we develop in this section a foam interpretation of Soergel bimodules themselves and not only of morphisms between them.

DEFINITION 4.6. — Let Γ be a braid-like \underline{k}_1 -MOY graph- \underline{k}_0 . We set $F_{\Gamma}^D(\Gamma)$ to be the free R_N -module generated by the set of rooted Γ -foams modded out by \sim -equivalence. If F is in $2\text{-Hom}_{\text{DLF}_k}(\Gamma_0; \Gamma_1)$, we denote by $F_{\Gamma}^D(F)$ the map $F_{\Gamma}^D(\Gamma_0) \rightarrow F_{\Gamma}^D(\Gamma_1)$ induced by F . It is a map of graded R_N -modules. Given two objects \underline{k}_0 and \underline{k}_1 of DLF_k , this defines a functor F_{Γ}^D from $1\text{-Hom}_{\text{DLF}_k}(\underline{k}_0; \underline{k}_1)$ to the category of graded R_N -modules.

LEMMA 4.7. — Let $\underline{k} = (k_1; \dots; k_r)$ be a finite sequence of positive integers. The space $F_{\Gamma}^D(\underline{k}; I)$ has a natural structure of R_N -algebra. As an algebra it is isomorphic to $A_{\underline{k}}$.

Proof. — The algebra structure is induced by concatenation of disk-like $(\underline{k}; I)$ -foams along the standard tree. The unit is the standard tree times the interval, with each facets trivially decorated. Note, that there is an isomorphism of R_N -algebras:

$$A_{\underline{k}} \cong \bigotimes_{i=1}^r R_N[x_1; \dots; x_{k_i}]^{S_{k_i}}.$$

It is convenient to use this description of $A_{\underline{k}}$ to define the isomorphism between $A_{\underline{k}}$ and $F_{\Gamma}^D(\underline{k}; I)$. For $\mathbf{P} = P_1 \otimes \dots \otimes P_r$ a pure tensor in $A_{\underline{k}}$, we define $\mathbf{P}(\Gamma)$ to be the (\sim) -equivalence class of the standard tree times the interval with decorations $P_1; \dots; P_r$ on its leaves and trivial decorations on the other facets. This is clearly an algebra morphism and it is surjective thanks to Lemma 3.30. We now focus on injectivity. Both $F_{\Gamma}^D(\underline{k}; I)$ and $A_{\underline{k}}$ have natural structures of A_k -modules. For $A_{\underline{k}}$ this comes from the injection of A_k in $A_{\underline{k}}$. For $F_{\Gamma}^D(\underline{k}; I)$, this comes by decorating the “root” facet, that is, the facet which bounds the edge $\underline{k}; I$ (which it self is in s_b). The map $\mathbf{P}(\Gamma)$ respects these structures of A_k -modules because the dots migration identity (3.3) is part of the \sim -equivalence (see Remark 3.20). Thanks to identity (3.4) (which is as well compatible with the \sim -equivalence) used $(k-1)$ times, we know that $F_{\Gamma}^D(\underline{k}; I)$ is free of rank $[k_1 \ k_2 \ \dots \ k_r]$. The algebra $A_{\underline{k}}$ is as well a free A_k -module of rank $[k_1 \ k_2 \ \dots \ k_r]$. This is enough to conclude that $\mathbf{P}(\Gamma)$ is indeed an isomorphism.

COROLLARY 4.8. — Let Γ be a braid-like \underline{k}_1 -MOY graph- \underline{k}_0 . The space $F_{\Gamma}^D(\Gamma)$ has a natural structure of $A_{\underline{k}_1}$ -module- $A_{\underline{k}_0}$. Let us define r_s and r_m the two Laurent polynomials by the formulas:

$$r_s(\Gamma) := \prod_{\substack{v \in 2V(\Gamma) \text{ split} \\ v \text{ of type } (a; b; a+b)}} \begin{bmatrix} a+b \\ a \end{bmatrix} \quad \text{and} \quad r_m(\Gamma) := \prod_{\substack{v \in 2V(\Gamma) \text{ merge} \\ v \text{ of type } (a; b; a+b)}} \begin{bmatrix} a+b \\ a \end{bmatrix}.$$

The space $F_{\Gamma}^D(\Gamma)$ is a free $A_{\underline{k}_1}$ -module of graded rank $r_m(\Gamma) q^{-s(\Gamma)}$ and a free module- $A_{\underline{k}_0}$ of graded rank $r_s(\Gamma) q^{-s(\Gamma)}$.

Proof. — The algebras $A_{\underline{k}_0}$ and $A_{\underline{k}_1}$ are isomorphic to $F_{\mathbb{Z}}^D(\underline{k}_0 - 1)$ and $F_{\mathbb{Z}}^D(\underline{k}_1 - 1)$ and the action of $A_{\underline{k}_0}$ and $A_{\underline{k}_1}$ are given by concatenating disk-like $(\underline{k}_0 - 1)$ -foam and $(\underline{k}_1 - 1)$ along s_r and s_c . The statement about the freeness and the rank follows directly from Lemma 2.12, Remark 2.13 and the fact that the identities (3.1) and (3.4) holds in the \mathbb{Z} -equivalence setting (see Remark 3.20.)

REMARK 4.9


(1) Note that this corollary implies, that for any braid-like \underline{k}_1 -MOY graph \underline{k}_0 , we have:

$$r_s(\) \dim_q^{\mathbb{O}} A_{\underline{k}_1} = r_m(\) \dim_q^{\mathbb{O}} A_{\underline{k}_0} :$$

where $\dim_q^{\mathbb{O}} W$ ($2 \mathbb{Z}[[q]]$) denotes the graded dimension of a W as a graded \mathbb{O} -vector space provided each graded piece is finite-dimensional.

(2) Recall that $F_{\mathbb{Z}}^D(\underline{k}_0 - 1)$ and $F_{\mathbb{Z}}^D(\underline{k}_1 - 1)$ are spanned by decorated versions of the interval times the standard \underline{k}_0 -tree and the standard \underline{k}_1 -tree respectively. This implies that the action of $A_{\underline{k}_0}$ and $A_{\underline{k}_1}$ on $F_{\mathbb{Z}}^D(\)$ can be thought of as multiplying decoration on facets adjacent to the standard \underline{k}_0 -tree (contained in the square s_c) and on the standard \underline{k}_1 -tree (contained in the square s_r) respectively.

COROLLARY 4.10. — *Let us consider \underline{m} the braid-like \underline{k}_1 -MOY graph \underline{k}_0 and \underline{s} the braid-like \underline{k}_0 -MOY graph \underline{k}_1 given by*



and

Then $F_{\mathbb{Z}}^D(\underline{s})$ isomorphic to $A_{\underline{k}_0}$ as a $A_{\underline{k}_1}$ -module- $A_{\underline{k}_0}$ and $F_{\mathbb{Z}}^D(\underline{m})$ isomorphic to $A_{\underline{k}_0}$ as a graded $A_{\underline{k}_0}$ -module- $A_{\underline{k}_1}$. This makes sense, since $A_{\underline{k}_1}$ is a sub-algebra of $A_{\underline{k}_0}$.

Proof. — We only prove the statement for \underline{m} (the proof for \underline{s} is similar). Denote ℓ the length of \underline{k}_0 and let us fix F a rooted tree-like \underline{m} -foam. We want to define a map from $A_{\underline{k}_0}$ to $F_{\mathbb{Z}}^D(\underline{m})$. An element of $A_{\underline{k}_0}$ is a finite sum of elements of the form $P := \prod_{i=1}^{\ell} P_i$ where P_i is a symmetric polynomial on k_0^i variables. Define $\psi(P)$ to be the foam F with decorations of that facets adjacent to the leaves of the \underline{k}_0 -tree given by the P 's. Pictorially, for $\underline{k}_0 = (a; b; c)$ and $\underline{k}_1 = (a + b; c)$, this looks like Figure 10.

This map is clearly a bimodule map and it has degree $\ell - s(\underline{m})$. Thanks to Lemma 3.30, the map is surjective. We conclude with Corollary 4.8 which implies that up to an overall grading shift of $\ell - s(\underline{m})$, $F_{\mathbb{Z}}^D(\underline{m})$ and $A_{\underline{k}_0}$ have the same graded dimension.




FIGURE 10.

LEMMA 4.11. — Let b and t be two braid-like \underline{k}_1 -MOY graphs- \underline{k}_0 and F be an element of $2\text{-hom}_{\text{DLF}}(b; t)$. The linear map $F_7^D(F)$ is a map of $A_{\underline{k}_1}$ -modules- $A_{\underline{k}_0}$.

Proof. — The structures of $A_{\underline{k}_1}$ -modules- $A_{\underline{k}_0}$ of $F_7^D(b)$ and $F_7^D(t)$ is given by multiplying the decoration of facets of foams adjacent to the \underline{k}_0 -tree (contained in the square s_b) and the \underline{k}_1 -tree (contained in the square s_t). Recall that the boundary foam F in the square s_b (resp. in the square s_t) is \underline{k}_0 -1 (resp. \underline{k}_1 -1). Hence stacking F over a rooted b -foam does not change the nature of the boundary in the squares s_b and s_t . In particular, acting with $A_{\underline{k}_1}$ or $A_{\underline{k}_0}$ before or after stacking F produces the same foam. Hence the actions of $A_{\underline{k}_1}$ and $A_{\underline{k}_0}$ commute with $F_7^D(F)$.

LEMMA 4.12. — Let α be a braid-like \underline{k}_1 -MOY graph- \underline{k}_0 and β be a \underline{k}_2 -MOY graph- \underline{k}_1 . Then we have the following isomorphism of $A_{\underline{k}_2}$ -module- $A_{\underline{k}_0}$:

$$F_7^D(\alpha \underline{k}_1 \beta) \simeq F_7^D(\alpha) \otimes_{A_{\underline{k}_1}} F_7^D(\beta):$$

Proof. — We have an $A_{\underline{k}_1}$ -bilinear morphism of $A_{\underline{k}_2}$ -module- $A_{\underline{k}_0}$ from $F_7^D(\alpha) \otimes_{A_{\underline{k}_1}} F_7^D(\beta)$ (where $1^k = (1; \dots; 1)$) to $F_7^D(\alpha \underline{k}_1 \beta)$ given by concatenating foams along the standard tree for \underline{k}_1 . This induce a map $\phi: F_7^D(\alpha) \otimes_{A_{\underline{k}_1}} F_7^D(\beta) \rightarrow F_7^D(\alpha \underline{k}_1 \beta)$. We now prove that ϕ is bijective.

The surjectivity is easy. Indeed, thanks to Lemmas 3.30 and 3.32 every element of $F_7^D(\alpha \underline{k}_1 \beta)$ is a linear combination of decorated tree-like foams F_i and we can choose the shape of these tree-like foams. Hence we can suppose that at the locus where α and β are glued together, the intersection of the foams F_i with a vertical plane are equal to the standard trees. Hence every F_i is in the image of ϕ and their sum as well.

To conclude, we argue with graded \mathbb{Q} -dimensions since every graded piece is finite-dimensional. We have:

$$\begin{aligned} \dim_{\mathbb{Q}}^{\circ} (F_{\mathbb{Z}}^D(\underline{k}_1) \otimes_{A(\underline{k})} F_{\mathbb{Z}}^D(\underline{k}_2)) &= r_m(\underline{k}_1) r_s(\underline{k}_2) \dim_{\mathbb{Q}}^{\circ} A_{\underline{k}_1} q^{s(\underline{k}_1)} q^{-s(\underline{k}_2)} \\ &= r_m(\underline{k}_1) r_m(\underline{k}_2) \dim_{\mathbb{Q}}^{\circ} A_{\underline{k}_2} q^{s(\underline{k}_1) - s(\underline{k}_2)} \\ &= r_m(\underline{k}_1 + \underline{k}_2) \dim_{\mathbb{Q}}^{\circ} A_{\underline{k}_2} q^{s(\underline{k}_1) - s(\underline{k}_2)} \\ &= \dim_{\mathbb{Q}}^{\circ} (F_{\mathbb{Z}}^D(\underline{k}_1 + \underline{k}_2)) : \end{aligned}$$

REMARK 4.13. — Note that from this lemma, we can re-obtain the fact that the bimodule $B(\underline{k})$ is well-defined. See Remark 4.5.

DEFINITION 4.14. — For every k , we denote by BimS_k the 2-category of singular Soergel bimodules of level k . More precisely:

- (1) The objects of BimS_k are finite sequences of positive integers which sum up to k .
- (2) The category of 1-morphisms from \underline{k}_0 to \underline{k}_1 is the smallest abelian full subcategory of $A_{\underline{k}_1}$ -module- $A_{\underline{k}_0}$ containing the $A_{\underline{k}_1}$ -module- $A_{\underline{k}_0}$ $B(\underline{k})$ for any braid-like MOY graph \underline{k} . Note that thanks to Corollary 4.8, all objects of this category are projective (and therefore free) as $A_{\underline{k}_1}$ -modules and as modules- $A_{\underline{k}_0}$, and are finitely generated for both of these structures.

From Lemmas 4.7 and 4.12 and Corollaries 4.8 and 4.10, we deduce the following proposition:

PROPOSITION 4.15. — We have a 2-functor

$$\begin{aligned} F_{\mathbb{Z}}^D : \text{DLF}_k &\rightarrow \text{BimS}_k \\ \underline{k} &\mapsto \underline{k} \\ \mathbb{Z} &\mapsto F_{\mathbb{Z}}^D(\underline{k}) \\ F &\mapsto F_{\mathbb{Z}}^D(F) : \end{aligned}$$

which factorizes through the 2-category $\overline{\text{DLF}}_k$.

Actually, based on evidence given by Stošić [Sto08], we conjecture the following:

CONJECTURE 4.16. — The 2-functor $F_{\mathbb{Z}}^D$ induces an equivalence of 2-categories between $\overline{\text{DLF}}_k$ and BimS_k .

4.4. HOCHSCHILD HOMOLOGY. — If A is an algebra and M an A -module- A , the Hochschild homology of A with coefficients in M is denoted by $\text{HH}(A; M)$.

LEMMA 4.17. — Let \underline{k} be a braid-like \underline{k}_1 -MOY graph- \underline{k}_0 and \underline{k}^0 be a braid-like \underline{k}_0 -MOY graph- \underline{k}_1 , then $\text{HH}(A_{\underline{k}_0}; B(\underline{k}^0, \underline{k}_1))$ and $\text{HH}(A_{\underline{k}_1}; B(\underline{k}, \underline{k}_0^0))$ are canonically isomorphic.

Proof. — This follows from Corollary 4.8 and Lemma 4.12. Let us write $M = F_1^D(\cdot)$ and $M^0 = F_1^D(\cdot^0)$. Let $C(M)$ be a projective resolution of M as A_{k_1} -module- A_{k_0} . Since M^0 is projective as module- A_{k_1} , $C(M) \otimes_{A_{k_0}} M^0$ is a projective resolution of $M \otimes_{A_{k_0}} M^0$ as A_{k_1} -module- A_{k_1} . Similarly, $M^0 \otimes_{A_{k_1}} C(M)$ is a projective resolution of $M^0 \otimes_{A_{k_1}} M$ as A_{k_0} -module- A_{k_0} . We have a canonical isomorphism of chain complexes

$$A_{k_0} \otimes_{A_{k_0}}^{en} (M^0 \otimes_{A_{k_1}} C(M)) \simeq A_{k_1} \otimes_{A_{k_1}}^{en} (C(M) \otimes_{A_{k_0}} M^0).$$

The result follows because $HH(A_{k_0}; B(\cdot^0_{k_1}))$ and $HH(A_{k_1}; B(\cdot_{k_0}^0))$ are the homology groups of these two chain complexes.

PROPOSITION 4.18. — *Let \underline{k} be a braid-like k -MOY graph- k and denote by $\widehat{\cdot}$ its closure and by $F_1^T(\widehat{\cdot})$ the space of vinyl $\widehat{\cdot}$ -foams modulo 1-equivalence. The space $HH_0(A_{\underline{k}}; F_1^D(\cdot))$ is canonically isomorphic to $F_1^T(\widehat{\cdot})$.*

Proof. — Closing up rooted \cdot -foams into vinyl $\widehat{\cdot}$ -foams provides a well-defined map $\cdot : F_1^D(\cdot) \rightarrow F_1^T(\widehat{\cdot})$. The action of $A_{\underline{k}}$ can be seen as concatenating foams (see proof of Corollary 4.10). Hence, \cdot factors through $F_1^D(\cdot) = [A_{\underline{k}}; F_1^D(\cdot)]$ which is equal to $HH_0(A_{\underline{k}}; F_1^D(\cdot))$. We now denote by \cdot the induced map from $HH_0(A_{\underline{k}}; F_1^D(\cdot))$ to $F_1^T(\widehat{\cdot})$. This map is surjective thanks to Lemma 3.30. Instead of proving that \cdot is injective, we prove that spaces $HH_0(A_{\underline{k}}; F_1^D(\cdot))$ and $F_1^T(\widehat{\cdot})$ have the same graded dimension (and each of their homogeneous parts is finite-dimensional). Thanks to Lemma 4.17, $HH_0(A_{\underline{k}}; F_1^D(\cdot))$ only depends on $\widehat{\cdot}$. Hochschild homology is compatible with direct sum of bimodules in the sense that:

$$HH(A; M \oplus N) \simeq HH(A; M) \oplus HH(A; N):$$

Hence, thanks to Proposition 2.19, it is enough to prove the statement for $\widehat{\cdot}$ a collection of circles. If $\widehat{\cdot}$ is a collection of circles labeled by $\underline{k} := (k_1; \dots; k_r)$, then we have $F_1^D(\cdot) = A_{\underline{k}}$. Lemma B.1 implies that the space $HH_0(A_{\underline{k}}; F_1^D(\cdot))$ is isomorphic to $A_{\underline{k}}$. On the other hand, $F_1^T(\widehat{\cdot})$ is isomorphic to $A_{\underline{k}}$ as well thanks to Lemma 4.2.

5. ONE QUOTIENT AND TWO APPROACHES

5.1. A FOAMY APPROACH

5.1.1. Evaluation of vinyl foams

NOTATION 5.1. — The set of Young diagrams with at most a rows and at most b columns is denoted by $T(a; b)$ and the set of Young diagrams with at most a rows is denoted by $T(a; 1)$. The rectangular Young diagram with a rows and b columns is denoted by $(a; b)$.

NOTATION 5.2. — Recall that R_N denotes the ring of symmetric polynomials with coefficients in \mathbb{Q} .

(1) Denote the graded algebra $R_N[x_1, \dots, x_k]^{\mathbb{S}_k}$ by A_k , by $J_{N;k}$ the ideal of $R_N[x_1, \dots, x_k]$ generated by

$$\left\{ \prod_{i=1}^N (x_j - T_i) \mid j = 1, \dots, k \right\};$$

Note that elements of this set are indeed symmetric in the T . Denote by $M_{N;k}$ the R_N -algebra

$$A_k / (J_{N;k} \setminus A_k);$$

seen as an R_N -module. The indeterminates x have degree 2, just like the indeterminates T appearing in the definition of R_N (end of Section 3.4).

(2) If $\lambda = (\lambda_1, \dots, \lambda_k)$ is a Young diagram with at most k rows, define $\mathbf{x}^\lambda := \prod_{i=1}^k x_i^{\lambda_i}$. Denote by $m^\lambda(x_1, \dots, x_k)$ the symmetric polynomial $\sum_{\theta \in \text{distinct}} \mathbf{x}^{\theta \circ \lambda}$, where θ runs over all *distinct* permutations of λ . Denote by $\tilde{m}^\lambda(x_1, \dots, x_k)$ the symmetric polynomial $\sum_{\theta \in \text{all}} \mathbf{x}^{\theta \circ \lambda}$, where θ runs over *all* permutations of λ . The family $(m^\lambda)_{\lambda \in \mathcal{Y}(k;1)}$ is a \mathbb{Z} -basis of the ring of symmetric polynomials in k variables with coefficients in \mathbb{Z} (see [Mac15]). The family $(\tilde{m}^\lambda)_{\lambda \in \mathcal{Y}(k;1)}$ is a \mathbb{Q} -basis of the ring of symmetric polynomials in k variables with coefficients in \mathbb{Q} .

LEMMA 5.3. — *The R_N -module $M_{N;k}$ is free and has a basis given by images in $M_{N;k}$ of*

$$(m^\lambda(x_1, \dots, x_k))_{\lambda \in \mathcal{Y}(k;N-1)}$$

seen as element of A_k .

Proof. — The R_N -module $M_{N;k}$ is isomorphic to

$$(R_N[x_1, \dots, x_k] = J_{N;k})^{\mathbb{S}_k};$$

Indeed, the R_N -linear maps

$$A_k \xrightarrow{\pi} R_N[x_1, \dots, x_k] \twoheadrightarrow R_N[x_1, \dots, x_k] = J_{N;k} \twoheadrightarrow (R_N[x_1, \dots, x_k] = J_{N;k})^{\mathbb{S}_k}$$

is surjective because if $P + J_{N;k} \in (R_N[x_1, \dots, x_k] = J_{N;k})^{\mathbb{S}_k}$, one can assume that P is itself \mathbb{S}_k -invariant since the ideal $J_{N;k}$ is \mathbb{S}_k -invariant. The kernel of this morphism is $J_{N;k} \setminus A_k$, hence it induces an isomorphism of R_N -modules between $M_{N;k}$ and $(R_N[x_1, \dots, x_k] = J_{N;k})^{\mathbb{S}_k}$.

The R_N -module $(R_N[x_1, \dots, x_k] = J_{N;k})^{\mathbb{S}_k}$ is isomorphic to

$$\text{Sym}^k \left(R_N[x] / \left(\prod_{i=1}^N (x - T_i) \right) \right);$$

The R_N -module $R_N[x] / (\prod_{i=1}^N (x - T_i))$ is a free R_N module of rank N and has a natural R_N -basis given by $(x^i)_{0 \leq i \leq N-1}$. Therefore, $(m^\lambda(x_1, \dots, x_k))_{\lambda \in \mathcal{Y}(k;N-1)}$ is a R_N -basis of $(R_N[x_1, \dots, x_k] = J_{N;k})^{\mathbb{S}_k}$.

Denote $\mu_{N;k}$ the composition of the projection from A_k to $M_{N;k}$ with $\nu_{N;k}$. Let F be a vinyl S_k -foam- S_k . In \mathbb{R}^3 , we can cap and cup F with two disks labeled by k , to obtain a (non-vinyl) foam $\text{cl}(F)$.

DEFINITION 5.6. — The *equivariant symmetric evaluation* of a vinyl S_k -foams- S_k F is given by:

$$\langle\langle F \rangle\rangle_N := \mu_{N;k}(\text{hcl}(F)_{i_k});$$

where h_{i_k} denotes the sl_k -evaluation of closed foams (see Definition 3.14).

REMARK 5.7. — We can make the symmetric evaluation more explicit: let F be a vinyl S_k -foam- S_k , then $\langle\langle F \rangle\rangle_N$ is equal to the coefficient of $m_{(k;N-1)}$ (in the basis (m)) of $\text{hcl}(F)_{i_k}$. Alternatively it is equal to $k!$ times the $\tilde{m}_{(k;N-1)}$ -coefficient in the basis (\tilde{m}) of $\text{hcl}(F)_{i_k}$.

EXAMPLE 5.8. — Using Example 5.4, one can easily compute the evaluation of some simple vinyl foams.

(1) Suppose F is the foam $S_1 \times [0;1]$ decorated by x , then the foam $\text{cl}(F)$ is a sphere labeled by 1 decorated by x and $\text{hcl}(F)_{i_1} = X_1$. Hence

$$\langle\langle F \rangle\rangle_N = \nu_{N;1}(x) = \sum_{\substack{P \\ \sum_{i=1}^N n_i = N+1 \\ n_1, \dots, n_N > 0}} (-1)^{1+\sum_{i=1}^N (i-1)n_i} \frac{(\sum_{i=1}^N n_i)!}{\prod_{i=1}^N n_i!} \prod_{i=1}^N e_i^{n_i}(T_1; \dots; T_N);$$

(2) If F is a foam $S_k \times [0;1]$ decorated by a polynomial \tilde{m} , then

$$\langle\langle F \rangle\rangle_N = (-1)^{k(k+1)=2} \nu_{N;k}(\tilde{m});$$

The basis (\tilde{m}) is convenient because of the following lemma which is the key ingredient for the proof of monoidality of our construction.

LEMMA 5.9. — Let k_1 and k_2 be two non-negative integers, λ (resp. μ) be a Young diagram with at most k_1 (resp. k_2) rows and A be a set of $k_1 + k_2$ variables. We have

$$\sum_{\substack{A_1 \sqcup A_2 = A \\ \# A_1 = k_1 \\ \# A_2 = k_2}} \tilde{m}(A_1)\tilde{m}(A_2) = \tilde{m}(\lambda \cup \mu);$$

where $\lambda \cup \mu$ is Young diagram corresponding to the union of the partition of λ and μ . In particular, for any integer N , $\nu_{N;k} = \nu_{(k_1+k_2;N)}$ if and only if $\nu_{N;k_1} = \nu_{(k_1;N)}$ and $\nu_{N;k_2} = \nu_{(k_2;N)}$.


Lemma 5.9 implies a nice behavior of the evaluation with respect to disjoint union as illustrated in the following example.

EXAMPLE 5.10. — Let k_1 and k_2 be two non-negative integers, λ (resp. μ) be a Young diagram with at most k_1 (resp. k_2) rows. Denote by F (resp. G) the foam $S_{k_1} \times [0;1]$

(resp. $S_{k_2} [0;1]$) decorated by \tilde{m} (resp. \tilde{m}). From Example 5.8, one deduces:

$$\begin{aligned} \langle\langle F \rangle\rangle_N &= (-1)^{k_1(k_1+1)=2} \langle\langle N; k_1(\tilde{m}) \rangle\rangle = (-1)^{k_1(k_1+1)=2} k_1! \prod_{i=1}^{k_1} \langle\langle N; 1(X^i) \rangle\rangle \text{ and} \\ \langle\langle G \rangle\rangle_N &= (-1)^{k_2(k_2+1)=2} \langle\langle N; k_2(\tilde{m}) \rangle\rangle = (-1)^{k_2(k_2+1)=2} k_2! \prod_{i=1}^{k_2} \langle\langle N; 1(X^i) \rangle\rangle. \end{aligned}$$

Set $k = k_1 + k_2$ and consider H the $(S_k; S_k)$ -vinyl foam obtained by pre-composing $F [G$ by $Y_k^{k_1; k_2}$ and post-composing it by $\lambda_{k_1; k_2}^k$ where:




In other words, H is vinyl foam equal to a digon labeled by $k_1; k_2$ and k times S^1 decorated by \tilde{m} and \tilde{m} . The foam H is 1-equivalent to $S_k [0;1]$ decorated by \tilde{m} . Hence, one has:

$$\begin{aligned} \langle\langle H \rangle\rangle_N &= (-1)^{k(k+1)=2} \langle\langle N; k(\tilde{m}) \rangle\rangle \\ &= (-1)^{k(k+1)=2} k! \prod_{i=1}^{k_1} \langle\langle N; 1(X^i) \rangle\rangle \prod_{j=1}^{k_2} \langle\langle N; 1(X^j) \rangle\rangle \\ &= (-1)^{k(k+1)=2} (-1)^{k_1(k_1+1)=2} (-1)^{k_2(k_2+1)=2} \binom{k}{k_1} \langle\langle F \rangle\rangle_N \langle\langle G \rangle\rangle_N \\ &= (-1)^{k_1 k_2} \binom{k}{k_1} \langle\langle F \rangle\rangle_N \langle\langle G \rangle\rangle_N. \end{aligned}$$

The following lemma proves that this evaluation does not really depend on k .

LEMMA 5.11. — *Let F be a vinyl S_k -foam- S_k , then it is 1-equivalent to the foam $S_k [0;1]$ decorated with $\langle\langle (-1)^{k(k+1)=2} \text{cl}(F) \rangle\rangle_k$. The decoration makes sense since $\langle\langle (-1)^{k(k+1)=2} \text{cl}(F) \rangle\rangle_k$ is a symmetric polynomial in k variables.*

Proof. — Let us denote by T the foam



We need to show that T and F are N -equivalent for all N . If $N < k$, this is clear, since all S_k -foams- S_k are N -equivalent to 0. If $N = k$ this is clear as well since

the identity (3.2) has a very simple form in this case: we can apply it on the top and on the bottom of F and T , the result follows immediately. Let us now pick an integer N greater than k . Thanks to the definition of F_N (and its monoidality, see Corollary 3.18), the N -equivalence is equivalent to saying that for any $(G_1; \widehat{G}_2)$ in $\text{Hom}_{\text{Foam}}(S_k; ?) \cong \text{Hom}_{\text{Foam}}(?; S_k)$, we have:

$$h_{G_2} F G_1 i_N = h_{G_2} T G_1 i_N:$$

If $N = k$, the result holds by hypothesis. Since F is vinyl, for any sl_N -coloring c of $G_2 \cup F \cup G_1$, the induced coloring on the two circles which form the boundary of F are the same, hence it induces a coloring c^∂ on $G_2 \cup T \cup G_1$. Since the difference of the Euler characteristics of monochrome and bichrome surfaces as well as the parity of the difference of numbers of positive circles can be computed locally, the quantity

$$\frac{h_{G_2 \cup F \cup G_1; c} i_N}{h_{G_2 \cup T \cup G_1; c^\partial} i_N}$$

only depends on the restrictions of c and c^∂ to F and T . Since F and T are vinyl, there are exactly k pigments appearing in the restrictions of c and c^∂ to F and T . Hence if these pigments are $1; \dots; k$ and if we sum over all colorings c of $G_2 \cup F \cup G_1$, which induce c^∂ on $G_2 \cup T \cup G_1$ we obtain (thanks to the case $k = N$):

$$\sum_{c \text{ induces } c^\partial} \frac{h_{G_2 \cup F \cup G_1; c} i_N}{h_{G_2 \cup T \cup G_1; c^\partial} i_N} = 1:$$

Since permuting the pigments boils down to permuting the variables $x_1; \dots; x_N$ (see [RW17, Lem. 2.16]), we obtain:

$$\sum_{c \text{ coloring of } G_2 \cup F \cup G_1} h_{G_2 \cup F \cup G_1; c} i_N = \sum_{c^\partial \text{ coloring of } G_2 \cup T \cup G_1} h_{G_2 \cup T \cup G_1; c^\partial} i_N:$$

5.1.2. *Universal construction.* — We will use the evaluation defined in Definition 5.6 and a universal construction à la [BHMV95] in order to define a functor $S_{k;N} : \text{TLF}_k \rightarrow \mathcal{C}$, where \mathcal{C} is the category of \mathbb{Z} -graded finitely generated projective R_N -modules.

If Γ is a vinyl graph of level k , we consider the free graded R_N -module generated by $\text{Hom}_{\text{TLF}_k}(S_k; \Gamma) \cong q^{k(N-1)}$. We mod this space out by

$$\bigcap_{G \supseteq \Gamma} \text{Ker} \left(\text{Hom}_{\text{TLF}_k}(S_k; \Gamma) \otimes_{R_N} \langle \langle G \cup F \rangle \rangle_N \right):$$

We define $S_{k;N}(\Gamma)$ to be this quotient. The definition of $S_{k;N}$ on morphisms follows naturally.

5.1.3. *Categorified identities*

PROPOSITION 5.12. — *The R_N -module $S_{k;N}(S_k)$ is isomorphic to $M_{N;k} q^{k(N-1)}$ (see Notation 5.2). In particular, we have*

$$\text{rk}_q^{R_N}(S_{k;N}(S_k)) = \begin{bmatrix} k + N & 1 \\ & k \end{bmatrix}:$$

Proof. — Define $\text{ev} : A_k \rightarrow S_{k;N}(S_k)$ the R_N -linear map which maps any symmetric polynomial in k variables P to the cylinder $S_k \times [0;1]$ decorated by P . Thanks to Lemma 5.11, this map is surjective. By the very definition of the equivariant symmetric evaluation $J_{N;k} \setminus A_k$ is in the kernel of this map. Hence it induces a R_N -linear map $M_{N;k} \rightarrow S_{k;N}(S_k)$ denoted by ev^0 .

The map ev^0 is injective. Indeed, let x be a non-zero element of $M_{N;k}$. Since $M_{N;k}$ is a Frobenius algebra there exists y in $M_{N;k}$ such that $\langle xy \rangle \neq 0$. Let X and Y be two R_N -linear combinations of vinyl S_k -foams- S_k representing $\text{ev}^0(x)$ and $\text{ev}^0(y)$ in $S_{k;N}(S_k)$. By definition, $\langle X, Y \rangle_N = \langle xy \rangle \neq 0$. Hence $\langle \text{ev}^0(x), \text{ev}^0(y) \rangle \neq 0$ and $\text{ev}^0(x) \neq 0$. It follows that ev^0 is an isomorphism. Note that it is homogeneous of degree $(N-1)k$.

From the definition of the evaluation of vinyl foams, we immediately deduce that the identities of Section 3.2 which can be expressed by vinyl foams are still valid. This gives the following proposition.

PROPOSITION 5.13. — *Let k be a non-negative integer and N be a positive integer, then the functor $S_{k;N}$ satisfies the following local relations:*

$$(5.1) \quad S_{k;N} \left(\begin{array}{c} i \quad j \quad k \\ \swarrow \quad \searrow \quad \nearrow \\ \downarrow \\ i+j+k \end{array} \right) = S_{k;N} \left(\begin{array}{c} i \quad j \quad k \\ \swarrow \quad \nearrow \quad \searrow \\ \downarrow \\ i+j+k \end{array} \right)$$

$$(5.2) \quad S_{k;N} \left(\begin{array}{c} m+n \\ \uparrow \quad \downarrow \\ m \quad n \\ \downarrow \quad \uparrow \\ m+n \end{array} \right) = S_{k;N} \left(\begin{array}{c} \uparrow \\ m+n \end{array} \right) \begin{bmatrix} m+n \\ m \end{bmatrix}$$

$$(5.3) \quad S_{k;N} \left(\begin{array}{ccc} m & n+k & n \\ \uparrow & \uparrow & \uparrow \\ n+k & m & m+k \\ \leftarrow & \rightarrow & \leftarrow \\ n & m & m+k \end{array} \right) = \bigoplus_{j=\max(0; m-n)}^m S_{k;N} \left(\begin{array}{ccc} m & n+k & n \\ \uparrow & \uparrow & \uparrow \\ m & j & n+j \\ \leftarrow & \rightarrow & \leftarrow \\ n & m & m+k \end{array} \right) \begin{bmatrix} k \\ j \end{bmatrix}$$


$$(5.4) \quad S_{k;N} \left(\begin{array}{ccc} k & r & r+k \\ \uparrow & \uparrow & \uparrow \\ k & r & r+k \\ \leftarrow & \rightarrow & \leftarrow \\ k+s & r+s & r+s \end{array} \right) = S_{k;N} \left(\begin{array}{ccc} k & r & r+k \\ \uparrow & \uparrow & \uparrow \\ k+s & r+s & r+s \end{array} \right) \begin{bmatrix} r+s \\ s \end{bmatrix}$$

5.1.4. *Monoidality.* — The category TLF_k does not have a notion of disjoint union, hence for a fixed k we cannot have monoidality of the functor $S_{k;N}$. However, if we consider the disjoint union TLF_N of the categories TLF_k for k in N , then we can speak about disjoint union, and this obviously endows TLF_N with a structure of a monoidal category. The empty vinyl graph seen as an object of TLF_0 is the monoidal unit. Note that in this category $\tau_1 \tau_2$ is in general not isomorphic to $\tau_2 \tau_1$.

We consider the functor $S_{N;N}$: it is given by the functor $S_{k;N}$ on TLF_k for all $k \in N$. In order to fix notations, set

$$\sqcup_{k_1, k_2} : \text{TLF}_{k_1} \times \text{TLF}_{k_2} \rightarrow \text{TLF}_{k_1+k_2}$$

which sends pairs of objects (resp. morphisms) onto their (rescaled) disjoint union. This is illustrated on object below:



PROPOSITION 5.14. — Let τ_1 (resp. τ_2) be a vinyl graph of level k_1 (resp. k_2). Suppose that $S_{k_1;N}(\tau_1)$ and $S_{k_2;N}(\tau_2)$ are free R_N -modules, then $S_{k_1+k_2;N}(\sqcup_{k_1, k_2}(\tau_1; \tau_2))$ is isomorphic to $S_{k_1;N}(\tau_1) \otimes_{R_N} S_{k_2;N}(\tau_2)$. In particular, $S_{k_1+k_2;N}(\sqcup_{k_1, k_2}(\tau_1; \tau_2))$ is a free R_N -module.

Proof. — Let us fix two vinyl graphs τ_1 and τ_2 of level k_1 and k_2 . We denote $\sqcup_{k_1, k_2}(\tau_1; \tau_2)$ by τ and k_1+k_2 by k . We will define a map γ_{k_1, k_2}^k from $S_{k_1;N}(\tau_1) \otimes_{R_N} S_{k_2;N}(\tau_2)$ to $S_{k_1+k_2;N}(\tau)$. It is enough to define γ_{k_1, k_2}^k on pure tensors. Let v_1 (resp. v_2) be an element of $S_{k_1;N}(\tau_1)$ (resp. $S_{k_2;N}(\tau_2)$). We can suppose that v_1 is represented by a τ_1 -foam- S_{k_1} F_1 and v_2 by a τ_2 -foam- S_{k_2} F_2 . We define $\gamma_{k_1, k_2}^k(v_1 \otimes v_2)$ to be the element of $S_{k;N}(\tau)$ obtained by re-scaling F_1 and F_2 , taking their disjoint union (this gives an element of $\text{Hom}_{\text{TLF}_k}(\sqcup_{k_1, k_2}(S_{k_1}; S_{k_2}); \tau)$) and pre-composing it with the foam

$$\gamma_k^{k_1, k_2} := \begin{array}{c} \begin{array}{cc} k_1 & k_2 \\ \swarrow & \searrow \\ & \downarrow \\ & k_1 + k_2 \end{array} \end{array} \quad S^1:$$

We extend this definition linearly. We now need to show that:

- (1) this is well-defined,
- (2) this is an isomorphism,

(1) In order to prove that γ_{k_1, k_2}^k is well-defined, we only need to show that if for all vinyl τ_2 -foam- S_{k_2} F_2 (resp. τ_1 -foam- S_{k_1} F_1) and R_N -linear combination of vinyl τ_1 -foams- S_{k_1} $\sum_i a_i F_1^i$ (resp. τ_2 -foams- S_{k_2} $\sum_j b_j F_2^j$) representing 0 in $S_{k_1;N}(\tau_1)$ (resp. in

$S_{k_2;N}(\mathbb{Z})$, we have $\sum_i a_i \langle [F_1^i] [F_2] \rangle = 0$ (resp. $\sum_i a_i \langle [F_1^i] [F_2] \rangle = 0$).
 By symmetry we only prove

$$\sum_i a_i \langle [F_1^i] [F_2] \rangle = 0:$$


The square brackets stand for “the element of the appropriate graded R_N -module represented by this foam”.

Suppose that the element $\sum_i a_i [F_1^i]$ is equal to 0. This means that for any vinyl S_{k_1} -foam- \mathbb{Z} G_1 , $\sum_i a_i \langle G_1 [F_1^i] \rangle_N = 0$. In other words, the coefficient of $\tilde{m}^{(k_1;N-1)}$ (in the base $(\tilde{m}^{\mathbb{Z}})_{\mathcal{ZT}(k_1;N-1)}$) of $\sum_i a_i \langle \text{cl}(G_1 [F_1^i]) \rangle_k$ is equal to 0.

We want to prove that for any vinyl S_k -foam- \mathbb{Z} G ,

$$\sum_i a_i \langle G (F_1^i \# F_2) Y_k^{k_1;k_2} \rangle_N = 0:$$

Thanks to Lemmas 3.38 and 3.40, we might suppose that G is tree-like and that it can be obtained by re-scaling the disjoint union of a tree-like S_{k_1} -foam- \mathbb{Z} G_1 and a S_{k_2} -foam- \mathbb{Z} G_2 composed with $\lambda_{k_1;k_2}^k$ (which is the foam $Y_k^{k_1;k_2}$ turned upside down). Thanks to Lemma 5.11, for all i , $G_1 [F_1^i]$ is 1-equivalent to $S_{k_1} [0;1]$ decorated with $\langle \text{cl}(G_1 [F_1^i]) \rangle_{k_1}$ and $G_2 [F_2]$ is 1-equivalent to $S_{k_2} [0;1]$ decorated with $\langle \text{cl}(G_2 [F_2]) \rangle_{k_2}$. Hence $\sum_i a_i G (F_1^i \# F_2) Y_k^{k_1;k_2}$ is 1-equivalent to



Following Example 5.10, one deduces that for all i :

$$\langle G (F_1^i \# F_2) Y_k^{k_1;k_2} \rangle_N = (-1)^{k_1 k_2} \binom{k}{k_1} \langle G_1 [F_1^i] \rangle_N \langle G_2 [F_2] \rangle_N;$$

and therefore

$$\sum_i a_i \langle G (F_1^i \# F_2) Y_k^{k_1;k_2} \rangle_N = (-1)^{k_1 k_2} \binom{k}{k_1} \langle G_2 [F_2] \rangle_N \sum_i a_i \langle G_1 [F_1^i] \rangle_N = 0:$$

(2) Let F_1 (resp. F_2) be a trivially decorated tree-like \mathbb{Z} -foam- S_{k_1} (resp. \mathbb{Z} -foam- S_{k_2}). The foam F obtained by pre-composing $F_1 \# F_2$ with the foam $Y_k^{k_1;k_2}$ described in the construction of $\sum_i a_i$ is in the image of $\sum_i a_i$ as well as its decorated version with non-trivial decoration on leaves. Thanks to Lemma 3.38 and 3.40 the R_N -vector space $S_{k;N}(\mathbb{Z})$ is spanned by elements represented by foams of this type. This proves the surjectivity.

To prove injectivity, we first pick bases $(B_i^1)_{i \in I}$ and $(B_j^2)_{j \in J}$ of $S_{k_1;N}(\mathbb{Z})$ and $S_{k_2;N}(\mathbb{Z})$ and their dual bases $(B_i^1)_{i \in I}$ and $(B_j^2)_{j \in J}$. All these elements can be

represented by (R_N -linear combinations of) tree-like foams. Denote these (R_N -linear combinations of) tree-like foams by F_i^1, F_j^2, F_i^{-1} and F_j^{-2} . Suppose that

$$\left[\sum_{\substack{i \in I \\ j \in J}} a_{ij} (F_i^1 \mp F_j^2) \gamma_k^{k_1, k_2} \right] = 0:$$

Let us fix an i_0 in I and a j_0 in J . The hypothesis implies that

$$\sum_{\substack{i \in I \\ j \in J}} a_{ij} \left\langle \lambda_{k_1, k_2}^k (B_{i_0}^1 \mp B_{j_0}^2) (F_i^1 \mp F_j^2) \gamma_k^{k_1, k_2} \right\rangle_N = 0:$$

In the previous expression, $\langle \rangle_N$ has been R_N -linearly extended to makes sense on R_N -linear combinations of foams. Thanks to Lemma 5.9, the $\tilde{m}_{(k; N-1)}$ -coefficient in the base $(\tilde{m})_{2T(k; 1)}$ of

$$\sum_{\substack{i \in I \\ j \in J}} a_{ij} \left\langle \text{cl}(\lambda_{k_1, k_2}^k (B_{i_0}^1 \mp B_{j_0}^2) (F_i^1 \mp F_j^2) \gamma_k^{k_1, k_2}) \right\rangle_k$$

is equal to $a_{i_0 j_0}$. However, this coefficient should be equal to 0. This prove injectivity.


THEOREM 5.15. — *The functor $S_{N;N}$ is monoidal.*

Proof. — First of all, since the only vinyl foam of level 0 is the empty set, it is clear that $S_{N;N}(\mathbf{1}_{\text{TLF}_N}) = R_N = \mathbf{1}_C$ ($\mathbf{1}$ denotes the unital object of monoidal categories). We need to construct a natural isomorphism $(\cdot; \cdot)_{1; 2 \text{ob}(\text{TLF}_N)}$ from $S_{N;N}(\cdot)_{R_N}$ to $S_{N;N}(\bigsqcup(\cdot; \cdot))$.

This isomorphism is provided by (the proof of) Proposition 5.14. In order to use it, we only need to show that for any vinyl graph Γ , $S_{N;N}(\mathbf{1}_{\text{TLF}_N})$ is a free R_N -module.

If Γ is a collection of circles we can argue by induction on the number of circles. If Γ consists of only one circle, we can use Proposition 5.12. If it consists of more than one circle, we can use the induction hypothesis and Proposition 5.14. We deduce the general case from the case of collection of circles thanks to Proposition 2.19 and Proposition 5.13.

REMARK 5.16. — Formally we should have checked the compatibility of associators. Since we did not write down them explicitly, we cannot be very precise here. However, this compatibility trivially holds because the foams



are 1-equivalent, (because they are both tree-like with only trivial decorations (see Lemma 3.40)).

We can now prove that the functor $S_{N;N}$ categorifies the symmetric MOY calculus.

THEOREM 5.17. — *The functor $S_{N;N} : \text{TLF}_N \rightarrow \mathcal{C}$ satisfies for every vinyl graph Γ ,*

$$\text{rk}_q^{R_N}(S_{N;N}(\Gamma)) = \langle \langle \Gamma \rangle \rangle_N :$$

Proof. — We have already seen in the proof of Theorem 5.15, that the R_N -module $S_{N;N}(\Gamma)$ is free for every vinyl graph Γ . Thanks to Theorem 5.15 and Propositions 5.12 and 5.13 we obtain that the function $\text{rk}_q^{R_N}(S_{N;N}(\Gamma))$ satisfies identities (2.12), (2.13), (2.14), (2.18) and (2.19). We conclude by Theorem 2.18.

5.2. AN ALGEBRAIC APPROACH

5.2.1. Hochschild and Koszul homologies. — Koszul homology has been formalized in [BLS18]. If R is a unital commutative ring, A an R -algebra and M a A -module- A , it associates with the pair $(A; M)$ a sequence $\mathcal{KH}(A; M)$ of R -modules in a functorial way. If A is Koszul (and this will be our case), then $\mathcal{KH}(A; M) = \mathcal{HH}(A; M)$. We do not aim to discuss Koszul homology in details, we refer to [BLS18] and references therein for a nice presentation. We will use Koszul homology instead of Hochschild homology because it enables to have more structure: in fact an extra differential.

NOTATION 5.18. — In what follows, $\underline{k} = (k_1; \dots; k_r)$ is a finite sequence of positive integers of level k , $A_{\underline{k}}$ is the polynomial algebra $R_N[x_1; \dots; x_k]^{\mathbb{S}_{\underline{k}}}$ and A_k denotes $R_N[x_1; \dots; x_k]^{\mathbb{S}_k}$. Note that $A_{\underline{k}}$ is a polynomial algebra over R_N . For i in $\{1; \dots; r\}$ and j in $\{1; \dots; k_i\}$, we set $e_j^{(i)}$ to be the j th elementary symmetric polynomial in variables $x_{r_i+1}; \dots; x_{r_i+k_i}$, where $r_i = \sum_{t=1}^{i-1} k_t$. It is standard that we have (see for example [Lan02, Chap. IV, §6]):

$$A_{\underline{k}} = R_N[e_1^{(1)}; \dots; e_{k_1}^{(1)}; e_1^{(2)}; \dots; e_{k_2}^{(2)}; \dots; e_1^{(r)}; \dots; e_{k_r}^{(r)}] \quad A_{1^k} :$$

where $A_{1^k} := A_{(1; \dots; 1)} = R_N[x_1; \dots; x_k]$.

DEFINITION 5.19. — The *Koszul resolution* of $A_{\underline{k}}$ is the complex

$$C(A_{\underline{k}}) := \bigotimes_{i=1}^r \bigotimes_{j=1}^{k_i} \left(R_N[e_j^{(i)}] \rightarrow R_N[e_j^{(i)}] q^{2j} \binom{e_j^{(i)} - 1}{j} \rightarrow R_N[e_j^{(i)}] \rightarrow R_N[e_j^{(i)}] \right) :$$

The homological degree of $C(A_{\underline{k}})$ is called the *H-degree*.

It is convenient to think of this complex in this way: let $V_{\underline{k}}$ be the R_N -module generated by $(e_j^{(i)})_{\substack{i=1; \dots; r \\ j=1; \dots; k_i}}$ (with $(e_j^{(i)})$ having degree $2j$). Then $C(A_{\underline{k}}) = A_{\underline{k}} \otimes V_{\underline{k}} \otimes A_{\underline{k}}$ with the differential:

$$d^k : C(A_{\underline{k}}) \rightarrow C(A_{\underline{k}})$$

$$a \otimes v_1 \wedge \dots \wedge v_r \otimes b \mapsto \sum_{i=1}^r (-1)^{i+1} (a v_i \otimes v_1 \wedge \dots \wedge \widehat{v}_i \wedge \dots \wedge v_r \otimes b - a \otimes v_1 \wedge \dots \wedge \widehat{v}_i \wedge \dots \wedge v_r \otimes v_i b) :$$

It is standard that it is a projective resolution of $A_{\underline{k}}$ as $A_{\underline{k}}$ -module- $A_{\underline{k}}$. Hence, for any $A_{\underline{k}}$ -module- $A_{\underline{k}}$ M , we have:

$$HH(A_{\underline{k}}; M) \cong H(C(A_{\underline{k}}) \otimes_{A_{\underline{k}}} M) =: KH(A_{\underline{k}}; M):$$

From now on, when speaking about Hochschild homology, we mean Hochschild homology computed in this way. Of course this precision is irrelevant when we only look at the homology groups. But we will shortly introduce an extra differential d_N on $C(A_{\underline{k}})$. It will equip the Hochschild homology with a structure of chain complex. As far as we understand, the differential d_N can be thought as an equivariant version of the extra differential introduced by Cautis in [Cau17]. Some proofs are postponed to Appendix B.

5.2.2. *An extra differential.* — Let N be a positive integer.

NOTATION 5.20. — We denote by D_N the following derivation on A_{1^k} :

$$D_{1^k}^N: A_{1^k} \rightarrow A_{1^k}$$

$$P(x_1; \dots; x_k) \mapsto \sum_{i=1}^k \prod_{j=1}^N (x_i - T_j) @_{x_i} P(x_1; \dots; x_k):$$

We denote by $d_{\underline{k}}^N$ the following map $A_{\underline{k}}$ -linear- $A_{\underline{k}}$ map on $C(A_{\underline{k}})$:

$$d_{\underline{k}}^N: C(A_{\underline{k}}) \rightarrow C(A_{\underline{k}})$$

$$1 \otimes v_1 \wedge \dots \wedge v_{k-1} \mapsto \sum_{i=1}^k (-1)^{i+1} D_k^N(v_i) \otimes v_1 \wedge \dots \wedge \widehat{v}_i \wedge \dots \wedge v_k: 1:$$

note that this is H -homogeneous of degree -1 and q -homogeneous of degree $2(N-1)$. When the context is clear we will drop the subscript \underline{k} .

LEMMA 5.21. — For any finite sequence of positive integers \underline{k} , the map $d_{\underline{k}}^N$ anti-commutes with $d^{\underline{k}}$ and is a differential on $C(A_{\underline{k}})$.

Since we have this extra structure, we need a refined version of Lemma 4.17. The following lemma should be compared to [Cau17, Lem. 6.2].

LEMMA 5.22. — Let $\widehat{\Gamma}$ be a braid-like \underline{k}_1 -MOY graph- \underline{k}_0 and Γ be a braid-like \underline{k}_0 -MOY graph- \underline{k}_1 , then the complexes

$$(HH(A_{\underline{k}_0}; F_{\Gamma}^D(\widehat{\Gamma}_{\underline{k}_1})); d_{\underline{k}_0}^N) \text{ and } (HH(A_{\underline{k}_1}; F_{\Gamma}^D(\Gamma_{\underline{k}_0})); d_{\underline{k}_1}^N)$$

are isomorphic.

This is proved in Appendix B where we restrict to the special case where $\widehat{\Gamma}$ contains only one vertex. Note that this is actually enough to conclude in general. We define an explicit homotopy equivalence $\widehat{\cdot}$ between two complexes computing the Hochschild homology. Finally we prove that $\widehat{\cdot} \circ d_{\underline{k}_0}^N \circ d_{\underline{k}_1}^N \circ \widehat{\cdot}$ is null-homotopic. Note that the previous lemma shows that the complex $(HH(A_{\underline{k}}; F_{\Gamma}^D(\widehat{\Gamma})); d^N)$ only depends on the vinyl graph $\widehat{\Gamma}$.

In order to make the differential d_N of q -degree 0, we shift $\mathbb{H}_i(A_{\underline{k}}; M)$ by $2i(N-1)$ in q -degree. Note that this adjustment does not change $\mathbb{H}_0(A_{\underline{k}}; M)$. We denote this normalization by $\mathbb{H}^N(A_{\underline{k}}; M)$.

The following proposition should be compared to [Wed19, Lem. 3.23].

PROPOSITION 5.23. — *Let Γ be a vinyl graph. The homology of $(\mathbb{H}^N(A_{\underline{k}}; F_{\Gamma}^D(\cdot)); d^N)$ (denoted by $\mathbb{H}^N(\Gamma)$) is concentrated in H -degree 0.*

Proof. — Thanks to Que-élec-Rose's algorithm (see Theorem 2.18), it is enough to show the statement when Γ is a collection of circles. Suppose that Γ is a collection of circles labeled by \underline{k} . The result follows from the regularity of the sequence $(D^N(e_j^{(i)}))_{1 \leq i \leq N}$ seen as polynomials in x_1, \dots, x_k with coefficients in R_N . The regularity of this sequence follows from that of the sequences $(D^N(e_j^{(i)}))_{1 \leq j \leq k_i}$ for every i because they involve different sets of variables. Hence, from now on, we can suppose that $N = 1$ and we write $D^N(e_j)$ instead of $D^N(e_j^{(1)})$. The regularity of a sequence is equivalent to the fact that the set Z of common zeroes of these polynomials is 0-dimensional. Note that $fT_1, \dots, T_N g^k$ is a subset of Z . One shall see that $Z = fT_1, \dots, T_N g^k$.

Let us consider the ideal I of $R_N[x_1, \dots, x_k]^{S_k}$ generated by the polynomials

$$P_r = \sum_{i=1}^k \prod_{j=1}^N (x_i - T_j) x_i^r \quad \text{for } 0 \leq r \leq k-1.$$

Note that I is equal to the ideal generated by $(D^N(e_j))_{1 \leq j \leq k}$.

Let $(y_1, \dots, y_k) \in R_N^k$ be a common zero of the polynomials $(P_r)_{0 \leq r \leq k-1}$. Some of the y 's might be equal. Let s be the cardinal of $f y_1, \dots, y_k g$. By symmetry, one can assume that the first y_1, \dots, y_s are pairwise different. For $i \in f 1, \dots, s g$, let us write $m_i = f a \in f 1, \dots, k g j_a = y_i g$. For $0 \leq r \leq s-1$, one has:

$$P_r(y_1, \dots, y_k) = \sum_{i=1}^s m_i \prod_{j=1}^N (y_i - T_j) y_i^r = 0;$$

in other words:

$$\begin{pmatrix} 1 & 1 & \dots & 1 \\ y_1 & y_2 & \dots & y_s \\ \vdots & \vdots & \ddots & \vdots \\ y_1^{s-1} & y_2^{s-1} & \dots & y_s^{s-1} \end{pmatrix} \begin{pmatrix} m_1 \prod_{j=1}^N (y_1 - T_j) \\ m_2 \prod_{j=1}^N (y_2 - T_j) \\ \vdots \\ m_s \prod_{j=1}^N (y_s - T_j) \end{pmatrix} = 0;$$

This implies that for every $i \in f 1, \dots, s g$, we have $y_i \in f T_1, \dots, T_N g$. Hence $Z = f T_1, \dots, T_N g^k$.

REMARK 5.24. — The previous proof remains valid over $R_N - T_i \neq z_i \mathbb{C}$, for any choice of z 's. It is adapted from the elegant proof of [CKW09, Prop. 2.9] which treats the case where all the z_i are 0.

In next section, we will need a primality result concerning the polynomials $(D^N(e_j^{(i)}))_{\substack{1 \leq i \leq k \\ 1 \leq j \leq k_i}}$. As in the previous proof, we start with the case $k = 1$.

LEMMA 5.25. — For $1 \leq i \leq k$, let e_i be the i th elementary symmetric polynomial in $x_1; \dots; x_k$. The polynomials $(D^N(e_i))_{1 \leq i \leq k}$ are pairwise co-prime in $A_k (= A_{1^k}^S)$.

Proof. — We work by induction on k . For $1 \leq i \leq k$, we write $P_{i;k} := D^N(e_i)$ with e_i symmetric in k variables and $P_{i;k}^0 := D^N(e_i)_{j_{T_1} = \dots = T_N = 0}$. For $k = 1$, there is nothing to show. For $k = 2$, we have:

$$P_{1,2}^0 = x_1^N + x_2^N \quad \text{and} \quad P_{2,2}^0 = x_1 x_2^N + x_2 x_1^N = x_1 x_2 (x_1^{N-1} + x_2^{N-1})$$

which are co-prime since $x_1^{N-1} + x_2^{N-1}$ and $x_1^N + x_2^N$ are co-prime as polynomial in x_1 with coefficients in $\mathbb{Q}[x_2]$. If Q is a non-trivial homogeneous element of A_k which divides $P_{1,2}$ and $P_{2,2}$, then $Q_{j_{T_1} = \dots = T_N = 0}$ is of degree 0. Since Q is not equal to 0 it has degree 0 which proves that $P_{1,2}$ and $P_{2,2}$ are co-prime. Suppose now that $k > 3$.

If P is a polynomial in A_k , $P(x_k = 0)$ denotes the polynomial of A_{k-1} obtained by specializing the variable x_k to 0 in P . From the very definition of D^N , we have:

$$P_{i;k}^0 = \sum_{j=1}^k x_j^N e_{i-1}(x_1; \dots; \widehat{x}_j; \dots; x_k):$$

Hence if $1 \leq i \leq k-1$, $P_{i;k}^0(x_k = 0) = P_{i;k-1}^0$. Let $1 \leq i_1 < i_2 \leq k-1$. Suppose that a polynomial Q in A_{1^k} divides $P_{i_1;k}$ and $P_{i_2;k}$. The polynomial $Q^0 := Q_{j_{T_1} = \dots = T_N = 0}$ is homogeneous and $Q(x_k = 0)$ divides $P_{i_1;k-1}^0$ and $P_{i_2;k-1}^0$. By induction, we know that $Q^0(x_k = 0)$ has degree 0. This implies that Q^0 and therefore Q has degree 0. Hence $P_{i_1;k}$ and $P_{i_2;k}$ are co-prime.

It remains to show that for $1 \leq i \leq k-1$, $P_{i;k}$ and $P_{k;k}$ are co-prime. Let Q be a polynomial of A_k which divides $P_{i;k}$ and $P_{k;k}$.

$$P_{k;k}^0 = x_1 x_2 \dots x_k (x_1^{N-1} + \dots + x_k^{N-1}) = e_k(x_1; \dots; x_k) (x_1^{N-1} + \dots + x_k^{N-1}):$$

The polynomial e_k is prime in A_k and does not divide $P_{i;k}$, since $P_{i;k}(x_k = 0)$ is not equal to 0. Hence Q^0 divides $x_1^{N-1} + \dots + x_k^{N-1} =: p_{k;N-1}$. Since Q^0 divides $p_{k;N-1}$ and $P_{i;k}^0$, it divides (in A_k)

$$P_{i;k} - p_{k;N-1} x_k e_{i-1}(x_1; \dots; x_{k-1}):$$

Hence its x_k -degree is at most equal to 1. Since Q^0 is symmetric in the x_i , if it does not have degree equal to 0, it is a multiple of $x_1 + \dots + x_k = e_1$. But for e_1 to divide $p_{k;N-1}$, one must have $k = 2$ and $N-1$ odd. But we supposed $k > 3$. Hence Q^0 has degree 0. This implies that Q has degree 0 and finally that $P_{i;k}$ and $P_{i_2;k}$ are co-prime.

COROLLARY 5.26. — The polynomials $(D^N(e_j^{(i)}))_{\substack{1 \leq i \leq k \\ 1 \leq j \leq k_i}}$ are pairwise co-prime in A_k .

Proof. — First note that $D^N(e_j^{(i)})$ is a polynomial in the variables $x_{r_i+1}; \dots; x_{r_i+k_i}$ and $T_1; \dots; T_N$. It is homogeneous and has degree $N-1+j$ which is bigger than 1.

Hence for $i_1 \neq i_2$, $1 \leq j_1 \leq k_{i_1}$ and $1 \leq j_2 \leq k_{i_2}$, if $D^N(e_{j_1}^{(i_1)})$ and $D^N(e_{j_2}^{(i_2)})$ would have a non-trivial common divisor, they would have an homogeneous divisor in the variables T . However $D^N(e_{j_1}^{(i_1)})$ is not divisible by any non-trivial homogeneous polynomial in the variables T , because evaluating all these variables to 0 in $D^N(e_{j_1}^{(i_1)})$ does not give the 0 polynomial. The case $i_1 = i_2$ follows from Lemma 5.25.

NOTATION 5.27. — If $\widehat{\Gamma}$ is a vinyl graph of level k , denote $T_N(\widehat{\Gamma})$, the space

$$HN_0(HH^N(A_{\underline{k}}; F_{\mathbb{T}}^D(\widehat{\Gamma})))q^{-k(N-1)}$$

for a braid-like \underline{k} -MOY graph- \underline{k} whose closure is equal to $\widehat{\Gamma}$. This is legitimate thanks to Proposition 5.23.

5.3. WHEN ALGEBRA MEETS FOAMS. — The aim of this section is to compare $T_N(\widehat{\Gamma})$ and $S_{N;N}(\widehat{\Gamma})$, namely to prove that these spaces are isomorphic.

Let us consider a braid-like \underline{k} -MOY graph- \underline{k} and denote $\widehat{\Gamma}$ the closure of Γ . We know, thanks to Proposition 4.18, that there is a canonical isomorphism from $F_{\mathbb{T}}^T(\widehat{\Gamma})$ to $HH_0(A_{\underline{k}}; B(\widehat{\Gamma}))$. The space $S_{N;N}(\widehat{\Gamma})$ is a quotient of $F_{\mathbb{T}}^T(\widehat{\Gamma})q^{-k(N-1)}$ while the space $T_N(\widehat{\Gamma})$ is a quotient of $HH_0(A_{\underline{k}}; B(\widehat{\Gamma}))q^{-k(N-1)}$ (thanks to Proposition 5.23). Using the isomorphism φ , we can think of $T_N(\widehat{\Gamma})$ and $S_{N;N}(\widehat{\Gamma})$ as being both quotients of $HH_0(A_{\underline{k}}; B(\widehat{\Gamma}))q^{-k(N-1)}$. The rest of the section is devoted to proving the following proposition:

PROPOSITION 5.28. — *The spaces $T_N(\widehat{\Gamma})$ and $S_{N;N}(\widehat{\Gamma})$ are isomorphic.*

Thanks to the Quelelec-Rose algorithm and Proposition 4.15, it is enough to prove the statement when $\widehat{\Gamma}$ is a collection of circles labeled by \underline{k} . Since the spaces $T_N(\widehat{\Gamma})$ and $S_{N;N}(\widehat{\Gamma})$ are both quotients of $HH_0(A_{\underline{k}}; B(\widehat{\Gamma}))q^{-k(N-1)}$ which is itself isomorphic to $A_{\underline{k}}q^{-k(N-1)}$, let us write $T_N(\widehat{\Gamma}) = A_{\underline{k}}q^{-k(N-1)} = I_1$ and $S_{N;N}(\widehat{\Gamma}) = A_{\underline{k}}q^{-k(N-1)} = I_2$. With these notations, we only need to show that the spaces I_1 and I_2 of $A_{\underline{k}}q^{-k(N-1)}$ are equal.

LEMMA 5.29. — *The space I_1 is generated by the polynomials $(D^N(e_j^{(i)}))_{\substack{1 \leq i \leq N \\ 1 \leq j \leq k_i}}$. Forgetting about the action of the variables T , it is a graded vector space. Its graded dimension over \mathbb{Q} is equal to:*

$$\dim_{\mathbb{Q}}^{\text{gr}} I_1 = q^{-k(N-1)} \left(1 \prod_{b=1}^k \prod_{i=1}^{k_b} (1 - q^{2(i+N-1)}) \right) \dim_{\mathbb{Q}} A_{\underline{k}};$$

Proof. — The first statement is obvious. The second one follows from the fact that the polynomials $(D^N(e_j^{(i)}))_{\substack{1 \leq i \leq N \\ 1 \leq j \leq k_i}}$ are pairwise co-prime. Indeed this implies that

$$\begin{aligned} \langle D^N(e_{j_1}^{(i_1)}) \rangle_{A_{\underline{k}}} \setminus \langle D^N(e_{j_2}^{(i_2)}) \rangle_{A_{\underline{k}}} \setminus \dots \setminus \langle D^N(e_{j_a}^{(i_a)}) \rangle_{A_{\underline{k}}} \\ = \langle D^N(e_{j_1}^{(i_1)}) D^N(e_{j_2}^{(i_2)}) \dots D^N(e_{j_a}^{(i_a)}) \rangle_{A_{\underline{k}}}; \end{aligned}$$

Since the polynomial $D^N(e_j^{(i)})$ is homogeneous of degree $2(N + j - 1)$, we have:

$$\dim_q \langle D^N(e_{j_1}^{(i_1)}) \ D^N(e_{j_2}^{(i_2)}) \ \dots \ D^N(e_{j_a}^{(i_a)}) \rangle_{A_{\underline{k}}} = q^{-k(N-1)} (\dim_q^\square A_{\underline{k}}) \prod_{i=1}^a q^{2(j_i + N - 1)}.$$

The space I_1 is the sum of all spaces $\langle D^N(e_j^{(i)}) \rangle_{A_{\underline{k}}}$. This implies:

$$\dim_q^\square I_1 = q^{-k(N-1)} \left(1 \prod_{b=1}^{k_b} \prod_{i=1}^{k_b} (1 - q^{2(i+N-1)}) \right) \dim_q^\square A_{\underline{k}}.$$

The proof of Proposition 5.28, follows from next lemma.

LEMMA 5.30. — *The spaces I_1 and I_2 are equal.*

Proof. — It is clear that I_1 is in I_2 because the polynomials $(D^N(e_j^{(i)}))_{\substack{1 \leq i \leq k_b \\ 1 \leq j \leq k_i}}$ are all in I_2 (this follows from the definition of the evaluation of vinyl foams). We know thanks to Lemma 5.3 that $S_{N;N}$ associates with a circle labeled k_b a free R_N -module of graded rank

$$\prod_{i=1}^{k_b} \frac{q^{-i(N-1)} q^{i+N-1}}{q^{-i} q^i} = q^{-k_b(N-1)} \prod_{i=1}^{k_b} \frac{1 - q^{2(i+N-1)}}{1 - q^{2i}}.$$

Thanks to the monoidality of this functor, we obtain that the graded rank of $S_{N;N}(\)$ is equal to

$$q^{-k(N-1)} \prod_{b=1}^{k_b} \prod_{i=1}^{k_b} \frac{1 - q^{2(i+N-1)}}{1 - q^{2i}}.$$

But $A_{\underline{k}} q^{-k(N-1)}$ has a graded rank over R_N equal to:

$$q^{-k(N-1)} \prod_{b=1}^{k_b} \prod_{i=1}^{k_b} \frac{1}{1 - q^{2i}}.$$

Hence we have:

$$\begin{aligned} \dim_q^\square I_2 &= q^{-k(N-1)} \dim_q^\square A_{\underline{k}} \cdot q^{-k(N-1)} \prod_{b=1}^{k_b} \prod_{i=1}^{k_b} (1 - q^{2(i+N-1)}) \dim_q^\square A_{\underline{k}} \\ &= q^{-k(N-1)} \left(1 \prod_{b=1}^{k_b} \prod_{i=1}^{k_b} (1 - q^{2(i+N-1)}) \right) \dim_q^\square A_{\underline{k}} \\ &= \dim_q^\square I_1. \end{aligned}$$

6. LINK HOMOLOGIES

In this section, we define the symmetric Khovanov-Rozansky homology on diagrams of braid closures and prove that they are indeed links invariants. The definition is of a purely foamy nature. However for proving the invariance we need to use the dictionary developed in Section 4. We derive the invariance of the symmetric homologies from the invariance of the triply graded homology [KR08]. We use the description of this

homology as Hochschild homology of complexes of Soergel bimodules due to Khovanov and Rouquier [Kho07, Rou17]. We show that the extra differential introduced in Section 5.2.2, is compatible with their construction. Finally, we prove that when taking the homology with respect to this extra differential, one gets the same link homology as the one obtained by applying the foamy functor of the previous sections. This link homology categorifies the Reshetikhin-Turaev link invariant associated with q -symmetric powers of the standard representation of $U_q(\mathfrak{sl}_N)$. We call it the *symmetric Khovanov-Rozansky homology*.

6.1. THE CHAIN COMPLEXES. — The idea of the construction is somewhat classical. We follow the normalization used in [Ras15]. Let D be a diagram of a braid-closure of level k , and $\text{Cross}(D)$ be its set of crossings. For $x \in \text{Cross}(D)$ we define a finite set I_x by the following rules:

$$\begin{aligned}
 &\text{if } x = \begin{array}{c} \nearrow^m \searrow^n \\ \nwarrow^m \nearrow^n \end{array} \text{ and } m \leq n \text{ then } I_x = \{m, \dots, n-1, 0\}g; \\
 &\text{if } x = \begin{array}{c} \nearrow^m \searrow^n \\ \nwarrow^m \nearrow^n \end{array} \text{ and } m > n \text{ then } I_x = \{n, \dots, m-1, 0\}g; \\
 &\text{if } x = \begin{array}{c} \nearrow^m \searrow^n \\ \nwarrow^m \nearrow^n \end{array} \text{ and } m \leq n \text{ then } I_x = \{0, 1, \dots, m\}g; \\
 &\text{if } x = \begin{array}{c} \nearrow^m \searrow^n \\ \nwarrow^m \nearrow^n \end{array} \text{ and } m > n \text{ then } I_x = \{0, 1, \dots, n\}g;
 \end{aligned}$$

In the first two cases, we say that x is of type $(m; n; +)$ and in the last two cases of type $(m; n; -)$. If x is a crossing and i is an element of I_x we define

$$(x; i; x; i) = \begin{cases} (m+i; i; i; m) & \text{if } x \text{ is of type } (m; n; +), \text{ and } m \leq n \\ (n+i; i; i; n) & \text{if } x \text{ is of type } (m; n; +), \text{ and } m > n, \\ (i; m; m; i) & \text{if } x \text{ is of type } (m; n; -), \text{ and } m \leq n \\ (i; n; n; i) & \text{if } x \text{ is of type } (m; n; -) \text{ and } m > n. \end{cases}$$

We set $I(D)$ to be $\prod_{x \in \text{Cross}(D)} I_x$ and call the elements of $I(D)$ the *states of D* . With every state $s = (s_x)_{x \in \text{Cross}(D)}$ of D we associate a vinyl graph D_s of level k by replacing every crossing x of D according to the rules given by Figure 11.

If s is a state, we define

$$|s\rangle = \sum_{x \in \text{Cross}(D)} |x; s_x\rangle \quad \text{and} \quad \langle s| = \sum_{x \in \text{Cross}(D)} \langle x; s_x|$$

and we set D_s to sit in topological degree $-|s\rangle$ and to be shifted in q -degree by $|s\rangle$.

If $s = (s_x)_{x \in \text{Cross}(D)}$ and $s^\flat = (s_x^\flat)_{x \in \text{Cross}(D)}$ are two states which are equal on all but one of their coordinate x , for which $s_x^\flat = s_x + 1$, we write $(s \flat s^\flat)$ (or $(s \flat_x s^\flat)$ to be precise). In this case, we define $F_{D; s \flat s^\flat}$ to be the vinyl D_{s^\flat} -foam- D_s , which is the identity everywhere but in a neighborhood of x , where it is given by Figure 12.






FIGURE 11.

REMARK 6.1. — If x is of type $(1; 1; +)$ or $(1; 1; -)$, the set I_x has two elements and the foam $F_{D; s!} \times s^0$ simplifies upon removing 0-labeled facets:



We define an hyper-rectangle $R(D)$ of graded R_N -modules. The vertices of this hyper-rectangle are labeled by states and the edges by pair of states $(s; s')$ for which $s \neq s'$. With every state $s = (s_x)_{x \in 2}$ we associated the graded R_N -module $V_s := S_{k; N}(D_s)q^s$ and we declare that it has homological degree $-s$.

With every edge $(s \neq s')$, we associate the map $d_{s! s'} := S_{k; N}(F_{D; s! s'}) : V_s \rightarrow V_{s'}$. One easily checks that all these maps are q -homogeneous of degree 0 (thanks to the degree shift q^s) and increase the homological degree by 1. Hence we call them *pre-differentials*.



if x is of type $(m; n; +)$ and $m < n$, if x is of type $(m; n; -)$ and $n < m$,

if x is of type $(m; n; +)$ and $m > n$, if x is of type $(m; n; -)$ and $n < m$.

FIGURE 12. It is worth noting that this has q -degree 1.

All squares in $R(D)$ commute because of the TQFT nature of the functor $S_{k;N}$. Furthermore, if the composition of two pre-differentials $d_{s^0|s^0}$ and $d_{s^0|s^0}$ does not fit into a square in $R(D)$, this means that we have $(s^0|s^0)$ and $(s^0|s^0)$ for the same x in \mathcal{F} . In this case $d_{s^0|s^0} d_{s^0|s^0} = 0$ because the foams of Figure 13 are $\mathbb{1}$ -equivalent to 0. Indeed, given a closure F of one of this two foams, one can gather the colorings of F in canceling pairs. Given a coloring c of F , the two facets labeled 1 must carry different pigments, say i and j . One can locally exchange the pigments i and j . This produces a new coloring c^0 and the contributions of c and c^0 in the evaluation formula (Definition 3.14) cancel each others. Hence if we add⁽⁸⁾ some signs

⁽⁸⁾There are many ways to do it, but all possibilities produce isomorphic chain complexes. In order to get functoriality of the construction announced in the introduction, one should be careful in this choice. This boils down to endowing the set \mathcal{F} with a total order as detailed in [ETW18]. A more systematical construction is given in [Bla10] making use of the exterior algebra generated by \mathcal{F} . This last approach works only for uncolored links, but can be easily adapted to the colored case.




FIGURE 13.

to the pre-differential to turn the commutativity of squares into anti-commutativity we can flatten the hyper-rectangle $R(D)$ and obtain a complex of graded R_N -modules $S^0(D)$.

Finally we set $\widehat{S}^0(D)$ to be equal to $S^0(D)q^{o(D)}$, where

$$o(D) = \sum_{x^2} o_x \quad \text{and} \quad o_x = \begin{cases} m(m+N-1) & \text{if } x \text{ is of type } (m; m; +), \\ m(m+N-1) & \text{if } x \text{ is of type } (m; m; -), \\ 0 & \text{else.} \end{cases}$$

THEOREM 6.2. — *The homology of $\widehat{S}^0(D)$ is a link invariant which categorifies the symmetric Reshetikhin-Turaev invariant.*

NOTATION 6.3. — In what follows, we deal with three different differentials: the Hochschild one (d_H), the topological one (d_T) and the extra one (d^N). We denote by $\mathcal{H}(\cdot)$ (resp. $\mathcal{H}T(\cdot)$, $\mathcal{H}N(\cdot)$), the homology taken with respect to d_H (resp. d_T , d^N) and by $\mathcal{H}NT(\cdot)$ the homology taken with respect to the total complex⁽⁹⁾ built out of the bicomplex with bi-differentials ($d^N; d_T$). The Hochschild homology is computed using the Koszul complex (as explained in Section 4.4 and Appendix B). Moreover, for the Hochschild homology, we will often drop the algebra in the notation: writing $\mathcal{H}(M)$ instead of $\mathcal{H}(A; M)$. As explained in Section 5.2.2, we denote by \mathcal{H}^N the Hochschild homology with an additional q -degree shift making the extra differential of q -degree 0.

Note that homological degree for $\mathcal{H}\mathcal{H}$ and $\mathcal{H}N$ coincide. We denote by $[]_H$ and $[]_T$ grading shifts⁽¹⁰⁾ with respect to the H and the T -degree. When considering the total complex built out of the bicomplex with bi-differentials ($d^N; d_T$), the homological degree shift is denoted by $[]_{TH}$. Let us recall that grading shifts with respect to the q -degree are denoted by q .


⁽⁹⁾Note that d^N is a differential of chain complex (it has H -degree -1) while d_T is a differential of cochain complex (it has T -degree $+1$), the total complex we consider is a complex of cochain complex: the total homological degree is equal to the T -degree minus the H -degree.

⁽¹⁰⁾We use the homological convention for grading shifts: If $V = \sum_i V_i$, we have $(V[k])_i = V_{i-k}$.

Proof of Theorem 6.2. — The fact the graded Euler characteristic of $\widehat{S}^{\cdot}(\cdot)$ is indeed the symmetric Reshetikhin-Turaev invariant follows from Theorem 5.17 and from the construction of the hyper-rectangle which is clearly designed to categorify the identities (2.10) and (2.11). In order to prove invariance, we proceed in two steps:

(1) First, we prove that if D is a diagram of a knotted vinyl graph (that is, a diagram of a knotted MOY graph which satisfies the same condition on tangent than vinyl graph), then the homotopy type of complex $S^{\cdot}(D)$ (up to a q -grading shift) only depends on the isotopy type of knotted graph in the annulus.

(2) Then we prove that we have a stabilization property. Namely, we will show, that for any diagram of knotted braid-like \underline{k} -MOY graph- \underline{k} , the complexes associated with the following three diagrams



have the same homology. We write \underline{k}^+ (resp. \underline{k}^-) for the two knotted braid-like \underline{k}^0 -MOY graph- \underline{k}^0 obtained from \underline{k}^0 by adding one strand labeled by 1 on the right and a positive (resp. negative) crossing on the top of it.

Note that in our stabilization, we only deal with a strand labeled by 1. Thanks to a trick due to Mackaay-Stošić-Vaz [MSV11] (see as well [Wu14] and [Cau17, Fig. 1]), this implies (together with the homotopy equivalences (6.1)) that we actually get the stabilization property for any labels. Using this trick requires that we actually deal with knotted vinyl graphs in step (1) (and not only with links). Finally thanks to Markov theorem, we can conclude that $S^{\cdot}(D)$ is a link invariant.

$$(6.1) \quad q^{-ab} S \left(\begin{array}{c} a+b \\ \uparrow \\ \circlearrowleft \\ \swarrow \quad \searrow \\ a \quad b \end{array} \right) = S \left(\begin{array}{c} a+b \\ \uparrow \\ \circlearrowright \\ \swarrow \quad \searrow \\ a \quad b \end{array} \right) = q^{ab} S \left(\begin{array}{c} a+b \\ \uparrow \\ \circlearrowright \\ \swarrow \quad \searrow \\ a \quad b \end{array} \right) :$$

The proof of step (1) is quite standard. One first consider the case where all strands involved in the braid relations have label 1. This case is treated in terms of foams in [Vaz08, Figs. 5.14 & 5.16]. The case with strands of arbitrary labels follows from the first case and the invariance under the so-called *fork slide moves* (see Figure 14).

Indeed, one can twist (see Figure 15) each strands and use the fork slide moves and the 1-labeled braid relation to deduce the arbitrary labeled braid relations. See [WW17, MSV09, Wu14] for similar arguments. Proof of invariance for the fork slide move using foam is quite standard see for instance [QR16, Proof of Prop. 4.10] or [ETW18, §3.3]. But let us briefly sketch how it works on the fork slide move on the top left move of Figure 14. We consider the diagram on the left-hand side of the move and its associated complex in Figure 16. The space on the first line can be




FIGURE 14. Fork slide moves



FIGURE 15. Blistening an edge of label i .




FIGURE 16. For simplicity we dropped the symbols S_N .




FIGURE 17.





FIGURE 18.

decomposed using the isomorphisms (5.1) and (5.3). The space on the second line can be decomposed using isomorphisms (5.1) and (3.4). This gives the diagrams in Figure 17. One can check that the vertical maps injective on the second term of the direct sum. We can use them to simplify the complex. The complex is homotopy equivalent to that in Figure 18. The second line turns out to be exact (thanks to isomorphism (5.3)) except on the rightmost (corresponding to $j = 0$) term if $n < m$. If the line is exact we can simplify and obtain that the complex is homotopy equivalent to that corresponding to the right-hand side diagram of the for slide move we are interested in.

If $n < m$, we can cancel out the second line except the rightmost term which becomes:



Hence the whole complex is homotopy equivalent to that corresponding to the right-hand side diagram of the for slide move we are interested in.

The invariance under braid relations can as well be deduced from the algebraic setting using Soergel bimodules, see [WW17]. The proof of step (2) is more involved. We need to use the dictionary between Soergel bimodules and vinyl foams developed in Section 5 and a stabilization result which holds for Soergel bimodules.

First note that if D has braid index equal to k ,

$$S(D) = HT(HN(HH^N(B(R(D))))))q^{-k(N-1)};$$

thanks to Proposition 5.28. We claim that we have

$$HNT(HH^N(B(R(D)))) = HT(HN(HH^N(B(R(D))))):$$

We consider $HH^N(B(R(D)))$. It is a bi-complex which we temporarily denote by C . Proposition 5.23 tells us that $HN(C)$ is concentrated in H -degree equal to 0. This implies that the spectral sequence $E(C)$ induced by the bi-complex structure with $HT(HN(C))$ on the second page has only trivial differentials on this page. This spectral sequence converges to $HNT(C)$ because the bi-complex is bounded. Hence, we have

$$HNT(HH^N(B(R(D)))) = E^1(C) = E^2(C) = HT(HN(HH^N(B(R(D))))):$$

The stabilization result for Soergel bimodules is given by Lemma 6.5 and its proof occupies Section 6.2. It tells us that:

$$HT(HH^N(B(\dots)))[1]_T[1]_Hq^{-1} = HT(HH^N(B(\dots))) + HT(HH^N(B(\dots)))q^1:$$

Using HH^N instead of HH and collapsing the T -grading and the H -grading to their difference, we get:

$$HT(HH^N(B(\ +)))q^{2(N-1)-1} \simeq HT(HH^N(B(\ 0))) \simeq HT(HH^N(B(\ -)))q^1:$$

Moreover these isomorphisms preserve the extra-differential d^N . We shift by $q^{-k(N-1)}$. This proves that

$$\begin{aligned} HN(HT(HH^N(B(\ +))))q^{-N}q^{(k+1)(N-1)} \simeq HN(HT(HH^N(B(\ 0))))q^{-k(N-1)} \\ \simeq HN(HT(HH^N(B(\ -))))[1]_{TH}q^Nq^{(k+1)(N-1)}. \end{aligned}$$

Since homological degrees are all finite, there is a spectral sequence with

$$HN(HT(HH A_{\underline{k}}; B(\)))$$

on the second page converging to $HNT(HH A_{\underline{k}}; B(\))$. The previous isomorphisms descend to the spectral sequences and to their limits. Hence we have:

$$\begin{aligned} HNT(HH^N A_{\underline{k}}; B(\ +))q^Nq^{(k+1)(N-1)} \simeq HNT(HH^N(A_{\underline{k}}; B(\ 0)))q^{-k(N-1)} \\ \simeq HNT(HH^N(A_{\underline{k}}; B(\ -)))q^{-N}q^{(k+1)(N-1)}. \end{aligned}$$

This implies that $S(D_+)q^N \simeq S(D_0) \simeq S(D_-)q^{-N}$, and finally that

$$\widehat{S}(D_+) \simeq S(D_0) \simeq \widehat{S}(D_-):$$

REMARK 6.4. — Note that the previous proof can be adapted to disk-like foams up to $\mathbb{1}$ -equivalence. Then it turns out to be a rewriting of the invariance of the triply graded homology very close to [Rou17] (see also [KR08, Ras15]).

6.2. STABILIZATION. — The aim of this section is to prove the stabilization move for Soergel bimodules. This basically follows from Rouquier [Rou17]. However, since for further use we need to be careful with some additional structures going on, we repeat the proof. Note, however, that the framework in which the results of [Rou17] are stated is more general⁽¹¹⁾ than ours.

Let k be a positive integer and $\underline{k}^{11} := (k_1; \dots; k_{\cdot-1}; 1; 1)$ be a finite sequence of positive integers of level k . We define $\underline{k}^2 := (k_1; \dots; k_{\cdot-1}; 2)$ to be the same sequence where the last two 1s has been merged $\underline{k}^1 := (k_1; \dots; k_{\cdot-1}; 1)$ to be the same sequence where the last 1 has been dropped, and finally $\underline{k}^0 := (k_1; \dots; k_{\cdot-1})$ to be the same sequence where the last two 1s has been dropped. We consider the algebras $A^{11} := A_{\underline{k}^{11}}$, $A^2 := A_{\underline{k}^2}$ and $A^1 := A_{\underline{k}^1}$.

We consider M a complex of A^1 -modules- A^1 which is projective as an A^1 -module and as a module- A^1 , and we define $MI := M \otimes_{R_N[x_k]}$. It has a natural structure complex of A^{11} -modules- A^{11} and these modules are projective as A^1 -modules and as a modules- A^1 . We denote by $\text{the } A^{11}\text{-module-}A^{11} A^{11} \otimes_{A^2} A^{11} q^{-1}$. Note that with the

⁽¹¹⁾ Namely, he deals with arbitrary Coxeter groups, while we only consider the type A .

notations of Section 4, A^1 is the Soergel bimodule associated with $\begin{matrix} 1 & 1 \\ & \searrow \swarrow \\ & 2 \\ & \swarrow \searrow \\ 1 & 1 \end{matrix}$ and the A^{11} -module- A^{11} A^{11} is the Soergel bimodule associated with $\begin{matrix} 1 \\ \downarrow \\ 1 \end{matrix}$ $\begin{matrix} 1 \\ \uparrow \\ 1 \end{matrix}$. Finally we consider two morphisms of A^{11} -modules- A^{11} .

$$s: A^1 \otimes A^{11} \rightarrow A^{11} \otimes A^1 \quad \text{and} \quad m: A^{11} \otimes A^1 \rightarrow A^1 \otimes A^{11}$$

The notation may seem confusing: m is for merge and s is for split. We define

$$F := 0 \rightarrow A^{11} q^{-1} \rightarrow 0; \\ F^{-1} := 0 \rightarrow A^{11} q^{-1} \rightarrow 0;$$

In F^{-1} and F , $A^{11} q^{-1}$ is in T -degree 0, $A^{11} q^{-1}$ in T -degree 1 and $A^{11} q$ in T -degree -1 .

Note that in the language of Section 4, m and s are the maps induced by the foams given in Remark 6.1.

The following lemma should be compared to [Cau17, Lem. 6.3].

LEMMA 6.5. — *The homology of the complexes*

$$(HH(A^1; M); d_T); \quad (HH(A^{11}; MI_{A^{11}} F)[1]_T[1]_{Hq^{-1}}; d_T); \\ (HH(A^{11}; MI_{A^{11}} F^{-1})q^1; d_T)$$

are isomorphic as triply graded R_N -modules. Moreover we can choose the isomorphisms to commute with the extra-differential d^N .

Proof. — We will deal with complexes carrying three differential: the Hochschild differential d_H , the topological differential d_T and the additional differential d^N .

$$C(A^{11}) = C(A^1) \otimes_{R_N} X;$$

where

$$X := R_N[x_k] \xrightarrow{R_N[x_k]q^2 \xrightarrow{x_k \ 1 \ 1 \ x_k} R_N[x_k]} R_N[x_k] \xrightarrow{R_N[x_k]} R_N[x_k],$$

$\xrightarrow{x_k^N @ x_k}$

Hence

$$C(A^{11})_{(A^{11})^{\text{en}}} (MI_{A^{11}} F)^{\vee} = C(A^1)_{(A^1)^{\text{en}}} (M_{A^1} (X \otimes_{R_N[x_k]^{\text{en}}} F));$$

where B^{en} denotes the algebra $B \otimes_R B^{\text{opp}}$ for any R -algebra. We have, focusing on $X \otimes_{R_N[x_k]^{\text{en}}} F$,

$$X \otimes_{R_N[x_k]^{\text{en}}} F \xrightarrow{x_k \ 1 \ 1 \ x_k} \begin{matrix} q^2 & \xrightarrow{s} & A^{11}q \\ \downarrow & & \downarrow 0 \\ & & A^{11}q^{-1} \\ & \xrightarrow{s} & \end{matrix} :$$

In \mathcal{C} , the elements $2(x_k \ 1 \ 1 \ x_k)$ and $(x_k \ x_k \ 1) \ 1 \ 1 \ (x_k \ x_k \ 1)$ are equal. Hence, we have the following exact sequence of bicomplexes (for d_T and d_H) of graded A^{11} -modules- A^{11} in Figure 19. Note that this is *not* a sequence of tri-complexes: it does




FIGURE 19. Here A^{11} is equal to A^{11} as a A^{11} -module and has a right A^{11} -action twisted by the transposition $x_k \ 1$ which exchanges x_k and x_{k-1} .

not respect the differential d^N . The dotted arrows represent the part of d^N appearing in X . In each topological degree, this sequence splits as a sequence of complexes (for the Hochschild differential) of

$$(p(x_1; \dots; x_{k_2})x_{k-1}^i x_k^j) = p(x_1; \dots; x_{k_2})x_{k-1}^i x_k^j$$

We now take the homology with respect to the Hochschild differentials.⁽¹²⁾ It implies, that for each i in \mathbb{N} , that we have the sequence of graded complexes:

$$0 \rightarrow \mathbb{H}_i(C(A^1)_{(A^1)^{en}}(M_{A^1} Y_1)) \rightarrow \mathbb{H}_i(A^{11}; M_{A^1} F) \rightarrow \mathbb{H}_i(C(A^1)_{(A^1)^{en}}(M_{A^1} Y_2)) \rightarrow 0$$

⁽¹²⁾Here, we abuse a little bit the appellation Hochschild differential: in Y_1 and Y_2 the vertical arrows are part of the Hochschild differential, while the horizontal ones are part of the topological differential.

We have a short exact sequence of complexes of A^{11} -modules- A^{11} :


$$0 \rightarrow A^{11}q^{a \nabla a(x_k \ x_k \ 1) \ 1 \ a \ (x_k \ x_k \ 1)} \rightarrow A^{11}q^{1} \rightarrow 0;$$

This implies that for all i , the complex $HH_i(C(A^1)_{(A^1)^{en}}(M_{A^1} Y_2))$ is homotopically trivial (with respect to d_T). It follows that:

$$HT(HH_i(A^{11}; MI \ F^{-1})) \cong HT(HH_i(C(A^1)_{(A^1)^{en}}(M_{A^1} Y_1))):$$

This implies that the part of the differential d^N coming from X is equal to 0 in $HT(HH_i(A^{11}; MI \ F^{-1}))$, and therefore that this isomorphism commutes with d^N . The complex $(HH_i(C(A^1)_{(A^1)^{en}}(M_{A^1} Y_1)); d_T)$ has the same homology as the complex $(HH_i(C(A^1)_{(A^1)^{en}}(M_{A^1} A^1 q^1)) [+1]_H; d_T)$ which itself is equal to $HH_i(A^1; M) [+1]_H q^1$.

The argument for F^{-1} is similar, with the following short exact sequence of bi-complexes:




6.3. AN EXAMPLE OF COMPUTATION. — In this section, we compute the homology of the positive uncolored Hopf link in the non-equivariant setting (i.e., evaluating the variables T on 0).

First remark (see Remark 6.1) that the pre-differential given by the foam $F_{D;S^1 \times S^0}$ for x of type $(1;1;+)$ (resp. of type $(1;1;-)$) is always surjective (resp. injective). This implies in particular that if an uncolored braid diagram D has n crossings, then the

homological length (for the topological degree) of $H(\widehat{S}(D))$ is at most n . If D contains both positive and negative crossings then $H(\widehat{S}(D))$ has length at most $n - 1$.

The complex of resolutions of the positive uncolored Hopf link is given by:



In this picture, the red labels are meant to represent a basis. For a given diagram γ , pick a \mathbb{Z} -tree-like foam and decorate it with the elements given by red sets. The blue quantum numbers are meant to represent the quantum dimension of the spaces.

Observe that the two spaces in the middle are isomorphic and that the pre-differential on the left are equal up to this isomorphism, and therefore have the same kernel. On the other hand, the pre-differential on the right are surjective. Taking care of the different grading shifts, this gives:

$$H_i(\widehat{S}(D_n)) = H_i(S(D_n))q^{-Nn} = \begin{cases} \mathbb{Q}[N][N+1]q^{2N+1} & \text{if } i = 0, \\ \mathbb{Q}[N]q^{-(N+1)} & \text{if } i = 1, \\ 0 & \text{else.} \end{cases}$$

REMARK 6.6

(1) Note in particular that the symmetric uncolored homology of links is not trivial for $N = 1$ (for which the corresponding polynomial invariant is always 1). We expect that a simple combinatorial description of this homology is achievable since the polynomial invariant of MOY graph has an especially simple form for $N = 1$ (see Lemma 2.8).

(2) For $N = 2$, this gives a different homology than the Khovanov homology and odd Khovanov homology.

APPENDIX A. QUANTUM LINK INVARIANTS AND REPRESENTATIONS OF $U_q(\mathfrak{gl}_N)$

In this appendix we provide details about the relation between the graphical MOY calculi defined in Section 2.1 and the representations of $U_q(\mathfrak{gl}_N)$. The aim is to give explicit definitions the Reshetikhin-Turaev functors which associate with a MOY graph an intertwiner of $U_q(\mathfrak{gl}_N)$ -representations. We will define two such functors: one sends an edge labeled k onto the identity of ${}^k V_q$ where V_q is the standard representation of $U_q(\mathfrak{sl}_N)$. This yields the *exterior* MOY calculus.⁽¹³⁾ The other one associates with such an edge the identity of $\text{Sym}_q^k V_q$. This yields what we call the *symmetric* MOY calculus.

A.1. THE QUANTUM GROUP $U_q(\mathfrak{gl}_N)$. — We first recall the definition of $U_q(\mathfrak{gl}_N)$. For a general presentation of quantum groups we refer to [Lus10].

DEFINITION A.1. — Let N be a positive integer. The *quantum general linear algebra* $U_q(\mathfrak{gl}_N)$ is the associative, unital $\mathbb{C}(q)$ -algebra generated by L_i, L_i^{-1}, F_j and E_j , with $1 \leq i \leq N$ and $1 \leq j \leq N - 1$ subject to the relations

$$\begin{aligned} L_i L_j &= L_j L_i; & L_i L_i^{-1} &= L_i^{-1} L_i = 1; \\ L_i F_i &= q^{-1} F_i L_i; & L_{i+1} F_i &= q F_i L_{i+1}; & L_i E_i &= q E_i L_i; & L_{i+1} E_i &= q^{-1} E_i L_{i+1}; \\ L_j F_i &= F_i L_j; & L_j E_i &= E_i L_j & \text{for } j &\neq i; i + 1, \\ E_i F_j &- F_j E_i &= &_{ij} \frac{L_i L_{i+1}^{-1} - L_i^{-1} L_{i+1}}{q - q^{-1}}; \\ [2]_q F_i F_j F_i &= F_i^2 F_j + F_j F_i^2 & \text{if } ji - jj = 1, \\ [2]_q E_i E_j E_i &= E_i^2 E_j + E_j E_i^2 & \text{if } ji - jj = 1, \\ E_i E_j &= E_j E_i; & F_i F_j &= F_j F_i & \text{if } ji - jj > 1. \end{aligned}$$

PROPOSITION A.2. — Defining $\iota : U_q(\mathfrak{gl}_N) \rightarrow U_q(\mathfrak{gl}_N)^2, S : U_q(\mathfrak{gl}_N)^{\text{opp}} \rightarrow U_q(\mathfrak{gl}_N)$ and $\epsilon : U_q(\mathfrak{gl}_N) \rightarrow \mathbb{C}(q)$ to be the $\mathbb{C}(q)$ algebra maps defined by:

$$\begin{aligned} (L_i^{-1}) &= L_i^{-1} L_i^{-1}; & S(L_i^{-1}) &= L_i^{-1}; & \epsilon(L_i^{-1}) &= 1; \\ (F_i) &= F_i^{-1} + L_i^{-1} L_{i+1}^{-1} F_i; & S(F_i) &= L_i L_{i+1}^{-1} F_i; & \epsilon(F_i) &= 0; \\ (E_i) &= E_i^{-1} + L_i L_{i+1}^{-1} E_i; & S(E_i) &= E_i L_i^{-1} L_{i+1}; & \epsilon(E_i) &= 0; \end{aligned}$$

endow $U_q(\mathfrak{gl}_N)$ with a structure of Hopf algebras with antipode. Furthermore the category of finite-dimensional $U_q(\mathfrak{gl}_N)$ -modules is braided.

⁽¹³⁾This is the “classical” MOY calculus. MOY stands for Murakami-Ohtsuki-Yamada who gave the identities described in Section 2.1 in [MOY98].

PROPOSITION A.3. — We define V_q to be an N th dimensional $C(q)$ -vector space with basis $(b_i)_{i=1,\dots,N}$. The formulas:

$$\begin{aligned} L_i b_j &= q b_j; & L_i^{-1} b_j &= q^{-1} b_j; & L_i^{-1} b_j &= b_j \quad j \notin i; \\ E_{i-1} b_i &= b_{i-1} & E_i b_j &= 0; & & \text{if } i \notin j-1, \\ F_i b_i &= b_{i+1} & F_i b_j &= 0; & & \text{if } i \notin j, \end{aligned}$$

endow V_q with a structure of $U_q(\mathfrak{gl}_N)$ -modules.

Following [ST19] we now consider the tensor algebra $T V_q$. This algebra is naturally graded and endowed with an action of $U_q(\mathfrak{gl}_N)$ which preserve the grading (i.e., for every integer a , $T^a V_q$ is a $U_q(\mathfrak{gl}_N)$ -submodule of $T V_q$). We consider two two-sided ideals $E^2 V_q$ and $S^2 V_q$ inside this algebra $T V_q$:

$$\begin{aligned} E V_q &:= \langle h q b_i - b_j - b_j \text{ for } i < j \rangle \\ \text{and } S V_q &:= \langle h b_m - b_m, b_i - b_j + q b_j - b_j \text{ for all } m \text{ and for } i < j \rangle \end{aligned}$$

Since these two ideals are homogeneous the quotient

$${}^q V_q := T V_q / S V_q \quad \text{and} \quad \text{Sym}_q V_q := T V_q / E V_q$$

inherits a grading from $T V_q$. One easily checks that $E^2 V_q$ and $S^2 V_q$ are stable under the action of $U_q(\mathfrak{gl}_N)$ over $T V_q$. This implies that for any non-negative integer a , ${}^a V_q$ and $\text{Sym}_q^a V_q$ inherit structure of $U_q(\mathfrak{gl}_N)$ -modules. One can show that for every integer a , ${}^a V_q$ and $\text{Sym}_q^a V_q$ are simple modules. The image of a pure tensor $x_1 \otimes \dots \otimes x_a$ is denoted by

$$x_1 \wedge \dots \wedge x_a \text{ in } {}^a V_q \quad \text{and by } x_1 \otimes \dots \otimes x_a \text{ in } \text{Sym}_q^a V_q.$$

The $C(q)$ vector space ${}^a V_q$ has dimension $\binom{N}{a}$ and is spanned by the vectors

$$(b_{i_1} \wedge b_{i_2} \wedge \dots \wedge b_{i_a})_{1 \leq i_1 < i_2 < \dots < i_a \leq N}.$$

If $1 \leq i_1 < i_2 < \dots < i_a \leq N$ and $I = i_1 \dots i_a g$, we write $b_I = b_{i_1} \wedge b_{i_2} \wedge \dots \wedge b_{i_a}$.

DEFINITION A.4. — Let X be a set. A *multi-subset* of X is a map $Y : X \rightarrow \mathbb{N}$. If $\sum_{x \in X} Y(x) < \infty$, the multi-subset Y is said to be *finite* and the sum is its *cardinal* (denoted by $\#Y$). If x is an element of X , the number $Y(x)$ is the *multiplicity* of x in Y .

The $C(q)$ vector space $\text{Sym}_q^a V_q$ has dimension $\binom{N+a-1}{a}$ and is spanned by the vectors

$$(b_{i_1} \otimes b_{i_2} \otimes \dots \otimes b_{i_a})_{1 \leq i_1 \leq i_2 \leq \dots \leq i_a \leq N}.$$

If $1 \leq i_1 \leq i_2 \leq \dots \leq i_a \leq N$ and I is the multi-subset of $I_N := \{1, \dots, N\}^a$, we write $b_I^{\otimes} = b_{i_1} \otimes b_{i_2} \otimes \dots \otimes b_{i_a}$.

A.2. EXTERIOR MOY CALCULUS. — In this subsection we work in the full subcategory $U_q(\mathfrak{gl}_N)\text{-mod}$ of finite-dimensional $U_q(\mathfrak{gl}_N)$ -modules generated (as a monoidal category) by the modules a_qV_q for $0 \leq a \leq N$ and their duals. We define a few morphisms:

$$\begin{aligned}
 & \begin{array}{ccc}
 {}^a_qV_q & {}^b_qV_q & \xrightarrow{a;b} {}^{a+b}_qV_q \\
 b_I & b_J \xrightarrow{!} & \begin{cases} q^{jJ < Ij} b_{I \cup J} & \text{if } I \setminus J = \emptyset, \\ 0 & \text{else;} \end{cases}
 \end{array} & \begin{array}{ccc}
 {}^{a+b}_qV_q & \xrightarrow{Y_{a;b}} & {}^a_qV_q & {}^b_qV_q \\
 b_K \xrightarrow{!} & \sum_{I \cup J = K} q^{I < Jj} b_I & b_J;
 \end{array} \\
 \\
 C(q) & \begin{array}{ccc}
 \begin{array}{c} \lceil_a \\ \hline \end{array} & \begin{array}{c} {}^a_qV_q \\ \hline \end{array} & \begin{array}{c} ({}^a_qV_q) \\ \hline \end{array} \\
 1 \xrightarrow{!} & \sum_{\#I=a} b_I & b_I;
 \end{array} & \begin{array}{ccc}
 ({}^a_qV_q) & \begin{array}{c} {}^a_qV_q \\ \hline \end{array} & \begin{array}{c} \setminus_a \\ \hline \end{array} C(q) \\
 f \times \xrightarrow{!} & f(x);
 \end{array} \\
 \\
 C(q) & \begin{array}{ccc}
 \begin{array}{c} \lceil_a \\ \hline \end{array} & \begin{array}{c} ({}^a_qV_q) \\ \hline \end{array} & \begin{array}{c} {}^a_qV_q \\ \hline \end{array} \\
 1 \xrightarrow{!} & \sum_{\#I=a} q^{jI < INj + jIN < Ij} b_I & b_I;
 \end{array} & \begin{array}{ccc}
 {}^a_qV_q & \begin{array}{c} ({}^a_qV_q) \\ \hline \end{array} & \begin{array}{c} \setminus_a \\ \hline \end{array} C(q) \\
 b_I & b_J \xrightarrow{!} & q^{iI < INj} b_I & b_J;
 \end{array}
 \end{aligned}$$

We should explain what $jJ < Ij$ and $jI < Jj$ mean here. If A and B are two subsets of an ordered set C , we define

$$jA < Bj := \#f(a; b) \geq A \cup B \text{ and } Bja < bg;$$

Using the Reshetikhin-Turaev functor one can interpret any MOY graph as a morphism in $U_q(\mathfrak{gl}_N)\text{-mod}$. Using identities (2.1) and (2.2) we can extend this interpretation to MOY graph with crossings.

A.3. SYMMETRIC MOY CALCULUS. — In this subsection we work in the full subcategory $U_q(\mathfrak{gl}_N)\text{-mod}_S$ of finite-dimensional $U_q(\mathfrak{gl}_N)$ -modules generated (as a monoidal category) by the modules $\text{Sym}_q^a V_q$ for a in \mathbb{N} and their duals. We define a few morphisms:

$$\begin{aligned}
 & \begin{array}{ccc}
 \text{Sym}_q^a V_q & \text{Sym}_q^b V_q & \xrightarrow{a;b} \text{Sym}_q^{a+b} V_q \\
 b_I & b_J \xrightarrow{!} & q^{jJ < Ij} b_{I \cup J};
 \end{array} & \begin{array}{ccc}
 \text{Sym}_q^{a+b} V_q & \xrightarrow{Y_{a;b}} & \text{Sym}_q^a V_q & \text{Sym}_q^b V_q \\
 b_K \xrightarrow{!} & \sum_{I \cup J = K} [I; J]_q q^{jJ < Ij} b_I & b_J;
 \end{array} \\
 \\
 C(q) & \begin{array}{ccc}
 \begin{array}{c} \lceil_a \\ \hline \end{array} & \begin{array}{c} \text{Sym}_q^a V_q \\ \hline \end{array} & \begin{array}{c} (\text{Sym}_q^a V_q) \\ \hline \end{array} \\
 1 \xrightarrow{!} & \sum_{\#I=a} q^{iI < Jj} b_I & (b_I);
 \end{array} & \begin{array}{ccc}
 (\text{Sym}_q^a V_q) & \begin{array}{c} \text{Sym}_q^a V_q \\ \hline \end{array} & \begin{array}{c} \setminus_a \\ \hline \end{array} C(q) \\
 f \times \xrightarrow{!} & f(x);
 \end{array} \\
 \\
 C(q) & \begin{array}{ccc}
 \begin{array}{c} \lceil_a \\ \hline \end{array} & \begin{array}{c} (\text{Sym}_q^a V_q) \\ \hline \end{array} & \begin{array}{c} \text{Sym}_q^a V_q \\ \hline \end{array} \\
 1 \xrightarrow{!} & \sum_{\#I=a} q^{jI < INj + jIN < Ij} (b_I) & b_I;
 \end{array} & \begin{array}{ccc}
 \text{Sym}_q^a V_q & \begin{array}{c} (\text{Sym}_q^a V_q) \\ \hline \end{array} & \begin{array}{c} \setminus_a \\ \hline \end{array} C(q) \\
 b_I & (b_J) \xrightarrow{!} & q^{iI < INj} b_I & b_J;
 \end{array}
 \end{aligned}$$

We should explain what $jJ < Ij, jI < Jj$, and $[I; J]$ mean here. If A and B are two multi-subsets of an ordered set X , we define

$$jA < Bj := \prod_{x < y \in 2X} A(x)B(y); \quad [A; B] = \prod_{x \in 2X} \begin{bmatrix} A(x) + B(x) \\ A(x) \quad B(x) \end{bmatrix};$$

Using the Reshetikhin-Turaev functor one can interpret any MOY graph as a morphism in $U_q(\mathfrak{gl}_N)\text{-mod}_S$. Using identities (2.10) and (2.11) we can extend this interpretation to MOY graph with crossings. Note that this is *not* consistent with the exterior MOY calculus: the braiding has been changed for its inverse.

APPENDIX B. KOSZUL RESOLUTIONS OF POLYNOMIAL ALGEBRAS

In this appendix we recall the definition of the Koszul resolution of polynomial algebras. Then we describe a way to construct other differentials on the Koszul complex which anti-commute with the Koszul differential.

For an introduction to Koszul resolution see [Lod98, §3.4] and [Kas04, BLS18].

B.1. KOSZUL RESOLUTIONS. — Let R be an unitary commutative ring and V a free R -module of rank k . Let us fix an ordered basis (x_1, \dots, x_k) . We denote by A the symmetric tensor algebra SV and we will think of A as the polynomial algebra $R[x_1, \dots, x_n]$. The algebra V is naturally graded (we speak of H -grading): the non-zero elements of V seen in V have H -degree equal to 1. Let $C(A)$ be the A -module- A $A \oplus V \oplus A$. It inherits an H -grading from V . We consider the following endomorphism of A -module- A on $(C(A))$:

$$d: C(A) \rightarrow C(A)$$

$$1 \oplus v_1 \wedge \dots \wedge v_{i-1} \cdot \frac{1}{(i-1)!} \sum_{j=1}^i (-1)^{i+1} (v_j \wedge v_1 \wedge \dots \wedge \widehat{v}_i \wedge \dots \wedge v_{i-1} \cdot 1 - 1 \wedge v_1 \wedge \dots \wedge \widehat{v}_i \wedge \dots \wedge v_{i-1} \cdot v_j):$$

LEMMA B.1. — *The map d is a differential on $C(A)$, and $(C(A), d)$ is a projective resolution of A as an A -module- A . The complex $C(A)$ is called the Koszul resolution of A .*

Proof. — If V has dimension 1, then $A = R[x_1]$. It is easy to check, that the short exact sequence

$$0 \rightarrow A \xrightarrow{x_1} A \xrightarrow{x_1} A \xrightarrow{m} A \rightarrow 0$$

is exact. This proves that

$$0 \rightarrow A \xrightarrow{x_1} A \xrightarrow{x_1} A \rightarrow 0$$

is a projective resolution of A . If V is k dimensional, then $C(A)$ is by definition the complex

$$C(R[x_1]) \rightarrow C(R[x_2]) \rightarrow \dots \rightarrow C(R[x_k]);$$

and we deduce the result from the one-dimensional case.

REMARK B.2. — The vector space V could be graded, we speak of q -grading. In this case we suppose that the basis $(x_1; \dots; x_k)$ is homogeneous. The H -grading of $C(A)$ is not influenced by the q -grading. On the contrary, q -grading of V induces a q grading on A . Hence if all the x_i 's have positive q -degree, we have:

$$\text{rk}_q^R A = \prod_{i=1}^k \frac{1}{q^{\deg_q x_i}}.$$

B.2. AN HOMOTOPY EQUIVALENCE. — We consider the algebra $A := A_{1^k} = R[x_1; \dots; x_k]$. For consistency with the rest of the paper, we set the indeterminate x_i 's to have q -degree equal to 2. If $\underline{k} = (k^1; \dots; k^r)$ is a finite sequence of positive integers of level k , we set $A_{\underline{k}} := A^{\mathbb{S}_{\underline{k}}}$, where $\mathbb{S}_{\underline{k}}$ is equal to $\mathbb{S}_{k^1} \times \dots \times \mathbb{S}_{k^r}$ and acts naturally on the indeterminates $x_1; \dots; x_k$. Note that the algebras $A_{\underline{k}}$ can be thought of as polynomial algebras as well. One only need to consider some appropriate elementary symmetric polynomials.

Let us fix two finite sequences $\underline{k}_1 = (k_1^1; \dots; k_1^r)$ and $\underline{k}_2 = (k_2^1; \dots; k_2^s)$ of positive integers of levels k . We suppose furthermore, that \underline{k}_2 is obtained from \underline{k}_1 by merging two of consecutive elements of \underline{k}_1 . For instance:

$$\underline{k}_1 = (2; 3; 1; 1; 5; 4) \quad \text{and} \quad \underline{k}_2 = (2; 4; 1; 5; 4):$$

For simplicity of notations we will actually suppose that $\underline{k}_1 = (a; b)$ and $\underline{k}_2 = (a + b)$ with $a; b > 1$. In what follows we will be interested in $A^1 := A_{\underline{k}_1}$ and $A^2 := A_{\underline{k}_2}$.

Since A^2 is a sub-algebra of A^1 , A^1 can be consider as a A^2 -module- A^1 . As a module- A^1 it is free of rank 1, as a A^2 -module, it is free of rank $\binom{a+b}{a}$. Hence it is both a projective module- A^1 and a projective A^2 -module. This implies that

$$C^1 := A^1 \xrightarrow{A^1} C(A^1) \xrightarrow{A^1} C(A^1) \quad \text{and} \quad C^2 := C(A^2) \xrightarrow{A^2} A^1$$

are both projective resolutions of A^1 as A^2 -module- A^1 . Hence we know that these two complexes are homotopic. We denote by d^1 (resp. d^2) the differential of C^1 (resp. C^2).

We want to give an explicit homotopy equivalence of complexes of A^2 -module- A^1 $\psi : C^2 \rightarrow C^1$. We consider the vector space $V_1 := \langle hf_1; \dots; f_a; g_1; \dots; g_b \rangle_{A^1}$ and $V_2 := \langle he_1; \dots; e_{a+b} \rangle_{A^1}$. The element f_i (resp. g_i , resp. e_i) is meant to represent the i th elementary symmetric polynomials in the first a variables (resp. the last b variables, resp. $a + b$ variables).

Thanks to some standard argument of homological algebra (see e.g. [Bro82, Chap. 1, Lem. 7.3]) we know that if ψ is a chain map such that

$$(B.1) \quad \begin{array}{ccc} C^2 & \longrightarrow & A^1 \\ \psi \downarrow & & \downarrow \text{Id}_{A^1} \\ C^1 & \longrightarrow & A^1 \end{array}$$

commutes, then ψ is an homotopy equivalence. By definition we have:

$$C^1 := A^1 \xrightarrow{V_1} A^1 \quad \text{and} \quad C^2 := A^2 \xrightarrow{V_2} A^1:$$

Since C^2 is a free A^2 -module- A^1 with a basis given by elements of the form

$$1_{A^2} \wedge e_{i_1} \wedge \dots \wedge e_{i_r} \wedge 1_{A^1};$$

with $0 \leq i_1 < i_2 < \dots < i_r$, we only need to define δ^i on these elements. For $r = 0$, we define:

$$\delta^0(1_{A^2} \otimes 1_R \otimes 1_{A^1}) = 1_{A^1} \otimes 1_R \otimes 1_{A^1}.$$

For $r = 1$, we define:

$$\delta^1(1_{A^2} \otimes e_i \otimes 1_{A^1}) = \sum_{j=1}^i f_{i-j} g_j \otimes 1_{A^1} + 1_{A^1} \otimes f_j g_{i-j};$$

with the convention that the j th elementary polynomial in c variables is equal to 0 whenever $j > c$. For $r > 1$ we set:

$$\delta^i(1_{A^2} \otimes e_{i_1} \wedge \dots \wedge e_{i_r} \otimes 1_{A^1}) = \prod_{h=1}^r \delta^i(1_{A^2} \otimes e_{i_h} \otimes 1_{A^1})$$

and extend this map $A^2 \otimes (A^1)^{\text{opp}}$ -linearly. In the last formula, the space $A \otimes V \otimes A$ is endowed with the algebra structures given by:

$$(a_1 \otimes v_1 \otimes b_1) (a_2 \otimes v_2 \otimes b_2) = a_1 a_2 \otimes v_1 \wedge v_2 \otimes b_1 b_2.$$

Note however that the product is ordered and that δ^i is *not* an algebra morphism.

PROPOSITION B.3. — *The map δ^i is a morphism of complexes of A^2 -modules- A^1 such that the diagram (B.1) commutes.*

Proof. — Thanks to the definition of $\delta^i(1_{A^2} \otimes 1_R \otimes 1_{A^1})$, the diagram (B.1) obviously commutes. It remains to show that δ^i is indeed a chain map. Thanks to the $A^2 \otimes (A^1)^{\text{opp}}$ -linearity, it is enough to consider elements of the form

$$1_{A^2} \otimes e_{i_1} \wedge \dots \wedge e_{i_r} \otimes 1_{A^1}.$$

Let us consider the case $r = 1$. We have

$$\delta^1(d^2(1 \otimes e_i \otimes 1)) = e_i \otimes 1_R \otimes 1 \otimes 1 \otimes 1_R \otimes e_i$$

and

$$\begin{aligned} d^1(\delta^1(1 \otimes e_i \otimes 1)) &= \sum_{j=1}^i (f_{i-j} g_j \otimes 1_R \otimes 1 \otimes f_{i-j} \otimes 1_R \otimes g_j) \\ &\quad + \sum_{j=1}^i (f_j \otimes 1_R \otimes g_{i-j} \otimes 1 \otimes 1_R \otimes f_j g_{i-j}) \\ &= \sum_{j=0}^i (f_j g_{i-j} \otimes 1_R \otimes 1 \otimes 1 \otimes 1_R \otimes f_j g_{i-j}) = e_i \otimes 1_R \otimes 1 \otimes 1 \otimes 1_R \otimes e_i. \end{aligned}$$

If $r > 1$, we have:

$$\begin{aligned} d^1(\delta^i(1 \otimes e_{i_1} \wedge \dots \wedge e_{i_r} \otimes 1)) &= d^1\left(\prod_{h=1}^r \delta^i(1 \otimes e_{i_h} \otimes 1)\right) \\ &= \sum_{k=1}^r (1)^{k+1} \left(\prod_{h=1}^{k-1} \delta^i(1 \otimes e_{i_h} \otimes 1)\right) d^1(\delta^i(1 \otimes e_{i_k} \otimes 1)) \left(\prod_{h=k+1}^r \delta^i(1 \otimes e_{i_h} \otimes 1)\right) \\ &= \sum_{k=1}^r (1)^{k+1} \left(\prod_{h=1}^{k-1} \delta^i(1 \otimes e_{i_h} \otimes 1)\right) (\delta^i(d^2(1 \otimes e_{i_k} \otimes 1))) \left(\prod_{h=k+1}^r \delta^i(1 \otimes e_{i_h} \otimes 1)\right) \\ &= \delta^i(d^2(1 \otimes e_{i_1} \wedge \dots \wedge e_{i_r} \otimes 1)). \end{aligned}$$

B.3. AN ADDITIONAL DIFFERENTIAL

NOTATION B.4. — Suppose $D : A \rightarrow A$ is a derivation on A which lets $A_{\underline{k}}$ stable for any finite sequence of positive \underline{k} integers of level k . We consider the endomorphisms $\delta_{\underline{k}}$ of $C(A_{\underline{k}})$ given by:

$$(B.2) \quad \delta_{\underline{k}} : C(A_{\underline{k}}) \rightarrow C(A_{\underline{k}}) \\ (v_1 \wedge \dots \wedge v_{i-1} \wedge v_{i+1} \wedge \dots \wedge v_r) \mapsto \sum_{i=1}^r (-1)^{i+1} D(v_i) v_1 \wedge \dots \wedge \widehat{v_i} \wedge \dots \wedge v_r$$

Both these maps are H -homogeneous of degree -1 . For simplifying notations we denote the maps on $C(A^1)$ and $C(A^2)$ by δ^1 and δ^2 .

If we fix a positive integer N and if $R = R_N$, an example of such a derivation is given by

$$D^N : A_{1^k} \rightarrow A_{1^k} \\ P(x_1, \dots, x_k) \mapsto \sum_{i=1}^k \prod_{j=1}^N (x_i - T_j)_{@x_i} P(x_1, \dots, x_k)$$

The maps defined with D^N by the formula (B.2) are denoted $d_{\underline{k}}^N$. These are precisely the differentials considered in Section 5.2.2.

LEMMA B.5. — For $i = 1, 2$, the map δ^i anti-commutes with d^i and is a differential on $C(A^i)$.

Proof. — The fact that δ^1 and δ^2 are differentials follows as usual from the signs which forces every square to anti-commute. To check that d^i and δ^i anti-commute is an easy computation. We assume $i = 1$. We have:

$$\begin{aligned} & \delta^1(d^1(v_1 \wedge \dots \wedge v_r)) \\ &= \sum_{1 \leq i_1 < i_2 \leq r} (-1)^{i_1+i_2+2} (v_{i_1} D(v_{i_2}) v_1 \wedge \dots \wedge \widehat{v_{i_1}} \wedge \dots \wedge \widehat{v_{i_2}} \wedge \dots \wedge v_r) \\ & \quad - D(v_{i_2}) v_1 \wedge \dots \wedge \widehat{v_{i_1}} \wedge \dots \wedge \widehat{v_{i_2}} \wedge \dots \wedge v_r - v_{i_1} \\ & \quad - D(v_{i_1}) v_{i_2} v_1 \wedge \dots \wedge \widehat{v_{i_1}} \wedge \dots \wedge \widehat{v_{i_2}} \wedge \dots \wedge v_r \\ & \quad + D(v_{i_2}) v_1 \wedge \dots \wedge \widehat{v_{i_1}} \wedge \dots \wedge \widehat{v_{i_2}} \wedge \dots \wedge v_r - v_{i_1}) \end{aligned}$$

and

$$\begin{aligned} & d^1(\delta^1(v_1 \wedge \dots \wedge v_r)) \\ &= \sum_{1 \leq i_1 < i_2 \leq r} (-1)^{i_1+i_2+2} (D(v_{i_1}) v_{i_2} v_1 \wedge \dots \wedge \widehat{v_{i_1}} \wedge \dots \wedge \widehat{v_{i_2}} \wedge \dots \wedge v_r) \\ & \quad - D(v_{i_1}) v_1 \wedge \dots \wedge \widehat{v_{i_1}} \wedge \dots \wedge \widehat{v_{i_2}} \wedge \dots \wedge v_r - v_{i_2} \\ & \quad - v_{i_1} D(v_{i_2}) v_1 \wedge \dots \wedge \widehat{v_{i_1}} \wedge \dots \wedge \widehat{v_{i_2}} \wedge \dots \wedge v_r \\ & \quad + D(v_{i_1}) v_1 \wedge \dots \wedge \widehat{v_{i_1}} \wedge \dots \wedge \widehat{v_{i_2}} \wedge \dots \wedge v_r - v_{i_2}) \\ &= \delta^1(d^1(v_1 \wedge \dots \wedge v_r)). \end{aligned}$$

We define $\delta : C^2 \rightarrow C^1$ to be the $A^2 = (A^1)^{opp}$ -linear map defined by:

$$(1 - e_{i_1} \wedge \dots \wedge e_{i_r}) = D \text{ Id}_V - \text{Id}_{A^1} (\delta^1(e_{i_1} \wedge \dots \wedge e_{i_r}))$$

Note that we have:

$$(1 - e_{i_1} \wedge \dots \wedge e_{i_n} - 1) = \sum_{k=1}^n \left(\prod_{h=1}^{k-1} (1 - e_{i_h} - 1) \right) (1 - e_{i_k} - 1) \left(\prod_{h=k+1}^n (1 - e_{i_h} - 1) \right);$$

This follows from the fact that D is a derivation. We can write a (complicated) explicit formula for d^l :

$$(1 - e_{i_1} \wedge \dots \wedge e_{i_n} - 1) = \sum_{j_1+j_1^0=i_1} \sum_{k=0}^i \sum_{\substack{A+B=f_1+\dots+f_i \\ A=f_{a_1} < \dots < a_k g \\ B=f_{b_1} < \dots < b_k \cdot g}} (1 - 1)^{jB < Aj} \\ D(f_{j_{a_1}} \dots f_{j_{a_k}}) g_{j_{a_1}^0} \wedge \dots \wedge g_{j_{a_k}^0} \wedge f_{j_{b_1}} \wedge \dots \wedge f_{j_{b_k}} g_{j_{b_1}^0} \dots g_{j_{b_k}^0};$$

where $jB < Aj := \#f(b; a) \geq A - Bj$ and $b < ag$.

LEMMA B.6. — We have the following identity:

$$(1 - e_i - 1)^2 = d_2 + d_1;$$

Proof. — Suppose $i = 1$. On the one hand, we have

$$\begin{aligned} (1 - e_i - 1)^2 &= D(e_i - 1) = \sum_{j=1}^i (f_{i-j} D(g_j) - 1 + D(f_j) - 1) g_{i-j} \\ &= \sum_{j=0}^i D(f_j) g_{i-j} - 1 + f_j D(g_{i-j}) - 1 \\ &= \sum_{j=1}^i D(f_j) g_{i-j} - 1 + D(f_j) - 1) g_{i-j} \\ &= \sum_{j=1}^i D(f_j) g_{i-j} - 1 + \sum_{j=1}^i D(f_j) - 1) g_{i-j}; \end{aligned}$$

On the other hand, we have:

$$d^2(1 - e_i - 1) = (e_i - 1) - (1 - e_i) = 0$$

and

$$\begin{aligned} d^1(1 - e_i - 1) &= d^1 \left(\sum_{j=1}^i D(f_j) - g_{i-j} \right) \\ &= \sum_{j=1}^i D(f_j) g_{i-j} - 1 + D(f_j) - 1) g_{i-j} \\ &= (1 - e_i - 1)^2; \end{aligned}$$

Suppose now that $i > 1$. We have:

$$\begin{aligned} (1 - e_{i_1} \wedge \dots \wedge e_{i_n} - 1) &= \sum_{h=1}^n (1 - e_{i_h} - 1) \prod_{k=1}^{h-1} (1 - e_{i_k} - 1) \prod_{k=h+1}^n (1 - e_{i_k} - 1) \\ &= \sum_{h=1}^n (1 - e_{i_h} - 1) (d^1) \prod_{k=h+1}^n (1 - e_{i_k} - 1); \end{aligned}$$

On the other hand, we have:

$$\begin{aligned}
 & d^2(1 \otimes e_{i_1} \wedge \dots \wedge e_{i_j} \otimes 1) \\
 &= \left(\sum_{h=1}^j (1)^{h+1} \prod_{k=1}^{h-1} (1 \otimes e_{i_k} \otimes 1) d^2(1 \otimes e_{i_h} \otimes 1) \prod_{k=h+1}^j (1 \otimes e_{i_k} \otimes 1) \right) \\
 &= \sum_{h=1}^j \sum_{j=1}^{h-1} (1)^{h+1} \prod_{k=1}^{j-1} (1 \otimes e_{i_k} \otimes 1) (1 \otimes e_{i_j} \otimes 1) \\
 &\quad \prod_{k=j+1}^{h-1} (1 \otimes e_{i_k} \otimes 1) (d^2(1 \otimes e_{i_h} \otimes 1)) \prod_{k=h+1}^j (1 \otimes e_{i_k} \otimes 1) \\
 &\quad + \sum_{h=1}^j \sum_{j=h+1}^j (1)^{h+1} \prod_{k=1}^{h-1} (1 \otimes e_{i_k} \otimes 1) (d^2(1 \otimes e_{i_h} \otimes 1)) \\
 &\quad \prod_{k=h+1}^{j-1} (1 \otimes e_{i_k} \otimes 1) (1 \otimes e_{i_j} \otimes 1) \prod_{k=j+1}^j (1 \otimes e_{i_k} \otimes 1) \\
 &= \sum_{h=1}^j \sum_{j=1}^{h-1} (1)^{h+1} \prod_{k=1}^{j-1} (1 \otimes e_{i_k} \otimes 1) (1 \otimes e_{i_j} \otimes 1) \\
 &\quad \prod_{k=j+1}^{h-1} (1 \otimes e_{i_k} \otimes 1) d^1(1 \otimes e_{i_h} \otimes 1) \prod_{k=h+1}^j (1 \otimes e_{i_k} \otimes 1) \\
 &\quad + \sum_{h=1}^j \sum_{j=h+1}^j (1)^{h+1} \prod_{k=1}^{h-1} (1 \otimes e_{i_k} \otimes 1) d^1(1 \otimes e_{i_h} \otimes 1) \\
 &\quad \prod_{k=h+1}^{j-1} (1 \otimes e_{i_k} \otimes 1) (1 \otimes e_{i_j} \otimes 1) \prod_{k=j+1}^j (1 \otimes e_{i_k} \otimes 1)
 \end{aligned}$$

and

$$\begin{aligned}
 & d^1(1 \otimes e_{i_1} \wedge \dots \wedge e_{i_j} \otimes 1) \\
 &= d^1 \left(\sum_{h=1}^j (1)^{h+1} \prod_{k=1}^{h-1} (1 \otimes e_{i_k} \otimes 1) (1 \otimes e_{i_h} \otimes 1) \prod_{k=h+1}^j (1 \otimes e_{i_k} \otimes 1) \right) \\
 &= d^2((1 \otimes e_{i_1} \wedge \dots \wedge e_{i_j} \otimes 1) \\
 &\quad + \sum_{h=1}^j (1)^{h+1} \prod_{k=1}^{h-1} (1 \otimes e_{i_h} \otimes 1) d^1(1 \otimes e_{i_h} \otimes 1) \prod_{k=h+1}^j (1 \otimes e_{i_k} \otimes 1) \\
 &= d^2(1 \otimes e_{i_1} \wedge \dots \wedge e_{i_j} \otimes 1) \\
 &+ \sum_{h=1}^j (1)^{h+1} \prod_{k=1}^{h-1} (1 \otimes e_{i_k} \otimes 1) (1 \otimes e_{i_h} \otimes 1) \prod_{k=h+1}^j (1 \otimes e_{i_k} \otimes 1) \\
 &= d^2(1 \otimes e_{i_1} \wedge \dots \wedge e_{i_j} \otimes 1) + (1 \otimes e_{i_1} \wedge \dots \wedge e_{i_j} \otimes 1):
 \end{aligned}$$

APPENDIX C. A PINCH OF ALGEBRAIC GEOMETRY

Algebraic geometry has been a very useful guideline for the definition of the (exterior) Khovanov-Rozansky homologies. The exterior \mathfrak{sl}_N -invariant of the unknot labeled by k is equal to $\left[\begin{smallmatrix} N \\ k \end{smallmatrix} \right]$, which is the graded Euler characteristic of the cohomology ring of $\mathrm{Gr}_{\mathbb{C}}(k; N)$ the Grassmannian variety of k -spaces in \mathbb{C}^N up to an overall grading shift. Indeed the Frobenius algebra associated with the unknot labeled by k in the Khovanov-Rozansky homology is isomorphic to $H^*(\mathrm{Gr}_{\mathbb{C}}(k; N))$. This can be extended to an equivariant⁽¹⁴⁾ setting see [Kra10a, RW17].

The symmetric \mathfrak{sl}_N invariant of the unknot with a label k is equal to $\left[\begin{smallmatrix} N+k \\ k \end{smallmatrix} \right]$. This is (up to an overall grading shift) the graded Euler characteristic of $\mathrm{Gr}_{\mathbb{C}}(k; k+N-1)$. However, this does not seem to be the correct point of view when categorifying the symmetric \mathfrak{sl}_N -invariant.⁽¹⁵⁾ Indeed, in an equivariant setting, one expect to have a natural action of $\mathbb{Q}[T_1; \dots; T_N]$ or $\mathbb{Q}[T_1; \dots; T_N]^{\mathfrak{S}_N}$ on the Frobenius algebra associated with the unknot labeled by k . Instead, we believe that it is better to consider the space $S^k(\mathrm{Gr}_{\mathbb{C}}(1; N))$ of collections of k lines (counted with multiplicity) in \mathbb{C}^N :

$$S^k(\mathrm{Gr}_{\mathbb{C}}(1; N)) = (\mathrm{Gr}_{\mathbb{C}}(1; N))^k / \mathfrak{S}_k.$$

There is a natural action of GL_N on this space and its (non-equivariant) cohomology ring is isomorphic⁽¹⁶⁾ to that of $\mathrm{Gr}_{\mathbb{C}}(k; k+N-1)$:

THEOREM C.1 ([ES02, Th. 2.4]). — *There is a birational map $f : (\mathrm{Gr}_{\mathbb{C}}(1; N))^k \rightarrow \mathrm{Gr}_{\mathbb{C}}(k; k+N-1)$ such that the correspondence induced by the graph Γ_f of f induces an isomorphism*

$$H(\Gamma_f) : H(\mathrm{Gr}_{\mathbb{C}}(k; k+N-1); \mathbb{Q}) \xrightarrow{\cong} H(S^k(\mathrm{Gr}_{\mathbb{C}}(1; N)); \mathbb{Q})$$

of graded \mathbb{Q} -algebras.

As stated before, we are interested in the equivariant version of the cohomology ring of $S^k(\mathrm{Gr}(1; N))$.

The group $G := GL_N$ acts naturally on $\mathrm{Gr}(1; N)$ and diagonally on $S^k(\mathrm{Gr}(1; N))$. Since $H^*(BG; \mathbb{Q}) \cong \mathbb{Q}[T_1; \dots; T_N]^{\mathfrak{S}_N} =: R_N$ as a ring, the equivariant cohomologies of both $\mathrm{Gr}(1; N)$ and $S^k(\mathrm{Gr}(1; N))$ have structures of graded R_N -algebras (the degrees of the variables T_i are all equal to 2). There exists a presentation of $H_G(\mathrm{Gr}(1; N))$:

LEMMA C.2 ([Ful07, Lect. 3, Ex. 1.2]). — *The R_N -algebra $H_G(\mathrm{Gr}(1; N); \mathbb{Q})$ is isomorphic to $R_N[x] / J_{N,1}$, where x has degree 2 and $J_{N,1}$ is the ideal of $R_N[x]$ generated by $\prod_{i=1}^N (x - T_i)$.*

It follows immediately that $H_G(\mathrm{Gr}(1; N); \mathbb{Q})$ is a free graded R_N -module and that the family $(1; x; \dots; x^{N-1})$ forms an homogeneous R_N -basis of this space. The

⁽¹⁴⁾This motivates the term "equivariant" \mathfrak{sl}_N -homology.

⁽¹⁵⁾This remark is due to François Costantino.

⁽¹⁶⁾Note that if $N = 2$, these two varieties are actually isomorphic.

space $M_{N;k}$ considered in Section 5 is isomorphic to

$$\mathrm{Sym}^k(H_G(\mathrm{Gr}(1; N); \mathbb{Q}));$$

We believe that this is the same as

$$H_G(\mathrm{Sym}^k(\mathrm{Gr}(1; N); \mathbb{Q}))$$

but could not locate such a statement in the literature.

REFERENCES

- [BN05] D. BAR-NATAN – “Khovanov’s homology for tangles and cobordisms”, *Geom. Topol.* **9** (2005), p. 1443–1499.
- [BW08] A. BELIAKOVA & S. WEHRLI – “Categorification of the colored Jones polynomial and Rasmussen invariant of links”, *Canad. J. Math.* **60** (2008), no. 6, p. 1240–1266.
- [BLS18] R. BERGER, T. LAMBRE & A. SOLOTAR – “Koszul calculus”, *Glasgow Math. J.* **60** (2018), no. 2, p. 361–399, Addendum: *Ibid.* **61** (2019), no. 1, p. 249.
- [Bla10] C. BLANCHET – “An oriented model for Khovanov homology”, *J. Knot Theory Ramifications* **19** (2010), no. 2, p. 291–312.
- [BHMV95] C. BLANCHET, N. HABEGGER, G. MASBAUM & P. VOGEL – “Topological quantum field theories derived from the Kauffman bracket”, *Topology* **34** (1995), no. 4, p. 883–927.
- [Bro82] K. S. BROWN – *Cohomology of groups*, Graduate Texts in Math., vol. 87, Springer-Verlag, New York-Berlin, 1982.
- [Cau15] S. CAUTIS – “Clasp technology to knot homology via the affine Grassmannian”, *Math. Ann.* **363** (2015), no. 3-4, p. 1053–1115.
- [Cau17] ———, “Remarks on coloured triply graded link invariants”, *Algebraic Geom. Topol.* **17** (2017), no. 6, p. 3811–3836.
- [CK08a] S. CAUTIS & J. KAMNITZER – “Knot homology via derived categories of coherent sheaves. I. The $\mathrm{sl}(2)$ -case”, *Duke Math. J.* **142** (2008), no. 3, p. 511–588.
- [CK08b] ———, “Knot homology via derived categories of coherent sheaves. II: sl_m case”, *Invent. Math.* **174** (2008), no. 1, p. 165–232.
- [CKW09] A. CONCA, C. KRATTENTHALER & J. WATANABE – “Regular sequences of symmetric polynomials”, *Rend. Sem. Mat. Univ. Padova* **121** (2009), p. 179–199.
- [CH15] B. COOPER & M. HOGANCAMP – “An exceptional collection for Khovanov homology”, *Algebraic Geom. Topol.* **15** (2015), no. 5, p. 2659–2707.
- [CK12] B. COOPER & V. KRUSHKAL – “Categorification of the Jones-Wenzl projectors”, *Quantum Topol.* **3** (2012), no. 2, p. 139–180.
- [Dow17] N. DOWLIN – “A categorification of the HOMFLY-PT polynomial with a spectral sequence to knot Floer homology”, 2017, arXiv: 1703.01401.
- [EST17] M. EHRIG, C. STROPPEL & D. TUBBENHAUER – “The Blanchet-Khovanov algebras”, in *Categorification and higher representation theory*, Contemp. Math., vol. 683, American Mathematical Society, Providence, RI, 2017, p. 183–226.
- [ETW18] M. EHRIG, D. TUBBENHAUER & P. WEDRICH – “Functoriality of colored link homologies”, *Proc. London Math. Soc. (3)* **117** (2018), no. 5, p. 996–1040.
- [ES02] E. J. ELIZONDO & V. SRINIVAS – “Some remarks on Chow varieties and Euler-Chow series”, *J. Pure Appl. Algebra* **166** (2002), no. 1, p. 67–81.
- [FSS12] I. FRENKEL, C. STROPPEL & J. SUSSAN – “Categorifying fractional Euler characteristics, Jones-Wenzl projectors and $3j$ -symbols”, *Quantum Topol.* **3** (2012), no. 2, p. 181–253.
- [Ful07] W. FULTON – “Equivariant cohomology in algebraic geometry”, Eilenberg lectures, 2007, <https://people.math.osu.edu/anderson.2804/eilenberg/>.
- [GGS18] E. GORSKY, S. GUKOV & M. STOŠIĆ – “Quadruply-graded colored homology of knots”, *Fund. Math.* **243** (2018), no. 3, p. 209–299.
- [Kas04] C. KASSEL – “Homology and cohomology of associative algebras — A concise introduction to cyclic homology”, Notes of a course given in the Advanced School on Non-commutative Geometry at ICTP, 2004, 37 pages, <http://www-irma.u-strasbg.fr/~kassel/Kassel-ICTPnotes2004.pdf>.

- [Kau13] L. H. KAUFFMAN – *Knots and physics*, 4th ed., World Scientific, Singapore, 2013.
- [Kho04] M. KHOVANOV – “ $sl(3)$ link homology”, *Algebraic Geom. Topol.* **4** (2004), p. 1045–1081.
- [Kho05] ———, “Categorifications of the colored Jones polynomial”, *J. Knot Theory Ramifications* **14** (2005), no. 1, p. 111–130.
- [Kho07] ———, “Triply-graded link homology and Hochschild homology of Soergel bimodules”, *Internat. J. Math.* **18** (2007), no. 8, p. 869–885.
- [KR08] M. KHOVANOV & L. ROZANSKY – “Matrix factorizations and link homology. II”, *Geom. Topol.* **12** (2008), no. 3, p. 1387–1425.
- [Kra10a] D. KRASNER – “Equivariant $sl(n)$ -link homology”, *Algebraic Geom. Topol.* **10** (2010), no. 1, p. 1–32.
- [Kra10b] ———, “Integral HOMFLY-PT and $sl(n)$ -link homology”, *Internat. J. Math. Math. Sci.* (2010), article ID 896879 (25 pages).
- [Lan02] S. LANG – *Algebra*, third ed., Graduate Texts in Math., vol. 211, Springer-Verlag, New York, 2002.
- [LQR15] A. D. LAUDA, H. QUEFFELEC & D. E. V. ROSE – “Khovanov homology is a skew Howe 2-representation of categorified quantum sl_m ”, *Algebraic Geom. Topol.* **15** (2015), no. 5, p. 2517–2608.
- [Lod98] J.-L. LODAY – *Cyclic homology*, second ed., Grundlehren Math. Wiss., vol. 301, Springer-Verlag, Berlin, 1998.
- [Lus10] G. LUSZTIG – *Introduction to quantum groups*, Modern Birkhäuser Classics, Birkhäuser/Springer, New York, 2010, Reprint of the 1994 edition.
- [Mac15] I. G. MACDONALD – *Symmetric functions and Hall polynomials*, second ed., Oxford Classic Texts in the Physical Sciences, The Clarendon Press, Oxford University Press, New York, 2015.
- [MSV09] M. MACKAAY, M. STOŠIĆ & P. VAZ – “ $sl(N)$ -link homology ($N > 4$) using foams and the Kapustin-Li formula”, *Geom. Topol.* **13** (2009), no. 2, p. 1075–1128.
- [MSV11] ———, “The 1;2-coloured HOMFLY-PT link homology”, *Trans. Amer. Math. Soc.* **363** (2011), no. 4, p. 2091–2124.
- [MW18] M. MACKAAY & B. WEBSTER – “Categorified skew Howe duality and comparison of knot homologies”, *Adv. Math.* **330** (2018), p. 876–945.
- [MS09] V. MAZORCHUK & C. STROPPEL – “A combinatorial approach to functorial quantum sl_k knot invariants”, *Amer. J. Math.* **131** (2009), no. 6, p. 1679–1713.
- [MOY98] H. MURAKAMI, T. OHTSUKI & S. YAMADA – “Homfly polynomial via an invariant of colored plane graphs”, *Enseign. Math. (2)* **44** (1998), no. 3-4, p. 325–360.
- [QR16] H. QUEFFELEC & D. E. V. ROSE – “The sl_n foam 2-category: a combinatorial formulation of Khovanov-Rozansky homology via categorical skew Howe duality”, *Adv. Math.* **302** (2016), p. 1251–1339.
- [QR18] ———, “Sutured annular Khovanov-Rozansky homology”, *Trans. Amer. Math. Soc.* **370** (2018), no. 2, p. 1285–1319.
- [QRS18] H. QUEFFELEC, D. E. V. ROSE & A. SARTORI – “Annular evaluation and link homology”, 2018, arXiv: 1802.04131.
- [Ras15] J. RASMUSSEN – “Some differentials on Khovanov-Rozansky homology”, *Geom. Topol.* **19** (2015), no. 6, p. 3031–3104.
- [RW17] L.-H. ROBERT & E. WAGNER – “A closed formula for the evaluation of sl_n -foams”, 2017, To appear in *Quantum Topology*, arXiv: 1702.04140.
- [RT16] D. E. V. ROSE & D. TUBBENHAUER – “Symmetric webs, Jones-Wenzl recursions, and q -Howe duality”, *Internat. Math. Res. Notices* **2016** (2016), no. 17, p. 5249–5290.
- [RW16] D. E. V. ROSE & P. WEDRICH – “Deformations of colored sl_N link homologies via foams”, *Geom. Topol.* **20** (2016), no. 6, p. 3431–3517.
- [Rou17] R. ROUQUIER – “Khovanov-Rozansky homology and 2-braid groups”, in *Categorification in geometry, topology, and physics*, Contemp. Math., vol. 684, American Mathematical Society, Providence, RI, 2017, p. 147–157.
- [Roz14] L. ROZANSKY – “An infinite torus braid yields a categorified Jones-Wenzl projector”, *Fund. Math.* **225** (2014), no. 1, p. 305–326.

- [ST19] A. SARTORI & D. TUBBENHAUER – “Webs and q -Howe dualities in types BCD”, *Trans. Amer. Math. Soc.* **371** (2019), no. 10, p. 7387–7431.
- [Soe92] W. SOERTEL – “The combinatorics of Harish-Chandra bimodules”, *J. reine angew. Math.* **429** (1992), p. 49–74.
- [Sto08] M. STOŠIĆ – “Hochschild homology of certain Soergel bimodules”, 2008, arXiv: 0810. 3578.
- [Str04] C. STROPPEL – “A structure theorem for Harish-Chandra bimodules via coinvariants and Golod rings”, *J. Algebra* **282** (2004), no. 1, p. 349–367.
- [SS14] C. STROPPEL & J. SUSSAN – “Categorified Jones-Wenzl projectors: a comparison”, in *Perspectives in representation theory*, Contemp. Math., vol. 610, American Mathematical Society, Providence, RI, 2014, p. 333–351.
- [Sus07] J. SUSSAN – “Category \mathcal{O} and $\mathfrak{sl}(k)$ link invariants”, PhD Thesis, Yale University, Ann Arbor, MI, 2007, arXiv: math/0701045.
- [TVW17] D. TUBBENHAUER, P. VAZ & P. WEDRICH – “Super q -Howe duality and web categories”, *Algebraic Geom. Topol.* **17** (2017), no. 6, p. 3703–3749.
- [Vaz08] P. VAZ – “A categorification of the quantum $\mathfrak{sl}(n)$ -link polynomials using foams”, PhD Thesis, Universidade do Algarve, 2008, arXiv: 0807. 2658.
- [Web17] B. WEBSTER – *Knot invariants and higher representation theory*, Mem. Amer. Math. Soc., vol. 250, no. 1191, American Mathematical Society, Providence, RI, 2017.
- [WW17] B. WEBSTER & G. WILLIAMSON – “A geometric construction of colored HOMFLYPT homology”, *Geom. Topol.* **21** (2017), no. 5, p. 2557–2600.
- [Wed19] P. WEDRICH – “Exponential growth of colored HOMFLY-PT homology”, *Adv. Math.* **353** (2019), p. 471–525.
- [Wil11] G. WILLIAMSON – “Singular Soergel bimodules”, *Internat. Math. Res. Notices* **2011** (2011), no. 20, p. 4555–4632.
- [Wu13] H. WU – “Colored Morton-Franks-Williams inequalities”, *Internat. Math. Res. Notices* **2013** (2013), no. 20, p. 4734–4757.
- [Wu14] ———, “A colored $\mathfrak{sl}(N)$ homology for links in S^3 ”, *Dissertationes Math. (Rozprawy Mat.)* **499** (2014), p. 1–217.
- [Yon11] Y. YONEZAWA – “Quantum $(\mathfrak{sl}_n; \wedge V_n)$ link invariant and matrix factorizations”, *Nagoya Math. J.* **204** (2011), p. 69–123.

Manuscript received 13th July 2018

accepted 16th March 2020

LOUIS-HADRIEN ROBERT, Université de Genève
2–4 rue du lièvre, 1227 Genève, Switzerland
Cogitamus Laboratory
E-mail : louis-hadrien.robert@uni.ge.ch
Url : <https://www.uni.ge.ch/math/folks/robert/>

EMMANUEL WAGNER, Univ Paris Diderot, IMJ-PRG, UMR 7586 CNRS
F-75013, Paris, France
Université de Bourgogne Franche-Comté, IMB, UMR 5584
21000 Dijon, France
Cogitamus Laboratory
E-mail : emmanuel.wagner@imj-prg.fr
Url : <http://wagner.perso.math.cnrs.fr/index.html>