



# Journal de l'École polytechnique

## *Mathématiques*

Camille LAURENT & Matthieu LÉAUTAUD

**On uniform observability of gradient flows in the vanishing viscosity limit**

Tome 8 (2021), p. 439-506.

[http://jep.centre-mersenne.org/item/JEP\\_2021\\_\\_8\\_\\_439\\_0](http://jep.centre-mersenne.org/item/JEP_2021__8__439_0)

© Les auteurs, 2021.

*Certains droits réservés.*



Cet article est mis à disposition selon les termes de la licence  
LICENCE INTERNATIONALE D'ATTRIBUTION CREATIVE COMMONS BY 4.0.  
<https://creativecommons.org/licenses/by/4.0/>

L'accès aux articles de la revue « Journal de l'École polytechnique — Mathématiques » (<http://jep.centre-mersenne.org/>), implique l'accord avec les conditions générales d'utilisation (<http://jep.centre-mersenne.org/legal/>).

Publié avec le soutien  
du Centre National de la Recherche Scientifique



Publication membre du  
Centre Mersenne pour l'édition scientifique ouverte  
[www.centre-mersenne.org](http://www.centre-mersenne.org)

## ON UNIFORM OBSERVABILITY OF GRADIENT FLOWS IN THE VANISHING VISCOSITY LIMIT

BY CAMILLE LAURENT & MATTHIEU LÉAUTAUD

**ABSTRACT.** — We consider a transport equation by a gradient vector field with a small viscous perturbation  $\varepsilon \geq 0$ . We study uniform observability (resp. controllability) properties in the (singular) vanishing viscosity limit  $\varepsilon \rightarrow 0^+$ , that is, the possibility of having a uniformly bounded observation constant (resp. control cost). We prove with a series of examples that in general, the minimal time for uniform observability may be much larger than the minimal time needed for the observability of the limit equation  $\varepsilon = 0$ . We also prove that the two minimal times coincide for positive solutions. The proofs rely on a semiclassical reformulation of the problem together with (a) Agmon estimates concerning the decay of eigenfunctions in the classically forbidden region [HS84] (b) fine estimates of the kernel of the semiclassical heat equation [LY86].

**RÉSUMÉ** (Sur l'observabilité uniforme des flots de gradient dans la limite de viscosité évanescence)

Nous considérons l'équation de transport par un champ de gradient avec une petite perturbation visqueuse  $\varepsilon \geq 0$ . Nous étudions la propriété d'observabilité (resp. de contrôlabilité) uniforme dans la limite (singulière) de viscosité évanescence  $\varepsilon \rightarrow 0^+$ , c'est-à-dire la possibilité d'avoir une constante d'observabilité (resp. un coût du contrôle) uniforme. Nous prouvons avec une série d'exemples que le temps minimal pour l'observabilité uniforme peut être bien plus grand que le temps minimal pour l'équation limite  $\varepsilon = 0$ . Nous montrons aussi que les deux temps minimaux coïncident pour les solutions positives. Les preuves reposent sur une reformulation semiclassique du problème ainsi que (a) des estimées d'Agmon de décroissance des fonctions propres dans la zone classiquement interdite [HS84] (b) des estimées fines du noyau de la chaleur semiclassique [LY86].

### CONTENTS

1. Introduction and main results.....	440
2. Preliminaries: duality, conjugation of gradient flows and (GCC).....	454
3. General lower bounds without geometric assumption.....	461
4. Surfaces of revolution.....	469
5. Uniform time of observability for positive solutions.....	488
Appendix. About the distances.....	499
References.....	503

**MATHEMATICAL SUBJECT CLASSIFICATION (2020).** — 93B07, 93B05, 35B25, 35F05, 35K05, 93C73.

**KEYWORDS.** — Transport equation, gradient flow, vanishing viscosity limit, parabolic equation, minimal control time, semiclassical Schrödinger operator.

The authors are partially supported by the Agence Nationale de la Recherche under grants SRGI ANR-15-CE40-0018 (for C.L.), SALVE ANR-19-CE40-0004 (for M.L.) and ISDEEC ANR-16-CE40-0013 (for C.L. and M.L.).

## 1. INTRODUCTION AND MAIN RESULTS

1.1. INTRODUCTION. — Given a smooth connected compact manifold  $M$  without boundary (the case of a bounded domain of  $\mathbb{R}^n$  is also discussed in Section 1.3 below), a smooth real valued vector field  $X$  on  $M$  and a real valued potential  $q(x)$ , we consider the question of observability/detectability for the autonomous transport equation

$$(1.1) \quad \begin{cases} (\partial_t + X + q)u = 0, & \text{in } \mathbb{R} \times M, \\ u|_{t=0} = u_0, & \text{on } M, \end{cases}$$

from an observation (open) set  $\omega \subset M$  through the time interval  $(0, T)$ . More precisely, the question is whether there exists a constant  $C_0 = C_0(T, \omega) > 0$  such that

$$(1.2) \quad C_0^2 \int_0^T \int_\omega |ju(t, x)|^2 ds(x) dt > ku(T)k_{L^2(M)}^2,$$

for all  $u_0 \in L^2(M)$  and  $u$  solution of (1.1).

Here,  $ds(x)$  denotes any positive density measure<sup>(1)</sup> on  $M$ , and the  $L^2$  norm is defined accordingly. The observability question (1.2) is naturally solved by introducing an appropriate Geometric Control Condition (recall  $\partial M = \emptyset$ ): we say that  $(M, X, \omega, T)$  satisfies (GCC) if for all  $x \in M$ , there is  $t \in (0, T)$  such that  $\phi_{-t}(x) \in \omega$ , where  $(\phi_t)_{t \in \mathbb{R}}$  denotes the flow of  $X$  (see Section 2.3 for precise statements and proofs). We also say that  $(M, X, \omega)$  satisfies (GCC) if  $(M, X, \omega, T)$  does for some  $T > 0$ ; and if so, we denote by  $T_{\text{GCC}}(M, X, \omega)$  the infimum of times for which  $(M, X, \omega, T)$  satisfies (GCC).

On the other hand, endowing  $M$  with a Riemannian metric  $g$ , one may want to investigate the observability question for the viscously damped transport equation:

$$(1.3) \quad \begin{cases} (\partial_t + X + q + \varepsilon \Delta_g)u = 0, & \text{in } \mathbb{R}^+ \times M, \\ u|_{t=0} = u_0, & \text{on } M, \end{cases}$$

from the same observation set  $(0, T) \times \omega$ . The question is whether there exists a constant  $C_0(T, \varepsilon) > 0$  such that

$$(1.4) \quad C_0(T, \varepsilon)^2 \int_0^T \int_\omega |ju(t, x)|^2 ds(x) dt > ku(T)k_{L^2(M)}^2,$$

for all  $u_0 \in L^2(M)$  and  $u$  solution of (1.3),

(and one may then choose the Riemannian volume density  $ds(x) = d\text{Vol}_g(x)$  without changing the problem). For fixed  $\varepsilon > 0$ , Equation (1.3) is of parabolic type and the observability inequality (1.4) is known to hold for any open set  $\omega \subsetneq M$  and  $T > 0$ , see [F196] (see also [LR95] and its variant in [Léa10]). Of course, in such results, the

<sup>(1)</sup>See e.g. [Lee13, Chap. 16 p. 427]: given a local chart  $(U_\kappa, \kappa)$  of  $M$ , we have  $\int_{U_\kappa} u ds = \int_{\kappa(U_\kappa)} u \kappa^{-1}(y) \varphi^\kappa(y) dy$  for an appropriate smooth positive function  $\varphi^\kappa$ , and for any  $u \in C_c^0(U_\kappa)$ .

observability constant  $C_0(T, \varepsilon)$  in (1.4) depends *a priori* on  $\varepsilon$ . For many different reasons (some of them described in Section 1.2 below), it is interesting to investigate the behavior of the observability constant  $C_0(T, \varepsilon)$  in the vanishing viscosity limit  $\varepsilon \rightarrow 0^+$ . This problem was first studied in the one-dimensional setting by Coron and Guerrero in [CG05], and later extended to any dimension by Guerrero and Lebeau [GL07]. Their main result in this direction can be formulated (in the present geometric context, see the remark preceding Proposition 5.7) as follows.

**THEOREM 1.1** (Guerrero-Lebeau [GL07]). — *Given an open set  $\omega \subset M$ , the following two results hold.*

– [GL07, Th.1] *Assume  $(M, X, \bar{\omega}, T)$  does not satisfy (GCC). Then there is  $C, \varepsilon_0 > 0$  such that any constant  $C_0(T, \varepsilon)$  in (1.4) satisfies  $C_0(T, \varepsilon) > \exp(C/\varepsilon)$  for  $\varepsilon \in (0, \varepsilon_0)$ .*

– [GL07, Th.3] *Assume  $(M, X, \omega)$  satisfies (GCC). Then there is  $T_{\text{unif}}(\omega) > T_{\text{GCC}}(M, X, \omega)$  and  $K_0 > 0$  such that for all  $T > T_{\text{unif}}(\omega)$ , (1.4) holds with  $C_0(T, \varepsilon) \leq K_0$  for all  $\varepsilon \leq 1$ .*

Note that the results in [GL07] are even more general since time-dependent vector fields are allowed and the boundary-value problem is also considered (with Dirichlet boundary conditions). We also refer to our Proposition 1.12 below for a more precise lower bound of the constant  $C$  when (GCC) is not satisfied.

Note that if (1.4) holds for some  $T_0$  and constant  $C_0(T_0, \varepsilon)$ , then it also holds for all times  $T > T_0$  with the same constant  $C_0(T_0, \varepsilon)$ . In [GL07], the question of the minimal time  $T_{\text{unif}}(\omega)$ , more precisely defined by

$$T_{\text{unif}}(\omega) = \inf \left\{ T > 0 \text{ for which there exist } K_0, \varepsilon_0 > 0 \right. \\ \left. \text{such that (1.4) holds with } C_0(T, \varepsilon) \leq K_0 \text{ for all } \varepsilon \in (0, \varepsilon_0) \right\},$$

and its link with the minimal observation time  $T_{\text{GCC}}(M, X, \omega)$  associated to the limit problem (1.1) is left open. In particular, the formulation of the results in [GL07] (see e.g. Th.2 and the discussion thereafter in that reference) suggests the possible existence of a universal constant  $K > 1$  such that

$$(1.5) \quad T_{\text{unif}}(\omega) \leq K T_{\text{GCC}}(M, X, \omega).$$

The present article investigates this question in a very particular case, namely assuming the vector field  $X$  is a gradient vector field, i.e.,  $X = \nabla_g f$  for a function  $f \in W^{2,1}(M; \mathbb{R})$  (note that the gradient is taken with respect to the Riemannian metric  $g$ ). Hence, Equation (1.3) becomes

$$(1.6) \quad \begin{cases} (\partial_t - \nabla_g f - \nabla_g \cdot q - \varepsilon \Delta_g)u = 0, & \text{in } \mathbb{R}^+ \times M, \\ u|_{t=0} = u_0, & \text{on } M, \end{cases}$$

Here, given two vector fields  $Y_1$  and  $Y_2$ , we have denoted  $Y_1 \cdot Y_2 = g(Y_1, Y_2)$  or  $(Y_1 \cdot Y_2)(x) = Y_1(x) \cdot Y_2(x) = g_x(Y_1(x), Y_2(x))$  for all  $x \in M$ . We denote similarly

$\|Y\|_g = \sqrt{\langle Y, Y \rangle_g}$  the associated Riemannian norm of a vector (or a vector field). Note that the vector field  $r_g f$  is canonically identified with the derivation  $r_g f = r_g$ .

In this context, the first consequence of our main results can be (loosely) stated as follows.

**THEOREM 1.2**

(1) *There are geometries  $(M, g)$  such that for all  $\Lambda > 0$ , one can find  $f \in C^1(M)$  and  $\omega$  open such that  $(M, r_g f, \omega)$  satisfies (GCC) and  $T_{\text{unif}}(\omega) > \Lambda T_{\text{GCC}}(M, r_g f, \omega)$ .*

(2) *There are  $(M, f, X, \omega)$  such that for all  $\Lambda > 0$ , one can find a metric  $g_\Lambda$  on  $M$  such that*

- $X = r_{g_\Lambda} f$ ,
- $(M, X, \omega)$  satisfies (GCC),
- $T_{\text{unif}}(\omega) > \Lambda T_{\text{GCC}}(M, X, \omega)$ .

In particular, Theorem 1.2 states that there is no  $K$  such that (1.5) holds for all  $(M, X, \omega)$ .

The second item in Theorem 1.2 stresses the importance of the viscosity one chooses. Namely, with the same vector field  $X$ , changing the metric  $g$ , that is the viscous perturbation, may change the minimal uniform observability time. We also obtain related results for domains of  $\mathbb{R}^n$  (see Section 1.3.1).

Our second main result in this setting concerns the uniform observability of *positive* solutions to (1.6). Recall that nonnegative data  $u_0 > 0$  give rise to positive solutions to (1.6). We define  $C_0^+(T, \varepsilon)$  the observability constant for positive solutions, that is for which (1.2) holds for all  $u_0 > 0$ , and accordingly set

$$(1.7) \quad T_{\text{unif}}^+(\omega) = \inf \{ T > 0 \text{ for which there exist } K_0, \varepsilon_0 > 0 \text{ such that (1.4) holds} \\ \text{for all } u_0 > 0, \text{ with } C_0^+(T, \varepsilon) \leq K_0 \text{ for all } \varepsilon \in (0, \varepsilon_0) \}.$$

**THEOREM 1.3 (Positive solutions).** — *For all  $f \in C^3(M; \mathbb{R})$ , and  $\omega \subset M$  such that  $(M, r_g f, \omega)$  satisfies (GCC), we have  $T_{\text{unif}}^+(\omega) = T_{\text{GCC}}(M, r_g f, \omega)$ .*

As usual, these uniform observability/non-observability results can be reformulated in terms of uniform controllability/non-controllability statements for an adjoint controlled equation, see Section 2.1.1 below.

**1.2. BACKGROUND AND MOTIVATION.** — Uniform controllability problems for singular perturbations of partial differential equations already appeared in the reference book of Lions [Lio88, Chap. 3]. In the context of transport/heat equation in vanishing viscosity limit, this study was initiated by Coron and Guerrero on the 1D problem with constant speed in [CG05], where the authors make a conjecture on the minimal time needed to achieve uniform controllability. Then, the estimates on this minimal time have been improved successively in [Gla10, Lis12, Lis14, Lis15]. We also refer to the articles [Mün18, AM19b, AM19a] proposing numerical experiments to find the optimal minimal time. Such uniform control properties in singular limits are also addressed for vanishing dispersion in [GG08] and for vanishing dispersion and viscosity in [GG09].

Whereas the one-dimensional problem with a constant vector field has received a lot of attention in the past fifteen years, there are very few results in higher dimension or for non-constant vector fields. Besides [GL07] we are only aware of the results of [BP20a] and [BP20b] for the flat Laplace operator and the vector field  $\partial_{x_1}$ , with several boundary conditions.

Note that controllability problems for nonlinear conservation laws in vanishing viscosity have also been studied in [GG07], [Léa12], and [Mar14].

There are several motivations for studying the vanishing viscosity limit. A first motivation comes from the theory of conservation laws, for which the vanishing viscosity criterion is a selection principle for the physical (called entropy) solution, see [Kru70] or [Daf00, Chap. 6]. It is therefore very natural, when considering control problems for conservation laws, to study the cost of the viscosity, that is, to determine if known controllability properties for the hyperbolic equation are still valid for the model with small viscosity, and how the size of the control evolves as the viscosity approaches 0. So far the only known results in this directions seem to be [GG07] and [Léa12].

Another important motivation for studying singular limits in control problems is the seek of controllability properties for the perturbed system itself. This is well-illustrated by the papers [Cor96, CF96, Cha09, CMS20], where the authors investigate the Navier-Stokes system with Navier slip or slip-with-friction boundary conditions. They use a global controllability result for the inviscid equation (in this case, the Euler equation) to deduce global approximate controllability of the Navier-Stokes system.

On the other hand, the study of gradient fields naturally arises as the simplest dynamical situation among all vector fields. The importance of gradient vector fields with a vanishing viscosity coefficient also appears in theoretical physics and differential topology, through the Witten-Hellier-Sjöstrand theory [Wit82, HS85]. See e.g. the monographs [Hel88, CFKS87]. In that theory, the operator  $\nabla_g f - \nabla_g \varepsilon \Delta_g$  (and its analogues acting on forms) is conjugated to a particular semiclassical Schrödinger operator, namely

$$(1.8) \quad P_\varepsilon = -\varepsilon^2 \Delta_g + \frac{j \nabla_g f \nabla_g}{4} + \frac{\varepsilon}{2} \Delta_g f,$$

sometimes called the Witten Laplacian. Topological properties of the couple  $(M, f)$  (e.g. the Morse inequalities, linking the number of critical points of the Morse function  $f$  with the Betti numbers of  $M$ ) are deduced from spectral properties of the Witten Laplacian. We also refer to [DR20] for the understanding of other links between the spectral theories of the Witten Laplacian and the vector field  $\nabla_g f - \nabla_g$  (in appropriate spaces linked to the dynamics of the gradient flow), in the semiclassical limit  $\varepsilon \rightarrow 0^+$ .

Viscous perturbations of gradient dynamics also arise naturally in molecular dynamics. Indeed, in  $\mathbb{R}^n$ , the operator  $\nabla f - \nabla \varepsilon \Delta$  is the infinitesimal generator of the so-called overdamped Langevin process

$$dX_t = \nabla f(X_t) dt + \sqrt{2\varepsilon} dB_t,$$

where  $X_t \geq \mathbb{R}^n$  and  $(B_t)_{t>0}$  is a standard Brownian motion of dimension  $n$ . This stochastic process is a classical model in statistical physics. It is used in particular for the simulation of molecular dynamics at low temperature (proportional to  $\varepsilon$ ), see [Cha43, SM79]. The possible convergence to equilibrium, as well as the so-called metastability phenomenon are closely related to the low-lying eigenvalues (and associated eigenfunctions) of  $r f r - \varepsilon \Delta$ , or equivalently of the Witten Laplacian (1.8). For a very precise asymptotic study of the exponentially small eigenvalues of this operator, we refer e.g. to [HKN04, Mic19] in the case of a compact manifold and [HN06, LP10, DGLLPN20] in the case with boundary (see also the references therein).

We finally remark that the above-mentioned works concerning the Witten Laplacian mostly study the behavior of the bottom of the spectrum  $P_\varepsilon$  (thus linked to critical points of  $f$ ). In the present work, we rely on a similar conjugation.

**1.3. MAIN RESULTS.** — As already seen in the end of Section 1.1, the results of this article go in two different directions.

In a first part (Section 3), we prove some general lower bounds on the time  $T_{\text{unif}}(\omega)$  for a general class of domains and vector fields. This implies in particular that the quite natural idea that  $T_{\text{unif}}(\omega)$  is  $T_{\text{GCC}}(M, X, \omega)$  or even  $KT_{\text{GCC}}(M, X, \omega)$  for a universal constant  $K$  is false in general. This might be interpreted by the fact that in the vanishing viscosity limit, some strong oscillations can be responsible for concentration phenomena. The latter are not only described by the flow of  $X$ , but other parameters where an Agmon distance plays a crucial role. We also study (in Section 4) the particular case of surfaces of revolutions where we obtain refined lower bounds. This analysis also shows that the global geometry of the Riemannian manifold we consider has an effect on the vanishing viscosity limit. In particular, with a fixed vector field, we show that the choice of the Laplacian  $\Delta_g$  can change drastically the time  $T_{\text{unif}}(\omega)$  of uniform controllability. This shows definitely that the flow of the vector field is not the unique parameter defining  $T_{\text{unif}}(\omega)$ .

In a second part (Section 5), our results go exactly in the opposite direction, but for a specific class of solutions, namely positive solutions. As announced in Theorem 1.3, we prove that  $T_{\text{unif}}(\omega)^+ = T_{\text{GCC}}(M, r_g f, \omega)$  for positive solutions. This shows that the dynamics of positive solutions are actually well represented by the sole flow of  $r_g f$ .

In both cases, using the change of unknown  $v = e^{f/2\varepsilon} u$ , see Section 2.2, the problem is reduced (modulo lower order terms, and in weighted spaces) to observability of solutions of a semiclassical heat equation

$$\varepsilon \partial_t v - \varepsilon^2 \Delta_g v + V v = 0,$$

where  $V = j r_g f_g^2 / 4$ . Note that most of the results we obtain are of interest for this particular question as well.

**1.3.1. A general lower bound.** — The first family of results in this paper concern the general setting of a compact connected Riemannian manifold  $(M, g)$ , with or without boundary  $\partial M$ , and the associated internal/boundary observability question. Namely,

we consider the parabolic–transport problem with small viscosity  $\varepsilon > 0$  and Dirichlet boundary conditions:

$$(1.9) \quad \begin{cases} (\partial_t - X - q - \varepsilon \Delta_g)u = 0, & \text{in } (0, T) \cap \text{Int}(M), \\ u = 0 & \text{on } (0, T) \cap \partial M, \\ u|_{t=0} = u_0, & \text{in } M. \end{cases}$$

Moreover, we assume that the vector field  $X$  is a gradient vector field for the metric  $g$ , that is: there is  $f \in W^{2,1}(M)$  (at least) such that

$$X = \text{r}_g f - \text{r}_g.$$

For the Dirichlet problem (1.9) as well as for the case  $\partial M = \emptyset$  discussed in Section 1.1, one may discuss the behavior of the observability constant (and in particular its possible uniform boundedness in the limit  $\varepsilon \rightarrow 0^+$ ) in the internal observability inequality (1.4). Also, a boundary observability problem for (1.9) can be formulated as follows (see [GL07] and Section 2.1.2 below). Given  $\theta \in C^1(\partial M)$ , there exist a constant  $C_0(T, \varepsilon) > 0$  such that

$$(1.10) \quad C_0(T, \varepsilon)^2 \int_0^T \int_{\partial M} \theta \varepsilon \partial_\nu u|_{\partial M}(t) k_{H^{1/2}(\partial M)}^2 dt > k u(T) k_{H_0^1(M)}^2,$$

for all  $u_0 \in H_0^1(M)$  and  $u$  solution of (1.9).

Here  $\partial_\nu$  denotes a unit normal (for the metric  $g$ ) vector field to  $\partial M$ . Then, the uniform observability question is whether  $C_0(T, \varepsilon)$  remains uniformly bounded in the limit  $\varepsilon \rightarrow 0^+$ , and the associated minimal uniform observation time is defined again by

$$T_{\text{unif}}(\theta) = \inf \{ T > 0 \text{ for which there exist } K_0, \varepsilon_0 > 0 \text{ such that (1.10) holds} \\ \text{for all } u_0 \in H_0^1(M), \text{ with } C_0(T, \varepsilon) \leq K_0 \text{ for all } \varepsilon \in (0, \varepsilon_0) \}.$$

Before going further, let us first give the definition of an analogue of the condition (GCC) in case of a manifold with boundary  $\partial M \neq \emptyset$  (called here Flushing Condition), as used in the Guerrero-Lebeau result [GL07]. For this, we need to extend  $(M, g)$  in a slightly bigger Riemannian manifold  $(\tilde{M}, \tilde{g})$ , i.e., such that  $M \subset \text{Int}(\tilde{M})$  and  $\tilde{g}$  any Riemannian metric on  $\tilde{M}$  such that  $\tilde{g}|_M = g$ . In the case of a bounded domain of  $\mathbb{R}^n$ , one may choose  $\tilde{M} = \mathbb{R}^n$ . We also extend  $f \in W^{2,1}(M)$  as a *compactly supported* function  $\tilde{f} \in W^{2,1}(\tilde{M})$  such that  $\tilde{f}|_M = f$ .

**DEFINITION 1.4.** — For  $x \in M$ , denote by  $\gamma_x$  the maximal solution to

$$\dot{\gamma}_x(t) = -\text{r}_{\tilde{g}} \tilde{f}(\gamma_x(t)), \quad \gamma_x(0) = x.$$

Note that this solution is defined globally in time since  $\tilde{f}$  is compactly supported.

We say that  $(M, \text{r}_g f, T)$  (resp.  $(M, \text{r}_g f, \omega, T)$ ) satisfies the Flushing Condition (FC) if for all  $x \in M$  there is  $t \in (0, T)$  such that  $\gamma_x(t) \notin M$  (resp.  $\gamma_x(t) \notin M$  or  $\gamma_x(t) \in \omega$ ). We also say that  $(M, \text{r}_g f)$  (resp.  $(M, \text{r}_g f, \omega)$ ) satisfies (FC) if there is  $T > 0$  such that  $(M, \text{r}_g f, T)$  (resp.  $(M, \text{r}_g f, \omega, T)$ ) does. We then define accordingly



the time  $T_{FC}(M, r_g f)$  (resp.  $T_{FC}(M, \omega, r_g f)$ ) as the infimum of  $T > 0$  for which this property holds.

Remark finally that these definitions do not depend on the extensions  $(\tilde{M}, \tilde{g})$  and  $\tilde{f}$ .

Guerrero-Lebeau [GL07] prove an analogue of Theorem 1.1 in the present setting (and for general vector fields), namely: if  $(M, r_g f)$  (resp.  $(M, r_g f, \omega)$ ) satisfies the Flushing Condition, given  $\theta$  any nontrivial observation function, then there is  $T_u > 0$  and  $K_0 > 0$  such that for all  $T > T_u$ , (1.10) holds (resp. (1.4) holds for all solutions to (1.9)), with  $C_0(T, \varepsilon) \leq K_0$  for all  $\varepsilon \leq 1$ .

Two important geometric quantities in our results are the potential associated to the function  $f$ , defined by

$$(1.11) \quad V(x) := \frac{\int r_g f(x) j_g^2}{4},$$

and the Agmon distance (see e.g. [Hel88, Chap. 3]) to the minimum of this potential, namely, with  $E_0 = \min_M V$ ,

$$(1.12) \quad \begin{aligned} d_A(x, y) &= \inf \left\{ \int_0^1 \sqrt{(V(\gamma(t)) - E_0)_+} |\dot{\gamma}(t)|_g dt, \gamma \in \mathbf{U}_1(x, y) \right\}, \\ \mathbf{U}_1(x, y) &= \{ \gamma \in W^{1,1}([0, 1]; M) ; \gamma(0) = x, \gamma(1) = y \}, \\ d_A(x) &= d_A(x, V^{-1}(E_0)) = \inf_{y \in V^{-1}(E_0)} d_A(x, y). \end{aligned}$$

Here  $V^{-1}(E_0)$  is the classically allowed region at the potential minimum,

$$(V(x) - E_0)_+ = \max(V(x) - E_0, 0),$$

and  $d_A(x)$  is the Agmon distance of  $x$  to the set  $V^{-1}(E_0)$  for the (pseudo-)metric  $(V - E_0)_+ g$ . Remark that the index  $(\cdot)_+$  is not needed at the bottom energy  $E_0 = \min_M V$ ; however, we keep it here since the definition (1.12) will also be useful for a general energy level.

Our main result in this general setting formulates as follows.

**THEOREM 1.5.** — *We assume that  $f \in W^{2,1}(M)$  (hence  $V \in W^{1,1}(M)$ ) and  $q \in L^1(M)$ . We let  $E_0 = \min_M V$ , set*

$$W_{E_0}(x) = \frac{f(x)}{2} + d_A(x),$$

*and fix  $\omega \in M$  (resp.  $\theta \in C^1(\partial M)$ ), and in the case of boundary estimates, we further assume  $f, q \in C^1(M)$ . For any  $\delta > 0$ , there is  $\varepsilon_0 > 0$  such that for all  $\varepsilon \in (0, \varepsilon_0)$  the observability inequality (1.4) (resp. (1.10)) with constant  $C_0(T, \varepsilon)$  implies*

$$\begin{aligned} C_0(T, \varepsilon) &> \exp \frac{1}{\varepsilon} \left( \min_{\bar{\omega}} W_{E_0} - \max_{V^{-1}(E_0)} \frac{f}{2} - E_0 T - \delta \right), \\ \text{resp. } C_0(T, \varepsilon) &> \exp \frac{1}{\varepsilon} \left( \min_{\text{supp } \theta} W_{E_0} - \max_{V^{-1}(E_0)} \frac{f}{2} - E_0 T - \delta \right). \end{aligned}$$

In particular, we have

$$(1.13) \quad E_0 T_{\text{unif}}(\omega) > \frac{\min_{\bar{\omega}} W_{E_0}}{\bar{\omega}} \max_{V^{-1}(E_0)} f/2,$$

$$(1.14) \quad \left( \text{resp. } E_0 T_{\text{unif}}(\theta) > \min_{\text{supp } \theta} W_{E_0} \max_{V^{-1}(E_0)} f/2 \right).$$

Note that the quantity in the right hand-side of (1.13)-(1.14) as well as  $E_0$  are invariant under the change  $f \mapsto f + C$  for  $C$  constant on  $M$ . This is consistent with the fact that the equations remain unchanged by such a modification of  $f$ . Note also that if  $E_0 = 0$  and  $V^{-1}(E_0) \setminus \omega = ?$  (resp.  $V^{-1}(E_0) \setminus \text{supp}(\theta) = ?$ ), a more precise version of this result (see Theorem 3.1 below) actually shows that  $C_0(T, \varepsilon) > e^{c/\varepsilon}$  for one  $c > 0$  and all time  $T > 0$  (in particular, uniform observability never holds). This is consistent with (and a particular case of) the Guerrero-Lebeau [GL07] result (first part of Theorem 1.1 above) for in this case,  $(M, r_g f, \omega)$  does not satisfy (GCC). Indeed, a point  $x_0 \in V^{-1}(E_0)$  satisfies  $r_g f(x_0) = 0$  and is thus a stationary point of the gradient dynamics.

We refer to Remark 3.5 concerning the additional smoothness assumption for the boundary estimate.

**THEOREM 1.6.** — Assume  $M = \bar{\Omega}$  where  $\Omega \subset \mathbb{R}^n$  is any smooth bounded connected open set endowed with  $g = \text{Eucl}$  the Euclidean metric and  $q \in L^1(M)$ . There exist  $\omega \subset \Omega$  and constants  $c_\omega, \delta > 0$  such that for any  $\lambda > 0$ , there is a function  $f_\lambda \in C^1(\bar{\Omega})$  such that:

- $(\bar{\Omega}, r f_\lambda, \omega)$  satisfies (FC) and  $T_{\text{FC}}(\bar{\Omega}, r f_\lambda, \omega) \leq \text{diam}(\Omega)$ ;
- $T_{\text{unif}}(\omega) > c_\omega \lambda$ ;
- $\lambda^2 \delta \leq k r f_\lambda k_{L^1(\Omega)}^2 \leq \lambda^2 \text{diam}(\Omega)^2 + n$ .

In particular, for all  $\Lambda > 0$ , there is  $f \in C^1(\bar{\Omega})$  such that

$$T_{\text{unif}}(\omega) > \Lambda T_{\text{FC}}(\bar{\Omega}, r f, \omega).$$

The result of Theorem 1.6 is already of interest in dimension one. In this case  $\Omega = (L, L)$ , the vector field we consider is  $f^\theta(x) \partial_x$  with  $f^\theta > 1$  on  $[L, L]$  and  $f^\theta(0) = 1$  and the observation set  $\omega$  is a neighborhood of the boundary point  $L$  (note that this would correspond to the case  $M < 0$  in the Coron-Guerrero problem with the notation of [CG05]). Note that the function  $f_\lambda$  in this result satisfies  $\max f_\lambda - \min f_\lambda \sim \lambda$ . As a consequence, one cannot even hope to have existence of a constant  $K > 0$  depending *only* on  $\min_{x \in [0, L]} f^\theta(x)$  (a uniform flushing time) such that  $T_{\text{unif}}(\omega) \leq K T_{\text{FC}}(\bar{\Omega}, r f_\lambda, \omega)$ . However, at this point, it does not seem hopeless that such a constant  $K$  depends only on  $k r f_\lambda k_{L^1(\Omega)}$ , at least for a fixed metric.

**REMARK 1.7.** — In the case  $\partial M = ?$ , Theorem 1.5 does not seem to suffice to construct functions  $f, \omega$  having  $T_{\text{unif}}(\omega)/T_{\text{GCC}}(M, r_g f, \omega)$  arbitrarily large. In a domain of  $\mathbb{R}^n$ , Theorem 1.5 is however enough to provide counterexamples.

Another drawback of Theorem 1.5 is that it does not produce any useful lower bound in case  $\omega$  is a whole neighborhood of  $\partial M$  (or in the boundary observation case from the whole boundary  $\partial M$ ).

We remedy these issues in the next section on surfaces of revolution.

1.3.2. *Lower bounds on surfaces of revolution.* — In Theorems 1.5 and 1.6 above, the lower bound of the minimal uniform observability time is essentially due to the contribution of the potential  $V(x) = j_{\Gamma_g} f(x) j_g^2 / 4$  (and the difference between its maximal and minimal values on  $M$ ). In this section, we consider a family of geometric settings, namely surfaces of revolution, for which the contribution of the geometry of  $(M, g)$  plays an important role. This leads in particular to explicit versions of Theorem 1.2.

The precise description of the geometry of the surfaces we consider is given in Section 4.1 and we only describe here features required to state the result. We may consider either:

- (1)  $M = S \subset \mathbb{R}^3$  a smooth compact surface diffeomorphic to the sphere  $S^2$ ;
- (2)  $M = S \subset \mathbb{R}^3$  a smooth compact surface diffeomorphic to the disk  $D$ ;
- (3)  $M = S \subset \mathbb{R}^3$  a smooth compact surface diffeomorphic to the cylinder  $[0, 1] \times S^1$ ;
- (4)  $M = S$  a smooth compact surface diffeomorphic to the torus  $T^2 = S^1 \times S^1$ .

We assume moreover that it has revolution invariance around an axis. In particular, the axis may intersect  $S$  in two points (sphere), one point (disk) or no point (cylinder or torus). Except near these points,  $S$  has a global coordinate chart  $(s, \theta) \in (0, L) \times S^1$  for some  $L > 0$ . In the first three cases, the surface is endowed with the metric  $g$  inherited from the Euclidean metric on  $\mathbb{R}^3$  which writes

$$(1.15) \quad g = ds^2 + R(s)^2 d\theta^2,$$

on account to the rotation invariance. Here the function  $R : (0, L) \rightarrow (0, \infty)$  describes the shape of  $S$  (distance to the revolution axis). In the torus case, we simply endow  $T^2$  with the metric (1.15).

We further assume that the function  $f$  and  $q$  are themselves rotationally invariant, that is  $f = f(s)$  and  $q = q(s)$  in these coordinates. In this setting (and as opposed to results presented in above Section 1.3.1), the relevant quantities for our analysis are the following.

We define for any  $c > 0$  (that can be chosen) the ( $\theta$ -invariant) effective potential

$$(1.16) \quad V_c(s) := \frac{c^2}{R(s)^2} + \frac{f^\theta(s)^2}{4}.$$

Note that, as opposed to the potential appearing in (1.11), this potential  $V_c$  is different from  $j_{\Gamma_g} f j_g^2 / 4 = f^\theta(s)^2 / 4$ . Moreover, it depends explicitly on the geometry (namely, on  $R$ ). We shall make the simplifying assumption that

$$(1.17) \quad V_c^{-1}(\min V_c) = f_{s_{\min}g}$$
 consists in a single point  $s_{\min}$ .

Note that in case  $S$  has a boundary, one may have  $s_{\min}$  at the boundary (see Section 4 for more precise statements). As in the previous section, we introduce the associated

Agmon distance, which simply writes in the first three cases:

$$(1.18) \quad d_A^c(s) = \left| \int_{s_{\min}}^s \sqrt{V_c(y) - V_c(s_{\min})} dy \right|.$$

In the fourth case  $S = S_L^1 \setminus S^1$ , an analog of (1.12) still makes sense on  $S_L^1$  when  $V_c$  is defined on  $S_L^1$ . We may choose a representation of  $S_L^1 = \mathbb{R}/L\mathbb{Z}$  in which  $s_{\min} = 0 + LZ$ , a definition of the Agmon distance then reads

$$(1.19) \quad d_A^c(s) = \min_{s_{\min} \leq LZ} \left[ \left| \int_{s_{\min}}^s \sqrt{V_c(y) - V_c(s_{\min})} dy \right| \right].$$

We also set

$$(1.20) \quad W^c(s) = d_A^c(s) + \frac{f(s)}{2}, \quad \text{for } s \in (0, L).$$

Then our main result in this geometric context can be (loosely) stated as follows.

**THEOREM 1.8.** — *Let  $c > 0$ , assume that  $V_c$  satisfies (1.17) and that, in the coordinates  $(s, \theta) \in (0, L) \times S^1$ , we have  $\omega = (0, \delta) \in S^1 \setminus [(L - \delta, L) \times S^1]$  (this can be rewritten in an intrinsic way on  $S$ ). Assuming the observability inequality (1.4) (resp. (1.10)) with constant  $C_0(T, \varepsilon, \omega)$  (resp.  $C_0(T, \varepsilon, fLg \setminus S^1)$ ), there is a sequence  $\varepsilon_k \searrow 0^+$  such that for any  $\delta > 0$ , there is  $k_0(\delta) > 0$  such that*

$$C_0(T, \varepsilon_k, \omega) > e^{(1/\varepsilon_k)(W_\omega^c - W_m^c - V_c(s_{\min})T - \delta)} \quad \text{for all } k > k_0(\delta), \quad W_\omega^c = \min_{\bar{\omega}} W^c,$$

$$\text{resp. } C_0(T, \varepsilon_k, fLg \setminus S^1) > e^{(1/\varepsilon_k)(W^c(L) - W_m^c - V_c(s_{\min})T - \delta)} \quad \text{for all } k > k_0(\delta),$$

where  $W_m^c = \inf_{(0, L)} W^c$ . In particular, we have

$$(1.21) \quad V_c(s_{\min})T_{\text{unif}}(\omega) > W_\omega^c - W_m^c,$$

$$(1.22) \quad V_c(s_{\min})T_{\text{unif}}(fLg \setminus S^1) > W^c(L) - W_m^c.$$

Theorem 1.8 differs from Theorem 1.5 in several respects. First notice that the potential appearing in Theorem 1.5 is  $f^\theta(s)^2/4$ , that is  $V_0(s)$  with the definition of  $V_c$  in (1.16). In particular, it does not depend on  $R$ : neither does its minimal value, nor the associated Agmon distance and function  $W^0$ . Therefore, in this very particular geometric context, the results of Theorem 1.5 do not depend on the geometry of  $R$ , and hence only formulate as a one-dimensional result in the  $s$  variable. As such, they do not care about the "transverse dynamics" in the  $\theta$ -variable. Theorem 1.8 overcomes this lack and shows that both have to be taken into account.

Another difference with the estimates of Theorem 1.5 is that  $\max_{V^{-1}(E_0)} f/2 = \max_{V^{-1}(E_0)} W$  is here replaced by  $\inf_{(0, L)} W^c$ . This improvement is due to the "one-dimensional" underlying framework (in which localization properties of eigenfunctions are better understood).

Again, we remark that the initial problem is invariant by changing  $f$  by a constant  $f + C_0$ . In Estimate (1.21), both the potential  $V_c$  and the quantity  $W_\omega^c - W_m^c$  are as well invariant by this change of the function  $f$ .

We now state three particular examples of application of Theorem 1.8. The latter imply Theorem 1.2.

**COROLLARY 1.9.** — *Assume  $S$  is a surface of revolution in  $\mathbb{R}^3$  diffeomorphic to  $S^2$  (resp.  $\mathbb{D}$ ) and such that  $R^{-1}(\max R)$  is a single point ( $R$  has a unique maximum). Denote by  $N, S$  (resp.  $N$  only) the north/south poles of  $S$ , which are the only two (resp. the unique) invariant points under the revolution symmetry. Then, for any  $\delta > 0$ , there exists  $f_\delta \in C^1(S)$  invariant by rotation such that with  $\omega_\delta = B_g(N, \delta) \cup B_g(S, \delta)$  (resp.  $\omega_\delta = B_g(N, \delta)$ ) we have*

(1)  $(S, r_g f_\delta, \omega_\delta)$  satisfies (GCC) (resp. satisfies (FC)) and

$$T_{\text{GCC}}(S, r_g f_\delta, \omega_\delta) = L \quad 2\delta \leq L = \text{dist}_g(N, S) \\ (\text{resp. } T_{\text{FC}}(S, r_g f_\delta, \omega_\delta) = L \quad \delta \leq L = \text{dist}_g(N, \partial S));$$

(2) for all  $c > 0$ , there is  $\delta_0, C > 0$  such that for all  $\delta \in (0, \delta_0)$

$$T_{\text{Unif}}(\omega_\delta) > \left( \frac{c^2}{R(s_{\min})^2} + \frac{1}{4} \right)^{-1} (c \log(1/\delta) - C).$$

This result proves the first item in Theorem 1.2. In particular, the limit  $\delta \rightarrow 0^+$  prevents from the existence of a universal constant  $K > 1$  such that

$$T_{\text{Unif}}(\omega) \leq K T_{\text{GCC}}(M, X, \omega).$$

Note that in this construction, the vector fields  $r_g f_\delta, r_g$  are rotationally invariant and independent of the metric  $g$ . Moreover, for  $\delta < \delta^0$ , the two functions  $f_\delta$  and  $f_{\delta^0}$  coincide on  $M \setminus \omega_{\delta^0}$ .

In our second result, the geometry is close to that of Corollary 1.9. However, we consider fixed  $\omega$  and  $f$  (and even a fixed vector field), but let the metrics  $g$  vary. We denote  $S_L^1 = \mathbb{R}/L\mathbb{Z}$  and  $S^1 = S_{2\pi}^1$ .

**COROLLARY 1.10.** — *Assume  $S = S_L^1 \times S^1$  with coordinates  $(s, \theta)$  and let  $f \in C^1(S_L^1)$ . Let  $\omega = I_\omega \times S^1$  with  $I_\omega$  a nonempty interval such that  $\overline{I_\omega} \subset S_L^1$ .*

*Then, there is a constant  $C > 0$  such that for any  $\delta \in (0, 1)$ , there exists a function  $R_\delta \in C^1(S_L^1; \mathbb{R}^+)$  such that*

(1) *the vector field  $r_{g_\delta} f - r_{g_\delta} = f^\theta(s) \partial_s$  (defined by (1.15) associated to  $R_\delta$ ) does not depend on the metric  $g_\delta$ , the triple  $(S, r_{g_\delta} f, \omega)$  satisfies (GCC) if and only if  $f^\theta \notin 0$  on  $S_L^1 \cap I_\omega$ , and, in this case, we have*

$$T_{\text{GCC}}(S, r_{g_\delta} f, I_\omega \times S^1) = \left| \int_{S_L^1 \cap I_\omega} \frac{ds}{f^\theta(s)} \right| < +\infty;$$

(2) *for the transport equation (1.6) with viscosity given by the metric  $g_\delta$  under the form (1.15) associated to  $R_\delta$ , we have*

$$T_{\text{Unif}}(I_\omega \times S^1) > C\delta^{-1/2};$$

(3)  $\delta^{1/2}(1 - C\delta) \leq \min_{S^1} R_\delta \leq \delta^{1/2}$  for all  $\delta \in (0, 1)$ .

This result implies the second item in Theorem 1.2.

Another application is given by the following result, which is an analogue of Corollary 1.10 for the boundary observability problem in the cylindrical geometry.

**COROLLARY 1.11.** — Assume  $S = [0, L] \times S^1$  (whence  $\partial S = (\{0\} \times S^1) \cup \{L\} \times S^1$ ), and let  $f \in C^1([0, L])$ . Then, for any  $\gamma > 2$ ,  $\delta_0 > 0$  there is a constant  $C > 0$  such that for all  $\delta \in (0, \delta_0]$ , there exists a function  $R_\delta : [0, L] \rightarrow \mathbb{R}^+$  such that

(1) the vector field  $X = r_{g_\delta} f \cdot r_{g_\delta} = f^\theta(s) \partial_s$  does not depend on the metric  $g_\delta$  (defined by (1.15) with  $R_\delta$ );

(2)  $(S, r_{g_\delta} f)$  satisfies (FC) (in the sense of Definition 1.4) if and only if  $f^\theta \not\equiv 0$  on  $[0, L]$  and in this case,

$$T_{\text{FC}}(M, r_{g_\delta} f) = \left| \int_0^L \frac{ds}{f^\theta(s)} \right| < +\infty;$$

(3) for the transport equation with viscosity and Dirichlet boundary conditions (1.6), and with metric  $g_\delta$  under the form (1.15) associated to  $R_\delta$ , we have

$$T_{\text{unif}}(\partial S) > \left( \max_{[0, L]} \frac{f^{\theta 2}}{4} \right)^{-1} \left( \frac{1}{\delta^{\gamma/2 - 1}} \frac{1}{\gamma/2 - 1} C \right);$$

$$(4) \begin{cases} (s + \delta)^{\gamma/2} (1 - C(s + \delta)) \in R_\delta(s) \in (s + \delta)^{\gamma/2} & \text{for } s \in [0, L/4], \\ (L - s + \delta)^{\gamma/2} (1 - C(L - s + \delta)) \in R_\delta(s) \in (L - s + \delta)^{\gamma/2} & \text{for } s \in [3L/4, L]. \end{cases}$$

Note the link between the asymptotic singularity of the metric  $R_\delta(s) \sim (s + \delta)^{\gamma/2}$  (i.e.,  $S$  becomes close to a “conical geometry” for small  $\delta$ ) and the blowup of the minimal time  $T_{\text{unif}} \sim 1/\delta^{\gamma/2 - 1}$ . Note also that the limit case  $\gamma = 2$ , all calculations can be done as well and lead to  $R_\delta(s) \sim s + \delta$  together with  $T_{\text{unif}} \sim \log \delta$ . This corresponds to the case where the geometry of the cylinder degenerates towards that of the disk, leading to the same blowup estimate as in Corollary 1.9.

**1.3.3. Observability for positive solutions.** — As already mentioned, our last result concerns the uniform observability question for positive solutions of (1.6), and is restricted to the case  $\partial M = \emptyset$ . We also assume  $f \in C^3$ . Note that if  $u_0 \in L^2(M; \mathbb{R}^+)$ , then the associated solution  $u$  to (1.6) satisfies  $u(t, x) > 0$  for a.e.  $(t, x) \in \mathbb{R}^+ \times M$  (see e.g. Th. 9 in §7.1 p. 369 together with Problem 7 in §7.5 in [Eva98], or Chap. III, Th. 7.1, p. 181 in [LSU68]).

We consider the observability inequality for nonnegative solutions:

$$(1.23) \quad C_0^+(T, \varepsilon)^2 \int_0^T \int_\omega |ju(t, x)|^2 ds(x) dt > ku(T) k_{L^2(M)}^2,$$

for all  $u_0 \in L^2(M; \mathbb{R}^+)$  and  $u$  solution of (1.6).

and the associated minimal time  $T_{\text{unif}}^+(\omega)$  of uniform observability for positive solution, already defined in (1.7). The main result we obtain in this context is the above Theorem 1.3, stating that  $T_{\text{unif}}^+(\omega) = T_{\text{GCC}}(M, r_g f, \omega)$ . As a byproduct of our analysis, we also obtain a lower estimate on the blow up of the control cost when the Geometric Control Condition is not satisfied. It involves the definition of a quantity

that roughly speaking, measures how two points are far from being the image of a trajectory at time  $t$ , namely

$$(1.24) \quad d_{r_g f}(x, y, t) = \frac{1}{4} \inf \left\{ \int_0^t |\dot{\gamma}(s)|_{r_g f(\gamma(s))}^2 ds; \gamma \in \mathbf{U}_t(x, y) \right\},$$

$$\mathbf{U}_t(x, y) = \{ \gamma \in W^{1,1}([0, t]; M); \gamma(0) = x, \gamma(t) = y \}.$$

Note that we have  $d_{r_g f}(x, y, t) \sim d(x, \phi_t(y))$  for bounded  $t$ , where  $d$  denotes the Riemannian distance and  $\phi_t$  the flow of the vector field  $r_g f$  (see Lemma A.2 where  $d_{r_g f}(x, y, t)$  is interpreted as a control cost from  $x$  to  $\phi_t(y)$  with time varying metric).

**PROPOSITION 1.12.** — *Assume that  $(M, r_g f, \bar{\omega})$  does not satisfy (GCC). Then, we have*

$$d_{([0, T], \bar{\omega})} := \sup_{y \in M} \inf_{x \in \bar{\omega}} \inf_{t \in [0, T]} d_{r_g f}(x, y, t) > 0.$$

Moreover, for any  $\delta > 0$ , there is  $\varepsilon_0 > 0$  such that for all  $\varepsilon \in (0, \varepsilon_0)$ , we have

$$(1.25) \quad C_0(T, \varepsilon) > C_0^+(T, \varepsilon) > e^{(d_{([0, T], \bar{\omega})} - \delta)/\varepsilon}.$$

Note that this exponential blowup is a refinement of the Guerrero-Lebeau [GL07] result (first part of Theorem 1.1 above). However, we provide here with a precise geometric rate (namely  $d_{([0, T], \bar{\omega})}$ ) quantifying this blowup phenomenon.

The proofs of Theorem 1.3 and Proposition 1.12 rely on estimates on the kernel of the associated equation. Note that kernel estimates have already been used in [Mil04] to prove lower bounds for the cost of controllability of the usual heat equation in the short time asymptotics, and in [LL21a] to prove observability of positive solutions to the heat equation with optimal constants.

**1.4. FURTHER REMARKS.** — In this section, we collect several remarks and comments related to our results.

(1) The principal interest of working with gradient vector fields  $X = r_g f$  is that the associated operator  $r_g f - \varepsilon \Delta_g$  can be conjugated to a *selfadjoint* Schrödinger operator (1.8). And the limit  $\varepsilon \rightarrow 0^+$  then corresponds to the semiclassical limit, which has been the object of many studies (see e.g. [Wit82, Sim83, HS84, HS85, Hel88, CFKS87, All98, HKN04, HN06]). This conjugation does no longer hold in case  $X$  is not a gradient vector field. One could also consider that giving counter-examples with gradient flows is “stronger” than general counter-examples. We do not know whether an analogue of Theorem 1.3 for positive solutions remains true for general vector fields  $X$ . This seems to be an interesting open problem.

(2) In the context of surfaces of revolution, as presented in Section 1.3.2, we only provide with lower bounds of  $T_{\text{unif}}$ . It would of course be interesting to obtain related upper bounds on this uniform control time. This would require to provide a very precise description of several spectral quantities (spectral gaps, localization of eigenfunctions at all energy levels...) for the semiclassical Schrödinger operator  $P_\varepsilon$  in (1.8), and seems to be a difficult question. See the companion paper [LL21b] for an upper bound in a related one-dimensional situation.

(3) The one-dimensional one-well problem is considered in [LL21b]. In this very particular situation, we are able to provide with

- improved lower bounds on the minimal time when compared to Theorem 1.5;
- an upper bound on the minimal time.

This requires the knowledge of precise information on the spectral gap and the localization of eigenfunctions at all energy levels  $E > E_0$  (whereas Theorem 1.5 is only concerned with the bottom energy level  $E_0$ ). See also the discussion at the beginning of Section 3.

(4) Notice that if one is not interested in null-controllability (i.e., driving the solution exactly to zero at time  $T$ ), but rather in approximate controllability with a reasonable cost (and a precision depending on the viscosity  $\varepsilon$ ), one might be satisfied by the following statement.

**PROPOSITION 1.13.** — *Suppose  $\partial M = ?$  and  $(M, X, \omega, T)$  satisfies (GCC) (resp.  $\partial M \notin ?$  and  $(M, X, \omega, T)$  satisfies (FC)). Then, there exist  $C, C_0 > 0$  such that for all  $y_0 \in L^2(M)$ ,  $\varepsilon \in (0, 1]$  there is  $h = h_\varepsilon \in L^2((0, T) \times \omega)$  with*

$$\|h_\varepsilon\|_{L^2((0, T) \times \omega)} \leq C \|y_0\|_{L^2(M)},$$

such that the associated solution to (2.1) satisfies

$$\|y(T)\|_{L^2(M)} \leq C e^{-C_0/\varepsilon} \|y_0\|_{L^2(M)}.$$

That is to say, one can drive the solutions  $e^{-C_0/\varepsilon}$  close to zero with a uniformly bounded cost. This result follows from Proposition 5.7 below (a particular case of [GL07, Prop. 3]) together with [LL21c, App.]. This can be particularly useful for numerical purposes, since  $e^{-C_0/\varepsilon} = 0$  numerically for  $\varepsilon$  small enough.

In the situation of Theorems 1.5 or 1.8, this means that for intermediate times  $T \geq (T_{\text{GCC}}(M, r_{\text{gf}}, \omega), T_{\text{Unif}}(\omega))$  (resp. for  $T \geq (T_{\text{FC}}(M, r_{\text{gf}}, \omega), T_{\text{Unif}}(\omega))$  if  $\partial M \notin ?$ ), controlling the solution  $e^{-C_0/\varepsilon}$  close to zero costs  $\sim 1$ , whereas controlling the solution exactly to zero costs  $\sim e^{C/\varepsilon}$ .

(5) Note that in the context of revolution surfaces of Section 1.3.2, we can prove a complementary result compared to [LL21a, Th. 1.9/Cor. 1.10]. The techniques used in Proposition 4.9 show that in all cases of Section 1.3.2, for any rotationally invariant set  $\omega$ , we have (with the notation of [LL21a], the constant  $K_{\text{eig}}(\omega)$  being the smallest constant  $K$  in the inequality  $\|k\psi_\lambda\|_{L^2(M)} \leq C e^{K \overline{\lambda}} \|k\psi_\lambda\|_{L^2(\omega)}$  where  $\Delta_g \psi_\lambda = \lambda \psi_\lambda$ )

$$K_{\text{eig}}(\omega) = R(s_{\min}) d_A(\omega), \quad \text{with } d_A(\omega) = \inf_{x \in \omega} d_A(x).$$

In [LL21a], we only proved  $K_{\text{eig}}(\omega) > R(s_{\min}) d_A(\omega)$  (and only in case  $S$  is diffeomorphic to a sphere). This result is close to that of Allibert [All98], which already proves this in case  $S$  is diffeomorphic to a cylinder and the function  $R$  has a single local maximum which is non-degenerate.



*Acknowledgements.* — Part of this research was done when the second author was in École Polytechnique, Centre de Mathématiques Laurent Schwartz UMR7640, 91128 Palaiseau cedex France.

We would like to thank Olivier Glass (who suggested Proposition 1.13), Franck Sueur and Dorian Le Peutrec for interesting discussions related to this work. We are also grateful to Jon Asier Bárcena-Petisco for his comments on a first version of the article. Finally we would like to thank both referees for their careful reading of the manuscript and their constructive remarks that helped to improve the presentation of the paper.

## 2. PRELIMINARIES: DUALITY, CONJUGATION OF GRADIENT FLOWS AND (GCC)

**2.1. UNIFORM CONTROLLABILITY PROBLEMS AND DUAL FORMULATION.** — In this section, we reformulate the above uniform observability questions in terms of uniform controllability statements. This follows essentially the classical duality approach, see [DR77] or [Cor07, Chap. 2.3].

**2.1.1. Duality between internal control and observation problems.** — In this section, we present the controllability problems associated to the above observability questions, and we briefly describe the duality between the control and the observation problems. We introduce the internal control problem

$$(2.1) \quad \begin{cases} (\partial_t + X + \operatorname{div}_g(X) - q - \varepsilon \Delta_g)y = \mathbf{1}_\omega h, & \text{in } (0, T) \cap M, \\ y = 0, & \text{on } (0, T) \cap \partial M, \\ y|_{t=0} = y_0, & \text{on } M. \end{cases}$$

Notice that, as opposed to (1.3), the operator appearing in these control problems is  $X = X - \operatorname{div}_g(X)$ , where the adjoint is taken in the space  $L^2(M, d\operatorname{Vol}_g)$ .

That the appropriate dual observation problem is (1.3) is a consequence of the following lemma.

**LEMMA 2.1 (Duality equation).** — *For all solutions  $u \in C^0([0, T]; L^2(M))$  of (1.3) on  $[0, T]$  with Dirichlet boundary conditions and all  $y \in C^0([0, T]; L^2(M))$  solution to (2.1) with  $h \in L^2((0, T) \cap M)$ , we have*

$$(2.2) \quad (u(T), y_0)_{L^2(M)} - (u_0, y(T))_{L^2(M)} + \int_0^T (\mathbf{1}_\omega u(t), h(T-t))_{L^2(M)} dt = 0.$$

Notice that one passes from the observed evolution to the controlled evolution by changing  $(X, q)$  into  $(-X, q - \operatorname{div}_g(X))$ . The interest of adding a potential term  $q(x)$  in these equations is that the free equation (1.9) and the controlled equation (2.1) then have the same form (i.e., the adjoint of a vector field is not a vector field but the adjoint of a first order differential operator is a first order differential operator).

**DEFINITION 2.2 (Controllability and cost).** — Given  $(\omega, \varepsilon, T)$ , we say that (2.1) is null-controllable from  $(\omega, T)$  if for any  $y_0 \in L^2(M)$ , there is  $h \in L^2((0, T) \cap M)$  such that the associated solution to (2.1) satisfies  $y(T) = 0$ . If (2.1) is null-controllable

from  $(\omega, T)$ , we define for  $y_0 \in L^2(M)$  the set  $U(y_0)$  of all such controls  $h \in L^2((0, T) \times M)$ , and the cost function

$$C_0(T, \varepsilon) := \sup_{\substack{y_0 \in L^2(M) \\ \|y_0\|_{L^2(M)} \leq 1}} \left\{ \inf_{h \in U(y_0)} \|h\|_{L^2((0, T) \times M)} \right\}.$$

As a corollary of Lemma 2.1, together with classical arguments (see e.g. [DR77] or [Cor07, Chap. 2.3]) we deduce the following statement.

**COROLLARY 2.3** (Observability constant = control cost). — *Given  $(\omega, \varepsilon, T)$ , Equation (2.1) is null-controllable from  $(\omega, T)$  if and only if the observability inequality (1.4) holds. Moreover, we then have  $C_0(T, \varepsilon) = C_0(T, \varepsilon)$ .*

As a consequence, all lower bounds on  $C_0(T, \varepsilon)$  formulated in Theorems 1.5 and 1.8 translate into lower bounds on  $C_0(T, \varepsilon)$ . The time  $T_{\text{unif}}(\omega)$  is equal to the minimal time of uniform *controllability*, and all lower bounds on the time  $T_{\text{unif}}(\omega)$  obtained in Theorems 1.5 and 1.8 and their corollaries apply.

The uniform observability result for positive solutions of the heat equation in Theorem 1.3 also has a controllability counterpart. This fact was indeed proved by Le Bal'c'H [LB20, Th. 4.1] for the classical heat equation. In the present context, the uniform observability estimate for positive solutions, associated to Theorem 1.3, implies the following controllability result.

**COROLLARY 2.4.** — *Let  $M$  be a compact Riemannian manifold with  $\partial M = \emptyset$ ,  $X = r_g f$  where  $f \in C^3(M)$ , and  $\omega \subset M$  an open subset. Assume that  $(M, r_g f, \omega)$  satisfies (GCC), and  $T > T_{\text{GCC}}(M, r_g f, \omega)$ . Then, there exist  $C, \varepsilon_0 > 0$  so that for any  $y_0 \in L^2(M)$  and  $0 < \varepsilon \leq \varepsilon_0$ , there exists a control  $h \in L^2([0, T], L^2(\omega))$  with*

$$\|h\|_{L^2([0, T], L^2(\omega))} \leq C \|y_0\|_{L^2(M)}$$

such that the solution of (2.1) satisfies  $y(T) > 0$ .

We refer to Section 5.4 for a proof.

**2.1.2. Duality between boundary control and observation problems.** — We now briefly discuss the boundary case and we refer to [GL07] for the details. The boundary control problem under interest is

$$(2.3) \quad \begin{cases} (\partial_t + X + \text{div}_g(X) - q - \varepsilon \Delta_g)y = 0, & \text{in } (0, T) \times \text{Int}(M), \\ y = \theta h, & \text{on } (0, T) \times \partial M, \\ y|_{t=0} = y_0, & \text{on } M, \end{cases}$$

where  $\theta \in C^1(\partial M; \mathbb{R})$  is meant to be a smooth version of  $1_\Gamma, \Gamma \subset \partial M$ . Solutions of (2.3) are defined in the sense of transposition, and a well-posedness statement can be written as follows.

LEMMA 2.5 (Guerrero-Lebeau [GL07] pp 1814-1815). — Assume  $X$  is a  $L^1$  vector field on  $M$  with  $\operatorname{div}_g(X) \in L^1(M)$ ,  $q \in L^1(M)$ , and let  $T > 0$ . Then, there exists  $C > 0$  such that for all  $y_0 \in H^{-1}(M)$ , all  $h \in L^2(0, T; H^{-1/2}(\partial M))$  and all  $\varepsilon > 0$ , there is a unique solution  $y$  to (2.3) in the sense of transposition, which satisfies  $y \in L^2((0, T) \times M) \setminus C^0([0, T]; H^{-1}(M)) \setminus H^1(0, T; H^{-2}(M))$  with

$$\|y\|_{L^2((0, T) \times M)} + \|y\|_{L^1(0, T; H^{-1}(M))} + \|\partial_t y\|_{L^2(0, T; H^{-2}(M))} \leq \frac{C}{\varepsilon} (\|y_0\|_{H^{-1}(M)} + \varepsilon^{1/2} \|h\|_{L^2(0, T; H^{-1/2}(\partial M))}).$$

Such solutions in particular solve the first equation of (2.3) in the sense of distributions.

DEFINITION 2.6 (Controllability and cost). — Given  $(\theta, \varepsilon, T)$ , we say that (2.3) is null-controllable from  $(\theta, T)$  if for any  $y_0 \in H^{-1}(M)$ , there is  $h \in L^2(0, T; H^{-1/2}(\partial M))$  such that the associated solution to (2.3) satisfies  $y(T) = 0$ . If (2.3) is null-controllable from  $(\theta, T)$ , we define for  $y_0 \in H^{-1}(M)$  the set  $U(y_0) \subset \mathcal{U}$  of all such controls  $h \in L^2(0, T; H^{-1/2}(\partial M))$ , and the cost function

$$C_0(T, \varepsilon) := \sup_{\substack{y_0 \in H^{-1}(M) \\ \|y_0\|_{H^{-1}(M)} \leq 1}} \left\{ \inf_{h \in U(y_0)} \|h\|_{L^2(0, T; H^{-1/2}(\partial M))} \right\}.$$

We now describe the link with the boundary observation problem (1.9). We start with the duality identity.

LEMMA 2.7 (Duality equation). — For all solutions  $u \in C^0([0, T]; H_0^1(M))$  of (1.9) on  $[0, T]$  and all  $y \in C^0([0, T]; H^{-1}(M))$  solution to (2.1) with  $h \in L^2(0, T; H^{-1/2}(M))$ , we have

$$\langle hu(T), y_0 \rangle_{H_0^1, H^{-1}} - \langle hu_0, y(T) \rangle_{H_0^1, H^{-1}} - \int_0^T \langle h\theta \varepsilon \partial_\nu u|_{\partial M}(t), h(T-t) \rangle_{H^{1/2}(\partial M), H^{-1/2}(\partial M)} dt = 0.$$

The proof is omitted here and only consists in an integration by parts for smooth solutions, and then a density argument. As in the internal case, classical duality arguments (see [DR77] or [Cor07, Chap. 2.3]) yield the following statement.

COROLLARY 2.8 (Observability constant = control cost). — Given  $(\theta, \varepsilon, T)$ , Equation (2.3) is null-controllable from  $(\theta, T)$  if and only if the observability inequality (1.10) holds. Moreover, we then have  $C_0(T, \varepsilon) = C_0(T, \varepsilon)$ .

Again, all lower bounds on  $C_0(T, \varepsilon)$  formulated in Theorems 1.5 and 1.8 translate into lower bounds on  $C_0(T, \varepsilon)$ . The time  $T_{\text{unif}}(\theta)$  is equal to the minimal time of uniform controllability, and all lower bounds on the time  $T_{\text{unif}}(\theta)$  obtained in Theorems 1.5 and 1.8 and their corollaries apply.

2.2. THE VANISHING VISCOSITY LIMIT FOR GRADIENT FLOWS. CONJUGATION AND REFORMULATION

We focus in this article on the very particular case (1.6) where  $X$  is a gradient vector field (with respect to the same metric  $g$  defining the viscous perturbation  $\varepsilon\Delta_g$ ) of a weight function  $f : M \rightarrow \mathbb{R}$ , that is  $X = r_g f - r_g$ . In this case, it is classical (see e.g. [Wit82, HS85]) that the operator  $\varepsilon\Delta_g - r_g f - r_g - q$  can be conjugated to a “semiclassical selfadjoint operator”. Here,  $X \cdot Y$  is the inner product of the two vector fields  $X$  and  $Y$  given by the metric  $g$  in  $TM$ .

The first basic computation is the following:

$$e^{-f/2\varepsilon} \Delta_g e^{f/2\varepsilon} = \Delta_g + \frac{1}{\varepsilon} (r_g f - r_g) + \frac{jr_g f j_g^2}{4\varepsilon^2} + \frac{\Delta_g f}{2\varepsilon}.$$

We denote by

$$(2.4) \quad \frac{1}{\varepsilon^2} P_\varepsilon := \Delta_g + \frac{jr_g f j_g^2}{4\varepsilon^2} + \frac{\Delta_g f}{2\varepsilon} - \frac{q}{\varepsilon}, \quad \text{that is} \quad P_\varepsilon := \varepsilon^2 \Delta_g + \frac{jr_g f j_g^2}{4} + \varepsilon q_f,$$

where  $q_f = (\Delta_g f / 2) - q$ . The above computation implies that

$$(2.5) \quad e^{-f/2\varepsilon} \left( \frac{1}{\varepsilon^2} P_\varepsilon \right) e^{f/2\varepsilon} = \Delta_g - \frac{1}{\varepsilon} (r_g f - r_g) - \frac{q}{\varepsilon}.$$

The interest of this conjugation is that the operator  $P_\varepsilon$  is selfadjoint in  $L^2(M, d\text{Vol}_g)$  endowed with domain  $D(P_\varepsilon) = H^2(M) \setminus H_0^1(M)$ . Henceforth, the operator

$$\Delta_g - (1/\varepsilon)(r_g f - r_g) - q/\varepsilon$$

is also selfadjoint in  $L^2(M, e^{f/2\varepsilon} d\text{Vol}_g)$ . Let us now reformulate the uniform observability problem (1.4) in terms of the heat equation involving the operator  $P_\varepsilon$  defined in (2.4).

Note that the constant coefficient one-dimensional problem introduced in [CG05] enters the “gradient flow” setting with  $M = (0, L) \subset \mathbb{R}$ ,  $g = 1$ ,  $\Delta_g = \partial_x^2$ ,  $q = 0$ ,  $f = Mx$  for  $M \geq \mathbb{R}$ , and thus  $r_g f - r_g = M\partial_x$ . In that context, this form together with its formulation (2.4) have already been used in [CG05, Gla10, Lis14, Lis15].

LEMMA 2.9. — Given  $T_0, C_0, \varepsilon > 0$  and a function  $u$ , the following statements are equivalent.

(1) The function  $u$  solves

$$(2.6) \quad \begin{cases} (\partial_t - r_g f - r_g - q - \varepsilon\Delta_g)u = 0, & \text{in } (0, T_0) \times \text{Int}(M), \\ u = 0 & \text{on } (0, T_0) \times \partial M, \end{cases}$$

$$(2.7) \quad \text{resp.} \quad \|ku(T_0)\|_{L^2(M)}^2 \leq C_0^2 \int_0^{T_0} \|ku\|_{L^2(\omega)}^2 dt,$$

$$(2.8) \quad \text{resp.} \quad \|ku(T_0)\|_{H_0^1(M)}^2 \leq C_0^2 \int_0^{T_0} \|k\theta\varepsilon\partial_\nu u|_{\partial M}\|_{H^{1/2}(\partial M)}^2 dt.$$

(2) The function  $v(t, x) = e^{f(x)/2\varepsilon}u(t, x)$  solves

$$(2.9) \quad \begin{cases} \varepsilon \partial_t v + P_\varepsilon v = 0, & \text{in } (0, T_0) \quad \text{Int}(M), \\ v = 0 & \text{on } (0, T_0) \quad \partial M, \end{cases}$$

$$(2.10) \quad \text{resp.} \quad \|e^{f/2\varepsilon}v(T_0)\|_{L^2(M)}^2 \subset C_0^2 \int_0^{T_0} \|e^{f/2\varepsilon}v\|_{L^2(\omega)}^2 dt,$$

$$(2.11) \quad \text{resp.} \quad \|e^{f/2\varepsilon}v(T_0)\|_{H_0^1(M)}^2 \subset C_0^2 \int_0^{T_0} \|\theta e^{f/2\varepsilon} \varepsilon \partial_\nu v|_{\partial M}\|_{H^{1/2}(\partial M)}^2 dt.$$

(3) The function  $w(t, x) = v(t/\varepsilon, x) = e^{f(x)/2\varepsilon}u(t/\varepsilon, x)$  solves

$$(2.12) \quad \begin{cases} \partial_t w + \frac{1}{\varepsilon^2} P_\varepsilon w = 0, & \text{in } (0, T_0) \quad \text{Int}(M), \\ w = 0 & \text{on } (0, T_0) \quad \partial M, \end{cases}$$

$$\text{resp.} \quad \|e^{f/2\varepsilon}w(\varepsilon T_0)\|_{L^2(M)}^2 \subset C_0^2 \int_0^{\varepsilon T_0} \|e^{f/2\varepsilon}w\|_{L^2(\omega)}^2 dt,$$

$$(2.13) \quad \text{resp.} \quad \|e^{f/2\varepsilon}w(\varepsilon T_0)\|_{L^2(M)}^2 \subset C_0^2 \int_0^{\varepsilon T_0} \|\theta e^{f/2\varepsilon} \varepsilon \partial_\nu w\|_{H^{1/2}(\partial M)}^2 dt.$$

*Proof.* — Start e.g. with  $u$  satisfying (2.6). Using (2.5) and the definition of  $P_\varepsilon$ , Equation (2.6) rewrites equivalently as

$$\partial_t u + e^{f/2\varepsilon} \frac{1}{\varepsilon} P_\varepsilon e^{f/2\varepsilon} u = 0, \quad t \geq [0, T_0].$$

The function  $v = e^{f/2\varepsilon}u$  then satisfies (2.9) (and conversely). Setting  $w(t, x) = v(t/\varepsilon, x)$  then satisfies (2.12), and conversely.

The proof that (2.8), (2.11), (2.13) uses additionally that, on account to the Dirichlet boundary condition, we have  $\partial_\nu(e^{f/2\varepsilon}u)|_{\partial M} = e^{f/2\varepsilon} \partial_\nu u|_{\partial M}$ .

**2.3. (GCC) AND CONTROLLABILITY OF THE LIMIT EQUATION  $\varepsilon = 0$ .** — In this section, we characterize the observability inequality (1.2) for solutions of (1.1) in terms of the Geometric Control Condition (GCC). In this section,  $M$  is always assumed without boundary.

We denote by  $(\phi_t)_{t \in \mathbb{R}}$  the flow of the vector field  $X$ , namely

$$(2.14) \quad \dot{\phi}_t(x) = X(\phi_t(x)), \quad \phi_0(x) = x \in M.$$

This flow is globally defined on account to the compactness of  $M$ . We consider the following definition of the geometric control condition in the manifold  $M$  for the vector field  $X$  and the set  $\omega$ , which we denote (GCC).

**DEFINITION 2.10.** — Let  $M$  be a compact manifold without boundary,  $X$  a Lipschitz vector field on  $M$ ,  $\omega \subset M$ ,  $\chi \in C^0(M)$ ,  $I \subset \mathbb{R}$  and interval and  $T > 0$ . We say that

- $(M, X, \omega, I)$  satisfies (GCC) if for all  $x \in M$ , there is  $t \in I$  such that  $\phi_{-t}(x) \in \omega$ ;
- $(M, X, \omega, T)$  satisfies (GCC) if  $(M, X, \omega, (0, T))$  satisfies (GCC);
- $(M, X, \omega)$  satisfies (GCC) if there is  $T > 0$  such that  $(\omega, T)$  satisfies (GCC);
- $(M, X, \chi, T)$  satisfies (GCC) if  $(M, X, f_\chi \neq 0g, T)$  satisfies (GCC);
- $(M, X, \chi)$  satisfies (GCC) if  $(M, X, f_\chi \neq 0g)$  satisfies (GCC).

In this section, the manifold  $M$  is fixed. To lighten notation, we omit the dependence on  $M$  in  $(M, X, \omega, I)$  and we simply write  $(X, \omega, I)$  instead of  $(M, X, \omega, I)$  (with a similar notation for the other definitions).

Note in particular that an open set  $\omega$  satisfying (GCC) must contain all singular points of the vector field  $X$  (i.e., all points  $x \in M$  such that  $X(x) = 0$ ). We now provide with different reformulations of this property.

**LEMMA 2.11.** — *Let  $M$  be a compact manifold and  $X$  a Lipschitz vector field on  $M$ . Given  $\omega \subset M$  and  $T > 0$ , the following properties are equivalent:*

- (1)  $(X, \omega, T)$  satisfies (GCC);
- (2)  $\bigcup_{t \in (0, T)} \phi_t(\omega) = M$ ;
- (3)  $\bigcup_{t \in (0, T)} \phi_{T-t}(\omega) = \phi_T(M) = M$  after having applied  $\phi_T$ , which itself is equivalent to Item (2). Equivalence between Item (4) and Item (3) finally follows from the fact that the flow of  $-X$  is  $(\phi_{-t})_{t \in \mathbb{R}}$ .
- (4)  $(-X, \omega, T)$  satisfies (GCC).

*Proof.* — The definition of  $(X, \omega, T)$  satisfying (GCC) is equivalent to: for all  $x \in M$ , there is  $t \in (0, T)$  such that  $x \in \phi_t(\omega)$ . Equivalence between the Items (1) and (2) follows. Item (3) is equivalent to  $\bigcup_{t \in (0, T)} \phi_{T-t}(\omega) = \phi_T(M) = M$  after having applied  $\phi_T$ , which itself is equivalent to Item (2). Equivalence between Item (4) and Item (3) finally follows from the fact that the flow of  $-X$  is  $(\phi_{-t})_{t \in \mathbb{R}}$ .

**PROPOSITION 2.12.** — *Assume  $M$  is a compact manifold,  $X$  is a Lipschitz vector field on  $M$ ,  $ds$  is a positive density on  $M$ , and  $q \in L^1(M)$ . Given  $\omega \subset M$ ,  $\chi \in C^0(M)$  and  $T > 0$ , the following statements hold true:*

- (1) *If  $(X, \omega, T)$  satisfies (GCC), then the observability inequality (1.2) for solutions of (1.1) is true.*
- (2) *The observability inequality (1.2) for solutions of (1.1) implies that  $(X, \bar{\omega}, [0, T])$  satisfies (GCC).*
- (3) *The observability inequality*

$$(2.15) \quad C_0^2 \int_0^T \int_M j\chi(x)u(t, x)^2 ds(x) dt > ku(T)k_{L^2(M)}^2,$$

*for all  $u_0 \in L^2(M)$  and  $u$  solution of (1.1)*

*holds true if and only if  $(X, \chi, T)$  satisfies (GCC).*

- (4) *In all the above observability statements,  $ku(T)k_{L^2(M)}^2$  can be equivalently replaced by  $ku(0)k_{L^2(M)}^2$ .*

The proof below is inspired by that in [DL09, LL16] for the wave equation. It is constructive and would also yield a characterization of the HUM control operator (see e.g. [HKL15] or [Léa18, §1.2] for more on controllability/stabilization properties for transport equations).

*Proof.* — First notice that for  $u_0 \in L^2(M)$ , the unique solution to (1.1) is explicitly given by

$$u(t, x) = e^{\int_0^t q(\phi_t^{-1}(x)) d\tau} u_0(\phi_t(x)) \in C^0(\mathbb{R}; L^2(M)).$$

A first direct consequence is the existence of a constant  $C_{T,q} > 1$  such that

$$C_{T,q}^{-1} \|ku(0)\|_{L^2(M)}^2 \leq \|ku(T)\|_{L^2(M)}^2 \leq C_{T,q} \|ku(0)\|_{L^2(M)}^2, \quad \text{for all solutions to (1.1),}$$

which proves Item (4). Next, we write the observation term in (1.2) (the same holds for (2.15) if we replace  $1_\omega$  by  $\chi$ ) as

$$\begin{aligned} \int_0^T \int_\omega |ju(t, x)|^2 ds(x) dt &= \int_0^T \int_M 1_\omega(x) |ju(t, x)|^2 ds(x) dt \\ &= \int_0^T \int_M 1_\omega(x) u_0^2(\phi_t(x)) e^{2\int_0^t q(\phi_t^{-1}(x)) d\tau} ds(x) dt. \end{aligned}$$

Using the change of variable  $y = \phi_t(x)$  (see e.g. [Lee13, Prop. 16.42 p432]), we obtain

$$\int_0^T \int_\omega |ju(t, x)|^2 ds(x) dt = \int_0^T \int_M 1_\omega(\phi_t^{-1}(y)) u_0^2(y) e^{2\int_0^t q(\phi_t^{-1}(y)) d\tau} (\phi_t^{-1} ds)(y) dt$$

(note that  $\text{div}_{ds}(X)$  is defined by  $\frac{d}{dt}(\phi_t ds)|_{t=0} = \text{div}_{ds}(X) ds$ , so that this expression simplifies slightly in case  $\text{div}_{ds}(X) = 0$ ). Using that the density is positive on the compact  $M$ , we get the existence of  $C_T > 1$  such that

$$C_T^{-1} ds(y) \leq e^{2\int_0^t q(\phi_t^{-1}(y)) d\tau} (\phi_t^{-1} ds)(y) \leq C_T ds(y) \quad \text{uniformly for } (t, y) \in [0, T] \times M.$$

As a consequence, we obtain

$$\begin{aligned} C_T^{-1} \int_0^T \int_M 1_\omega(\phi_t^{-1}(y)) u_0^2(y) ds(y) dt &\leq \int_0^T \int_M 1_\omega(x) |ju(t, x)|^2 ds(x) dt \\ &\leq C_T \int_0^T \int_M 1_\omega(\phi_t^{-1}(y)) u_0^2(y) ds(y) dt. \end{aligned}$$

Hence setting

$$\begin{aligned} g_{\omega,T}(y) &:= \int_0^T 1_\omega(\phi_t^{-1}(y)) dt \in L^1(M) \\ \text{resp. } g_{\chi,T}(y) &:= \int_0^T \chi^2(\phi_t^{-1}(y)) dt \in C^0(M), \end{aligned}$$

we deduce

$$(2.16) \quad \begin{aligned} C_T^{-1} \int_M g_{\omega,T}(y) u_0^2(y) ds(y) &\leq \int_0^T \int_M 1_\omega(x) |ju(t, x)|^2 ds(x) dt \\ &\leq C_T \int_M g_{\omega,T}(y) u_0^2(y) ds(y). \end{aligned}$$

Recalling that  $\omega$  is an open set and  $M$  compact, together with Definition 2.10, we deduce that if  $(X, \omega, T)$  satisfies (GCC), then we have the existence of  $c > 0$  such that  $g_{\omega,T}(y) > c$  for a.e.  $y \in M$ . The lower bound in (2.16) then implies the observability inequality (1.2), and Item (1) follows.

Concerning Item (2), if  $(X, \bar{\omega}, [0, T])$  does not satisfy (GCC), then there is a point  $x_0 \in M$  such that  $\phi_{-t}(x_0) \setminus \bar{\omega} = \emptyset$  for all  $t \in [0, T]$ . The set  $\bar{\omega} \cap [0, T]$  being compact, there is a neighborhood  $U$  of  $x_0$  such that  $\phi_{-t}(U) \setminus \bar{\omega} = \emptyset$  for all  $t \in [0, T]$ . Setting  $u_0 = 1_U$ , we have on the one hand that  $k_{u_0} k_{L^2(M)} > 0$ . On the other hand, we have  $1_{\omega} \phi_{-t}(y) = 0$  for all  $y \in U$  and  $t \in (0, T)$ . This implies that  $g_{\omega, T} u_0 = g_{\omega, T}(y) 1_U = 0$  a.e. and, according to the upper bound in (2.16), that  $\int_0^T \int_{\omega} |ju(t, x)|^2 ds(x) dt = 0$ . This contradicts (2.16) and concludes the proof of Item (2).

Finally, the proof of Item (3) is split in two parts. That  $(X, \chi, T)$  satisfies (GCC) implies the observability inequality (2.15) follows as in the proof of Item (1). Now assume that  $(X, \chi, T)$  does not satisfy (GCC). Then there is a point  $x_0 \in M$  such that  $\phi_{-t}(x_0) \setminus \bar{\chi} \cap \text{supp } g = \emptyset$  for all  $t \in (0, T)$ . Hence, we have

$$g_{\chi, T}(x_0) = \int_0^T \chi^2(\phi_{-t}(x_0)) dt = 0.$$

We now choose a sequence of continuous real-valued initial data  $(u_0^n)_{n \in \mathbb{N}}$  such that  $k_{u_0^n} k_{L^2(M)} = 1$  and  $(u_0^n)^2(x) ds(x) \rightharpoonup \delta_{x_0}$  in the sense of measures on  $M$ . The fact that  $g_{\chi, T}$  is continuous on  $M$  together with the upper bound in (2.16) implies that, denoting by  $u^n$  the solution of (1.1) with initial datum  $u_0^n$ , we have

$$\int_0^T \int_M |j\chi(x) u^n(t, x)|^2 ds(x) dt \leq C_T \int_M g_{\chi, T}(y) (u_0^n)^2(y) ds(y) \\ \leq C_T h \delta_{x_0}, g_{\chi, T} \in \text{Meas}, C^0 = 0,$$

which contradicts the observability inequality (2.15), and concludes the proof of Item (3).

### 3. GENERAL LOWER BOUNDS WITHOUT GEOMETRIC ASSUMPTION

In this section, we consider a general manifold (with or without boundary)  $M$ , and prove the lower bound for the minimal time of uniform controllability provided in Theorem 1.5. We also give a proof of Theorem 1.6 as a corollary. To do this, we use the semiclassical reformulation (2.9)-(2.10) of the problem in Lemma 2.9, as well as exponential decay properties of eigenfunctions of the operator  $P_\varepsilon$ . We rely on the Helffer-Sjöstrand theory as developed in [HS84, Hel88, DS99]. All results presented in this section apply as well for the semiclassical heat equation.

The result of Theorem 1.5 is stated for a potential minimum. However, we shall prove a seemingly more general result, at any energy level in  $V(M)$ . We shall then explain why this latter result is not more general, and how it can be improved in dimension one. We recall the definition of  $V$  in terms of  $f$  in (1.11), and define the classically allowed region at any energy level  $E$ :

$$K_E = \{x \in M; V(x) \leq E\}.$$



We then define the Agmon distance (see e.g. [Hel88, Chap. 3]) to the set  $K_E$  at the energy level  $E$ :

$$(3.1) \quad \begin{aligned} d_{A,E}(x, y) &= \inf \left\{ \int_0^1 \sqrt{(V(\gamma(t)) - E)_+} |\dot{\gamma}(t)| dt ; \gamma \in \mathbf{U}_1(x, y) \right\}, \\ \mathbf{U}_1(x, y) &= \{ \gamma \in W^{1,1}([0, 1]; M) ; \gamma(0) = x, \gamma(1) = y \}, \\ d_{A,E}(x) &= \inf_{y \in K_E} d_{A,E}(x, y). \end{aligned}$$

That is to say,  $d_{A,E}(x)$  is the distance of  $x$  to the set  $K_E$  for the (pseudo-)metric  $(V - E)_+ g$ . Here again  $(V(x) - E)_+ = \max(V(x) - E, 0)$ . We will use, as in (1.12) the notation  $d_A = d_{A,E_0}$  where  $E_0 = \min_M V$  for the Agmon distance at the bottom energy. Note that  $d_{A,E}$  vanishes identically on  $K_E$  (and only on this set). Finally, an important function in the estimates below is given by

$$(3.2) \quad W_E(x) = d_{A,E}(x) + \frac{f(x)}{2}.$$

We shall prove in this section the following result.

**THEOREM 3.1.** — *Assume the observability estimate (2.7) (resp. the boundary observability inequality (2.8)) for all solutions to (2.6) with constant  $C_0 = C_0(T_0, \varepsilon)$ . Then, for all  $E \in V(M) = [\min_M V, \max_M V]$  and all  $\delta > 0$ , there is  $\varepsilon_0 > 0$  such that we have for all  $\varepsilon \in (0, \varepsilon_0)$*

$$C_0(T_0, \varepsilon) > \exp \frac{1}{\varepsilon} \left( \min_{\bar{\omega}} W_E - \max_{K_E} W_E - \delta - ET_0 \right),$$

*in the internal observation case, and*

$$C_0(T_0, \varepsilon) > \exp \frac{1}{\varepsilon} \left( \min_{\bar{\Gamma}} W_E - \max_{K_E} W_E - \delta - ET_0 \right),$$

*in the boundary observation case.*

*In particular, we have for all  $E \in V(M)$ , for each respective case,*

$$\begin{aligned} T_{\text{unif}}(\omega) &> \frac{1}{E} \left( \min_{\bar{\omega}} W_E - \max_{K_E} W_E \right), \\ T_{\text{unif}}(\Gamma) &> \frac{1}{E} \left( \min_{\bar{\Gamma}} W_E - \max_{K_E} W_E \right). \end{aligned}$$

Theorem 1.5 is then the particular case  $E = E_0 = \min_M V$  in Theorem 3.1. Unfortunately, the function  $E \mapsto (1/E)(\min_{\bar{\omega}} W_E - \max_{K_E} W_E)$  is a decreasing function of  $E$ . Indeed,

– the sets  $K_E$  are increasing in  $E$ , hence the function  $E \mapsto \max_{K_E} W_E = \max_{K_E} f/2$  increases;

–  $E \mapsto d_{A,E}(x)$  is decreasing in  $E$ , hence the function  $E \mapsto \min_{\bar{\omega}} W_E$  decreases.

Therefore, the estimate of Theorem 3.1 simply reduces to that for  $E = E_0$ , that is Theorem 1.5 in the introduction. This comes from the fact that the estimate involving the term  $\max_{K_E} W_E$  is very rough (see Section 3.2 below for a more precise discussion). This can be improved in the one-dimensional one-well case, see [LL21b].

Moreover, it is interesting to notice that the proof of Theorem 3.1 is not more involved than the direct proof of Theorem 1.5, and we shall re-use part of it in case of revolution surfaces in Section 4.

**3.1. EIGENFUNCTIONS OF SEMICLASSICAL SCHRÖDINGER OPERATORS.** — In this section, we collect classical results concerning eigenfunctions of semiclassical Schrödinger operators, and some of their decay properties. Recall that

$$P_\varepsilon = \varepsilon^2 \Delta_g + \frac{j\Gamma_g f_g^2}{4} + \varepsilon q_f = \varepsilon^2 \Delta_g + V + \varepsilon q_f$$

is defined in (2.4). We first need to prove existence of eigenfunctions near any energy level.

**LEMMA 3.2 (Existence of eigenfunctions).** — *Assume  $V \in W^{1,1}(M)$  and  $q_f \in L^1(M)$  are both real valued. For all  $E \in V(M) = [\min_M V, \max_M V]$  and all  $\varepsilon \in (0, 1]$ , there is  $E_\varepsilon = E + O(\varepsilon^{2/3})$  and  $\psi_\varepsilon \in H^2(M) \setminus H_0^1(M)$  such that  $P_\varepsilon \psi_\varepsilon = E_\varepsilon \psi_\varepsilon$ .*

Note that the  $O(\varepsilon^{2/3})$  precision is relatively poor, and can be improved in different situations (e.g. if there is a critical point of  $V$  at energy  $E$ ). These refinements are however not needed here.

*Proof.* — The proof consists in constructing a (very rough) quasimode. Assume first that  $E$  is reached by an interior point, i.e., there is  $x_0 \in \text{Int}(M)$  such that  $V(x_0) = E$ . We then only work in a local chart near  $x_0$ , centered at  $x_0$  (hence we work in  $\mathbb{R}^n$  in a neighborhood of 0).

We take a cutoff function  $\chi \in C_c^1(\mathbb{R})$  such that  $\chi = 1$  in a neighborhood of 0. We set  $u^\varepsilon(x) = \varepsilon^{-n/3} \chi(\varepsilon^{-2/3} jx)$ , so that  $u^\varepsilon$  is smooth and moreover supported in the chart for all  $\varepsilon < \varepsilon_0$  with  $\varepsilon_0$  sufficiently small, and thus  $u^\varepsilon \in H^2(M) \setminus H_0^1(M)$ . Notice also that

$$k u^\varepsilon k_{L^2(M)}^2 = \int j u^\varepsilon(x)^2 \sqrt{jg}(x) dx = \int \chi^2(jy) \sqrt{jg}(\varepsilon^{2/3} y) dy = c_0 + O(\varepsilon^{2/3}),$$

with  $c_0 > 0$ . We now estimate  $(P_\varepsilon - E)u^\varepsilon$ . For this, we first have  $k \varepsilon q_f u^\varepsilon k_{L^2} \in C\varepsilon$ . Second, we always have the rough estimate  $V(x) - E = V(x) - V(0) = O(jx)$  so that we have

$$(3.3) \quad k(V - E)u^\varepsilon k_{L^2}^2 \in C \int jx^2 \chi^2(\varepsilon^{-2/3} jx) \frac{dx}{\varepsilon^{2n/3}} \in C\varepsilon^{4/3}.$$

Third, we have

$$\begin{aligned} \|\varepsilon^2 \Delta_g u^\varepsilon\|_{L^2}^2 &= \left\| \frac{\varepsilon^2}{\sqrt{jg}} \sum_{i,j} \partial_i (g^{ij} \sqrt{jg} \partial_j u^\varepsilon) \right\|_{L^2}^2 \\ &\in C\varepsilon^4 \sum_{i,j} \int \left[ \partial_i (g^{ij} \sqrt{jg} x_j / jx) \varepsilon^{-2/3} \chi^\theta(\varepsilon^{-2/3} jx) \right. \\ &\quad \left. + (g^{ij} \sqrt{jg} x_j / jx) \frac{x_i}{jx} \varepsilon^{-4/3} \chi^{\theta\theta}(\varepsilon^{-2/3} jx) \right]^2 \frac{dx}{\varepsilon^{2n/3}} \\ (3.4) \quad &\in C\varepsilon^{4/3}. \end{aligned}$$

Combining the above four estimates yields the existence of  $D, \varepsilon_0 > 0$  such that for all  $\varepsilon < \varepsilon_0$ , we have,

$$k(P_\varepsilon - E)u^\varepsilon k_{L^2} \leq C\varepsilon^{2/3} \leq D\varepsilon^{2/3} k u^\varepsilon k_{L^2}.$$

Hence, if  $E \notin \text{Sp}(P_\varepsilon)$ , this implies  $\|(P_\varepsilon - E)^{-1}\|_{L^2 \rightarrow L^2} > (D\varepsilon^{2/3})^{-1}$ . Finally, the operator  $P_\varepsilon$  being selfadjoint, we have, for  $z \in \mathbb{C} \setminus \text{Sp}(P_\varepsilon)$ ,  $k(P_\varepsilon - z)^{-1}k_{L^2 \rightarrow L^2} = 1/d(z, \text{Sp}(P_\varepsilon))$ , so that, if  $E \notin \text{Sp}(P_\varepsilon)$ ,

$$\frac{1}{d(E, \text{Sp}(P_\varepsilon))} > (D\varepsilon^{2/3})^{-1}.$$

In any case, this implies  $d(E, \text{Sp}(P_\varepsilon)) \leq D\varepsilon^{2/3}$ , and using that the spectrum of  $P_\varepsilon$  is purely pointwise, this proves the sought result.

Assume now that  $E$  is not reached by an interior point, i.e.,  $E \notin V(\text{Int}(M))$ . This means in particular that there is  $x_0 \in \partial M$  such that  $V(x_0) = E$ . Then, we again work in a local chart near  $x_0$ , centered at  $x_0$ . In this chart,  $M$  is given by  $\mathbb{R}^{n-1} \times \mathbb{R}$  and  $x_0$  by 0. We denote  $(x^0, x_n) \in \mathbb{R}^{n-1} \times \mathbb{R}$  local coordinates. We then take  $\chi$  as above and further define  $\tilde{\chi} \in C_c^\infty(\mathbb{R})$ , non-identically vanishing, such that  $\text{supp}(\tilde{\chi}) \subset (-1, 0)$ . We define  $u^\varepsilon(x) = \varepsilon^{-n/3} \tilde{\chi}(\varepsilon^{-2/3} x_n) \chi(\varepsilon^{-2/3} |x^0|)$ . One can check that all above properties of  $u^\varepsilon$  are still satisfied, and in particular (3.3)–(3.4). In addition, we have  $\text{supp} u^\varepsilon \subset \mathbb{R}^{n-1} \times \mathbb{R}$ , and thus  $u^\varepsilon \in H^2(M) \setminus H_0^1(M)$ . The remainder of the proof then follows the same as in the first case.

**REMARK 3.3.** — Note that near a noncritical value of  $V$ , or near the boundary of  $M$ , the appropriate local model is  $-\varepsilon^2 \partial_x^2 - x$ . Considering concentrating quasimodes of the form  $\chi(x/\varepsilon^\alpha)$  leads to

$$\varepsilon^2 \partial_x^2 (\chi(x/\varepsilon^\alpha)) = \frac{\varepsilon^2}{\varepsilon^{2\alpha}} \chi''(x/\varepsilon^\alpha), \quad \text{and} \quad x \chi(x/\varepsilon^\alpha) = \varepsilon^\alpha (x/\varepsilon^\alpha) \chi(x/\varepsilon^\alpha).$$

Henceforth, the right scaling is given by  $2 - 2\alpha = \alpha$ , that is  $\alpha = 2/3$ . The quasimode we construct are then  $O(\varepsilon^{2/3})$ . If one wants to obtain a better remainder, one could replace  $\chi$  by an Airy function, as one should replace  $\chi$  by a Hermite function in the case of bottom of potential (in which case the precision of the quasimode is improved). Also, the remainder  $\varepsilon^{2/3}$  is actually the worst possible case.

The next result states the decay estimates for eigenfunctions in the classically forbidden region, and is a consequence of so-called Agmon estimates (see [HS84, Hel88, DS99]). Here, it is a particular case of [Hel88, Prop. 3.3.1 & 3.3.4]. Note that with respect to [Hel88, Prop. 3.3.1], our operator  $P_\varepsilon$  contains an additional term, namely multiplication by  $\varepsilon q_f$ . However, this contribution is of lower order and can be absorbed in the proof of the Agmon estimates.

**THEOREM 3.4** (Decay of eigenfunctions in the classically forbidden region)

Assume  $V \in W^{1,1}(M)$  and  $q_f \in L^1(M)$ . Let

$$E \in V(M) = [\min_M V, \max_M V]$$

and assume  $\psi_\varepsilon \in H^2(M) \setminus H_0^1(M)$  and  $E_\varepsilon$  satisfy

$$(3.5) \quad P_\varepsilon \psi_\varepsilon = E_\varepsilon \psi_\varepsilon, \quad k\psi_\varepsilon k_{L^2(M)} = 1, \quad \max(E_\varepsilon - E, 0) = o(1) \text{ as } \varepsilon \rightarrow 0^+.$$

Then for all  $\delta > 0$ , there exist  $C = C(\delta)$ ,  $\varepsilon_0 = \varepsilon_0(\delta) > 0$  such that, for all  $\varepsilon < \varepsilon_0$ , we have

$$(3.6) \quad \|e^{1/\varepsilon d_{A,E}} \psi_\varepsilon\|_{L^2(M)} \leq C e^{\delta/\varepsilon}.$$

Assuming further that  $V, q_f \in C^1(M)$ , we have  $\psi_\varepsilon \in C^1(M)$  and for all  $\delta > 0$  and all smooth vector field  $Y$  on  $M$ , there exist  $C = C(\delta)$ ,  $\varepsilon_0 = \varepsilon_0(\delta) > 0$  such that, for all  $0 < \varepsilon < \varepsilon_0$ , we have

$$(3.7) \quad |j\psi_\varepsilon(x)| + |j(Y\psi_\varepsilon)(x)| \leq C e^{-(1/\varepsilon)(d_{A,E}(x) - \delta)}, \quad \text{for all } x \in M.$$

**REMARK 3.5.** — Note that the smoothness assumption  $V, q_f \in C^1(M)$  (as well as the smoothness assumptions in Theorem 1.5) is essentially only used in [HS84, Hel88, DS99] to perform elliptic regularity estimates so that the pointwise estimate (3.7) makes sense. A finer (much less demanding) regularity assumption can be formulated.

As a direct corollary, we have that most of the norm of  $\psi_\varepsilon$  is near  $K_E$ , see [Hel88, Cor. 3.3.2].

**COROLLARY 3.6** (Most of the norm is in the classically allowed region)

Let  $E \in V(M) = [\min_M V, \max_M V]$  and assume  $\psi_\varepsilon, E_\varepsilon$  satisfy (3.5). For any open set  $U$  containing  $K_E$ , there is  $\delta, \varepsilon_0 > 0$  such that for all  $0 < \varepsilon < \varepsilon_0$ , we have

$$k\psi_\varepsilon k_{L^2(U)}^2 = 1 + O(e^{-\delta/\varepsilon}).$$

### 3.2. ROUGH LOCALIZATION OF EIGENFUNCTIONS, AND A PROOF OF THEOREM 3.1

From the decay estimates in the classically forbidden region (Theorem 3.4) and the rough localization of the  $L^2$  mass of eigenfunctions (Corollary 3.6), together with the existence of eigenfunctions at any energy level (Lemma 3.2), we may now deduce a proof of Theorem 3.1. Recall that  $W_E$  is defined in (3.2). We first prove the following proposition, from which Theorem 3.1 will follow.

**PROPOSITION 3.7.** — Let  $E \in V(M) = [\min_M V, \max_M V]$  and assume  $\psi_\varepsilon \in H^2(M) \setminus H_0^1(M)$  and  $E_\varepsilon$  satisfy (3.5). Then for all  $\delta > 0$ , there exists  $\varepsilon_0 = \varepsilon_0(\delta) > 0$  such that, for any open set  $\omega \subset M$  and for all  $\varepsilon < \varepsilon_0$ , we have

$$(3.8) \quad \|e^{f/2\varepsilon} \psi_\varepsilon\|_{L^2(M)} > e^{-(1/\varepsilon)(\max_{K_E} W_E + \delta)},$$

$$(3.9) \quad \|e^{f/2\varepsilon} \psi_\varepsilon\|_{L^2(M)} \leq e^{(1/\varepsilon)(\min_M W_E - \delta)},$$

$$(3.10) \quad \|e^{f/2\varepsilon} \psi_\varepsilon\|_{L^2(\omega)} \leq e^{(1/\varepsilon)(\min_{\bar{\omega}} W_E - \delta)}.$$

Assuming also that  $V, q_f \in C^1(M)$  and  $\Gamma \subset \partial M$ , we have

$$(3.11) \quad \|e^{f/2\varepsilon} \partial_\nu \psi_\varepsilon\|_{L^2(\Gamma)} \leq e^{(1/\varepsilon)(\min_\Gamma W_E - \delta)}.$$

Note that Estimate (3.8) is very rough, due to our lack of knowledge on the localization of  $\psi_\varepsilon$  in the classically allowed region  $K_E$ . In the one-dimensional one-well case, this bound can be refined, see the companion paper [LL21b].

We first prove Proposition 3.7, and then deduce a proof of Theorem 3.1.

*Proof of Proposition 3.7.* — First, setting  $U_\delta = \{x \in M; f(x) < \max_{K_E} f + \delta\}$ , we have

$$\begin{aligned} \|e^{f/2\varepsilon}\psi_\varepsilon\|_{L^2(M)} &> \|e^{f/2\varepsilon}\psi_\varepsilon\|_{L^2(U_\delta)} > e^{-(1/\varepsilon)(\max_{K_E} f + \delta/2)} \|\psi_\varepsilon\|_{L^2(U_\delta)} \\ &> \frac{1}{2} e^{-(1/\varepsilon)(\max_{K_E} f + \delta/2)}, \end{aligned}$$

after having used Corollary 3.6 in the last inequality for  $\varepsilon < \varepsilon_0(\delta)$ . This proves (3.8), recalling that  $d_{A,E} = 0$  on  $K_E$ .

Second, notice that (3.9) is a consequence of (3.10). Third, to prove (3.10), we use (3.6) as follows

$$\begin{aligned} \|e^{f/2\varepsilon}\psi_\varepsilon\|_{L^2(\omega)} &\leq \|e^{f/2\varepsilon} e^{-(1/\varepsilon)d_{A,E}} e^{(1/\varepsilon)d_{A,E}} \psi_\varepsilon\|_{L^2(\omega)} \\ &\leq e^{-(1/\varepsilon)(\min_{\bar{\omega}}(f/2 + d_{A,E}))} \|e^{(1/\varepsilon)d_{A,E}} \psi_\varepsilon\|_{L^2(\omega)} \\ &\leq e^{-(1/\varepsilon)(\min_{\bar{\omega}}(f/2 + d_{A,E}))} C e^{\delta/\varepsilon}. \end{aligned}$$

Finally, to prove (3.11), we proceed similarly using (3.7) (instead of the sole (3.6)) with  $Y = \partial\nu$ . We have

$$\|e^{f/2\varepsilon}\partial\nu\psi_\varepsilon\|_{L^2(\Gamma)} \leq \|e^{f/2\varepsilon} C e^{-(1/\varepsilon)(d_{A,E} - \delta)}\|_{L^2(\Gamma)} \leq C e^{-(1/\varepsilon)(\min_{\Gamma}(f/2 + d_{A,E}) - \delta)},$$

which concludes the proof of the proposition.

*Proof of Theorem 3.1 from Proposition 3.7 and Lemma 3.2.* — We use the reformulation in Lemma 2.9 and consider the observability estimate (2.10) for solutions to the evolution equation (2.9).

More precisely, we select  $E \geq V(M)$ , and we let  $v_\varepsilon$  be the solution to (2.9) associated to the initial condition  $v_\varepsilon(0) = \psi_\varepsilon$ , where  $\psi_\varepsilon$  is given by Lemma 3.2. That is to say,  $v_\varepsilon(t, x) = e^{-(E_\varepsilon/\varepsilon)t}\psi_\varepsilon(x)$ . We estimate both sides of (2.10).

Firstly, using (3.8), we have

$$\begin{aligned} \|e^{f/2\varepsilon}v_\varepsilon(T_0)\|_{L^2(M)} &= e^{-(E_\varepsilon/\varepsilon)T_0} \|e^{f/2\varepsilon}\psi_\varepsilon\|_{L^2(M)} \\ &> \frac{1}{2} e^{-(E_\varepsilon/\varepsilon)T_0} e^{-(1/\varepsilon)(\max_{K_E} f + \delta/2)}, \end{aligned}$$

for  $\varepsilon < \varepsilon_0(\delta)$ . Recalling that  $E_\varepsilon = E + O(\varepsilon^{2/3})$ , this implies the existence of  $\varepsilon_0(\delta) > 0$  such that for  $\varepsilon < \varepsilon_0(\delta)$ ,

$$\|e^{f/2\varepsilon}v_\varepsilon(T_0)\|_{L^2(M)} > \frac{1}{2} e^{-(1/\varepsilon)(ET_0 + \max_{K_E} f + \delta)},$$

Secondly, concerning the internal case, using (3.10), we have

$$\begin{aligned} \int_0^{T_0} \|e^{-f/2\varepsilon} v_\varepsilon\|_{L^2(\omega)}^2 dt &= \int_0^{T_0} e^{-2(E_\varepsilon/\varepsilon)t} \|e^{-f/2\varepsilon} \psi_\varepsilon\|_{L^2(\omega)}^2 dt \\ &= \frac{\varepsilon}{2E_\varepsilon} (1 - e^{-2(E_\varepsilon/\varepsilon)T_0}) \|e^{-f/2\varepsilon} \psi_\varepsilon\|_{L^2(\omega)}^2 \leq C e^{-(2/\varepsilon)(\min_{\overline{\omega}} W_E - \delta)}. \end{aligned}$$

As a consequence of these two estimates, applying (2.10) implies

$$e^{-(2/\varepsilon)(ET_0 + \max_{K_E} W_E + \delta)} \leq C_0(T_0, \varepsilon)^2 C e^{-(2/\varepsilon)(\min_{\overline{\omega}} W_E - \delta)},$$

which concludes the proof.

In the case of observation from part of the boundary, we simply replace the above use of (3.10) by that of (3.11).

**3.3. AN EXPLICIT COUNTER-EXAMPLE FOR A DOMAIN OF  $\mathbb{R}^n$ .** — The purpose of this section is to prove Theorem 1.6. Here  $M = \overline{\Omega}$  where  $\Omega \subset \mathbb{R}^n$  is an open set, endowed with the Euclidean metric. For  $\delta > 0$ , we may assume, up to an appropriate translation of the domain  $\Omega$ , that there is  $\eta > 0$  such that

$$(3.12) \quad \overline{B}(0, \eta) \subset \Omega.$$

$$(3.13) \quad \omega = (\Omega \setminus \{x_1 > 0, x_2 > 0, \dots, x_n > 0\}), \quad \text{and} \quad B(0, \eta) \setminus \omega = ?.$$

We let  $f_\lambda(x_1, \dots, x_n)$  be defined as follows:

$$(3.14) \quad f_\lambda(t) := \int_0^t \sqrt{\lambda^2 s^2 + 1} ds \quad \text{and} \quad f_\lambda(x_1, \dots, x_n) := \sum_{i=1}^n f_\lambda(x_i).$$

With this definition, the associated gradient vector field and potential are given respectively by

$$\begin{aligned} r f_\lambda(x_1, \dots, x_n) &= \sum_{i=1}^n f_\lambda'(x_i) e_i = \sum_{i=1}^n \sqrt{\lambda^2 x_i^2 + 1} e_i, \\ V_\lambda(x) &= \frac{j r f_\lambda^2}{4} = \frac{\lambda^2 j x^2 + n}{4}, \end{aligned}$$

where  $(e_1, \dots, e_n)$  denotes the canonical basis of  $\mathbb{R}^n$ .

The proof of Theorem 1.6 now directly follows from the following Lemmas 3.8 and 3.9, when taking  $\lambda$  large enough.

**LEMMA 3.8.** — *In the above setting, recalling (3.12), (3.13) and that  $f_\lambda$  is defined in (3.14), we have  $T_{\text{unif}}(\omega) > \lambda \eta^2 / n$  for all  $\lambda > 0$ .*

**LEMMA 3.9.** — *In the above setting, recalling (3.12), (3.13) and that  $f_\lambda$  is defined in (3.14),  $(\overline{\Omega}, r f_\lambda, \omega)$  and  $(\overline{\Omega}, r f_\lambda)$  both satisfy (FC) and we have for all  $\lambda > 0$*

$$(3.15) \quad \begin{aligned} T_{\text{FC}}(\overline{\Omega}, r f_\lambda, \omega) &\leq T_{\text{FC}}(\overline{\Omega}, r f_\lambda) \leq \min_{\substack{v \in \mathbb{R}^n, |v_j|=1 \\ v_{e_i} > 0.8 e_i}} \left( \max_{x \in \overline{\Omega}} x \vee \min_{x \in \overline{\Omega}} x \right) \\ &\leq \text{diam}(\Omega). \end{aligned}$$

*In particular, both are bounded by a constant uniformly in  $\lambda > 0$ .*

We recall that (FC) and  $T_{FC}$  are defined in this context in Definition 1.4. Notice that the quantity  $\max_{x \in \bar{\Omega}} x \cdot \nu - \min_{x \in \bar{\Omega}} x \cdot \nu$  represents the minimal Euclidean distance between two parallel hyperplanes (normal to  $\nu$ ) such that  $\Omega$  is contained between the two hyperplanes.

*Proof of Lemma 3.8.* — The minimum of  $V_\lambda$  is reached at  $x_{\min} = 0$  and  $V_\lambda(x_{\min}) = n/4$ . The Agmon distance (3.1) at the bottom energy  $V_\lambda(x_{\min})$  can be explicitly computed for points  $x \in \bar{B}(0, \eta) \cap \Omega \cap \omega$ . Indeed, for  $x \in \Omega$  we have

$$\sqrt{V_\lambda(x) - V_\lambda(x_{\min})} = \frac{\lambda}{2} |x|$$

and thus for  $x \in \bar{B}(0, \eta)$ ,

$$\begin{aligned} d_A(x) &= \frac{\lambda}{2} \inf \left\{ \int_0^1 |\dot{\gamma}(t)|^2 dt, \gamma \in W^{1,1}([0, 1]; \mathbb{M}), \gamma(0) = 0, \gamma(1) = x \right\} \\ &= \frac{\lambda |x|^2}{2} \int_0^1 t dt = \frac{\lambda |x|^2}{4}, \end{aligned}$$

where, by symmetry arguments, we have noticed that the straight line  $\gamma(t) := xt$  reaches the infimum for  $x \in \bar{B}(0, \eta)$ . Recalling that  $V_\lambda(0) = n/4$  and  $f(0) = 0$ , application of (1.13) in Theorem 1.5 implies

$$(3.16) \quad \frac{n}{4} T_{\text{unif}}(\omega) = V_\lambda(0) T_{\text{unif}}(\omega) > \min_{\bar{\omega}} (f/2 + d_A).$$

By a connectedness argument, for any  $x \in \bar{\omega}$ , we have from (3.13)

$$d_A(x) > \min_{y \in \partial B(0, \eta)} d_A(y) > \lambda \eta^2 / 4.$$

Moreover, the condition (3.13) together with the definition of  $f_\lambda$  in (3.14) imply that  $f_\lambda(x) > 0$  for  $x \in \bar{\omega}$ . We thus have

$$\min_{\bar{\omega}} (f_\lambda/2 + d_A) > \min_{\bar{\omega}} (d_A) > \frac{\lambda \eta^2}{4}.$$

Combined with (3.16), this concludes the proof of the lemma.

*Proof of Lemma 3.9.* — Notice first that given  $\nu \in \mathbb{R}^n$  with  $|\nu| = 1$  such that  $e_i \cdot \nu > 0$  for all  $i \in \{1, \dots, n\}$ , we have for all  $x \in \bar{\Omega}$ ,

$$(3.17) \quad r f_\lambda(x) \cdot \nu = \sum_{i=1}^n \sqrt{\lambda^2 x_i^2 + 1} e_i \cdot \nu > \sum_{i=1}^n e_i \cdot \nu > \sqrt{\sum_{i=1}^n (e_i \cdot \nu)^2} = |\nu| = 1.$$

Second, following Definition 1.4, we extend  $f_\lambda$  as a smooth compactly supported function  $\tilde{f}_\lambda$  in  $\mathbb{R}^n$ . Given  $x \in \bar{\Omega}$ , we denote by  $\gamma_x$  the maximal (global) solution to  $\dot{\gamma}_x(t) = r \tilde{f}_\lambda(\gamma_x(t))$  with  $\gamma_x(0) = x$ , defined in  $\mathbb{R}^n$  for  $t \in \mathbb{R}$ .

Given  $\nu \in \mathbb{R}^n$  with  $|\nu| = 1$  such that  $e_i \cdot \nu > 0$ , Estimate (3.17) thus implies that  $\dot{\gamma}_x(t) \cdot \nu = r \tilde{f}_\lambda(\gamma_x(t)) \cdot \nu \in [1, \infty)$ . Integrating this between 0 and  $t$ , we obtain  $\gamma_x(t) \cdot \nu \geq x \cdot \nu + t$ . Assuming that  $T > \max_{y \in \bar{\Omega}} y \cdot \nu - \min_{y \in \bar{\Omega}} y \cdot \nu$  thus implies

$$\gamma_x(T) \cdot \nu \geq x \cdot \nu + T < \left( \max_{y \in \bar{\Omega}} y \cdot \nu - \min_{y \in \bar{\Omega}} y \cdot \nu \right),$$

that is

$$x \vee \gamma_x(T) \vee > \max_{y \in \bar{\Omega}} y \vee \min_{y \in \bar{\Omega}} y \vee.$$

Since  $x \geq \bar{\Omega}$ , this implies  $\gamma_x(T) \not\leq \bar{\Omega}$ . This holds true for any  $x \geq \bar{\Omega}$ . Recalling the definition of (FC) in Definition 1.4, we have obtained that this condition is satisfied by both  $(\bar{\Omega}, r f_\lambda, T)$  and  $(\bar{\Omega}, r f_\lambda, \omega, T)$ . Moreover, given the definition of  $T_{FC}$  as an infimum, we have also obtained that

$$T_{FC}(\bar{\Omega}, r f_\lambda, \omega) \leq T_{FC}(\bar{\Omega}, r f_\lambda) \leq \max_{y \in \bar{\Omega}} y \vee \min_{y \in \bar{\Omega}} y \vee.$$

Since this holds for all  $v$ , we have proved (3.15), which concludes the proof of the lemma.

#### 4. SURFACES OF REVOLUTION

4.1. GENERAL SETTING. — In this section we introduce the geometric setting for the results presented in Section 1.3.2. We are concerned with a revolution surface  $S \subset \mathbb{R}^3$  being either

- (1) Case 1: diffeomorphic to a sphere  $S^2$  (in which case  $\partial S = \emptyset$ );
- (2) Case 2: diffeomorphic to a disk (in which case  $\partial S$  is a circle embedded in  $\mathbb{R}^3$ );
- (3) Case 3: diffeomorphic to a cylinder  $[0, 1] \times S^1 \subset \mathbb{R}^3$  (in which case  $\partial S$  consists in two disjoint circles embedded in  $\mathbb{R}^3$ , and belonging to two parallel hyperplanes).

We follow [LL21a] and [Bes78, Chap. 4B p95] for the precise geometric description of such manifolds. At some places, we also consider the case of the torus  $T^2$ , endowed with a metric invariant in one direction. This setting does not strictly speaking enter the framework of the present section, but is much simpler to describe (and we thus do not mention it in the present section).

*Definition and differentiable structure.* — We assume that  $(S, g)$  is an embedded 2D submanifold of  $\mathbb{R}^3$  (endowed with the induced Euclidean structure), having  $S^1 = (\mathbb{R}/2\pi\mathbb{Z}) \times SO(2)$  as an effective isometry group. The action of  $S^1$  on  $S$ , denoted by  $\theta \curvearrowright \mathbb{R}_\theta$  (such that  $\mathbb{R}_\theta S = S$ ) has:

(1) exactly two fixed points denoted by  $N, S \in S$  called North and South poles in Case 1; we write  $\partial_N S = fNg$  and  $\partial_S S = fSg$ ;

(2) exactly one fixed point denoted by  $N \in S$  called North pole in Case 2; in this case, we write  $\partial_N S = fNg$ , and  $\partial_S S = \partial S$  has a single connected component (called "south boundary") which is also invariant by  $\mathbb{R}_\theta$ ;

(3) no fixed point in Case 3; in this case  $\partial S$  has two connected components denoted  $\partial_N S$  and  $\partial_S S$  (called "north and south boundaries") which are both invariant by  $\mathbb{R}_\theta$ .

We denote by  $P$  the set of poles, that is  $P = \{N, S\}g$  in Case 1,  $P = fNg$  in Case 2 and  $P = \emptyset$  in Case 3 and set

$$(4.1) \quad U = S \setminus P.$$



We now describe a nice parametrization of  $(S, g)$ , and, in particular, useful coordinates on the set  $U$ . We set  $L = \text{dist}_g(\partial_N S, \partial_S S)$  and denote by

$$(4.2) \quad I_L = (0, L) \text{ in Case 1, } I_L = (0, L] \text{ in Case 2, } I_L = [0, L] \text{ in Case 3.}$$

We let  $\gamma_0$  be a geodesic curve of  $S$  joining  $N$  (resp.  $\partial_N S$  in case 3) to  $S$  (resp.  $\partial_S S$  in Cases 2 and 3). Note in particular that  $\text{length}(\gamma_0) = L$ . For any  $\theta \in S^1$ , the isometry  $R_\theta$  transforms the geodesic  $\gamma_0$  into  $R_\theta(\gamma_0)$ , which is another geodesic joining  $N$  (resp.  $\partial_N S$ ) to  $S$  (resp.  $\partial_S S$ ). For every  $m \in U$  (defined in (4.1)), there exists a unique  $\theta \in S^1$  such that  $m$  belongs to  $R_\theta(\gamma_0)$ . The geodesic  $R_\theta(\gamma_0)$  can be parametrized by arclength

$$\begin{aligned} \rho : [0, L] &\rightarrow R_\theta(\gamma_0), \quad \rho(0) \in \partial_N S, \quad \rho(L) \in \partial_S S, \\ s = \text{dist}_g(\rho(s), \partial_N S) &= L - \text{dist}_g(\rho(s), \partial_S S), \end{aligned}$$

and there exists a unique  $s \in I_L$  such that  $\rho(s) = m$ . We use  $(s, \theta)$  as a parametrization of  $U \subset S$ :

$$\begin{aligned} \zeta : U = S \cap P &\rightarrow I_L \times S^1 \\ m &\mapsto \zeta(m) = (s, \theta). \end{aligned}$$

In Case 3,  $P = \emptyset$  and thus the whole surface  $S = U$  is diffeomorphic to the cylinder  $I_L \times S^1$  via  $\zeta$ . In Cases 2 and 1, we further need to describe coordinate charts around the poles. In cases 1 and 2, we define another exponential chart  $(U_N, \zeta_N)$  centered at the pole  $N$  by

$$\begin{aligned} U_N &= \text{fNg}[\zeta^{-1}((0, L/2) \times S^1)] = B_g(N, L/2) \subset S, \\ \zeta_N : U_N &\rightarrow B_{\mathbb{R}^2}(0, L/2), \quad \zeta_N(N) = 0. \end{aligned}$$

with the transition map

$$\begin{aligned} \zeta_N \circ \zeta^{-1} : \zeta(U \setminus U_N) = (0, L/2) \times S^1 &\rightarrow B_{\mathbb{R}^2}(0, L/2) \cap \text{fNg} \\ (s, \theta) &\mapsto (s \cos(\theta), s \sin(\theta)). \end{aligned}$$

In Case 1, we add similarly a last exponential chart  $(U_S, \zeta_S)$  centered at the pole  $S$  by

$$\begin{aligned} U_S &= \text{fSg}[\zeta^{-1}((L/2, L) \times S^1)] = B_g(S, L/2) \subset S, \\ \zeta_S : U_S &\rightarrow B_{\mathbb{R}^2}(0, L/2), \quad \zeta_S(S) = 0, \end{aligned}$$

with the transition map

$$\begin{aligned} \zeta_S \circ \zeta^{-1} : \zeta(U \setminus U_S) = (L/2, L) \times S^1 &\rightarrow B_{\mathbb{R}^2}(0, L/2) \cap \text{fSg} \\ (s, \theta) &\mapsto ((L - s) \cos(\theta), (L - s) \sin(\theta)). \end{aligned}$$

We shall need the following notation. For a subset  $J \subset I_L$ , we denote by

$$\mathcal{C}_J = \zeta^{-1}(J \times S^1) = \{m = \zeta^{-1}(s, \theta) \in U; s \in J\} \subset U \subset S$$

the  $R_\theta$  invariant set which projects down to  $J$ . We will also extend this definition to sets  $J \subset [0, L]$  by adding the point  $N$  if  $0 \in J$  (in Cases 1 and 2) and the point  $S$  if  $L \in J$  (in Case 1).

*Riemannian structure and operators involved.* — On the cylinder  $I_L \times S^1$ , the metric  $g$  is given by

$$(4.3) \quad (\zeta^{-1})^* g = ds^2 + R(s)^2 d\theta^2,$$

for some smooth function  $R : I_L \rightarrow \mathbb{R}^+$  (the function  $R$  can be interpreted as the Euclidean distance in  $\mathbb{R}^3$  of the point parametrized on  $S$  by  $s$  to the symmetry axis, see e.g. [LL21a, §3]). Since  $g$  is a smooth metric on  $S$ , [Bes78, Prop. 4.6] gives that  $R$  extends to a  $C^1$  function  $[0, L] \rightarrow \mathbb{R}^+$  satisfying moreover

$$(4.4) \quad R(0) = 0, \quad R'(0) = 1, \quad R^{(2p)}(0) = 0 \quad \text{for any } p \geq \mathbb{N}, \quad \text{in Cases 1 and 2,} \\ R(L) = 0, \quad R'(L) = -1, \quad R^{(2p)}(L) = 0 \quad \text{for any } p \geq \mathbb{N}, \quad \text{in addition in Case 1.}$$

For other parametrizations of surfaces of revolution, or direct computations on the sphere  $S^2$  and the disk  $D$ , we refer to [LL21a, §3].

**EXAMPLE 4.1.** — In particular, we consider here the following three examples:

- the unit sphere of  $\mathbb{R}^3$  is given by case 1  $L = \pi$ ,  $s \in (0, \pi)$ ,  $R(s) = \sin(s)$  (and the maximum of  $R$  is reached at  $s_0 = \pi/2$ )
- the unit disk of  $\mathbb{R}^2$  is given by case 2 with  $L = 1$  and  $R(s) = s$ ;
- flat cylinder of length  $L > 0$  and radius  $R_0 > 0$  is given by case 3 with  $R(s) = R_0$ .

In these coordinates, the Riemannian volume form is hence  $R(s) ds d\theta$ , the Riemannian gradient of a function is

$$(4.5) \quad \nabla_g u = \partial_s u \frac{\partial}{\partial s} + \frac{1}{R(s)^2} \partial_\theta u \frac{\partial}{\partial \theta}, \quad \text{with } g(\nabla_g u, \nabla_g u) = j \partial_s u j^2 + \frac{1}{R(s)^2} j \partial_\theta u j^2,$$

and the Laplace-Beltrami operator is given by

$$\Delta_{s,\theta} = \frac{1}{R(s)} \partial_s R(s) \partial_s + \frac{1}{R(s)^2} \partial_\theta^2.$$

We define by  $L^2(S) := L^2(S, d\text{Vol}_g)$  the space of square integrable functions, which is also invariant by the action of  $(R_\theta)_{\theta \in \mathbb{S}^1}$ . We will sometime also use the same definition for  $L^2(C_J) := L^2(C_J, d\text{Vol}_g)$  for  $J \in [0, L]$ .

Another important operator is the infinitesimal generator  $X_\theta$  of the group  $(R_\theta)_{\theta \in \mathbb{S}^1}$ , defined, for  $u \in C^1(S)$ , by

$$(4.6) \quad X_\theta u = \lim_{\vartheta \rightarrow 0} \vartheta^{-1} (u \circ R_\vartheta - u).$$

In the chart  $(U, \zeta)$ , the action of  $R_\theta$  is given by  $(\zeta^{-1})^* R_\theta(u, \theta^0) = (u, \theta^0 + \theta)$ , so that  $(\zeta^{-1})^* X_\theta = \partial_\theta$ . It is proved in [LL21a, §3.2] that  $X_\theta$  is a smooth vector field on  $S$ . Note also that  $X_\theta(N) = X_\theta(S) = 0$  and that its norm is given by  $\sqrt{g(X_\theta, X_\theta)}(s, \theta) = R(s)$  (in the coordinates of  $U$ ).

Now, remark that  $(R_\theta)_{\theta \in \mathbb{S}^1}$  acts as a (periodic) one-parameter unitary group on  $L^2(S)$  by  $f \mapsto f \circ R_\theta$ . The Stone Theorem (see e.g. [RS80, Th. VIII-8, p. 266]) hence implies that its infinitesimal generator is  $iA$ , where  $A$  is a selfadjoint operator on  $L^2(S)$  with domain  $D(A) \subset L^2(S)$ . Since  $iAf = X_\theta f$  for  $f \in C^1(S)$  (which is dense in  $D(A)$ ) according to (4.6), we have that  $A$  is the selfadjoint extension of  $X_\theta/i$ .

From now on, we slightly abuse the notation and still denote  $X_\theta/i$  for its selfadjoint extension  $A$ .

*Gradient vector field and conjugated operator.* — We finally introduce a function  $f : S \rightarrow \mathbb{R}$ , at least of class  $W^{2,1}$  to define the gradient flow. Throughout this section, we assume that  $X_\theta f = 0$ , i.e.,  $f$  is invariant by rotation and the same property holds for  $g$ . In the coordinates of  $U$ , we shall thus simply write  $f = f(s)$ . These regularity assumptions can be written in the coordinate of  $U$  by

$$(4.7) \quad \begin{aligned} & s \mapsto f(s) \in W^{2,1}(0, L), \text{ with} \\ & f'(0) = 0 \text{ (in Cases 1 and 2), and } f'(L) = 0 \text{ (in Case 1).} \end{aligned}$$

We may now define as in (2.4) the conjugated operator  $P_\varepsilon$  as

$$(4.8) \quad \begin{aligned} P_\varepsilon &= \varepsilon^2 \Delta_g + \frac{\int g f g^2}{4} + \varepsilon \frac{\Delta_g f}{2} \quad \varepsilon q \\ &= \varepsilon^2 \left( \frac{1}{R(s)} \partial_s R(s) \partial_s + \frac{1}{R(s)^2} \partial_\theta^2 \right) + \frac{\int f^\theta(s) f^2}{4} + \frac{\varepsilon}{2} \frac{1}{R(s)} \partial_s (R(s) f^\theta(s)) \quad \varepsilon q, \end{aligned}$$

where the second writing, in the coordinates of  $U$ , uses the invariance of  $f$ . Note that the last term in this expression acts as a multiplication operator by a function in  $L^1(S)$  with size  $\varepsilon$ . We shall often consider it as a lower order term, and keep the shorter notation  $\Delta_g f$  in place of  $(1/R(s)) \partial_s (R(s) f^\theta(s))$ .

Since both  $g$  and  $f$  are invariant by the action of  $R_\theta$ , we have

$$(4.9) \quad [X_\theta, P_\varepsilon] = 0.$$

Moreover,  $P_\varepsilon$  is selfadjoint in  $L^2(S, d\text{Vol}_g)$  with domain  $H^2(S) \setminus H_0^1(S)$  ( $= H^2(S)$  in Case 1), and has compact resolvent. Therefore, the operators  $P_\varepsilon$  and  $X_\theta$  share a common basis of eigenfunctions (see e.g. [LL21a, §3.2] for a proof). If  $\lambda \in \mathbb{R}$  is an eigenvalue of  $P_\varepsilon$ , then (in the coordinates of  $U$ ) the associated eigenfunction can be written as  $e^{ik\theta} v(s)$  with  $k \in \mathbb{Z}$ ,  $v \in H_{\text{loc}}^2(I_L) \setminus L^2((0, L), R(s) ds)$  satisfying

$$(4.10) \quad \frac{\varepsilon^2}{R(s)} \partial_s (R(s) \partial_s v) + \varepsilon^2 \frac{k^2}{R(s)^2} v + \left( \frac{\int f^\theta(s) f^2}{4} + \varepsilon q f \right) v = \lambda v,$$

together with  $v(L) = 0$  in Case 2 and  $v(0) = v(L) = 0$  in Case 3.

Restoring the dependence of the eigenvalues in the parameter  $\varepsilon$ , we call the normalized eigenfunctions of  $P_\varepsilon$ :  $\varphi_{k,n}^\varepsilon = e^{ik\theta} v_{k,n}^\varepsilon(s)$  with eigenvalues  $\lambda_{k,n}^\varepsilon$ , where  $n \in \mathbb{N}$ . In particular, for all  $\varepsilon > 0$ , we can write  $L^2(S) = \bigoplus_{(k,n) \in \mathbb{Z} \times \mathbb{N}} \text{span}(\varphi_{k,n}^\varepsilon)$ .

We further denote

$$L_k^2 = \ker(X_\theta - ik) = \{ \varphi \in L^2(S) ; \varphi|_U = e^{ik\theta} f(s), f \in L^2((0, L), R(s) ds) \},$$

and  $H_k^2 = D(P_\varepsilon) \setminus L_k^2 = H^2(S) \setminus H_0^1(S) \setminus L_k^2$ . The commutation property (4.9) implies that for all  $\varepsilon > 0$ ,  $P_\varepsilon H_k^2 \subset L_k^2$ , so we can define the operator

$$(4.11) \quad P_\varepsilon^{(k)} = P_\varepsilon|_{L_k^2}, \quad \text{with domain } H_k^2,$$

which is selfadjoint. This can be seen for instance directly on the simultaneous diagonalization which implies an isometry  $L^2(S) \rightarrow \ell^2(\mathbb{Z} \times \mathbb{N})$  where  $L_k^2 \rightarrow \ell^2(\mathbb{N})$ ;  $n \in \mathbb{N}$ .

as a closed subspace of  $\ell^2(Z \setminus N)$ . The fact that  $P_\varepsilon$  has compact resolvent implies that this is also the case for  $P_\varepsilon^{(k)}$ . With a slight abuse of notation, we shall still denote by  $P_\varepsilon^{(k)}$  the one-dimensional operator  $(\zeta^{-1}) P_\varepsilon^{(k)} \zeta$  defined on  $I_L$ , namely

$$(4.12) \quad P_\varepsilon^{(k)} w = \frac{\varepsilon^2}{R(s)} \partial_s(R(s) \partial_s w) + \left( \frac{\varepsilon^2 k^2}{R(s)^2} + \frac{f^\theta(s) j^2}{4} + \varepsilon q_f \right) w.$$

**4.2. THE CONDITIONS (GCC) AND (FC) ON SURFACES OF REVOLUTION.** — In this section, we characterize the conditions (GCC) (see Definition 2.10 if  $\partial M = ?$ ) and (FC) (see Definition 1.4 if  $\partial M \neq ?$ ) in the above very particular geometry, and further assuming that the observation region  $\omega$  is rotationally invariant as well. In case  $\partial M \neq ?$ , we consider the two cases of internal and boundary observation, and describe the associated minimal times  $T_{FC}$ .

**PROPOSITION 4.2.** — *Let  $\delta > 0$  and recall that  $f$  is assumed to be  $\theta$ -invariant.*

(1) *In Case 1, we consider the set  $\omega = B_g(N, \delta) \cup B_g(S, \delta)$ ; then  $(S, r_g f, \omega)$  satisfies (GCC) if and only if*

$$f^\theta(s) \neq 0 \text{ for all } s \in [\delta, L - \delta] \text{ and } T_{GCC}(S, r_g f, \omega) = \left| \int_\delta^{L-\delta} \frac{ds}{f^\theta(s)} \right|.$$

(2) *In Case 2, we consider the set  $\omega = B_g(N, \delta)$ ; then  $(S, r_g f, \omega)$  satisfies (FC) if and only if*

$$f^\theta(s) \neq 0 \text{ for all } s \in [\delta, L] \text{ and } T_{FC}(S, r_g f, \omega) = \left| \int_\delta^L \frac{ds}{f^\theta(s)} \right|.$$

(3) *In Case 3,  $(S, r_g f)$  satisfies (FC) if and only if*

$$f^\theta(s) \neq 0 \text{ for all } s \in [0, L] \text{ and } T_{FC}(S, r_g f) = \left| \int_0^L \frac{ds}{f^\theta(s)} \right|.$$

Note that in Case 2, the situations  $f^\theta > 0$  and  $f^\theta < 0$  play two different roles (see the proof below). Indeed, in case  $f^\theta > 0$ , all trajectories of  $\dot{s} = -f^\theta$  enter the controlled region  $\omega$ , whereas, in case  $f^\theta < 0$ , all trajectories of  $\dot{s} = -f^\theta$  flow out of the domain  $S$  through  $\partial S$  (without passing into  $\omega$ ). However, the definition of (FC) in Definition 1.4 does not make a distinction between these two situations.

*Proof.* — We only prove the second item; the other two items are proved similarly. According to Definition 1.4, (4.5) and the  $\theta$ -invariance of  $f$ , it suffices to check under which conditions the solutions to  $\dot{s}(t) = -f^\theta(s(t))$  all enter  $\omega = B_g(N, \delta)$  (resp. all exit  $S$ , that is satisfy  $s(T) > L$ ). If there is  $s_0 \in [\delta, L]$  such that  $f^\theta(s_0) = 0$ , then the associated solution satisfies  $s(t) = s_0 \in [\delta, L]$  for all  $t \geq 0$ , and  $(S, r_g f, \omega)$  does not satisfy (FC).

If  $f^\theta > 0$  on  $[\delta, L]$ , then  $s(t)$  is decreasing, and for any  $\sigma_0, \sigma_1 \geq 0$ , one has

$$\int_{s(\sigma_0)}^{s(\sigma_1)} \frac{ds}{f^\theta(s)} = \sigma_0 - \sigma_1.$$

The longest trajectory that does not enter  $\omega$  is such that  $s(0) = L$  and  $s(T) = \delta$  so that  $T = \int_{\delta}^L ds/f^{\theta}(s)$ . This proves

$$T_{\text{FC}}(S, r_g f, \omega) = \int_{\delta}^L \frac{ds}{f^{\theta}(s)}$$

in this case.

Finally, if  $f^{\theta} < 0$  on  $[\delta, L]$ , then  $s(t)$  is increasing, and for any  $\sigma_0, \sigma_1 \geq \mathbb{R}$ , one has

$$\int_{s(\sigma_0)}^{s(\sigma_1)} \frac{ds}{f^{\theta}(s)} = \sigma_0 - \sigma_1.$$

The longest trajectory that does not enter  $\omega$  is such that  $s(0) = \delta$  and  $s(T) = L$  so that  $T = \int_{\delta}^L ds/f^{\theta}(s)$ . This proves

$$T_{\text{FC}}(S, r_g f, \omega) = \int_{\delta}^L \frac{ds}{f^{\theta}(s)}$$

in this case, and hence the proposition.

**4.3. EXISTENCE OF EIGENFUNCTIONS.** — One may consider different asymptotic regimes in the parameters  $\varepsilon \searrow 0^+$  and  $k \searrow +1$ . Note that the case  $k$  bounded would correspond to the one-dimensional situation treated in the companion paper [LL21b]. Here, we shall consider the limit  $k \searrow +1$  and make the following choice of the parameter  $\varepsilon$ :

$$(4.13) \quad \varepsilon = \varepsilon_k = ck^{-1}$$

considered as a semiclassical parameter, where  $c > 0$  is a fixed parameter (i.e., which does not depend on  $k$ ) that will be chosen but fixed. All constants that appear below might depend on  $c$ . The analysis of the asymptotic of the constant involved as  $c \searrow 0$  (low level of rotation) or  $c \searrow 1$  (high level of rotation) would be interesting but would require much more work.

In view of (4.12), the choice (4.13) naturally leads to consider

$$(4.14) \quad s \nearrow L \quad V_c(s) := \frac{c^2}{R(s)^2} + \frac{f^{\theta}(s)^2}{4},$$

as the effective potential of the operator  $P_{\varepsilon_k}^{(k)}$  in the semiclassical limit  $\varepsilon = \varepsilon_k = ck^{-1} \searrow 0^+$ . In particular, the operator  $P_{\varepsilon}^{(k)}$  is now a semiclassical operator with small parameter  $\varepsilon_k$  and (4.12) can be rewritten

$$(4.15) \quad P_{\varepsilon_k}^{(k)} w = \frac{\varepsilon_k^2}{R(s)} \partial_s(R(s)\partial_s w) + (V_c(s) + \varepsilon_k q_f) w.$$

In the present section, we recall the existence of eigenfunctions (Analogue of Lemma 3.2 above) associated to any value of the effective potential  $V_c$ . More precisely, in the chosen regime (4.13), for any  $s_0 \geq I_L$  (recall that  $I_L$  is defined in (4.2)), we construct a sequence  $\psi_k$  such that  $P_{\varepsilon_k} \psi_k = (V_c(s_0) + r(k))\psi_k$ , with  $r(k) \searrow 0$  as  $k \searrow +1$ . As in Section 3.1, the precision  $r(k)$  might depend on whether  $V_c^{\theta}(s_0) \neq 0$  or  $V_c^{\theta}(s_0) = 0$  but we will only state the worst estimate, which is sufficient for our

needs. We shall later on prove localization properties of the  $\psi_k$ 's assuming further that  $V_c(s_0) = \min_{I_L} V_c$  (which is a global assumption). We recall the choice (4.13) and the definition (4.14).

**LEMMA 4.3** (Existence of eigenfunctions). — *For all  $s_0 \in I_L$ , there is  $k_0 > 0$  such that for all  $k \in \mathbb{N}, k > k_0$ , there exists  $\psi_k \in H^2(S) \setminus L_k^2$  in case 1 (resp.  $\psi_k \in H^2(S) \setminus H_0^1(S) \setminus L_k^2$  in Cases 2–3), and  $\mu_k \in \mathbb{R}$  such that*

$$\mu_k = V_c(s_0) + O(1/k^{2/3}) = \frac{c^2}{R(s_0)^2} + \frac{f^\theta(s_0)^2}{4} + O(1/k^{2/3}),$$

$$P_{\varepsilon_k} \psi_k = \mu_k \psi_k, \quad k \psi_k k_{L^2(S)} = 1, \quad \psi_k(s, \theta) = e^{ik\theta} \varphi_k(s),$$

with  $P_\varepsilon$  defined in (4.8) and  $\varepsilon_k$  in (4.13).

The proof is very similar to Lemma 3.2. Indeed, the study of the operator  $P_{\varepsilon_k}$  in  $L_k^2$  reduces to the 1D problem for the operator  $P_{\varepsilon_k}^{(k)}$  defined in (4.11) and (4.15). The proof consists first in constructing quasimodes exactly as in the proof of Lemma 3.2. Deducing existence of an exact eigenfunction from a quasimode requires the use of the right selfadjoint extension  $P_{\varepsilon_k}^{(k)}$ . This issue is however treated in detail in [LL21a, Lem. 3.6]. Note that  $s_0 \in I_L$  implies that it cannot be a pole so that  $R(s_0)^{-1}$  is finite.

**4.4. GEOMETRIC ASSUMPTIONS AND THE AGMON DISTANCE.** — The next step is to study the behavior of the eigenfunction  $\psi_k$  constructed in Lemma 4.3. This will require some global assumptions on the effective potential  $V_c$ . Recall that  $V_c$  is defined in (4.14), is continuous on  $I_L$ , and tends to  $+1$  near to the poles. Indeed, in Cases 1 and 2 we have for instance

$$V_c(s) \underset{s \rightarrow 0^+}{\sim} \frac{c^2}{R(s)^2} \underset{s \rightarrow 0^+}{\sim} \frac{c^2}{s^2} \underset{s \rightarrow 0^+}{\sim} +1,$$

as a consequence of (4.4) and (4.7) (and similarly when  $s \rightarrow L$  in Case 1). As a consequence,  $V_c$  admits a minimum on the interval  $I_L$ , which we denote by

$$V_{\min} = \min_{s \in I_L} V_c(s) \in \mathbb{R}_+.$$

In the following, we make Assumption (1.17) on the set where  $V_c$  reaches its global minimum.

**ASSUMPTION 4.4.** — *The set  $V_c^{-1}(V_{\min}) = f s_{\min} g \cap I_L$  consists in a single point.*

Note that this assumption is generic. Here it is not strictly needed to prove the main results, but simplifies the presentation and statements slightly. We again introduce the relevant Agmon distance at the minimal energy level  $V_{\min}$ , defined in the coordinates of  $U$  by the eikonal equation

$((d_A^c)^\theta(s))^2 - (V_c(s) - V_{\min}) = 0$ ,  $d_A^c(s_{\min}) = 0$ ,  $\text{sgn}((d_A^c)^\theta(s)) = \text{sgn}(s - s_{\min})$ , or, more explicitly, for  $s \in I_L$ , by (1.18). In view of the  $W^{2,1}$  regularity of  $f$  on  $S$  and the definition of  $V_c$  in (4.14), the function  $d_A^c$  is of class  $C^2$  away from  $s_{\min}$ ,  $0$  and  $L$ , and is locally Lipschitz on  $I_L$ . Note that this includes Lipschitz regularity up to the

boundary  $s = L$  in Case 2 and to both boundaries  $s = 0, L$  in Case 3. We also consider  $d_A^c$  as a  $\theta$ -invariant function on the surface  $S$ .

LEMMA 4.5 (Properties of  $d_A^c$ ). — *Under Assumption 4.4, we have  $d_A^c \in C^2(I_L \setminus \{s_{\min}\})$  together with*

$$(4.16) \quad d_A^c(s) = c \log(s) + O(1), \quad \text{as } s \rightarrow 0^+, \quad \text{in Cases 1 and 2,}$$

$$(4.17) \quad d_A^c(s) = c \log(L - s) + O(1), \quad \text{as } s \rightarrow L^-, \quad \text{in Case 1.}$$

*Proof.* — We only consider the asymptotics as  $s \rightarrow 0^+$ , that is, prove (4.16). The proof of (4.17) follows the same. Remark that according to (4.4), we have  $1/R(y) \sim 1$  as  $y \rightarrow 0^+$  with

$$R(s) = s + O(s^3), \quad \text{when } s \rightarrow 0^+.$$

As a consequence, since  $f \in W^{2,1}(I_L)$ , we have

$$V_c(s) = \frac{c^2}{s^2} + O(1), \quad \text{when } s \rightarrow 0^+.$$

With (1.18), we obtain

$$d_A^c(s) = \left| \int_{s_{\min}}^s \frac{c}{y} (1 + O(y)) dy \right| = c \log(s) + O(1) \quad \text{as } s \rightarrow 0^+.$$

4.5. UPPER BOUNDS FOR EIGENFUNCTIONS: AGMON ESTIMATES. — As far as upper bounds on  $\psi_k$  are concerned, we have the following Agmon type estimate.

PROPOSITION 4.6. — *Under Assumption 4.4, assume that  $\mu_k = V_c(s_{\min}) + r(k)$  with  $r(k) \rightarrow 0$  and  $\psi_k \in H^2(S) \setminus L_k^2(S)$  solves*

$$P_{\varepsilon_k} \psi_k = \mu_k \psi_k, \quad \text{on } S, \quad \psi_k|_{\partial S} = 0, \quad \|\psi_k\|_{L^2(S)} = 1,$$

with  $P_\varepsilon$  defined in (4.8) and  $\varepsilon_k$  in (4.13). Then for all  $0 < \delta < 1$ , there exist  $C = C(\delta)$ ,  $k_0 = k_0(\delta) > 0$  such that, for all  $k \in \mathbb{N}$ ,  $k > k_0$ , the following integral is well defined with the estimate

$$\int_S e^{2(1-\delta)d_A^c(m)/\varepsilon_k} |\psi_k|^2(m) d\text{Vol}_g(m) \leq C e^{2\delta/\varepsilon_k}.$$

Also, if  $\partial S \neq \emptyset$  (that is, in Cases 2 and 3), for all  $0 < \delta < 1$ , there exist  $C = C(\delta)$ ,  $k_0 = k_0(\delta) > 0$  such that, for all  $k \in \mathbb{N}$ ,  $k > k_0$ ,

$$(4.18) \quad \begin{aligned} k \partial_s \psi_k(L, \cdot) & \in C^0_{H^1(S^1)} \leq C e^{-2(1-\delta)d_A^c(L)/\varepsilon_k}, \\ k \partial_s \psi_k(0, \cdot) & \in C^0_{H^1(S^1)} \leq C e^{-2(1-\delta)d_A^c(0)/\varepsilon_k}, \end{aligned}$$

where the last estimate (at  $s = 0$ ) holds true in Case 3 only.

Note that given the asymptotic expansion of  $d_A^c$  in Lemma 4.5, this estimate implies that  $\psi_k$  vanishes strongly near the poles of  $S$ . The proof is made with classical Agmon type identity with some care with respect to the degeneracy at the poles. It is very similar to the one performed in [LL21a, Th. 3.9] and we omit it. Note that, as opposed to [LL21a, Th. 3.9], we do not assume here that the minimum be non-degenerate, and

only deduce an estimate with loss ( $\delta > 0$ ), which is sufficient for our needs. In the non-degenerate case, one can take  $\delta = 0$  in this estimate and replace the right hand-side by a polynomial bound of the type  $\varepsilon_k^M$ , see [LL21a, Th. 3.9].

The proof of the boundary estimate also requires a bootstrap argument to estimate higher  $H^s$  norms and the use of a trace estimate, see [Hel88, Prop. 3.3.1 & 3.3.4].

We obtain the following two direct Corollaries.

**COROLLARY 4.7.** — *Under the assumptions of Proposition 4.6, for all  $0 < \delta < 1$ , there exist  $C = C(\delta), k_0 = k_0(\delta) > 0$  such that, for all  $k \geq 2N, k > k_0$ , and for all rotationally invariant set  $\omega$ , we have*

$$\int_{\omega} j\psi_k^2(m) d\text{Vol}_g(m) \leq C e^{-(2/\varepsilon_k)((1-\delta)d_A^c(\omega)-\delta)}, \quad \text{with } d_A^c(\omega) = \inf_{s \in \omega} d_A^c(s).$$

*Proof.* — This is a direct consequence of the following estimate:

$$\begin{aligned} e^{2(1-\delta)d_A^c(\omega)/\varepsilon_k} \int_{\omega} j\psi_k^2(m) d\text{Vol}_g(m) &\leq \int_{\omega} e^{2(1-\delta)d_A^c(m)/\varepsilon_k} j\psi_k^2(m) d\text{Vol}_g(m) \\ &\leq \int_S e^{2(1-\delta)d_A^c(m)/\varepsilon_k} j\psi_k^2(m) d\text{Vol}_g(m) \\ &\leq C e^{2\delta/\varepsilon_k}, \end{aligned}$$

where we have used Proposition 4.6 in the last inequality.

**COROLLARY 4.8** (most of the norm is close to the minimum). — *Under the assumptions of Proposition 4.6, for any  $\rho > 0$ , there exists  $k_0 \geq 2N$  so that*

$$k\psi_k k_{L^2(\mathcal{C}_{(s_{\min}-\rho, s_{\min}+\rho)})}^2 > 1/2, \quad \text{for all } k \geq 2N, k > k_0.$$

*Proof.* — Applying Corollary 4.7 with  $\omega = S \cap \mathcal{C}_{(s_{\min}-\rho, s_{\min}+\rho)}$  ensures that for any  $\delta > 0$ , there is  $k_0 \geq 2N$  such that for all  $k > k_0$ , we have

$$\int_{\omega} j\psi_k^2(m) d\text{Vol}_g(m) \leq C(\delta) e^{-(2/\varepsilon_k)((1-\delta)d_A^c(\omega)-\delta)}, \quad \text{with } d_A^c(\omega) = \inf_{s \in \omega} d_A^c(s).$$

From Assumption 4.4, we have  $d_A^c(s) > 0$  for all  $s \in I_L \cap \mathcal{C}_{(s_{\min}-\rho, s_{\min}+\rho)}$ . That  $\omega$  is closed and does not contain  $s_{\min}$  implies that  $d_A^c(\omega) > 0$ . Then we fix  $\delta > 0$  small enough so that  $(1-\delta)d_A^c(\omega) - \delta > 0$ . There is  $k_0 \geq 2N$  such that we have  $k\psi_k k_{L^2(\omega)}^2 \rightarrow \infty$  for  $k > k_0$ , which implies the result.

**4.6. LOWER BOUNDS FOR EIGENFUNCTIONS: ALLIBERT ESTIMATES.** — In Corollary 4.7, we proved that the family of eigenfunctions  $\psi_k$  decays on  $\omega$  at least like  $e^{-d_A^c(\omega)/\varepsilon_k}$ . The purpose of this section is to prove the converse, i.e., that the  $\psi_k$ 's decay at most like  $e^{-d_A^c(\omega)/\varepsilon_k}$  up to  $\delta$  loss. This comes from the particular one-dimensional underlying context. We follow in this section the method of Allibert [All98]. More precisely, we prove the following estimates.

**PROPOSITION 4.9.** — *Under Assumption 4.4, assume further that*

$$(4.19) \quad \mu_k = V_c(s_{\min}) + r(k) \text{ with } r(k) \rightarrow 0,$$



and  $\psi_k \in H^2(S) \setminus L^2_k(S)$  solves

$$P_{\varepsilon_k} \psi_k = \mu_k \psi_k, \text{ on } S, \quad \psi_k j_{\partial S} = 0, \quad k \psi_k k_{L^2(S)} = 1, \quad \psi_k(s, \theta) = e^{ik\theta} \varphi_k(s),$$

with  $P_\varepsilon$  defined in (4.8) and  $\varepsilon_k$  in (4.13).

Then, for any  $\eta, \delta > 0$ , there exist  $k_0, C > 0$  so that

$$k \psi_k k_{L^2(\mathcal{C}_{B(s, \eta)})} > C e^{-(1/\varepsilon_k)(d_A^c(s) + \delta)}, \quad \text{for all } k > k_0 \text{ and } s \in I_L \text{ s.t. } \text{dist}(s, P) > \eta.$$

Note that  $\eta$  is a safety distance to the set of poles  $P$  defined in Section 4.1. The proof of Proposition 4.9 relies on two lemmas, which we give in the next section.

**4.6.1. Two preliminary lemmas.** — In this section, we assume that the assumptions of Proposition 4.9 are satisfied. In particular, the eigenfunctions under consideration are of the form  $\psi_k(s, \theta) = e^{ik\theta} \varphi_k(s)$ . We define the following “semiclassical energy densities” of the eigenfunctions  $\psi_k$ , for  $s \in I_L$  by

$$(4.20) \quad \begin{aligned} E_k(s) &:= \varepsilon_k^2 j \partial_s \psi_k j^2(s) + (V_c(s) - \mu_k + 1) j \psi_k j^2(s) \\ &= \varepsilon_k^2 j \varphi_k^0 j^2(s) + (V_c(s) - \mu_k + 1) j \varphi_k j^2(s), \end{aligned}$$

$$(4.21) \quad \begin{aligned} E_k^+(s) &:= \varepsilon_k^2 j \partial_s \psi_k j^2(s) + (V_c(s) - \mu_k) j \psi_k j^2(s) \\ &= \varepsilon_k^2 j \varphi_k^0 j^2(s) + (V_c(s) - \mu_k) j \varphi_k j^2(s). \end{aligned}$$

Note that according to elliptic regularity, we have  $\varphi_k \in H^2_{\text{loc}}(I_L)$  and, due to Sobolev embeddings,  $\varphi_k^0, E_k, E_k^+ \in C^0(I_L)$  and in particular  $E_k, E_k^+$  are defined everywhere on  $I_L$ . For  $s, t \in I_L$ , we define  $I_{s,t}$  to be the interval between the real numbers  $s$  and  $t$ , that is, either  $[s, t]$  or  $[t, s]$ . We also set

$$(4.22) \quad E_\alpha = f s \in I_L; j s - p j > \alpha, \text{ for all } p \in P, \text{ and } j s - s_{\min} j > \alpha g.$$

Recall that  $P$  is the set of poles defined in Section 4.1 (and is aimed at covering all Cases 1–3 at the same time) and  $s_{\min}$  is the point at which  $V_c$  reaches its minimum.

**LEMMA 4.10.** — Assume the assumptions of Proposition 4.9 and recall that  $E_k^+$  is defined in (4.21). Then, for any  $\alpha, \delta > 0$ , there exists  $k_0 > 0$  so that for all  $s, t \in I_L$  such that  $I_{s,t} \subset E_\alpha$ , we have

$$E_k^+(t) \leq e^{(2/\varepsilon_k)(j d_A^c(s) - d_A^c(t) j + \delta)} E_k^+(s), \quad \text{for all } k > k_0.$$

Lemma 4.10 provides with a Grönwall type estimate on the energy  $E_k^+$ , with a precise description of the constant, under the condition that we remain finitely away from  $s_{\min}$ . It is an analogue of [All98, Lem. 12] in our setting (with an additional uniform dependence).

Note that  $j d_A^c(s) - d_A^c(t) j = d_A^c(s, t)$  is the Agmon distance between  $s$  and  $t$  at the lowest energy level.

*Proof of Lemma 4.10.* — On the interval  $z \in I_{s,t} \subset E_\alpha$ , we differentiate  $E_k^+$ , yielding

$$(E_k^+)^0(z) = 2\varepsilon_k^2 \text{Re}(\overline{\varphi_k^0} \varphi_k^{00}) + V_c^0(z) \varphi_k^2 + 2(V_c(z) - \mu_k) \text{Re}(\varphi_k \overline{\varphi_k^0}).$$

We recall the choice of  $\varepsilon_k$  in (4.13) from the definition of  $P_{\varepsilon_k}$  in (4.8), and the definition of  $P_{\varepsilon_k}^{(k)}$  in (4.10)-(4.15) that we have

$$\mu_k \varphi_k = P_{\varepsilon_k} \varphi_k = \varepsilon_k^2 \varphi_k^{00} - \varepsilon_k^2 \frac{R^0}{R} \varphi_k^0 + V_c \varphi_k + \varepsilon_k q_f \varphi_k.$$

Replacing  $\varepsilon_k^2 \varphi_k^{00}$  in the above identity yields

$$\begin{aligned} (E_k^+)^0(z) &= 2(V_c(z) - \mu_k + \varepsilon_k q_f) \operatorname{Re}(\varphi_k \overline{\varphi_k^0}) - 2\varepsilon_k^2 \frac{R^0}{R} j\varphi_k^0 j^2 + V_c^0(z) j\varphi_k j^2 \\ &\quad + 2(V_c(z) - \mu_k) \operatorname{Re}(\varphi_k \overline{\varphi_k^0}) \\ (4.23) \quad &= (4(V_c(z) - \mu_k) + 2\varepsilon_k q_f) \operatorname{Re}(\varphi_k \overline{\varphi_k^0}) - 2\varepsilon_k^2 \frac{R^0}{R} j\varphi_k^0 j^2 + V_c^0(z) j\varphi_k j^2. \end{aligned}$$

First, using the continuity of  $V_c$  on  $I_L$  and the compactness of  $E_\alpha$  in  $I_L$ , we see that  $V_c$  reaches its minimum on  $E_\alpha$ . This, together with the fact that  $s_{\min} \notin E_\alpha$ , implies that  $C_\alpha^{-2} \subset V_c(s) - V_c(s_{\min}) \subset C_\alpha^2$  uniformly for  $s \in E_\alpha$ . Recalling (4.19), this yields the existence of  $k_0(\alpha)$  such that for  $k > k_0(\alpha)$ , we have  $V_c - \mu_k > 1/2C_\alpha > 0$  on  $E_\alpha$ .

We now estimate each of the terms in (4.23). We first have

$$\begin{aligned} j4(V_c - \mu_k) \operatorname{Re}(\varphi_k \overline{\varphi_k^0}) j &\subset 4\varepsilon_k^{-1} \sqrt{V_c - \mu_k} (\varepsilon_k j\varphi_k^0 j) \left( \sqrt{V_c - \mu_k} j\varphi_k j \right) \\ &\subset 2\varepsilon_k^{-1} \sqrt{V_c - \mu_k} [\varepsilon_k^2 j\varphi_k^0 j^2 + (V_c - \mu_k) j\varphi_k j^2] \\ &= 2\varepsilon_k^{-1} \sqrt{V_c - \mu_k} E_k^+. \end{aligned}$$

Moreover, according to (4.19), there exists a constant  $C_\alpha > 0$  such that we have  $\frac{1}{\sqrt{V_c - \mu_k}} \subset \sqrt{V_c - V_c(s_{\min})} + C_\alpha j r(k) j$  uniformly for  $s \in E_\alpha$ . Together with the above inequality, this implies

$$j4(V_c - \mu_k) \operatorname{Re}(\varphi_k \overline{\varphi_k^0}) j \subset \varepsilon_k^{-1} (2\sqrt{V_c - V_c(s_{\min})} + C_\alpha j r(k) j) E_k^+, \quad \text{on } E_\alpha.$$

Second, we have

$$|V_c^0 j\varphi_k j^2| = \frac{jV_c^0 j}{V_c - \mu_k} (V_c - \mu_k) j\varphi_k j^2 \subset 2C_\alpha k V_c^0 k_{L^1(E_\alpha)} E_k^+, \quad \text{on } E_\alpha.$$

Third, we have

$$j\varepsilon_k q_f \operatorname{Re}(\varphi_k \overline{\varphi_k^0}) j \subset C_\alpha E_k^+, \quad \text{on } E_\alpha.$$

Finally, since  $R^0/R$  is bounded on  $E_\alpha$ , we have

$$\left| \varepsilon_k^2 \frac{R^0}{R} j\varphi_k^0 j^2 \right| \subset C_\alpha \varepsilon_k^2 j\varphi_k^0 j^2 \subset C_\alpha E_k^+ \quad \text{on } E_\alpha.$$

Combining the last four estimates in (4.23) yields for another constant  $C_\alpha > 0$  and for all  $k > k_0(\alpha)$

$$|(E_k^+)^0(z)| \subset 2\varepsilon_k^{-1} \left[ \sqrt{V_c(z) - V_c(s_{\min})} + C_\alpha j r(k) j + C_\alpha \varepsilon_k \right] E_k^+(z), \quad \text{for all } z \in E_\alpha.$$

Applying the Grönwall Lemma on the interval  $I_{s,t}$  contained in  $E_\alpha$  yields

$$E_k^+(s) \subset e^{(2/\varepsilon_k) \left( \int_t^s \sqrt{V_c(z) - V_c(s_{\min})} dz + C_\alpha j r(k) j + C_\alpha \varepsilon_k \right)} E_k^+(t),$$

which is the sought result.

The next lemma is aimed at giving a rough Grönwall type estimate for the energy  $E_k$ , without precise constants. The interest of this less precise result is that it remains true close to the minimum  $s_{\min}$ . This allows to compensate the fact that Lemma 4.10 is not uniform when  $s$  is close to  $s_{\min}$ . Similarly as  $E_\alpha$  in (4.22), we define

$$F_\alpha = \{s \geq I_L; |j_s - p_j| > \alpha, \text{ for all } p \geq P_g\}.$$

LEMMA 4.11. — Assume the assumptions of Proposition 4.9 and recall that  $E_k$  is defined in (4.20). For any  $\alpha > 0$ , there exist  $C_\alpha, D_\alpha > 0$  so that for all  $s, t \geq F_\alpha$ , we have

$$E_k(s) \leq C_\alpha e^{(2/\varepsilon_k)D_\alpha s} E_k(t), \quad \text{for all } k > k_0.$$

Lemma 4.11 is an analogue of [All98, Lem. 11] in our setting. Recall that  $P$  is the set of poles defined in Section 4.1.

*Proof.* — The proof is quite close to that of Lemma 4.10. We only use the fact that there exists  $C_\alpha$  so that  $C_\alpha^{-1} \leq V_c - \mu_k + 1 \leq C_\alpha$  on  $F_\alpha$  if  $k > k_0(\alpha)$ . This gives a constant  $D_\alpha > 0$  such that

$$D_\alpha^{-1} (\varepsilon_k^2 j \varphi_k^0 j^2 + j \varphi_k j^2) \leq E_k \leq D_\alpha (\varepsilon_k^2 j \varphi_k^0 j^2 + j \varphi_k j^2) \quad \text{on } F_\alpha.$$

The same computation as in (4.23) gives

$$\begin{aligned} E_k^0 &= (E_k^+)^0 + 2 \operatorname{Re}(\varphi_k \overline{\varphi_k^0}) \\ &= (4(V_c - \mu_k) + r_k + 1) \operatorname{Re}(\varphi_k \overline{\varphi_k^0}) - 2\varepsilon_k^2 \frac{R^0}{R} j \varphi_k^0 j^2 + V_c^0(z) j \varphi_k j^2, \end{aligned}$$

with  $\sup_{F_\alpha} r_k \rightarrow 0$  as  $k \rightarrow +\infty$ . As a consequence, for  $k > k_0(\alpha)$ , we have constants  $C_\alpha^0, C_\alpha^{00}$  such that for all  $z \geq F_\alpha$

$$j(E_k^0)^0(z) \leq \varepsilon_k^{-1} C_\alpha^0 (\varepsilon_k^2 j \varphi_k^0 j^2 + j \varphi_k j^2) \leq \varepsilon_k^{-1} C_\alpha^{00} E_k(z),$$

which allows to conclude as in the proof of Lemma 4.10 above by Grönwall estimates.

4.6.2. *Proof of Proposition 4.9 from Lemmas 4.10 and 4.11.* — The sketch of the proof of Proposition 4.9 is as follows:

- (1) the total mass of  $\varphi_k$  is dominated by its mass near the minimum  $s_{\min}$  via Corollary 4.8;
- (2) the mass near  $s_{\min}$  is dominated by the energy at  $s_{\min} - \rho$  via Lemma 4.11 (with a small loss if  $\rho$  is small);
- (3) the energy at  $s_{\min} - \rho$  is dominated by the energy near  $s$  via Lemma 4.10 (with a geometric constant  $e^{(2/\varepsilon_k)d_A^c(s)}$ );
- (4) the energy near  $s$  is dominated by the  $L^2$  norm of  $\varphi_k$  (or  $\psi_k$ ) near  $s$  via elliptic regularity.

*Proof of Proposition 4.9.* — Without loss of generality, we can assume  $s < s_{\min} - \eta$ . Indeed, the case  $s > s_{\min} + \eta$  is treated similarly, and the case  $s \in [s_{\min} - \eta, s_{\min} + \eta]$  is a direct consequence of Corollary 4.8 applied for  $\rho = \eta/2$ .

Now, since  $s_{\min} \geq I_L$ , notice that we have necessarily  $d := \text{dist}(s_{\min}, P) > 0$ , and we may also assume that  $\delta < d/4$ . Lemma 4.11 can be applied with  $\alpha = d/4$  and produces some constants  $C = C_\alpha$ ,  $D = D_\alpha$ . Let us now choose

$$\rho = \rho_\delta = \min(\delta/(4D), d/4, \eta/2).$$

Application of Lemma 4.11 gives for any  $u \in (s_{\min} - \rho, s_{\min} + \rho)$  (so that  $\text{dist}(u, P) > \eta$ )

$$E_k(s_{\min} - \rho) > C^{-1} e^{-(4/\varepsilon_k)D\rho} E_k(u) > C^{-1} e^{-\delta/\varepsilon_k} E_k(u).$$

Integrating in  $u \in (s_{\min} - \rho, s_{\min} + \rho)$  gives

$$(4.24) \quad E_k(s_{\min} - \rho) > \frac{C}{2\rho} e^{-\delta/\varepsilon_k} \int_{s_{\min} - \rho}^{s_{\min} + \rho} E_k(u) \, du > C_1 e^{-\delta/\varepsilon_k},$$

where we have used  $E_k(u) > \frac{1}{2} j\varphi_k f^2(u)$  (for  $k$  large enough) and Corollary 4.8.

Taking  $y \in [s, s + \eta/2]$ , we still have  $y \in (s_{\min} - \rho, s_{\min} + \rho)$  from the definition of  $\rho$ . Choosing  $\alpha = \rho/2 \in \eta$ , we can check that  $[y, s_{\min} - \rho] \in E_\alpha$ , so that Lemma 4.10 applies on  $E_\alpha$  and gives

$$(4.25) \quad \begin{aligned} E_k^+(y) &> e^{-(2/\varepsilon_k)(jd_A^c(y) - d_A^c(s_{\min} - \rho)j + \delta)} E_k^+(s_{\min} - \rho) \\ &> e^{-(2/\varepsilon_k)(d_A^c(y) + \delta)} E_k^+(s_{\min} - \rho), \end{aligned}$$

where we have noticed

$$jd_A^c(y) - d_A^c(s_{\min} - \rho)j = d_A^c(y) - d_A^c(s_{\min} - \rho) \in d_A^c(y) - d_A^c(s_{\min}) = d_A^c(y).$$

Since  $s_{\min}$  is a strict minimum, there are constants  $k_0(\rho, \delta) = k_0(\delta, \eta) > 0$  and  $C(\rho, \eta) = C(\delta, \eta) > 0$  such that for  $k > k_0(\delta)$ ,

$$V_c(s_{\min} - \rho) - \mu_k = V_c(s_{\min} - \rho) - V_c(s_{\min}) + r(k) > C(\delta, \eta)^{-1}.$$

This implies

$$E_k(s_{\min} - \delta) = E_k^+ + j\varphi_k f^2 \in E_k^+ + C(\delta, \eta)(V_c - \mu_k)j\varphi_k f^2 \in (1 + C(\delta, \eta))E_k^+(s_{\min} - \delta),$$

where all functions are taken at the point  $(s_{\min} - \delta)$ . Combining this estimate together with (4.25) and (4.24) yields

$$\begin{aligned} E_k(y) &> E_k^+(y) > C e^{-(2/\varepsilon_k)(d_A^c(y) + \delta)} E_k^+(s_{\min} - \rho) > C e^{-(2/\varepsilon_k)(d_A^c(y) + \delta)} E_k(s_{\min} - \rho) \\ &> C e^{-(2/\varepsilon_k)(d_A^c(y) + 2\delta)} > C e^{-(2/\varepsilon_k)(d_A^c(s) + 2\delta)}, \end{aligned}$$

where  $C$  is a new constant depending only on  $\delta$  and  $\eta$ . Note that in the last inequality, we have used that  $d_A^c(s) > d_A^c(y)$ . Integrating for  $y \in [s, s + \eta/2]$ , we get

$$(4.26) \quad \int_{[s, s + \eta/2]} E_k(y) \, dy > C e^{-(2/\varepsilon_k)(d_A^c(s) + 2\delta)}.$$

Now, turning to the global manifold and recalling that  $\psi_k(s, \theta) = e^{ik\theta} \varphi_k(s)$ , (4.5) and (4.13), we have

$$\begin{aligned} \int_{\mathcal{C}_{[s, s+\eta/2]}} E_k(y) R(y) dy &= \varepsilon_k^2 \int_{\mathcal{C}_{[s, s+\eta/2]}} j r_g \psi_k f^2 d \text{Vol}_g \\ &+ \int_{\mathcal{C}_{[s, s+\eta/2]}} \left( \frac{j r_g f^2}{4} \mu_k + 1 \right) j \psi_k f^2 d \text{Vol}_g \\ (4.27) \quad &\leq C k \psi_k k_{H^1(\mathcal{C}_{[s, s+\eta/2]})}^2. \end{aligned}$$

Finally, an interpolation estimates together with  $P_{\varepsilon_k} \psi_k = \mu_k \psi_k$  and the definition of  $P_{\varepsilon_k}$  in (4.8) gives

$$\begin{aligned} k \psi_k k_{H^1(\mathcal{C}_{[s, s+\eta/2]})}^2 &\leq C_\eta \left( k \psi_k k_{L^2(\mathcal{C}_{[s, s+\eta/2]})}^2 + k \psi_k k_{L^2(\mathcal{C}_{[s-\eta, s+\eta]})} k \Delta_g \psi_k k_{L^2(\mathcal{C}_{[s-\eta, s+\eta]})} \right) \\ (4.28) \quad &\leq C_\eta \varepsilon_k^2 k \psi_k k_{L^2(\mathcal{C}_{[s-\eta, s+\eta]})}^2. \end{aligned}$$

Now combining (4.26)–(4.27)–(4.28) gives

$$k \psi_k k_{L^2(\mathcal{C}_{[s-\eta, s+\eta]})}^2 > C \varepsilon_k^2 e^{(2/\varepsilon_k)(d_A^c(s)+2\delta)}, \quad k > k_0(\delta, \eta).$$

Finally noticing that  $\varepsilon_k^2 > e^{-\delta/\varepsilon_k}$  for  $k > k_0(\delta)$  ends the proof of Proposition 4.9 up to replacing  $\delta$  by  $\delta/3$ .

#### 4.7. MINIMAL TIME FOR UNIFORM CONTROLLABILITY IN THE LIMIT $\varepsilon \rightarrow 0^+$

The main purpose of this section is the proof Theorem 1.8 and its corollaries.

4.7.1. *Proof of Theorem 1.8.* — Recall that we consider the following situation:

–  $S$  is a surface of revolution as described in Section 4.1,  $c > 0$  is a fixed constant. Moreover, Assumption (1.17) is fulfilled.

– For  $k \geq \mathbb{N}$ ,  $\varepsilon = \varepsilon_k = ck^{-1}$  and  $\psi_k$  is the set of solutions of  $P_{\varepsilon_k} \psi_k = \mu_k \psi_k$  defined in Lemma 4.3 associated to  $s_0 = s_{\min}$  is the minimum of  $V_c$ .

– the function

$$(4.29) \quad v_k(t, x) = e^{-(\mu_k/\varepsilon_k)t} \psi_k(x)$$

is the solution to (2.9), namely  $(\varepsilon_k \partial_t + P_{\varepsilon_k})v_k = 0$ ,  $v_k|_{\partial S} = 0$ , and  $v_k|_{t=0} = \psi_k$ . Here,  $\psi_k$  denotes the eigenfunction constructed in Section 4.3 above (in particular  $k \psi_k k_{L^2(S)} = 1$ ) and studied in Sections 4.5–4.6.

We now want to test Inequality (2.10) on  $v_k$ , and thus estimate both sides of this inequality. This is achieved in Lemmas 4.12 and 4.13. Theorem 1.8 is then a direct consequence of these two lemmas. We recall that  $V_c$  is defined in (1.16) and  $W^c$  in (1.20).

LEMMA 4.12. — *For any  $\delta > 0$ , there exist  $C, k_0 > 0$  such that for all  $k > k_0$  and all  $0 \leq T_0 \leq \delta^{-1}$ , we have*

$$\|e^{-f/2\varepsilon} v_k(T_0)\|_{L^2(S)} > C e^{(V_c(s_{\min})T_0 + W_m^c + \delta)/\varepsilon_k}, \quad W_m^c = \min_{I_L} W^c,$$

with  $v_k$  defined in (4.29) and  $W^c(s) = d_A^c(s) + f(s)/2$ .

*Proof.* — Note first that the function  $W^c = d_A^c + f/2$  is continuous on  $I_L$  and converges to  $+1$  close to the poles  $p \in P$ , according to the asymptotics of  $d_A^c$  in Lemma 4.5. Hence, it reaches its minimum in (at least) one point that we denote  $s_1 \in I_L$ , that is  $W^c(s_1) = W_m^c = \min_{I_L} W^c$ . We take  $0 < \eta < \text{dist}(s_1, P)$  small enough so that  $f(s) - f(s_1) \leq \delta$  for  $|s - s_1| \leq \eta$ . We have for  $k > k_0$  large enough

$$\begin{aligned} \|e^{-f/2\varepsilon} v_k(T_0)\|_{L^2(S)} &= e^{(\mu_k/\varepsilon_k)T_0} \|e^{-f/2\varepsilon} \psi_k\|_{L^2(S)} \\ &> e^{(\mu_k/\varepsilon_k)T_0} \|e^{-f/2\varepsilon} \psi_k\|_{L^2(\mathcal{C}_{(s_1-\eta, s_1+\eta)})} \\ &> e^{(\mu_k/\varepsilon_k)T_0} e^{(f(s_1)+\delta)/2\varepsilon_k} \|\psi_k\|_{L^2(\mathcal{C}_{(s_1-\eta, s_1+\eta)})} \\ &> C e^{(\mu_k/\varepsilon_k)T_0} e^{(f(s_1)+\delta)/2\varepsilon_k} e^{(1/\varepsilon_k)(d_A^c(s_1)+\delta)}, \end{aligned}$$

where we have used Proposition 4.9 for the last estimate. Since  $\mu_k \rightarrow V_c(s_{\min})$ , we have  $\mu_k T_0 \leq V_c(s_{\min})T_0 + \delta$  for  $k$  large enough, which, together with the above estimate, concludes the proof (up to changing  $3\delta$  into  $\delta$ ).

LEMMA 4.13. — For any  $\omega \subset S$  and  $\delta > 0$ , there exist  $C, k_0 > 0$  such that for all  $k > k_0$ , and all  $0 \leq T_0 \leq \delta^{-1}$ , we have

$$\begin{aligned} \int_0^{T_0} \|e^{-f/2\varepsilon} v_k(t)\|_{L^2(\omega)}^2 dt &\leq C e^{-2(W_\omega^c - \delta)/\varepsilon_k}, \quad W_\omega^c = \min_{\omega} W^c, \\ \int_0^{T_0} \|e^{-f/2\varepsilon} \partial_s v_k(t)|_{s=L}\|_{H^1(S^1)}^2 dt &\leq C e^{-2(W^c(L) - \delta)/\varepsilon_k}, \quad \text{in Cases 2 and 3,} \\ \int_0^{T_0} \|e^{-f/2\varepsilon} \partial_s v_k(t)|_{s=0}\|_{H^1(S^1)}^2 dt &\leq C e^{-2(W^c(L) - \delta)/\varepsilon_k}, \quad \text{in Case 3,} \end{aligned}$$

with  $v_k$  defined in (4.29).

*Proof.* — Since we need an upper bound, we can assume without loss of generality that  $\omega$  is invariant by rotation. Also,  $W_\omega^c$  is finite except in the trivial case  $\omega = P$ . Let  $\delta > 0$ .

We first estimate the contribution close to the poles in case  $P \notin \omega$ . There, the function  $\psi_k$  (hence  $v_k$ ) is supposed to be very small since  $d_A^c$  is large. More precisely, using the asymptotics of  $d_A^c$  close to  $P$  given by Lemma 4.5, there exists  $\tilde{\eta} > 0$  so that

$$(4.30) \quad \frac{d_A^c(s)}{2} > W_\omega^c + \frac{kfk_{L^1}}{2} + 1, \quad \text{for all } s \in N_{\tilde{\eta}} = fS \cap I_L; \text{dist}(s, P) < \tilde{\eta}g.$$

We start with the estimate

$$(4.31) \quad \int_0^{T_0} \|e^{-f/2\varepsilon} v_k\|_{L^2(\omega \setminus \mathcal{C}_{N_{\tilde{\eta}}})}^2 dt \leq e^{kfk_{L^1}/\varepsilon} k\psi_k k_{L^2(\mathcal{C}_{N_{\tilde{\eta}}})}^2 \int_0^{T_0} e^{-2(\mu_k/\varepsilon_k)t} dt.$$

We have

$$\int_0^{T_0} e^{-2(\mu_k/\varepsilon_k)t} dt = \frac{\varepsilon_k}{2\mu_k} \int_0^{T_0(2\mu_k/\varepsilon_k)} e^{-s} ds \leq C \frac{\varepsilon_k}{2\mu_k} \leq 1, \quad \text{for } k \text{ large.}$$

Corollary 4.7 applied to the set  $\mathcal{C}_{N_{\bar{\eta}}}$  and the constant  $\delta = 1/2$ , together with (4.30) implies

$$k\psi_k k_{L^2(\mathcal{C}_{N_{\bar{\eta}}})} \leq C e^{(1/\varepsilon_k)(W_\omega^c + kfk_{L^1}/2)}.$$

With (4.31), this gives

$$\int_0^{T_0} \|e^{f/2\varepsilon} v_k\|_{L^2(\omega \setminus \mathcal{C}_{N_{\bar{\eta}}})}^2 dt \leq C e^{kfk_{L^1}/\varepsilon} e^{(2/\varepsilon_k)(W_\omega^c + kfk_{L^1}/2)} = C e^{(2/\varepsilon_k)W_\omega^c},$$

which is the expected bound for this part.

Let us now treat the contribution of the norm away from the poles (which is the whole  $I_L$  in Case 3). Since  $f$  is uniformly continuous on  $[0, L]$  and  $d_A^c$  is uniformly continuous on the compact set  $N_{\bar{\eta}}^c := I_L \cap N_{\bar{\eta}}$ , there exists  $\eta > 0$  so that

$$(4.32) \quad s, s' \geq 2N_{\bar{\eta}}^c \text{ and } |s - s'| \leq \eta \Rightarrow |f(s) - f(s')| \leq \delta \text{ and } |d_A^c(s) - d_A^c(s')| \leq \delta.$$

We now select a finite sequence  $s_i \geq 2N_{\bar{\eta}}^c$ ,  $i = 1, \dots, N$  so that  $f|_{s_i \leq s \leq s_i + \eta}$  is a finite covering of  $N_{\bar{\eta}}^c$ . This property gives the estimate

$$\int_0^{T_0} \|e^{f/2\varepsilon} v_k\|_{L^2(\omega \setminus \mathcal{C}_{N_{\bar{\eta}}^c})}^2 dt \leq \int_0^{T_0} e^{2(\mu_k/\varepsilon_k)t} dt \sum_{i \geq 1} \|e^{f/2\varepsilon} \psi_k\|_{L^2(\mathcal{C}_{(s_i - \eta, s_i + \eta)})}^2,$$

where  $J = \{i = 1, \dots, N\}$ ;  $(s_i - \eta, s_i + \eta) \setminus \omega \subset \mathcal{C}_{N_{\bar{\eta}}}$ . Using  $|f(s) - f(s_i)| \leq \delta$  and  $|d_A^c(s) - d_A^c(s_i)| \leq \delta$  for  $s \in (s_i - \eta, s_i + \eta)$  and then Corollary 4.7 with some  $0 < \delta' \leq \delta/(1 + d_A^c(s_i))$  so that  $(1 - \delta')d_A^c(s_i) - \delta' > d_A^c(s_i) - \delta$ , we obtain

$$\begin{aligned} \|e^{f/2\varepsilon} \psi_k\|_{L^2(\mathcal{C}_{(s_i - \eta, s_i + \eta)})} &\leq e^{(f(s_i) + \delta)/2\varepsilon} k\psi_k k_{L^2(\mathcal{C}_{(s_i - \eta, s_i + \eta)})} \\ &\leq C e^{(f(s_i) + \delta)/2\varepsilon} e^{(1/\varepsilon_k)((1 - \delta')d_A^c(s_i) - \delta')} e^{\delta'/\varepsilon} \\ &\leq C e^{(W^c(s_i) - 4\delta)/\varepsilon_k}. \end{aligned}$$

We finally obtain

$$\int_0^{T_0} \|e^{f/2\varepsilon} v_k\|_{L^2(\omega \setminus \mathcal{C}_{N_{\bar{\eta}}^c})}^2 dt \leq C e^{8\delta/\varepsilon_k} \max_{i \geq 1} e^{2W^c(s_i)/\varepsilon_k}.$$

We remark from (4.32) that  $\min_{i \geq 1} W^c(s_i) > W_\omega^c - 2\delta$ . This finishes the proof since  $\delta$  is arbitrary and  $k$  can be chosen large enough.

Finally, the proof of the boundary estimates simply consists in replacing the use of Corollary 4.7 by that of the inequality (4.18).

We may now conclude the proof of Theorem 1.8 from Lemmas 4.12 and 4.13

*Proof of Theorem 1.8.* — Using Lemma 2.9, if uniform observability holds for  $T_0$ , then, we have the inequality

$$\|e^{f/2\varepsilon} v(T_0)\|_{L^2(S)}^2 \leq C_0^2 \int_0^{T_0} \|e^{f/2\varepsilon} v\|_{L^2(\omega)}^2 dt,$$

for any solution  $v$  of (2.9). In particular, this inequality holds true for the sequence  $v_k$  described above. So, combining Lemmas 4.12 and 4.13, we obtain

$$e^{(V_c(s_{\min})T_0 + W_m^c + \delta)/\varepsilon_k} \leq C_0 e^{(W_\omega^c - \delta)/\varepsilon_k}, \quad k > k_0(\delta), \quad \text{for all } k > k_0.$$

This implies  $V_c(s_{\min})T_0 > W_\omega^c - W_m^c - 2\delta$  when letting  $k \rightarrow +\infty$ , which gives the expected result since  $\delta$  is arbitrary. The proof of the boundary observability estimate (1.22) follows the same. We also notice that all the previous results also apply in the case with the alternative Definition 1.19 of the Agmon distance which is also Lipschitz with the same properties that we used.

**4.7.2. Proof of Corollaries of Theorem 1.8.** — Corollaries 1.9, 1.10 and 1.11, stated in the introduction, are significant examples of application of Theorem 1.8. In this section, we prove these three results.

*Proof of Corollary 1.9.* — We first consider the case  $S$  diffeomorphic to  $S^2$ , i.e., Case 1. The case  $S$  diffeomorphic to  $D$ , i.e., Case 2, is discussed at the end of the proof. We define  $f_\delta(s) = \int_0^s \chi_\delta(t) dt$  where  $\chi_\delta \in C^1((0, L); [0, 1])$ ,  $\chi_\delta(s) = 1$  in a neighborhood of  $[\delta, L - \delta]$ . Such a function  $f_\delta$  is constant near 0 and  $L$ , and hence can be extended by continuity as a  $C^1$  function on  $S$  (see e.g. [Bes78, Prop. 4.6]). We notice that  $f_\delta'(s) = \chi_\delta(s) = 1$  for  $s \in [\delta, L - \delta]$ , so the statements about  $T_{\text{GCC}}$  are direct consequences of Proposition 4.2.

Notice now that we have

$$V_c(s) := \frac{c^2}{R(s)^2} + \frac{f_\delta(s)^2}{4}.$$

Let us call  $s_{\min} \in (0, L)$  the unique point such that  $R(s_{\min}) = \max R$ , that is  $1/R(s_{\min})^2 = \min 1/R^2$ .

**CLAIM.** — For all  $c > 0$ , there is  $\delta_0 > 0$  such that for all  $\delta \in (0, \delta_0)$ , we have  $V_c(s_{\min}) = \min V_c$  and  $V_c^{-1}(V_c(s_{\min})) = f_{s_{\min}} \mathcal{G}$ .

To prove the claim, we let  $\delta_0 > 0$  be such that

$$s_{\min} \in [\delta_0, L - \delta_0], \quad \text{and} \quad \frac{c^2}{R^2(s)} > \frac{c^2}{R(s_{\min})^2} + \frac{1}{4} \text{ for } s \notin [\delta_0, L - \delta_0]$$

(note that  $\delta_0$  thus depends on  $c$ ). This is possible since  $R(s) \rightarrow 0$  as  $s \rightarrow 0^+$  and  $s \rightarrow L^-$ . Hence, recalling the definitions of  $\chi_\delta$  and  $V_c$ , for  $\delta < \delta_0$  we have

$$V_c(s_{\min}) = \frac{c^2}{R(s_{\min})^2} + \frac{1}{4}, \quad V_c(s) > \frac{c^2}{R(s_{\min})^2} + \frac{1}{4} \text{ for all } s \in [\delta_0, L - \delta_0] \cap f_{s_{\min}} \mathcal{G},$$

together with

$$V_c(s) > \frac{c^2}{R^2(s)} > \frac{c^2}{R(s_{\min})^2} + \frac{1}{4} \text{ for } s \notin [\delta_0, L - \delta_0].$$

As a consequence,  $V_c$  reaches its minimum at  $s_{\min}$  only, which proves the claim.



Now Assumption 4.4 is satisfied and  $s_{\min}$  does not depend on  $\delta$ . We compute the Agmon distance (1.18)

$$\begin{aligned} d_A^c(s) &= \left| \int_{s_{\min}}^s \sqrt{V_c(y) - V_c(s_{\min})} dy \right| \\ &= \left| \int_{s_{\min}}^s \sqrt{\frac{c^2}{R(y)^2} + \frac{j\chi_\delta(y)^2}{4} - \frac{c^2}{R(s_{\min})^2}} \frac{1}{4} dy \right|. \end{aligned}$$

Notice that

$$0 > \frac{j\chi_\delta(y)^2}{4} - \frac{1}{4} > -\frac{1}{4}$$

uniformly with respect to  $\delta$  so that the asymptotic expansion of Lemma 4.5 is valid uniformly in  $\delta$ : there are  $C, \gamma > 0$  such that for all  $\delta \geq (0, \delta_0)$ ,

$$\begin{aligned} jd_A^c(s) + c \log(s) &\leq C \quad \text{for } s \geq (0, \gamma], \\ jd_A^c(s) + c \log(L - s) &\leq C, \quad \text{for } s \geq [L - \gamma, L). \end{aligned}$$

Now we recall the definition of  $W^c = d_A^c + f_\delta/2$ , notice that  $0 \leq f_\delta^0 \leq 1$  so that  $0 \leq f_\delta(s) \leq L$  for  $s \geq I_L$ . As a consequence, using that  $d_A^c > 0$  and  $d_A^c(s_{\min}) = 0$ , we have for  $\delta \geq (0, \gamma]$ ,

$$\begin{aligned} W_m^c &= \min_{I_L} W^c = \min_{I_L} (d_A^c + f_\delta/2) \leq \frac{L}{2} + \min_{I_L} d_A^c \leq \frac{L}{2}, \\ W_\omega^c &= \min_{\bar{\omega}} W^c > \min_{\bar{\omega}} d_A^c > -c \log(\delta) - C \quad \text{for } \omega = B_g(N, \delta) \cup B_g(S, \delta). \end{aligned}$$

The bound (1.21) of Theorem 1.8 then yields

$$\left( \frac{c^2}{R(s_{\min})^2} + \frac{1}{4} \right) T_{\text{unif}}(\omega) > -c \log(\delta) - C - \frac{L}{2},$$

and hence concludes the proof in the case S di eomorphic to  $S^2$ , i.e., Case 1.

In the case S di eomorphic to D, i.e., Case 2, we instead define  $f_\delta(s) = \int_0^s \chi_\delta(t) dt$  where  $\chi_\delta \geq C^{-1} \mathbb{1}_{(0, L]}; [0, 1]$ ,  $\chi_\delta(s) = 1$  in a neighborhood of  $[\delta, L]$ . Then, the remainder of the proof is the same except that the minimum can be achieved at  $s = L$ , and all sets of the form  $[\delta_0, L - \delta_0]$  have to be replaced by  $[\delta_0, L]$  (i.e., only a neighborhood of zero is avoided, and not a neighborhood of  $L$ ).

*Proof of Corollary 1.10.* — For the sake of simplicity, we may identify

$$S_L^1 = [L/2, L/2] \quad \text{and} \quad I_\omega = (\alpha, \alpha) \quad \text{for } \alpha \geq (0, L/2).$$

We first choose  $\chi_\delta \geq C^{-1} \mathbb{1}_{(S_L^1)}$  even in this identification, and  $s_{\min} = L/2 \notin (\alpha, \alpha)$  such that

$$\begin{aligned} \chi_\delta &= \frac{1}{\delta} \text{ on } \left[ \frac{1}{2}(\alpha + L/2), \frac{1}{2}(\alpha + L/2) \right], \quad \chi_\delta(s_{\min}) = 1, \\ \chi_\delta^0(s) &= 0 \text{ if and only if } s \geq \left[ \frac{1}{2}(\alpha + L/2), \frac{1}{2}(\alpha + L/2) \right] \text{ or } s = s_{\min}. \end{aligned}$$

Note that  $\chi_\delta = 1/\delta$  in a neighborhood of  $(\alpha, \alpha)$ , and that for  $\delta < 1$ ,  $\chi_\delta$  reaches at  $s_{\min} = L/2$  a unique global minimum and in particular  $\chi_\delta > 1$  on  $S_L^1$ . We then set

$$V^\delta(s) := \chi_\delta + M, \quad M := \max_{S_L^1} f_\delta^0/4,$$

so that  $V^\delta(s) > f^\theta(s)^2/4$  on  $S_L^1$ , and

$$R_\delta(s) := (V^\delta(s) - f^\theta(s)^2/4)^{1/2}, \quad R_\delta \in C^1(S_L^1; \mathbb{R}^+),$$

where we have used that  $f \in C^1(S_L^1)$ . Notice that with these definitions, we have (where  $V_1$  denotes  $V_c$  with  $c = 1$ )

$$V_1(s) = \frac{1}{R_\delta(s)^2} + \frac{f^\theta(s)^2}{4} = V^\delta(s).$$

Notice then that  $V^\delta$  admits a unique global minimum at  $s_{\min} = L/2$ .

On the one hand, we have

$$(\chi_\delta(s) + M)^{-1/2} \leq R_\delta(s) \leq \chi_\delta^{-1/2}(s).$$

On the other hand, since  $\chi_\delta$  is even and with the appropriate definition (1.19), we have for  $s \in [0, L/2]$ ,

$$d_A^c(s) = \int_{L/2}^s \sqrt{V^\delta(y) - V^\delta(L/2)} dy = \int_{L/2}^s \sqrt{\chi_\delta(y) - 1} dy.$$

As a consequence, we have (recall  $\omega = (\alpha, \alpha) \in S^1$ )

$$W_m^c = \min_{[0, L]} W^c = \min_{[0, L]} (d_A^c + f/2) \leq (d_A^c + f/2)(L/2) = \frac{f(L/2)}{2},$$

$$W_\omega^c > \min_{\bar{\omega}} d_A^c + \min_{\bar{\omega}} f/2 = d_A^c(\alpha) + \min_{\bar{\omega}} f/2,$$

where

$$\begin{aligned} d_A^c(\alpha) &= \int_\alpha^{L/2} \sqrt{\chi_\delta(y) - 1} dy > \int_\alpha^{\frac{1}{2}(\alpha+L/2)} \sqrt{\chi_\delta(y) - 1} dy = \int_\alpha^{\frac{1}{2}(\alpha+L/2)} \sqrt{(1/\delta) - 1} dy \\ &> \frac{1}{2}((L/2) - \alpha)(\delta^{-1/2} - 1) \end{aligned}$$

for  $\delta < 1$ . Applying Theorem 1.8 concludes the proof of the corollary.

We next prove Corollary 1.11, stated in the introduction. The proof is close to that of Corollary 1.10.

*Proof of Corollary 1.11.* — We first take  $\chi \in C^1(\mathbb{R}_+; [0, 1])$  such that  $\chi = 1$  on  $[0, L/4]$  and  $\text{supp } \chi \subset [0, L/2]$ . Next, with

$$M := \max_{[0, L]} f^\theta^2/4,$$

we set

$$V^\delta(s) := \frac{\chi(s)}{(s + \delta)^\gamma} + (1 - \chi(s))(s - L/2)^2 + M, \quad \text{for } s \in [0, L/2],$$

$$V^\delta(s) := V^\delta(L - s) \quad \text{for } s \in [L/2, L].$$

This function is symmetric about  $L/2$ , smooth on  $[0, L]$ , and satisfies  $V^\delta(s) > f^\theta(s)^2/4$  on  $[0, L]$ . Hence, defining  $R_\delta$  as

$$R_\delta(s) := (V^\delta(s) - f^\theta(s)^2/4)^{1/2},$$

and using that  $f \geq C^{-1}([0, L])$ , we deduce that  $R_\delta \geq C^{-1}([0, L]; \mathbb{R}^+)$ . Notice that with these definitions, we have (where  $V_1$  denotes  $V_c$  with  $c = 1$ )

$$V_1(s) = \frac{1}{R_\delta(s)^2} + \frac{f^\delta(s)^2}{4} = V^\delta(s).$$

Notice then that  $V^\delta$  admits a unique global minimum at  $s_{\min} = L/2$ .

We have on the one hand that for all  $s \geq [0, L/4]$  and  $\delta \geq (0, \delta_0)$ ,

$$(s + \delta)^{\gamma/2} \frac{1}{\sqrt{1 + M(s + \delta)^\gamma}} = (V^\delta(s))^{-1/2} \circlearrowleft R_\delta(s) \circlearrowleft (V^\delta(s) - M)^{-1/2} = (s + \delta)^{\gamma/2},$$

which proves the Item 4 (on account to the symmetry  $V^\delta(s) := V^\delta(L - s)$ ).

On the other hand, recalling that  $s_{\min} = L/2$ , we have for  $s \geq [0, L/2]$

$$\begin{aligned} V_1(s_{\min}) &= V^\delta(L/2) = M, \\ d_A^c(s) &= \left| \int_{L/2}^s \sqrt{V^\delta(y) - V^\delta(s_{\min})} dy \right| \\ &= \left| \int_{L/2}^s \sqrt{\frac{\chi(s)}{(s + \delta)^\gamma} + (1 - \chi(s))(s - L/2)^2} dy \right|. \end{aligned}$$

As a consequence, we have

$$\begin{aligned} W_m^c &= \min_{[0, L]} W^c = \min_{[0, L]} (d_A^c + f/2) \circlearrowleft (d_A^c + f/2)(L/2) = \frac{f(L/2)}{2}, \\ W_{f \neq 0}^c &= W^c(0) > d_A^c(0) + \frac{f(0)}{2}, \end{aligned}$$

where

$$\begin{aligned} d_A^c(0) &= \int_0^{L/2} \sqrt{\frac{\chi(s)}{(s + \delta)^\gamma} + (1 - \chi(s))(s - L/2)^2} dy > \int_0^{L/4} \sqrt{\frac{1}{(s + \delta)^\gamma}} dy \\ &= \frac{\delta^{1 - \gamma/2}}{\gamma/2 - 1} - \frac{(\delta + L/4)^{1 - \gamma/2}}{\gamma/2 - 1} = \frac{\delta^{1 - \gamma/2}}{\gamma/2 - 1} + O(1), \quad \text{for } \gamma > 2. \end{aligned}$$

By symmetry of  $V^\delta$  about  $L/2$ , we also have  $d_A^c(L) = d_A^c(0)$ . Applying Theorem 1.8 concludes the proof of the corollary.

## 5. UNIFORM TIME OF OBSERVABILITY FOR POSITIVE SOLUTIONS

The proofs of Theorem 1.3 and Proposition 1.12 rely on fine estimates on the semiclassical heat kernel, which we borrow from [LY86]. The latter are first presented in Section 5.1. Then, in Section 5.2, we deduce  $L^1$  observability statements and finally conclude the proofs of Theorem 1.3 and Proposition 1.12 in Section 5.3. Throughout this section, we assume  $\partial M = ?$ .

5.1. ESTIMATES ON THE SEMICLASSICAL HEAT KERNEL. — The main tool we use to estimate the heat kernel in the semiclassical limit is the following theorem taken from Li-Yau [LY86] (see also [Sim83], for a similar result on  $\mathbb{R}^n$ ).

THEOREM 5.1 (Theorem 6.1 of [LY86]). — *Let  $M$  be a compact manifold without boundary. Suppose  $V_\varepsilon = V + \varepsilon q$  with  $V, q \in C^2(M)$ . For any  $\varepsilon > 0$ , we consider  $H_\varepsilon$ , the fundamental solution of*

$$\partial_t w - \Delta_g w + \frac{1}{\varepsilon^2} V_\varepsilon(x)w = 0, \quad \text{on } (0, +\infty) \times M.$$

Then, we have

$$(5.1) \quad \lim_{\varepsilon \downarrow 0} \varepsilon \log H_\varepsilon(x, y, \varepsilon t) = \rho(x, y, t),$$

where

$$(5.2) \quad \rho(x, y, t) = \inf \left\{ \int_0^t \frac{1}{4} |\dot{\gamma}(s)|_g^2 + V(\gamma(s)) ds; \gamma \in W^{1,1}([0, t]; M), \gamma(0) = x, \gamma(t) = y \right\}.$$

Moreover, the limit in (5.1) is uniform on any compact set of  $M^2 \times (0, +\infty)$ .

We recall that  $H_\varepsilon(x, y, t)$  is defined to be the unique solution to

$$(5.3) \quad \begin{cases} \left( \partial_t - \Delta_g + \frac{1}{\varepsilon^2} V_\varepsilon(x) \right) H_\varepsilon(x, y, t) = 0, & \text{for } (t, x) \in \mathbb{R}^+ \times M, \\ H_\varepsilon(x, y, t)|_{t=0} = \delta_{x=y}, & \text{for } x \in M, \end{cases}$$

where  $y \in M$  is fixed, and the differential operator  $\Delta_g + (1/\varepsilon^2)V_\varepsilon(x)$  acts in the  $x$ -variable. We also recall that  $H_\varepsilon(x, y, t)$  is (well-defined and) continuous in  $M^2 \times (0, +\infty)$  as soon as  $V_\varepsilon \in L^1(M)$ , see e.g. [Sim82, Th. B.7.1 (a<sup>000</sup>)], so that pointwise estimates like (5.1) make sense.

The statement of Theorem 5.1 is not strictly speaking a consequence of [LY86, Th. 6.1] for the following two reasons:

- the potential  $V_\varepsilon$  is assumed independent of  $\varepsilon$  in [LY86];
- the uniformity of the limit on any compact subset is not explicitly written in [LY86].

However, let us explain why the proof of [LY86, Th. 6.1] actually contains these two points. The limit (5.1) is proved in two steps, a lower bound and an upper bound. The lower bound

$$\lim_{\varepsilon \downarrow 0} \varepsilon \log H_\varepsilon(x, y, \varepsilon t) > \rho(x, y, t)$$

is obtained as a consequence of the explicit estimate ([LY86, Th. 2.1])

$$\varepsilon \log H_\varepsilon(x, y, \varepsilon t_1) \geq \varepsilon \log H_\varepsilon(x, y, \varepsilon t) + \varepsilon^2 A_{1/\varepsilon}(t - t_1) + \rho_{\alpha, R}(x, y, t - t_1),$$

for all  $0 < t_1 < t$ ,  $\alpha > 1$  and where the constant  $A_\lambda$  ( $\lambda = \varepsilon^{-1}$  in our context) only depends on

$$k \Delta_g V_\varepsilon k_{L^1(M)}, \quad k \Gamma_g V_\varepsilon k_{L^1(M)} \quad \text{and} \quad \alpha.$$

In particular, the limit  $\lim_{\varepsilon \rightarrow 0} \varepsilon A_{1/\varepsilon} = 0$  holds uniformly. The proof finally proceeds by taking the limit  $R \rightarrow +\infty$ , and then  $\alpha \rightarrow 1^+$  (in which case  $\rho_{\alpha, R}^{V_\varepsilon}(x, y, t - t_1) \rightarrow \rho^{V_\varepsilon}(x, y, t - t_1)$ , uniformly on compact sets; here  $\rho^{V_\varepsilon}$  denotes the function  $\rho$  defined by (5.2) with  $V$  replaced by  $V_\varepsilon$ ). Taking finally the limit  $\varepsilon \rightarrow 0^+$ , and then  $t_1 \rightarrow 0^+$ , and noticing that

$$\lim_{t_1 \rightarrow 0^+} \lim_{\varepsilon \rightarrow 0^+} \varepsilon \log H_\varepsilon(x, y, \varepsilon t_1) > 0$$

concludes the proof of the lower bound. This last argument relies only on a uniform upper bound on  $k_{V_\varepsilon} k_{L^1}(\mathbb{M})$  together with a comparison argument, namely Inequality [LY86, (6.3)]. It can be checked that the convergence is uniform on any compact set since it involves the asymptotics on the diagonal of kernels of heat equations on large balls and without potential, which are known to be uniform on compact sets, see for instance [Var67, Th. 4.6]. It only remains to notice that the limit  $\lim_{t_1 \rightarrow 0^+} \lim_{\varepsilon \rightarrow 0^+} \rho^{V_\varepsilon}(x, y, t - t_1) = \rho(x, y, t)$  is uniform on compact sets.

The upper bound

$$\lim_{\varepsilon \rightarrow 0} \varepsilon \log H_\varepsilon(x, y, \varepsilon t) \leq \rho(x, y, t)$$

follows from [LY86, Th. 3.3]. As for the lower bound, this result also furnishes an explicit and uniform bound involving another constant  $A_\lambda$  enjoying the same type of convergence properties as for the lower bound.

Note that we have chosen to use the estimate of the semiclassical limit of the kernel of [LY86] but it is likely that we could have obtained the observability inequality starting directly from Harnack type inequalities like [LY86, Th. 2.1], as we did in [LL21a].

Here, we are mostly interested in the case  $V(x) = \frac{1}{4} |r_g f(x)|_g^2$ . In this situation, we can reformulate the result in terms of the transport equation with vanishing viscosity (1.6).

We next define  $K_\varepsilon(x, y, t)$ , the fundamental solution of (1.6) on  $\mathbb{M}^2 \times (0, +\infty)$  by the unique solution to

$$(5.4) \quad \begin{cases} (\partial_t - r_g f - r_g - q - \varepsilon \Delta_g) K_\varepsilon(x, y, t) = 0, & \text{for } (t, x) \in \mathbb{R}^+ \times \mathbb{M}, \\ K_\varepsilon(x, y, t)|_{t=0} = \delta_{x=y}, & \text{for } x \in \mathbb{M}, \end{cases}$$

where  $y \in \mathbb{M}$  is fixed, and the differential operator  $\partial_t - r_g f - r_g - q - \varepsilon \Delta_g$  acts in the  $x$ -variable. Recall that for  $u_0 \in L^1(\mathbb{M})$ , the function

$$u(t, x) = \int_{\mathbb{M}} K_\varepsilon(x, y, t) u_0(y) d\text{Vol}_g(y)$$

is the unique solution of (1.6) on  $(0, +\infty) \times \mathbb{M}$  issued from  $u|_{t=0} = u_0$  (This uses the choice of volume form  $d\text{Vol}_g$  in the embedding  $L^1(\mathbb{M}) \hookrightarrow D^0(\mathbb{M})$ ).

**COROLLARY 5.2.** — *Let  $\mathbb{M}$  be a compact manifold without boundary. Suppose  $X = r_g f$  where  $f$  is a  $C^3$  function defined on  $\mathbb{M}$ . For any  $\varepsilon > 0$ , we consider  $K_\varepsilon$ , the fundamental*

solution of (1.6) on  $(0, +1) \times M$ . Then, we have

$$\lim_{\varepsilon \downarrow 0} \varepsilon \log K_\varepsilon(x, y, t) = d_{r_g f}(x, y, t),$$

with

$$(5.5) \quad d_{r_g f}(x, y, t) := \rho(x, y, t) + \frac{f(x) - f(y)}{2},$$

where  $\rho(x, y, t)$  is defined by (5.2) with  $V(x) = j r_g f_g^2/4$ . Moreover, the limit is uniform on any compact set of  $M^2 \times (0, +1)$ .

In particular, for any  $\delta > 0$  and any compact subset  $I \subset (0, +1)$ , there exists  $\varepsilon_0 > 0$  such that

$$(5.6) \quad e^{-(d_{r_g f}(x, y, t) + \delta)/\varepsilon} \leq K_\varepsilon(x, y, t) \leq e^{-(d_{r_g f}(x, y, t) - \delta)/\varepsilon}$$

for any  $(x, y, t) \in I$  and  $0 < \varepsilon \leq \varepsilon_0$ .

Note that the definition of  $d_{r_g f}$  in (5.5) is not the same as that given in (1.24) in the introduction. Equivalence between these two definitions is proved in Lemma A.2.

Note that although the kernel  $H_\varepsilon(x, y, t)$  is symmetric (with respect to the Riemannian volume measure  $d\text{Vol}_g$ ) since  $\Delta_g + \frac{1}{\varepsilon^2} V_\varepsilon$  is, this is no longer the case for the kernel  $K_\varepsilon(x, y, t)$  (since the operator in (1.6) is not symmetric in  $L^2(M, d\text{Vol}_g)$ ). Similarly,  $\rho(x, y, t)$  is symmetric whereas  $d_{r_g f}(x, y, t)$  is not.

*Proof.* — Setting  $u_y(t, x) = K_\varepsilon(x, y, t)$  and  $w_y(t, x) = e^{f(x)/2\varepsilon} u_y(t/\varepsilon, x)$ , we have from Equation (5.4) and Lemma 2.9 that  $w_y(t, x)$  solves (2.12). Moreover, we have

$$w_y(0, x) = e^{f(x)/2\varepsilon} u_y(0, x) = e^{f(x)/2\varepsilon} \delta_{x=y} = e^{f(y)/2\varepsilon} \delta_{x=y}.$$

This implies that  $w_y(t, x) = e^{f(y)/2\varepsilon} H_\varepsilon(x, y, t)$  where  $H_\varepsilon(x, y, t)$  is defined in (5.3) with  $V_\varepsilon = j r_g f_g^2/4 + \varepsilon(\Delta_g f/2 - q)$ . Finally, we have proved that

$$H_\varepsilon(x, y, \varepsilon t) = e^{f(x)/2\varepsilon} e^{-f(y)/2\varepsilon} K_\varepsilon(x, y, t),$$

and hence

$$\lim_{\varepsilon \downarrow 0} \varepsilon \log K_\varepsilon(x, y, t) = \frac{f(y) - f(x)}{2} + \lim_{\varepsilon \downarrow 0} \varepsilon \log H_\varepsilon(x, y, \varepsilon t) = \frac{f(y) - f(x)}{2} + \rho(x, y, t),$$

after having applied (5.1).

**5.2.  $L^1$  OBSERVABILITY ESTIMATES FOR POSITIVE SOLUTIONS.** — We first prove intermediate observability statements in  $L^1$ . The following elementary abstract lemma shows that concerning positive solutions, observability in a (possibly weighted)  $L^1$  norm is equivalent to the “observability of the kernel”.

**PROPOSITION 5.3.** — *Let  $M$  be a compact Borel space (on which we denote by  $dx$  a distinguished measure) and  $T > 0$ . Take  $K = K(x, y, t) \geq C^0(M \times M \times (0, T])$  be a nonnegative kernel. Assume further that*

$$(t, y) \mapsto \int_M K(x, y, t) dx$$

is uniformly bounded for  $(t, y) \in (0, T] \times M$ . Define for  $t \in (0, T]$  the operator

$$S(t) : \text{Meas}(M) \rightarrow C^0(M), \quad [S(t)\mu](x) = \int_M K(x, y, t) d\mu(y) = \langle \mu, K(x, \cdot, t) \rangle,$$

where  $\text{Meas}(M)$  denotes the space of Radon measures on  $M$ . Let  $w_1 \in L^1([0, T] \times \omega)$  and  $w_2 \in L^1(M)$  be two nonnegative weight functions.

Then, for all  $T, s, C_0 > 0$ , the following statements are equivalent

(1) Observability of positive measures:

$$(5.7) \quad \|w_2 [S(s)\mu]\|_{L^1(M)} \leq C_0 \|w_1 [S(\cdot)\mu]\|_{L^1([0, T] \times \omega)}$$

for any  $\mu \in \text{Meas}_+(M)$  nonnegative Radon measure.

(2) Observability of positive  $L^1$  functions:

$$\|w_2 [S(s)u_0]\|_{L^1(M)} \leq C_0 \|w_1 [S(\cdot)u_0]\|_{L^1([0, T] \times \omega)}$$

for any  $u_0 \in L^1(M, dx)$  with nonnegative value.

(3) Observability of Dirac distributions:

$$\|w_2 [S(s)\delta_y]\|_{L^1(M)} \leq C_0 \|w_1 [S(\cdot)\delta_y]\|_{L^1([0, T] \times \omega)}$$

for any  $y \in M$ .

(4) Observability of the kernel:

$$I_s(y) \leq C_0 O_T(y), \quad \text{for all } y \in M,$$

where

$$O_T(y) = \int_0^T \int_\omega w_1(t, y) K(x, y, t) dx dt, \quad \text{and} \quad I_s(y) = \int_M w_2(x) K(x, y, s) dx.$$

Note first that under the assumption of the theorem, both terms in (5.7) are well-defined. Indeed, the Tonelli theorem (all functions/measures are nonnegative) implies

$$\begin{aligned} \|w_1 [S(\cdot)\mu]\|_{L^1([0, T] \times \omega)} &= \int_0^T \int_\omega w_1(t, x) [S(t)\mu](x) dx dt \\ &= \int_0^T \int_\omega w_1(t, x) \int_M K(x, y, t) d\mu(y) dx dt \\ &= \int_0^T \int_M \left( \int_\omega w_1(t, x) K(x, y, t) dx \right) d\mu(y) dt \\ &\leq T \|w_1\|_{L^1([0, T] \times \omega)} \|\mu\|_{TV} \sup_{t \in [0, T]} \sup_{y \in M} \int_\omega K(x, y, t) dx < +\infty, \end{aligned}$$

by assumption. Here  $\|\mu\|_{TV}$  denotes the total variation of the measure  $\mu$ . Note also that  $u(t) = [S(t)\mu]$  is a continuous nonnegative function for any nonnegative measure  $\mu$  and  $t > 0$ . Remark that  $O_T(y)$  is essentially the observation of solutions starting from  $\delta_y$  while  $I_s(y)$  is the weighted norm of this solution at time  $s$ .

*Proof.* — Remark first that

$$O_T(y) = \int_0^T \int_{\omega} w_1(t, y) K(x, y, t) dx dt = \int_0^T \int_{\omega} w_1(t, y) h \delta_y, K(x, \cdot, t) i dx dt$$

$$= k w_1 [S(\cdot) \delta_y] k_{L^1([0, T] \times \omega)}$$

$$I_s(y) = \int_M w_2(x) K(x, y, s) dx = \int_M w_2(x) h \delta_y, K(x, \cdot, s) i dx = k w_2 [S(s) \delta_y] k_{L^1(M)},$$

so that Item (3), Item (4). Moreover, applying the Fubini Theorem, we get

$$k w_1 [S(\cdot) \mu] k_{L^1([0, T] \times \omega)} = \int_0^T \int_{x \in \omega} \int_{y \in M} w_1(t, y) K(x, y, t) d\mu(y) dx dt = h\mu, O_T i$$

$$k w_2 [S(s) \mu] k_{L^1(M)} = \int_{y \in M} \int_{x \in M} w_1(t, x) K(x, y, t) d\mu(y) dx dt = h\mu, I_s i.$$

Therefore, Item (1) is equivalent to  $h\mu, C_0 O_T - I_s i > 0$  for any  $\mu \in \text{Meas}_+(M)$  and Item (2) to  $\int_M u_0(y) (C_0 O_T - I_s)(y) dy > 0$  for any  $u_0 \in L^1_+(M, dx)$ . That Item (4), Item (1), Item (2) follows from the general fact that if  $f \in C^0(M)$ , one has

$$f > 0 \text{ on } M \iff h\mu, f i > 0 \text{ for all } \mu \in \text{Meas}_+(M)$$

$$\iff \int_M u_0(y) f(y) dy > 0 \text{ for all } u_0 \in L^1_+(M, dx).$$

We now give the  $L^1$  observability estimate for positive solutions to (1.6).

**PROPOSITION 5.4.** — Assume that  $T > T_{GCC}(M, r_g f, \omega)$ . Then, for any  $\delta, s > 0$ , there exists  $\varepsilon_0 > 0$  so that we have

$$(5.8) \quad k u(s) k_{L^1(M)} \leq e^{\delta/\varepsilon} k u k_{L^1([0, T] \times \omega)}, \quad \text{for all } \varepsilon \in (0, \varepsilon_0),$$

for any  $u_0 \in L^1(M)$  with non-negative values and  $u$  solution of (1.6).

Remark that we only use the case  $s = T$  below. It is however remarkable that the stronger result for  $s > 0$  small holds as well. This is linked to the  $L^1$  setting here. Note also that in  $L^1$ , we have a “converse inequality”, which we state for the sake of the comparison. Proposition 5.4 is proved afterward.

**LEMMA 5.5.** — Assume  $\partial M = ?$ . For all  $T > 0$ , there is  $C_T > 0$  such that for all  $u_0 \in L^1(M; \mathbb{R}^+)$  and  $u$  the associated solution of (1.6), we have

$$(5.9) \quad k u(t) k_{L^1(M)} \leq C_T k u(T) k_{L^1(M)}, \quad \text{for all } t \in [0, T] \text{ and } \varepsilon > 0.$$

In particular, this implies that one cannot hope to replace the loss  $e^{\delta/\varepsilon}$  by a gain  $e^{-\delta/\varepsilon}$  in (5.8).

*Proof of Lemma 5.5.* — Assume first that  $u_0 \in W^{2,1}(M)$  with  $u_0 > 0$  a.e. on  $M$ . Then, notice that  $u(t, x) > 0$  for a.e.  $(t, x) \in (0, T) \times M$ . Integrating (1.6) on  $M$ , we obtain after an integration by parts (using that  $\partial M = ?$ )

$$\frac{d}{dt} \int_M u - \int_M (\text{div}_g X - q) u = 0.$$



Since  $u > 0$ , this implies

$$\frac{d}{dt} \|ku(t, \cdot)\|_{L^1} > \gamma \|ku(t, \cdot)\|_{L^1} \quad \text{for all } t > 0,$$

with  $\gamma = \inf_M (\operatorname{div}_g X - q)$ . The Grönwall inequality yields (5.9). The conclusion for a general  $u_0 \geq L^1$  follows from a density argument.

We now turn to the proof of the  $L^1$  observability estimate of Proposition 5.4, which will use the following lemma.

**LEMMA 5.6.** — *Assume  $(M, r_g f, \omega, (\delta^0, T))$  satisfies (GCC) (see Definition 2.10) for some  $\delta^0 > 0$ . Then, for all  $\delta > 0$  there is  $C_\delta > 0$  such that for all  $y \in M$  there is an open set  $U_y \subset \omega \cap (\delta^0, T)$  with  $|U_y| > C_\delta$  and for all  $(x, t) \in U_y$ , we have  $d_{r_g f}(x, y, t) \leq \delta$ .*

*Proof of Lemma 5.6.* — The assumption (GCC) implies that for any  $y_0 \in M$ , there is  $t_{y_0} \in (\delta^0, T)$  and  $x_{y_0} = \phi_{-t_{y_0}}(y_0) \in \omega$  where  $(\phi_t)_{t \in \mathbb{R}}$  is the flow of  $r_g f$ . The trajectory  $\gamma(s) = \phi_s(x_{y_0})$  satisfies  $\dot{\gamma}(s) = r_g f(s)$  with  $\gamma(0) = x_{y_0}$  and  $\gamma(t_{y_0}) = y_0$  so that Proposition A.4 implies  $d_{r_g f}(x_{y_0}, y_0, t_{y_0}) = 0$ . In particular, we obtain that for any  $y_0 \in M$ , there is  $t_{y_0} \in (\delta^0, T)$  and  $x_{y_0} \in \omega$  such that  $d_{r_g f}(x_{y_0}, y_0, t_{y_0}) = 0$ .

By uniform continuity of  $d_{r_g f}$  (on the compact set  $M^2 \cap [\delta^0/2, T + \delta^0]$ ) together with the fact that  $\omega \cap (\delta^0, T)$  is open, there exists  $\nu_{y_0} > 0$  so that

$$B_g(x_{y_0}, \nu_{y_0}) \cap [t_{y_0} - \nu_{y_0}, t_{y_0} + \nu_{y_0}] \subset \omega \cap (\delta^0, T),$$

and for any  $y \in B_g(y_0, \nu_{y_0})$ ,  $x \in B_g(x_{y_0}, \nu_{y_0})$  and  $t \in [t_{y_0} - \nu_{y_0}, t_{y_0} + \nu_{y_0}]$ , we have  $d_{r_g f}(x, y, t) \leq \delta$ . By compactness, we can cover  $M$  by  $M = \bigcup_{i \in I} B_g(y_i, \nu_{y_i})$  where  $I$  is finite. Then, for any  $y \in M$ , there is  $i \in I$  such that  $y \in B_g(y_i, \nu_{y_i})$ , and the set

$$U_y := B_g(x_{y_i}, \nu_{y_i}) \cap [t_{y_i} - \nu_{y_i}, t_{y_i} + \nu_{y_i}]$$

satisfies the sought properties.

As a consequence of this lemma together with Corollary 5.2 and Proposition 5.3, we may now deduce a proof of the  $L^1$  observability estimate of Proposition 5.4.

*Proof of Proposition 5.4.* — Without any loss of generality, we can assume  $0 < \delta < s$ . According to Proposition 5.3, it is enough to study the “observability of the kernel”. Using Corollary 5.2, for any  $\delta > 0$ , there exists  $\varepsilon_0$  so that (5.6) holds for all  $(x, y) \in M^2$ ,  $t \in [\delta, \delta^{-1}]$ ,  $0 < \varepsilon \leq \varepsilon_0$ . Proposition 5.3 leads to compare

$$O_T(y) := \int_0^T \int_{x \in \omega} K_\varepsilon(x, y, t) dx dt \quad \text{with} \quad I_s(y) := \int_{x \in M} K_\varepsilon(x, y, s) dx.$$

From (5.6), the fact that  $K_\varepsilon(x, y, t) > 0$  for  $(x, y, t) \in M^2 \cap (0, 1)$ , and  $s > \delta$ , we deduce

$$O_T(y) > \int_{\delta^0}^T \int_{x \in \omega} e^{-(d_{r_g f}(x, y, t) + \delta)/\varepsilon} \quad \text{and} \quad I_s(y) \leq \int_{x \in M} e^{-(d_{r_g f}(x, y, s) - \delta)/\varepsilon},$$

where  $\delta^0 > 0$  is chosen sufficiently small so that  $(M, r_g f, \omega, (\delta^0, T))$  still satisfies (GCC), which is possible since  $T > T_{GCC}(M, r_g f, \omega)$ . Using Lemma 5.6 (where  $U_y$  and  $C_\delta$  are defined), we now have

$$(5.10) \quad O_T(y) > \int_{(x,t) \in 2U_y} e^{(d_{r_g f}(x,y,t) + \delta)/\varepsilon} dt dx > \int_{(x,t) \in 2U_y} e^{2\delta/\varepsilon} dt dx > C_\delta e^{2\delta/\varepsilon}.$$

Also, for any  $s > \delta$ , using that  $d_{r_g f} > 0$  (see Proposition A.4), we have

$$I_s(y) \leq \text{Vol}_g(M) e^{\delta/\varepsilon}.$$

When combined with (5.10), we obtain  $O_T(y) > C e^{3\delta/\varepsilon} I_s(y)$ . By Proposition 5.3, this gives (5.8) which concludes the proof of the proposition (up to changing  $4\delta$  into  $\delta$ ).

**5.3. FROM  $L^1$  TO  $L^2$  OBSERVABILITY ESTIMATES FOR POSITIVE SOLUTIONS.** — In this section, we conclude the proofs of Proposition 1.12 and Theorem 1.3. We first prove the negative result of Proposition 1.12 (uniform observability of positive solutions does not hold for  $T < T_{GCC}(M, r_g f, \omega)$ , with an exponential lower bound of the cost).

*Proof of Proposition 1.12.* — Let us check the first part of the proposition (geometric statement). Since  $(M, r_g f, \bar{\omega}, T)$  does not satisfy (GCC), there is  $y_0 \in \bar{M}$ , so that for all  $t \in [0, T]$ ,  $\phi_t(y_0) \notin \bar{\omega}$ . In particular, for any  $(t, x) \in [0, T] \times \bar{\omega}$ , we have  $\phi_t(x) \neq y_0$ , which implies  $d_{r_g f}(x, y_0, t) > 0$  by Proposition A.4. By compactness of  $[0, T] \times \bar{\omega}$ ,  $\inf_{x \in \bar{\omega}, t \in [0, T]} d_{r_g f}(x, y_0, t) > 0$ . Therefore,  $d_{([0, T], \bar{\omega})} > 0$  as expected.

For the second part, for any  $\delta > 0$ , select  $y_0 \in \bar{M}$  so that

$$\inf_{(t,x) \in [0, T] \times \bar{\omega}} d_{r_g f}(x, y_0, t) \leq d_{([0, T], \bar{\omega})} + \delta/8.$$

By uniform continuity of  $d_{r_g f}(x, y_0, t)$  defined on  $[0, 2T] \times \bar{\omega}$ , we can also find  $\eta > 0$  so that  $\inf_{(t,x) \in [0, T+\eta] \times \bar{\omega}} d_{r_g f}(x, y_0, t) > d_{([0, T], \bar{\omega})} - \delta/4$  and  $d_{r_g f}(x, y_0, \eta) \leq \delta/4$  for  $x \in B(y_0, \eta)$ . We take as initial datum  $u_0(x) = K_\varepsilon(x, y_0, \eta)$ , yielding  $u(t, x) = K_\varepsilon(x, y_0, t + \eta)$  (see the definition of  $K_\varepsilon$  in (5.4)) as the associated solution of (1.6). We have  $u_0 \in L^2(M)$  together with a lower bound coming from (5.6) with a sufficiently small  $\delta$  (replaced by  $\delta/4$ )

$$k u_0 k_{L^2(M)} > k u_0 k_{L^2(B(y_0, \eta))} > C(\eta) e^{-\delta/2\varepsilon}.$$

Concerning the observation term, we deduce from the upper bound in (5.6) that for  $\varepsilon$  small enough,

$$\begin{aligned} \int_0^T \int_{\bar{\omega}} j u(t, x)^2 dx dt &= \int_0^T \int_{\bar{\omega}} K_\varepsilon(x, y_0, t + \eta)^2 dx dt \\ &\leq \text{Vol}_g(\bar{\omega}) T \sup_{x \in \bar{\omega}, t \in [0, T+\eta]} e^{-2(d_{r_g f}(x, y_0, t) - \delta/4)/\varepsilon} \\ &\leq C e^{-2(d_{([0, T], \bar{\omega})} - \delta/2)/\varepsilon}. \end{aligned}$$

Applying the observability inequality (1.23) to  $u$  thus implies that

$$C_0^+(T, \varepsilon)^2 e^{-2(d_{([0, T], \mathbb{R})} - \delta/4)/\varepsilon} > C,$$

uniformly in  $\varepsilon \in (0, \varepsilon_0]$ , which concludes the proof of the proposition.

To conclude the proof of Theorem 1.3, we need the following dissipation result taken from Guerrero-Lebeau [GL07]. In that reference, it is written on an open subset  $\Omega \subset \mathbb{R}^n$  with the flat metric; however, it can be checked that the result also applies to the case of a Riemannian manifold  $(M, g)$  without boundary and with an additional potential  $q$ .

**PROPOSITION 5.7** (Proposition 3 of [GL07],  $m = 1$ ). — *Assume that  $\partial M = \emptyset$  and that  $(M, X, \omega, T)$  satisfies (GCC). Then, there exist  $C, C_0 > 0$  such that*

$$(5.11) \quad \|ku(T)\|_{L^2(M)}^2 \leq C \left( \int_0^T \int_{\omega} |ju|^2 dt dx + e^{-C_0/\varepsilon} \|ku(0)\|_{L^2(M)}^2 \right)$$

for all  $\varepsilon \in (0, 1]$  and any (not necessarily positive) solution  $u$  to (1.3) (and a fortiori for all solutions  $u$  to (2.6)). The same statement holds true if  $\partial M \neq \emptyset$  and  $(M, X, \omega, T)$  satisfies (FC).

We shall also need the following lemma in the proof of Theorem 1.3.

**LEMMA 5.8.** — *For any  $\delta^0 > \delta > 0$ , there exists  $\varepsilon_0 > 0$  so that*

$$(5.12) \quad \|ku(\delta^0)\|_{L^2(M)} \leq C \|ku(\delta^0)\|_{L^1(M)} \leq C e^{\delta/\varepsilon} \|ku(\delta)\|_{L^1(M)}$$

for any solution  $u$  to (2.6) and  $0 < \varepsilon \leq \varepsilon_0$ .

*Proof.* — Since the manifold is compact, we only need to prove the  $L^1$  bound which follows from the bound  $\|kK_\varepsilon(\cdot, \cdot, \delta^0 - \delta)\|_{L^1(M; M)} \leq C e^{\delta/\varepsilon}$  on the kernel. This estimate follows from Corollary 5.2 (e.g. (5.6) together with the fact that  $d_{r_g f} > 0$ , see Proposition A.4).

*Proof of Theorem 1.3.* — Inequality (1.25) of Proposition 1.12 directly yields

$$T_{\text{unif}}^+(\omega) > T_{\text{GCC}}(M, r_g f, \omega).$$

Note that it was mostly proved in Guerrero-Lebeau [GL07, Th. 1] since one can check that the counterexample they build is a nonnegative solution.

Now, we prove  $T_{\text{unif}}^+(\omega) \leq T_{\text{GCC}}(M, r_g f, \omega)$ . For any  $\delta > 0$  (we will later need  $2\delta < C_0$  where  $C_0$  is the constant in (5.11)), and for  $T > T_{\text{GCC}}(M, r_g f, \omega) + 2\delta$ , we prove the observability inequality for positive solution

$$(5.13) \quad \|ku(T)\|_{L^2(M)}^2 \leq C \int_0^T \int_{\omega} |ju|^2 dt dx.$$

The combination of (5.11) on the time interval  $(2\delta, T)$  together with (5.12) on the time interval  $(\delta, 2\delta)$  implies

$$(5.14) \quad \|ku(T)\|_{L^2(M)}^2 \leq C \left( \int_{2\delta}^T \int_{\omega} u^2 dt dx + e^{-C_0 \delta/\varepsilon} \|ku(\delta)\|_{L^1(M)}^2 \right).$$

Now, applying (5.8) with  $\lambda > 0$  such that  $\lambda T = \delta$ , we obtain

$$(5.15) \quad ku(\delta)k_{L^1(\mathbb{M})} \leq e^{\delta/\varepsilon} \int_0^T \int_{\omega} u \, dt \, dx.$$

Combining (5.14), (5.15) together with the Hölder inequality, we deduce

$$ku(T)k_{L^2(\mathbb{M})} \leq (C + Ce^{-(C_0 - 2\delta)/\varepsilon}) \int_0^T \int_{\omega} u^2 \, dt \, dx.$$

Choosing  $\delta \geq (0, C_0/2)$  implies (5.13) uniformly for  $\varepsilon \geq (0, \varepsilon_0(\delta)]$ , and hence concludes the proof of the theorem.

#### 5.4. FROM OBSERVABILITY OF POSITIVE SOLUTIONS TO A CONTROLLABILITY STATEMENT

This section is devoted to the proof of the controllability result of Corollary 2.4 from the observability of positive solutions. It relies on the following lemma. The result and its proof follow [LB20, Th. 4.1].

**LEMMA 5.9.** — *Let  $V$  be a closed convex set of  $L^2(\mathbb{M})$  with  $0 \in V$  and  $\tilde{V} = V$  so that  $\tilde{V} = V = v$  for any  $v \in V$ . Assume moreover that there exists  $C_V > 0$  so that*

$$(5.16) \quad C_V(T, \varepsilon)^2 \int_0^T \int_{\omega} |ju(t)|^2 ds(x) dt > ku(T)k_{L^2(\mathbb{M})}^2,$$

*for all  $u_0 \in V$  and  $u$  solution of (1.3).*

*Then, for any  $y_0 \in L^2(\mathbb{M})$  and  $0 < \varepsilon \leq \varepsilon_0$ , there exists a control*

$$h \in L^2([0, T], L^2(\mathbb{M}))$$

*with*

$$khk_{L^2([0, T], L^2(\mathbb{M}))} \leq C_V(T, \varepsilon) ky_0k_{L^2(\mathbb{M})},$$

*so that the solution of (2.1) satisfies  $(y(T), u_0)_{L^2(\mathbb{M})} > 0$  for any  $u_0 \in \tilde{V}$ .*

For the proof of Corollary 2.4, we apply this lemma to the sets  $\tilde{V} = V = L^2(\mathbb{M}; \mathbb{R}^+)$ . Notice that Lemma 5.9 also contains one implication (namely Observability  $\Rightarrow$  Controllability) in Corollary 2.3 when applied to  $\tilde{V} = V = L^2(\mathbb{M}; \mathbb{R})$ .

*Proof.* — For any  $\alpha > 0$ , we consider the functional  $J_\alpha$  defined for any  $u_0 \in V$  by

$$J_\alpha(u_0) = \frac{1}{2} \int_0^T \int_{\omega} |ju(t, x)|^2 dt \, dx + \frac{\alpha}{2} ku_0k_{L^2(\mathbb{M})}^2 + (u(T), y_0)_{L^2(\mathbb{M})},$$

where  $u$  is the solution of (1.3). The functional  $J_\alpha$  is continuous, convex and coercive. Therefore,  $J_\alpha$  admits a minimum  $u_{0, \alpha} \in V$  (see e.g. [ET74, Chap. II, Prop. 1.2]). The minimality condition gives (see e.g. [ET74, Chap. II, Proposition 2.1]) for any  $p_0 \in L^2(\mathbb{M})$  that can be written  $p_0 = v_0 - u_{0, \alpha}$ , with  $v_0 \in V$ ,

$$(5.17) \quad \int_0^T \int_{\omega} p(t, x)u_\alpha(t, x) dt \, dx + \alpha (p_0, u_{0, \alpha})_{L^2(\mathbb{M})} + (p(T), y_0)_{L^2(\mathbb{M})} > 0,$$

where  $p$  (resp.  $u_\alpha$ ) denotes the solution of (1.3) with  $p(0) = p_0$  (resp.  $u(0) = u_{0, \alpha}$ ).

Now, let  $y_\alpha$  be the solution of (2.1) with control function  $h_\alpha(t, x) = u_\alpha(T - t, x)$  and initial datum  $y_\alpha(0) = y_0$ . The duality equation (2.2) gives for any  $p_0 \in L^2(M)$  initial datum for  $p$  solution of (1.3)

$$\int_0^T \int_\omega p(t, x) u_\alpha(t, x) dt dx = (p_0, y_\alpha(T))_{L^2(M)} - (p(T), y_0)_{L^2(M)}.$$

Combined with (5.17), this implies

$$(5.18) \quad (p_0, y_\alpha(T))_{L^2(M)} + \alpha (p_0, u_{0,\alpha})_{L^2(M)} > 0,$$

for every  $p_0 \in \mathcal{V} \cap u_{0,\alpha}$ . This also holds for any  $p_0 \in \tilde{\mathcal{V}}$  since  $\tilde{\mathcal{V}} \cap \mathcal{V} = u_{0,\alpha}$  by assumption.

To obtain an estimate of the control, we apply (5.17) to  $p_0 = 0 \in u_{0,\alpha}$ . After an application of the Cauchy-Schwarz inequality, we have

$$\int_0^T \int_\omega j u_\alpha(t, x)^2 dt dx + \alpha k u_{0,\alpha} k_{L^2(M)}^2 \leq k u_\alpha(T) k_{L^2(M)} k y_0 k_{L^2(M)}.$$

The observability inequality (5.16) applies to  $u_{0,\alpha} \in \mathcal{V}$ , so that

$$\int_0^T \int_\omega j u_\alpha(t, x)^2 dt dx + \alpha k u_{0,\alpha} k_{L^2(M)}^2 \leq C_V(T, \varepsilon) \left( \int_0^T \int_\omega j u_\alpha(t, x)^2 dt dx \right)^{1/2} k y_0 k_{L^2(M)}.$$

We obtain successively

$$(5.19) \quad \int_0^T \int_\omega j u_\alpha(t, x)^2 dt dx \leq C_V(T, \varepsilon)^2 k y_0 k_{L^2(M)}^2,$$

$$(5.20) \quad \alpha k u_{0,\alpha} k_{L^2(M)}^2 \leq C_V(T, \varepsilon)^2 k y_0 k_{L^2(M)}^2.$$

We obtain that  $1_\omega u_\alpha$  is bounded in  $L^2([0, T] \times M)$  uniformly in  $\alpha > 0$ . Take a sequence  $\alpha_n \searrow 0$  so that  $1_\omega u_{\alpha_n} \rightharpoonup 1_\omega u$  in  $L^2([0, T] \times M)$ . The associated solutions  $y_{\alpha_n}$  with control  $1_\omega u_{\alpha_n}(T - t, x)$  is therefore bounded in  $L^1([0, T], L^2(M))$  and, again up to a subsequence, converges weakly- to a solution  $y$  of (2.1) with control  $h(t, x) = 1_\omega u(T - t, x)$  and initial datum  $y_0$ . Moreover, up to a subsequence, we can impose  $y_{\alpha_n}(T) \rightharpoonup y(T)$  in  $L^2(M)$ . Passing to the limit in (5.18) using (5.20), we finally obtain

$$(p_0, y(T))_{L^2(M)} > 0$$

for any  $p_0 \in \tilde{\mathcal{V}}$ . We finally get the expected estimate on  $h(t, x) = u(T - t, x)$  passing to the limit in (5.19).

We may now conclude the proof of Corollary 2.4 from Lemma 5.9.

*Proof of Corollary 2.4.* — We apply Lemma 5.9, with  $\tilde{\mathcal{V}} = \mathcal{V} = L^2(M; \mathbb{R}^+)$ . Note that the Lemma applies because for any  $v \in L^2(M; \mathbb{R}^+)$ ,  $L^2(M; \mathbb{R}^+) \cap L^2(M; \mathbb{R}^+) = v$ . Indeed, any  $u \in L^2(M; \mathbb{R}^+)$  can be written  $u = (u + v) - v \in L^2(M; \mathbb{R}^+) - v$  since  $u + v > 0$ . This gives a control  $h$  with the expected uniform bound and so that  $(y(T), u_0)_{L^2(M)} > 0$  for any  $u_0 \in L^2(M; \mathbb{R}^+)$ . This implies  $y(T) > 0$ .

## APPENDIX. ABOUT THE DISTANCES

In this appendix,  $(M, g)$  is a compact Riemannian manifold without boundary.

A.1. A GENERAL LEMMA. — We start with a general lemma.

LEMMA A.1. — Let  $V \geq W^{1,1}(M)$  with nonnegative value. Then, for all  $x, y \in M^2$ , we have

$$\begin{aligned} \frac{1}{2} \inf_{\gamma, t} \left\{ \int_0^t j\dot{\gamma}(s)j_g^2 + V(\gamma(s)) ds ; t > 0, \gamma \in \mathbf{U}_t(x, y) \right\} \\ = \inf_{\gamma, t} \left\{ \int_0^t j\dot{\gamma}(s)j_g \sqrt{V(\gamma(s))} ds ; t > 0, \gamma \in \mathbf{U}_t(x, y) \right\} \\ = \inf_{\gamma} \left\{ \int_0^1 j\dot{\gamma}(s)j_g \sqrt{V(\gamma(s))} ds ; \gamma \in \mathbf{U}_1(x, y) \right\}, \end{aligned}$$

where  $\mathbf{U}_t(x, y) = \{ \gamma \in W^{1,1}([0, t]; M) ; \gamma(0) = x, \gamma(t) = y \}$  for  $t > 0$ .

This lemma is particularly useful for  $V = (V - E)_+$  in which case the (pseudo-)distance defined is the Agmon distance at energy level  $E$ .

*Proof.* — We denote by  $d_1, d_2, d_3$  respectively the three (pseudo-)distances defined in the statement of the lemma. Then, we notice that the last two quantities are invariant by reparametrization, so that  $d_2 = d_3$  after a change of variable in the integral. Then, the inequality  $ab \leq \frac{1}{2}(a^2 + b^2)$  directly yields  $d_2 \leq d_1$ . Let us now prove the converse inequality, namely  $d_3 > d_1$ . For  $\varepsilon > 0$  there exist  $\delta > 0$  and a path  $\gamma : [0, 1] \rightarrow M$  such that

$$\int_0^1 j\dot{\gamma}(s)j_g \sqrt{V(\gamma(s)) + \delta} ds \leq d_3 + \varepsilon.$$

We can further assume  $j\dot{\gamma}(s)j_g > 0$  with the same estimate (indeed, defining a new parametrization  $\zeta$  by  $\gamma(t) = \zeta(\phi(t))$  with  $\phi(t) = \int_0^t j\dot{\gamma}(s)j_g ds$  even yields a Lipschitz reparametrization with constant positive speed, see e.g. [ABB20, Proof of Lem. 3.16]). Using an approximation argument, we can further assume that  $\gamma$  is smooth up to replacing  $\varepsilon$  by  $2\varepsilon$ . We now define the following reparametrization  $\tilde{\gamma}(s) = \gamma(\varphi(s))$ , where  $\varphi$  solves

$$\dot{\varphi}(s) = \frac{\sqrt{V(\gamma(\varphi(s))) + \delta}}{j\dot{\gamma}(\varphi(s))j_g} > \frac{\rho_\delta}{\max_{[0,1]} j\dot{\gamma}j_g} > 0, \quad \varphi(0) = 0,$$

so that  $j\dot{\tilde{\gamma}}(s)j_g = \sqrt{V(\tilde{\gamma}(s)) + \delta}$  for any  $s \in [0, \varphi^{-1}(1)]$ . In particular,

$$j\dot{\tilde{\gamma}}(s)j_g \sqrt{V(\tilde{\gamma}(s)) + \delta} = \frac{1}{2} (j\dot{\tilde{\gamma}}(s)j_g^2 + V(\tilde{\gamma}(s)) + \delta)$$

and

$$\begin{aligned} \frac{1}{2} \int_0^{\varphi^{-1}(1)} (j\dot{\tilde{\gamma}}(s)j_g^2 + V(\tilde{\gamma}(s)) + \delta) ds &= \int_0^{\varphi^{-1}(1)} j\dot{\tilde{\gamma}}(s)j_g \sqrt{V(\tilde{\gamma}(s)) + \delta} ds \\ &= \int_0^1 j\dot{\gamma}(s)j_g \sqrt{V(\gamma(s)) + \delta} ds \leq d_3 + 2\varepsilon, \end{aligned}$$

which gives  $d_1 \leq d_3 + 2\varepsilon$ , and concludes the proof of the lemma.

**A.2. EQUIVALENCE BETWEEN THE TWO DEFINITIONS OF  $d_{r_g f}$ .** — In this section, we prove equivalence between the two definitions of  $d_{r_g f}$ , respectively given in (1.24) in the introduction and in (5.5). We further give an equivalent quantity in terms of the Riemannian distance and the flow  $\phi_t$  of  $r_g f$ . The function  $f$  is assumed to be  $C^3$  throughout the section.

We recall that  $\rho$  is defined in (5.2) with  $V = jr_g f_j^2/4$ , that is to say

$$(A.1) \quad \rho(x, y, t) := \frac{1}{4} \inf \left\{ \int_0^t j\dot{\gamma}(s)j_g^2 + jr_g f(\gamma(s))j_g^2 ds ; \gamma \in \mathbf{U}_t(x, y) \right\},$$

with  $\mathbf{U}_t(x, y) = \{\gamma \in W^{1,1}([0, t]; M) ; \gamma(0) = x, \gamma(t) = y\}$ . Note that it is proved in [LY86, App.] that  $\rho$  is continuous on  $M^2 \times (0, +\infty)$  and, for all  $t > 0$  fixed, Lipschitz continuous as a function of  $(x, y) \in M^2$ . These quantities are related to the Agmon distance but in finite time, see Section A.3 below. Note that the quantity  $\rho$  is symmetric,  $\rho(x, y, t) = \rho(y, x, t)$ , and remains unchanged under the change of  $f$  by  $-f$ . This is not the case for  $d_{r_g f}$ .

**LEMMA A.2.** — *The function  $d_{r_g f}$  defined as*

$$(A.2) \quad d_{r_g f}(x, y, t) := \rho(x, y, t) + \frac{f(x) - f(y)}{2}$$

*is continuous on  $M^2 \times (0, +\infty)$  and, for all  $t > 0$  fixed, Lipschitz continuous as a function of  $(x, y) \in M^2$ . Moreover, we have*

$$(A.3) \quad d_{r_g f}(x, y, t) = \frac{1}{4} \inf \left\{ \int_0^t j\dot{\gamma}(s) + r_g f(\gamma(s))j_g^2 ds ; \gamma \in \mathbf{U}_t(x, y) \right\},$$

$$(A.4) \quad = \frac{1}{4} \inf \left\{ \int_0^t j\dot{\gamma}(s) - r_g f(\gamma(s))j_g^2 ds ; \gamma \in \mathbf{U}_t(x, y) \right\},$$

$$(A.5) \quad = \frac{1}{4} \inf \left\{ \int_0^t j\dot{\gamma}(s)j_{g_s}^2 ds ; \gamma \in W^{1,1}([0, t]; M), \right. \\ \left. \gamma(0) = x, \gamma(t) = \phi_{-t}(y) \right\},$$

where  $\mathbf{U}_t(x, y) = \{\gamma \in W^{1,1}([0, t]; M) ; \gamma(0) = x, \gamma(t) = y\}$ , and  $g_s$  is the time varying metric defined by  $jYj_{g_s} = jD\phi_s(Y)j_g$ .

In particular, for any  $T > 0$ , there exists some constant  $C_T > 0$  so that

$$C_T^{-1}d(x, \phi_{-t}(y)) \leq d_{r_g f}(x, y, t) \leq C_T d(x, \phi_{-t}(y)), \quad \text{for all } t \in [0, T],$$

where  $d$  denotes the Riemannian distance (associated to  $g$ ).

*Proof.* — The continuity property directly follows from that of  $\rho$  proved in [LY86, App.]. To prove (A.3), we remark that for any path  $\gamma$  so that  $\gamma(0) = y, \gamma(t) = x$ , we have

$$f(x) - f(y) = \int_0^t \frac{d}{ds}(f \circ \gamma)(t) dt = \int_0^t r_g f(\gamma(t)) \dot{\gamma}(t) dt.$$

In particular, from the definition of  $d_{r_g f}$  in (A.2) and  $\rho$  in (A.1), we can rewrite  $d_{r_g f}(x, y, t)$  as

$$\begin{aligned} d_{r_g f}(x, y, t) &= \frac{1}{4} \inf \left\{ \int_0^t j\dot{\gamma}(s)^2 + jr_g f(\gamma(s))j_g^2 + 2r_g f(\gamma(s)) \dot{\gamma}(s) ds ; \gamma \geq \mathbf{U}_t(x, y) \right\} \\ &= \frac{1}{4} \inf \left\{ \int_0^t j\dot{\gamma}(s) + r_g f(\gamma(s))j_g^2 ds ; \gamma \geq \mathbf{U}_t(x, y) \right\}. \end{aligned}$$

The statement (A.4) is obtained thanks to the change of path  $\tilde{\gamma}(s) = \gamma(t - s)$ .

Now, we compute  $d_{r_g f}(x, \phi_t(y), t)$  according to formula (A.4). To this aim, let  $\gamma \geq W^{1,1}([0, t]; M)$  so that  $\gamma(0) = x$ ,  $\gamma(t) = \phi_t(y)$ . Let  $\tilde{\gamma}(s) = \phi_{-s}(\gamma(s))$  so that

$$\dot{\gamma}(s) = r_g f(\gamma(s)) + D\phi_s(\dot{\tilde{\gamma}}(s)).$$

In particular,

$$j\dot{\gamma}(s) - r_g f(\gamma(s))j_g^2 = |D\phi_s(\dot{\tilde{\gamma}}(s))|_g^2 = |\dot{\tilde{\gamma}}(s)|_{g_s}^2.$$

It gives (A.5) since any path  $\gamma \geq W^{1,1}([0, t]; M)$  so that  $\gamma(0) = x$ ,  $\gamma(t) = y$  can be written  $\gamma(s) = \phi_s(\tilde{\gamma}(s))$  with  $\tilde{\gamma}(0) = x$ ,  $\tilde{\gamma}(t) = \phi_{-t}(y)$ , and conversely.

**A.3. FURTHER LINKS BETWEEN THE DIFFERENT DISTANCES.** — In this section, we relate the above quantities  $\rho(x, y, t)$ ,  $d_{r_g f}(x, y, t)$  with the Agmon distance to the bottom energy (see 1.12 for  $V = jr_g f(x)j_g^2/4$  and  $E_0 = \min_M V = 0$ ), that is to say

$$(A.6) \quad d_A(x, y) = \frac{1}{2} \inf \left\{ \int_0^1 jr_g f(\gamma(s))j_g \dot{\gamma}(s)j_g ds ; \gamma \geq \mathbf{U}_1(x, y) \right\},$$

with  $\mathbf{U}_1(x, y) = \{\gamma \geq W^{1,1}([0, 1]; M) ; \gamma(0) = x, \gamma(1) = y\}$  and the associated quantity (compare with the definition of  $d_{r_g f}$  in terms of  $\rho$  in (A.2))

$$(A.7) \quad W(x, y) := d_A(x, y) + \frac{f(x) - f(y)}{2}.$$

The results of this section are not explicitly used in the proofs of the main part of the paper; however we believe these links are interesting and enlightening. Indeed, they relate the quantity  $d_A(x, y)$  appearing in all general bounds of Section 3 together with the quantities  $\rho(x, y, t)$ ,  $d_{r_g f}(x, y, t)$  appearing in results of Section 5 concerning positive solutions.

**LEMMA A.3.** — For all  $(x, y) \geq M^2$ , we have

$$(A.8) \quad d_A(x, y) = \inf_{t>0} \rho(x, y, t),$$

$$(A.9) \quad W(x, y) = \inf_{t>0} d_{r_g f}(x, y, t).$$

Moreover, if  $r_g f(y) = 0$ , then we have  $d_A(x, y) = \lim_{t \rightarrow +\infty} \rho(x, y, t)$ .

*Proof.* — Equality in (A.8) is a consequence of Lemma A.1 applied to  $V = jr_g fj_g^2$ . Then, (A.9) is a direct consequence of the expression of  $W$  and  $d_{r_g f}$  in terms of  $d_A$  and  $\rho$  in (A.7), (A.2), together with (A.8).



Finally, if  $r_g f(y) = 0$ , then the function  $t \mapsto \rho(x, y, t)$  is non-increasing. Indeed, taking  $t_1 < t_2$ , from a path  $\gamma_1 : [0, t_1] \rightarrow M$  such that  $\gamma_1(0) = x$  and  $\gamma_1(t_1) = y$ , we can construct the path  $\gamma_2 : [0, t_2] \rightarrow M$  by  $\gamma_2(s) = \gamma_1(s)$  for  $s \in [0, t_1]$  and  $\gamma_2(s) = y$  for  $s \in [t_1, t_2]$ . This yields a path in  $W^{1,1}([0, t_2]; M)$  if  $\gamma_1 \in W^{1,1}([0, t_1]; M)$ , and thus the set of admissible paths on  $[0, t_2]$  is larger than the set of admissible paths on  $[0, t_1]$ . Since the contribution  $\int_{t_1}^{t_2} j\dot{\gamma}_2(s) \! \! \! /_g^2 ds = 0$  we deduce that  $\rho(x, y, t_2) \leq \rho(x, y, t_1)$ . This proves that the inf is actually a  $\lim_{t \rightarrow +\infty}$  in this case.

Note that related properties are proved in the appendix of [HS85]. For instance, [HS85, Lem. A2.2] with our notations can be loosely stated as follows: If  $W(x, y) = 0$ , then every minimizing geodesic of  $d_A$  is a generalized integral curve of  $r_g f$ .

Finally, we state a last result that explains that  $d_{r_g f}(x, y, t)$  measures how far  $x$  is the final state of a path of the vector field at time  $t$  and starting at  $y$ . Part of this result is contained in the last statement of Lemma A.2; we here give a different proof, which, we believe, is interesting in itself.

**PROPOSITION A.4.** — *With  $d_{r_g f}(x, y, t)$  defined in (5.5), we have for all  $(x, y, t) \in M^2 \times (0, +\infty)$ ,*

$$(1) \quad d_{r_g f}(x, y, t) > 0;$$

(2)  $d_{r_g f}(x, y, t) = 0$  if and only if there exists a trajectory of  $\dot{\gamma}(s) = r_g f(\gamma(s))$  with  $\gamma(0) = x, \gamma(t) = y$ , that is if and only if  $y = \phi_t(x)$ .

In particular,  $(M, r_g f, \omega, T)$  satisfies (GCC) if and only if for any  $x \in M$ , there exist  $y \in \omega$  and  $t \in (0, T)$  so that  $d_{r_g f}(x, y, t) = 0$ .

Recall that the flow  $(\phi_t)_{t \in \mathbb{R}}$  is defined in (2.14) and the Geometric Control Condition (GCC) is defined in Definition 2.10.

*Proof.* — Statement 1 follows from the definition of  $d_{r_g f}$  in (A.4). Let us now consider Statement (2). Assume first that there exists a trajectory of  $\dot{\gamma}(s) = r_g f(\gamma(s))$  with  $\gamma(0) = x$  and  $\gamma(t) = y$ . Then, by definition of the infimum in (A.4), this yields  $d_{r_g f}(x, y, t) \leq 0$  and hence  $d_{r_g f}(x, y, t) = 0$ . Conversely, assume  $d_{r_g f}(x, y, t) = 0$ . Take a minimizing sequence in (A.4), that is to say that  $\gamma_n \in W^{1,1}([0, t]; M)$  such that  $\gamma_n(0) = x, \gamma_n(t) = y$  and  $\dot{\gamma}_n \cdot r_g f(\gamma_n) = R_n$  (bounded continuous with values in the tangent bundle of  $M$ ) with

$$(A.10) \quad \int_0^t jR_n(s) \! \! \! /_g^2 ds \rightarrow 0.$$

Since  $r_g f$  is bounded on  $M$ , the sequence  $\int_0^t j\dot{\gamma}_n(s) \! \! \! /_g^2 ds$  is then uniformly bounded in  $\mathbb{R}$ . As a consequence, the sequence of paths  $(\gamma_n)_{n \in \mathbb{N}}$  is equicontinuous. From Ascoli's theorem, we may extract a subsequence (which we do not relabel)  $(\gamma_n)_{n \in \mathbb{N}}$  which converges strongly for the topology  $C^0([0, t]; M)$  to a limit  $\gamma \in C^0([0, t]; M)$ . The latter thus has the same endpoints  $\gamma(0) = x$  and  $\gamma(t) = y$ . It is solution of  $\dot{\gamma} = r_g f(\gamma)$  in

the distributional sense according to (A.10). Bootstrapping in the differential equation implies  $\gamma \in W^{1,1}([0, 1]; M)$  and  $\gamma$  is a strong solution to  $\dot{\gamma} = r_g f(\gamma)$ . This concludes the proof of Statement 2.

According to Lemma 2.11 Item (4), that  $(M, X, \omega, T)$  satisfies (GCC) is equivalent to the fact that for any  $x \in M$ , there exist  $t \in (0, T)$  such that  $y := \phi_t(x) \in \omega$ . As a consequence of Item (2), this is equivalent to having  $d_{r_g f}(x, y, t) = 0$ .

Note that an analogue statement for the function  $W$  in (A.9) is proved in [HS85, Lem. A2.2], and could also be deduced from Proposition A.4.

## REFERENCES

- [ABB20] A. AGRACHEV, D. BARILARI & U. BOSCAIN – *A comprehensive introduction to sub-Riemannian geometry*, Cambridge Studies in Advanced Math., vol. 181, Cambridge University Press, Cambridge, 2020.
- [All98] B. ALLIBERT – “Contrôle analytique de l’équation des ondes et de l’équation de Schrödinger sur des surfaces de révolution”, *Comm. Partial Differential Equations* **23** (1998), no. 9-10, p. 1493–1556.
- [AM19a] Y. AMIRAT & A. MÜNCH – “Asymptotic analysis of an advection-diffusion equation and application to boundary controllability”, *Asymptot. Anal.* **112** (2019), no. 1-2, p. 59–106.
- [AM19b] ———, “On the controllability of an advection-diffusion equation with respect to the diffusion parameter: asymptotic analysis and numerical simulations”, *Acta Math. Appl. Sinica (English Ser.)* **35** (2019), no. 1, p. 54–110.
- [BP20a] J. A. BÁRCENA-PETISCO – “Uniform controllability of a Stokes problem with a transport term in the zero-diffusion limit”, *SIAM J. Control Optim.* **58** (2020), no. 3, p. 1597–1625.
- [BP20b] ———, “Cost of null controllability for parabolic equations with vanishing diffusivity and a transport term”, 2020, HAL: hal-02455632.
- [Bes78] A. L. BESSE – *Manifolds all of whose geodesics are closed*, *Ergeb. Math. Grenzgeb.* (3), vol. 93, Springer-Verlag, Berlin-New York, 1978.
- [Cha43] S. CHANDRESEKHAR – “Stochastic problems in physics and astronomy”, *Rev. Modern Phys.* **15** (1943), p. 1–89.
- [Cha09] M. CHAPOULY – “On the global null controllability of a Navier-Stokes system with Navier slip boundary conditions”, *J. Differential Equations* **247** (2009), p. 2094–2123.
- [Cor96] J.-M. CORON – “On the controllability of the 2-D incompressible Navier-Stokes equations with the Navier slip boundary conditions”, *ESAIM Control Optim. Calc. Var.* **1** (1996), p. 35–75.
- [Cor07] ———, *Control and nonlinearity*, Math. Surveys and Monographs, vol. 136, American Mathematical Society, Providence, RI, 2007.
- [CF96] J.-M. CORON & A. V. FURSIKOV – “Global exact controllability of the 2-D Navier-Stokes equations on a manifold without boundary”, *Russian J. Math. Phys.* **4** (1996), p. 429–448.
- [CG05] J.-M. CORON & S. GUERRERO – “Singular optimal control: A linear 1-D parabolic-hyperbolic example”, *Asymptot. Anal.* **44** (2005), p. 237–257.
- [CMS20] J.-M. CORON, F. MARBACH & F. SUEUR – “Small-time global exact controllability of the Navier-Stokes equation with Navier slip-with-friction boundary conditions”, *J. Eur. Math. Soc. (JEMS)* **22** (2020), no. 5, p. 1625–1673.
- [CFKS87] H. L. CYCON, R. G. FROESE, W. KIRSCH & B. SIMON – *Schrödinger operators with application to quantum mechanics and global geometry*, Texts and Monographs in Physics, Springer-Verlag, Berlin, 1987.
- [Daf00] C. M. DAFERMOS – *Hyperbolic conservation laws in continuum physics*, Springer-Verlag, Berlin, 2000.

- [DR20] N. V. DANG & G. RIVIÈRE – “Pollicott-Ruelle spectrum and Witten Laplacians”, *J. Eur. Math. Soc. (JEMS)* (2020), to appear.
- [DL09] B. DEHMAN & G. LEBEAU – “Analysis of the HUM control operator and exact controllability for semilinear waves in uniform time”, *SIAM J. Control Optim.* **48** (2009), no. 2, p. 521–550.
- [DGLLPN20] G. DI GESÙ, T. LELIÈVRE, D. LE PEUTREC & B. NECTOUX – “The exit from a metastable state: Concentration of the exit point distribution on the low energy saddle points, part 1”, *J. Math. Pures Appl.* (9) **138** (2020), p. 242–306.
- [DS99] M. DIMASSI & J. SJÖSTRAND – *Spectral asymptotics in the semi-classical limit*, London Math. Society Lect. Note Series, vol. 268, Cambridge University Press, Cambridge, 1999.
- [DR77] S. DOLECKI & D. L. RUSSELL – “A general theory of observation and control”, *SIAM J. Control Optim.* **15** (1977), no. 2, p. 185–220.
- [ET74] I. EKELAND & R. TEMAM – *Analyse convexe et problèmes variationnels*, Dunod–Gauthier-Villars, Paris, 1974.
- [Eva98] L. C. EVANS – *Partial differential equations*, Graduate Studies in Math., American Mathematical Society, Providence, RI, 1998.
- [FI96] A. V. FURSIKOV & O. Y. IMANUVILOV – *Controllability of evolution equations*, Lect. Notes Series, vol. 34, Seoul National University Research Institute of Mathematics Global Analysis Research Center, Seoul, 1996.
- [Gla10] O. GLASS – “A complex-analytic approach to the problem of uniform controllability of a transport equation in the vanishing viscosity limit”, *J. Funct. Anal.* **258** (2010), p. 852–868.
- [GG07] O. GLASS & S. GUERRERO – “On the uniform controllability of the Burgers equation”, *SIAM J. Control Optim.* **46** (2007), p. 1211–1238.
- [GG08] ———, “Some exact controllability results for the linear KdV equation and uniform controllability in the zero-dispersion limit”, *Asymptot. Anal.* **60** (2008), p. 61–100.
- [GG09] ———, “Uniform controllability of a transport equation in zero dispersion-dispersion limit”, *Math. Models Methods Appl. Sci.* **19** (2009), p. 1567–1601.
- [GL07] S. GUERRERO & G. LEBEAU – “Singular optimal control for a transport-dispersion equation”, *Comm. Partial Differential Equations* **32** (2007), p. 1813–1836.
- [HKL15] D. HAN-KWAN & M. LÉAUTAUD – “Geometric analysis of the linear Boltzmann equation I. Trend to equilibrium”, *Ann. PDE* **1** (2015), no. 1, article no. 3 (84 pages).
- [Hel88] B. HELFFER – *Semi-classical analysis for the Schrödinger operator and applications*, Lect. Notes in Math., vol. 1336, Springer-Verlag, Berlin, 1988.
- [HKN04] B. HELFFER, M. KLEIN & F. NIER – “Quantitative analysis of metastability in reversible dispersion processes via a Witten complex approach”, *Mat. Contemp.* **26** (2004), p. 41–85.
- [HN06] B. HELFFER & F. NIER – *Quantitative analysis of metastability in reversible dispersion processes via a Witten complex approach: the case with boundary*, Mém. Soc. Math. France (N.S.), vol. 105, Société Mathématique de France, Paris, 2006.
- [HS84] B. HELFFER & J. SJÖSTRAND – “Multiple wells in the semiclassical limit. I”, *Comm. Partial Differential Equations* **9** (1984), no. 4, p. 337–408.
- [HS85] ———, “Puits multiples en mécanique semi-classique IV. Étude du complexe de Witten”, *Comm. Partial Differential Equations* **10** (1985), no. 3, p. 245–340.
- [Kru70] S. N. KRUŽKOV – “First order quasilinear equations with several independent variables. (russian)”, *Mat. Sb. (N.S.)* **81** (1970), p. 228–255.
- [LSU68] O. A. LADYŽENSKAJA, V. A. SOLONNIKOV & N. N. URAL’CEVA – *Linear and quasilinear equations of parabolic type*, Translations of Math. Monographs, vol. 23, American Mathematical Society, Providence, R.I., 1968.
- [LL16] C. LAURENT & M. LÉAUTAUD – “Uniform observability estimates for linear waves”, *ESAIM Control Optim. Calc. Var.* **22** (2016), no. 4, p. 1097–1136.
- [LL21a] ———, “Observability of the heat equation, geometric constants in control theory, and a conjecture of Luc Miller”, *Anal. PDE* (2021), to appear.

- [LL21b] ———, “On uniform controllability of 1D transport equations in the vanishing viscosity limit”, work in progress, 2021.
- [LL21c] ———, “The cost function for the approximate control of waves”, work in progress, 2021.
- [LB20] K. LE BALC’H – “Global null-controllability and nonnegative-controllability of slightly superlinear heat equations”, *J. Math. Pures Appl. (9)* **135** (2020), p. 103–139.
- [LP10] D. LE PEUTREC – “Small eigenvalues of the Neumann realization of the semiclassical Witten Laplacian”, *Ann. Fac. Sci. Toulouse Math. (6)* **19** (2010), no. 3-4, p. 735–809.
- [Léa10] M. LÉAUTAUD – “Spectral inequalities for non-selfadjoint elliptic operators and application to the null-controllability of parabolic systems”, *J. Funct. Anal.* **258** (2010), p. 2739–2778.
- [Léa12] ———, “Uniform controllability of scalar conservation laws in the vanishing viscosity limit”, *SIAM J. Control Optim.* **50** (2012), no. 3, p. 1661–1699.
- [Léa18] ———, “Sur quelques questions de prolongement unique, de propagation et de contrôle”, Habilitation à diriger des recherches, Université Paris Diderot, 2018, <http://leautaud.perso.math.cnrs.fr/files/HdR.pdf>.
- [LR95] G. LEBEAU & L. ROBBIANO – “Contrôle exact de l’équation de la chaleur”, *Comm. Partial Differential Equations* **20** (1995), p. 335–356.
- [Lee13] J. M. LEE – *Introduction to smooth manifolds*, 2nd ed., Graduate Texts in Math., vol. 218, Springer, New York, 2013.
- [LY86] P. LI & S.-T. YAU – “On the parabolic kernel of the Schrödinger operator”, *Acta Math.* **156** (1986), no. 3-4, p. 153–201.
- [Lio88] J.-L. LIONS – *Contrôlabilité exacte, perturbations et stabilisation de systèmes distribués 2. Perturbations*, Recherches en Math. Appliquées, vol. 9, Masson, Paris, 1988.
- [Lis12] P. LISSY – “A link between the cost of fast controls for the 1-D heat equation and the uniform controllability of a 1-D transport-diffusion equation”, *Comptes Rendus Mathématique* **350** (2012), no. 11-12, p. 591–595.
- [Lis14] ———, “An application of a conjecture due to Ervedoza and Zuazua concerning the observability of the heat equation in small time to a conjecture due to Coron and Guerrero concerning the uniform controllability of a convection-diffusion equation in the vanishing viscosity limit”, *Systems Control Lett.* **69** (2014), p. 98–102.
- [Lis15] ———, “Explicit lower bounds for the cost of fast controls for some 1-D parabolic or dispersive equations, and a new lower bound concerning the uniform controllability of the 1-D transport-diffusion equation”, *J. Differential Equations* **259** (2015), no. 10, p. 5331–5352.
- [Mar14] F. MARBACH – “Small time global null controllability for a viscous Burgers’ equation despite the presence of a boundary layer”, *J. Math. Pures Appl. (9)* **102** (2014), no. 2, p. 364–384.
- [Mic19] L. MICHEL – “About small eigenvalues of the Witten Laplacian”, *Pure Appl. Anal.* **1** (2019), no. 2, p. 149–206.
- [Mil04] L. MILLER – “Geometric bounds on the growth rate of null-controllability cost for the heat equation in small time”, *J. Differential Equations* **204** (2004), no. 1, p. 202–226.
- [Mün18] A. MÜNCH – “Numerical estimations of the cost of boundary controls for the equation  $y_t - \varepsilon y_{xx} + M y_x = 0$  with respect to  $\varepsilon$ ”, in *Recent advances in PDEs: analysis, numerics and control*, SEMA SIMAI Springer Ser., vol. 17, Springer, Cham, 2018, p. 159–191.
- [RS80] M. REED & B. SIMON – *Methods of modern mathematical physics I. Functional analysis*, 2nd ed., Academic Press, Inc., New York, 1980.
- [SM79] Z. SCHUSS & B. J. MATKOWSKY – “The exit problem: a new approach to diffusion across potential barriers”, *SIAM J. Appl. Math.* **36** (1979), no. 3, p. 604–623.
- [Sim82] B. SIMON – “Schrödinger semigroups”, *Bull. Amer. Math. Soc. (N.S.)* **7** (1982), no. 3, p. 447–526.
- [Sim83] ———, “Instantons, double wells and large deviations”, *Bull. Amer. Math. Soc. (N.S.)* **8** (1983), no. 2, p. 323–326.

- [Var67] S. R. S. VARADHAN – “Diffusion processes in a small time interval”, *Comm. Pure Appl. Math.* **20** (1967), p. 659–685.
- [Wit82] E. WITTEN – “Supersymmetry and Morse theory”, *J. Differential Geometry* **17** (1982), no. 4, p. 661–692 (1983).

Manuscript received 3rd March 2020

accepted 1st February 2021

CAMILLE LAURENT, CNRS UMR 7598 and Sorbonne Universités UPMC Univ Paris 06, Laboratoire Jacques-Louis Lions

F-75005, Paris, France

*E-mail* : laurent@ann.jussieu.fr

*Url* : <https://www.ljll.math.upmc.fr/~laurent/>

MATHIEU LÉAUTAUD, Laboratoire de Mathématiques d’Orsay, Université Paris-Sud, CNRS, Université Paris-Saclay

Bâtiment 307, 91405 Orsay Cedex France

*E-mail* : matthieu.leautaud@math.u-psud.fr

*Url* : <http://leautaud.perso.math.cnrs.fr/>