



Journal de l'École polytechnique

Mathématiques

Frédéric DÉGLISE, Jean FASEL, Fangzhou JIN, & Adeel A. KHAN

On the rational motivic homotopy category

Tome 8 (2021), p. 533-583.

http://jep.centre-mersenne.org/item/JEP_2021__8__533_0

© Les auteurs, 2021.

Certains droits réservés.



Cet article est mis à disposition selon les termes de la licence
LICENCE INTERNATIONALE D'ATTRIBUTION CREATIVE COMMONS BY 4.0.
<https://creativecommons.org/licenses/by/4.0/>

L'accès aux articles de la revue « Journal de l'École polytechnique — Mathématiques » (<http://jep.centre-mersenne.org/>), implique l'accord avec les conditions générales d'utilisation (<http://jep.centre-mersenne.org/legal/>).

Publié avec le soutien
du Centre National de la Recherche Scientifique



Publication membre du
Centre Mersenne pour l'édition scientifique ouverte
www.centre-mersenne.org

ON THE RATIONAL MOTIVIC HOMOTOPY CATEGORY

BY FRÉDÉRIC DÉGLISE, JEAN FASEL, FANGZHOU JIN
& ADEEL A. KHAN

ABSTRACT. — We study the structure of the rational motivic stable homotopy category over general base schemes. Our first class of results concerns the six operations: we prove absolute purity, stability of constructible objects, and Grothendieck–Verdier duality for $\mathrm{SH}_{\mathbb{Q}}$. Next, we prove that $\mathrm{SH}_{\mathbb{Q}}$ is canonically SL-oriented; we compare $\mathrm{SH}_{\mathbb{Q}}$ with the category of rational Milnor–Witt motives; and we relate the rational bivariant A^1 -theory to Chow–Witt groups. These results are derived from analogous statements for the minus part of $\mathrm{SH}[1/2]$.

RÉSUMÉ (Sur la catégorie A^1 -homotopique rationnelle). — Dans ce travail, nous étudions la structure de la catégorie A^1 -homotopique stable rationnelle sur une base arbitraire. Notre première famille de résultats concerne les six opérations : nous prouvons la pureté absolue, la stabilité des objets constructibles et la dualité de Grothendieck-Verdier pour cette catégorie. Dans un deuxième temps, nous prouvons que la catégorie A^1 -homotopique stable rationnelle est canoniquement SL-orientée et la comparons à la catégorie des motifs rationnels de Milnor-Witt. Cela permet de calculer les groupes d' A^1 -homotopie stable bivariants en termes des groupes de Chow-Witt supérieurs. Ces résultats s'obtiennent à partir d'énoncés analogues pour la partie négative de la catégorie A^1 -homotopique stable 2-localisée.

CONTENTS

1. Introduction.....	534
2. The key lemma.....	543
3. Duality and absolute purity.....	545
4. The dimensional homotopy t -structure.....	547
5. The minus A^1 -derived category and Witt motives.....	549
6. Rational stable homotopy and Milnor–Witt motives.....	554
7. SL-Orientations.....	555
8. Bivariant A^1 -theory and Chow–Witt groups.....	559

MATHEMATICAL SUBJECT CLASSIFICATION (2020). — 14F42, 19E15, 19G12, 11E81, 14C25, 14C35.

KEYWORDS. — Motivic homotopy, motivic cohomology, six operations, Chow–Witt groups, K-theory, hermitian K-theory.

F.D. received support from the French “Investissements d’Avenir” program, project ISITE-BFC (contract ANR-IS-IDEX-OOOB). F.J. is partially supported by the DFG Priority Programme SPP 1786 Project “Motivic filtrations over Dedekind domains” and Marc Levine’s ERC Advanced Grant QUADAG. A.K. was partially supported by SFB 1085 Higher Invariants, Universität Regensburg, and from the Simons Collaboration on Homological Mirror Symmetry.

Appendix A. Continuity in motivic ∞ -categories.....	564
Appendix B. Essentially of finite presentation morphisms.....	566
Appendix C. Absolute purity in equal characteristic.....	573
Appendix D. Absolute purity in hermitian K-theory.....	577
Appendix E. 2-primary unramified Witt ring.....	578
References.....	580

1. INTRODUCTION

SUMMARY OF RESULTS. — Let S be a scheme and $\mathrm{SH}(S)$ the stable motivic homotopy category. This paper is dedicated to the study of its rationalization $\mathrm{SH}(S)_{\mathbb{Q}}$.

The first relevant observation, due to Morel, is that after inverting 2, $\mathrm{SH}(S)$ splits into two parts:

$$\mathrm{SH}(S)[1/2] \simeq \mathrm{SH}(S)_+ \times \mathrm{SH}(S)_-.$$

In particular, one has a decomposition $\mathrm{SH}(S)_{\mathbb{Q}} \simeq \mathrm{SH}(S)_{\mathbb{Q},+} \times \mathrm{SH}(S)_{\mathbb{Q},-}$. The plus part is very well understood after the work of Cisinski–Déglise [CD19]:

THEOREM A. — *The motivic ∞ -category⁽¹⁾ $\mathrm{SH}_{\mathbb{Q},+}$ satisfies the following properties:*

(I) **Orientation.** *It admits a canonical orientation. In particular, there are canonical isomorphisms*

$$\mathrm{Th}_X(v) \simeq \mathbb{1}_X\langle n \rangle,$$

for every virtual vector bundle v of rank n over a scheme X . Here we write $\langle n \rangle = (n)[2n]$ (Tate twist by n and shift by $2n$) for an integer n .

(II) **Absolute purity.** *For any smoothable⁽²⁾ morphism $f : X \rightarrow S$ of regular noetherian schemes, there is a canonical isomorphism*

$$\mathbb{1}_X\langle d \rangle \simeq f^!(\mathbb{1}_S)$$

in $\mathrm{SH}(X)_{\mathbb{Q},+}$, where $d = \mathrm{rk}(T_{X/S})$ is the relative virtual dimension of f .

(III) **Finiteness.** *Over quasi-excellent schemes, each of the six operations preserves constructible objects.*

(IV) **Duality.** *For every separated morphism of finite type $f : X \rightarrow S$ with S quasi-excellent and regular, $f^!(\mathbb{1}_S)$ is a dualizing object of $\mathrm{SH}(X)_{\mathbb{Q},+}$.*

(V) **Comparison.** *There is a canonical equivalence of symmetric monoidal motivic ∞ -categories*

$$\mathrm{SH}_{\mathbb{Q},+} \simeq \mathbf{H}_M\mathbb{Q}\text{-mod},$$

where $\mathbf{H}_M\mathbb{Q}$ is the rational motivic cohomology spectrum.

⁽¹⁾This is a refinement of the notion of motivic triangulated category from [CD19]; see [Kha16, Kha21].

⁽²⁾A morphism is called *smoothable* if it admits a global factorization through a closed immersion and a smooth morphism.

(VI) Bivariant theory. Let S be a regular noetherian scheme equipped with the dimension function $\delta = -\text{codim}_S$. Then for every S -scheme X of finite type, there are canonical isomorphisms

$$\mathbf{H}_{2n,n}^{A^1}(X/S, \mathbb{Q})_+ := [\mathbb{1}_X \langle n \rangle, f^!(\mathbb{1}_S)]_{\text{SH}(X)_{\mathbb{Q},+}} \simeq \text{CH}_{\delta=n}(X) \otimes_{\mathbb{Z}} \mathbb{Q}$$

for every integer n , where $f : X \rightarrow S$ is the structural map and $\text{CH}_{\delta=n}(X)$ is the Chow group of algebraic cycles on X of δ -dimension n .⁽³⁾

Our goal in this paper is to demonstrate analogous properties for the minus part $\text{SH}_{\mathbb{Q},-}$ (and hence for $\text{SH}_{\mathbb{Q}}$ itself). In fact, we are able to prove these just after $Z[1/2]$ -linearization (i.e., for the minus part of the A^1 -derived category):

THEOREM B. — Consider the minus part of the A^1 -derived category, i.e., the $Z[1/2]$ -linear motivic ∞ -category

$$\mathbf{D}_{A^1,-} = \text{SH}_- \otimes_{\text{Spt}} \mathbf{D}(Z) \simeq (\text{SH} \otimes_{\text{Spt}} \mathbf{D}(Z))[1/2]_-,$$

where Spt is the ∞ -category of spectra, $\mathbf{D}(Z)$ the derived ∞ -category of Z -modules, and the tensor product is in the sense of [Lur12, §4.8]. More generally, let $\mathbf{D}_{A^1,-}(-, \Lambda) := \text{SH}_- \otimes_{\text{Spt}} \mathbf{D}(\Lambda)$ for any commutative ring Λ . Then $\mathbf{D}_{A^1,-}(-, \Lambda)$ satisfies the following properties:

(I) Weak orientation. The motivic ∞ -category $\mathbf{D}_{A^1,-}(-, \Lambda)$ admits a canonical SL-orientation. In particular, there are canonical isomorphisms

$$\text{Th}_X(v) \simeq \text{Tw}_X(\widetilde{\det}(v)),$$

for every virtual vector bundle v over a scheme X . Here $\widetilde{\det}(v)$ denotes the graded determinant, and for a graded line bundle (L, n) , we write $\text{Tw}_X(L, n) = \text{Th}_X(L) \langle n - 1 \rangle$.

(II) Absolute purity. For any smoothable morphism $f : X \rightarrow S$ of regular noetherian schemes, there is a canonical isomorphism

$$\text{Tw}_X(\widetilde{\omega}_{X/S}) \simeq f^!(\mathbb{1}_S)$$

in $\text{SH}(X)_-$, where $\widetilde{\omega}_{X/S}$ is the graded determinant of the virtual tangent bundle $T_{X/S}$ (see 7.10).

(III) Finiteness. Over quasi-excellent schemes, each of the six operations preserves constructible objects.

(IV) Duality. Let S be a quasi-excellent regular scheme and $K_S \in \text{SH}(S)_-$ a \otimes -invertible object. Then for every separated morphism of finite type $f : X \rightarrow S$, $f^!(K_S)$ is a dualizing object of $\text{SH}(X)_-$.

(V) Comparison. Assume Λ is a subring of \mathbb{Q} with $2 \in \Lambda^\times$. There is a canonical equivalence of Λ -linear symmetric monoidal motivic ∞ -categories

$$\mathbf{D}_{A^1,-}(-, \Lambda) \simeq \mathbf{H}_W \Lambda\text{-mod},$$

where $\mathbf{H}_W \Lambda$ is the Λ -linear homotopical Witt spectrum.

⁽³⁾Note that the dimension convention here differs from the usual one for Chow groups (see for example [Ful98]), as the integer δ may be negative. For more details about our convention, see Definition 8.5 or [BD17, Ex. 3.1.6].

(VI) Bivariant theory. Assume Λ is a subring of \mathbb{Q} with $2 \in \Lambda^\times$. Let S be a regular noetherian scheme equipped with the dimension function $\delta = -\text{codim}_S$. For every S -scheme X of finite type, there are canonical isomorphisms

$$H_{2n,n}^{A^1}(X/S, \Lambda_-) := [\mathbb{1}_X \langle n \rangle, f^!(\mathbb{1}_S)]_{D_{A^1,-}(X,\Lambda)} \simeq \widetilde{CH}_{\delta=n}(X) \otimes_{\mathbb{Z}} \Lambda$$

for every integer n , where $f : X \rightarrow S$ is the structural map and $\widetilde{CH}_{\delta=n}(X)$ is the Chow–Witt group of quadratic cycles on X of δ -dimension n .

In fact, the motivic ∞ -category SH_- also satisfies property (II), when formulated correctly, as well as properties (III) and (IV).

Combining Theorems A and B, we get:

COROLLARY C. — *The motivic ∞ -category $\text{SH}_{\mathbb{Q}}$ satisfies the following properties (where (I)–(IV) are formulated exactly as in Theorem B):*

- (I) Weak orientation.
- (II) Absolute purity.
- (III) Finiteness.
- (IV) Duality.
- (V) Comparison. *There is a canonical equivalence of \mathbb{Q} -linear symmetric monoidal motivic ∞ -categories*

$$\text{SH}_{\mathbb{Q}} \simeq \mathbf{H}_{\text{MW}}\mathbb{Q}\text{-mod},$$

where $\mathbf{H}_{\text{MW}}\mathbb{Q}$ is the rational Milnor–Witt motivic cohomology spectrum.

(VI) Bivariant theory. Let S be a regular noetherian scheme equipped with the dimension function $\delta = -\text{codim}_S$. Then for every S -scheme X of finite type, there are canonical isomorphisms

$$H_{2n,n}^{A^1}(X/S, \mathbb{Q}) := [\mathbb{1}_X \langle n \rangle, f^!(\mathbb{1}_S)]_{\text{SH}(X)_{\mathbb{Q}}} \simeq \widetilde{CH}_{\delta=n}(X) \otimes_{\mathbb{Z}} \mathbb{Q}$$

for every integer n , where $f : X \rightarrow S$ is the structural map and $\widetilde{CH}_{\delta=n}(X)$ is the Chow–Witt group of quadratic cycles on X of δ -dimension n .

We now discuss each of the properties (I)–(VI) in turn.

(I) **WEAK ORIENTATION.** — Orientations are the main point of difference we encounter from the classical case of $\text{SH}_{\mathbb{Q},+}$. In fact, $D_{A^1,-}$ and $\text{SH}_{\mathbb{Q}}$ do not admit (GL-)orientations in the sense of [CD19, Def. 2.4.38].

Before explaining the weaker notion of SL-orientation, we briefly recall the classical notion. For any motivic ∞ -category \mathcal{T} , there is a canonical map of presheaves of groupoids

$$\text{Th} = \text{Th}^{\mathcal{T}} : \underline{\mathbf{K}} \longrightarrow \underline{\mathbf{Pic}}(\mathcal{T}),$$

sending a virtual vector bundle v over a scheme S to the associated *Thom object*, a \otimes -invertible object $\text{Th}_S(v) \in \mathcal{T}(S)$. An *orientation* of \mathcal{T} amounts to the data of, for every virtual vector bundle v of rank n on a scheme X , a Thom isomorphism

$$\text{Th}_X(v) \simeq \langle n \rangle,$$

where $\langle n \rangle = (n)[2n]$.⁽⁴⁾ These isomorphisms are required to be functorial and compatible with exact triangles of perfect complexes. Note that this is the same data as that of a factorization of Th through the rank map $\underline{\mathbf{K}} \rightarrow Z$.

Now recall that an SL-structure on a vector bundle is a trivialization of its determinant, and let $\underline{\mathbf{K}}^{\text{SL}}$ denote the groupoid of SL-oriented virtual vector bundles. We define an SL-orientation of T as a factorization of the composite

$$\underline{\mathbf{K}}^{\text{SL}} \longrightarrow \underline{\mathbf{K}} \xrightarrow{\text{Th}} \underline{\text{Pic}}(T)$$

through the rank map $\underline{\mathbf{K}} \rightarrow Z$, where the first arrow is the forgetful map. In other words, we have Thom isomorphisms only for vector bundles equipped with SL-structures (and the isomorphism depends on the SL-structure). This notion is derived from the foundational work of Panin and Walter [PW19].

For any virtual vector bundle v of rank n , $v - \det(v)$ is a virtual vector bundle of rank $n - 1$ with a canonical SL-orientation. Thus if T is SL-oriented, we get canonical isomorphisms

$$\text{Th}(v) \simeq \text{Tw}(\det(v)),$$

where we adopt the notation $\text{Tw}(L) := \text{Th}_X(L)\langle n - 1 \rangle$ for any graded line bundle $(L, n) \in \underline{\text{Pic}}$.

The fact that $\text{SH}_{\mathbb{Q},+}$ is oriented (Theorem A(I)) is known by [CD19, 14.2.5 & Th. 16.2.13]. Our result states that $\text{D}_{\mathbb{A}^1,-}$ and $\text{SH}_{\mathbb{Q}}$ are canonically SL-oriented (see Theorem 7.8 and Corollary 7.9). In particular, this means that every rational motivic spectrum E admits a canonical SL-orientation. This is a motivic analogue of the fact that any rational topological spectrum corresponds to an “ordinary cohomology theory”, and in particular is complex-oriented (in the sense of Adams).

An important consequence concerns the *fundamental class*

$$\eta_f : \text{Th}_X(T_{X/S}) \longrightarrow f^!(\mathbb{1}_S)$$

associated to any smoothable lci morphism $f : X \rightarrow S$ in SH_- or $\text{SH}_{\mathbb{Q}}$ (see [DJK21, 4.3.4]). Namely, this morphism simplifies to

$$\eta_f : \text{Tw}_X(\tilde{\omega}_{X/Y}) \longrightarrow f^!(\mathbb{1}_S),$$

where $\tilde{\omega}_{X/S}$ is the graded determinant of the virtual tangent bundle $T_{X/S}$ (see 7.10 below). In particular, for a regular closed immersion $i : Z \rightarrow X$, the fundamental class determines a canonical element in twisted cohomology with support:

$$\eta_i \in H_Z^{2c-2,c-1}(X, \lambda_i)_{\mathbb{Q}} = [X/X - Z, \text{Tw}_X(\lambda_i, c)]_{T(X)_{\mathbb{Q}}},$$

where c is the codimension of i and λ_i the determinant of the normal bundle of i , which defines an element $(\lambda_i, c) \in \underline{\text{Pic}}$. In general, the above fundamental class lives in twisted \mathbb{A}^1 -bivariant theory, or rather its rational or minus part, and satisfies the classical properties of intersection theory, as explained in [DJK21] (compatibility with

⁽⁴⁾Recall that the *Tate sphere* is defined as $\mathbb{1}_X(1) := \text{Th}_X(\mathbb{A}_X^1)[-2]$; for any non-negative integer n , $\mathbb{1}_X(n) := \mathbb{1}_X(1)^{\otimes n}$, and if $n < 0$, $\mathbb{1}_X(n) := \mathbb{1}_X(-n)^{\otimes(-1)}$.

composition, transversal base change, excess intersection formula). Moreover, it induces the usual wrong-way functorialities (covariance of twisted cohomology with respect to proper smoothable lci morphisms, contravariance of bivariant theories with respect to smoothable lci morphisms).

(II)–(IV) **ABSOLUTE PURITY, FINITENESS AND DUALITY.** — These results are originally the main motivations of this paper. Grothendieck’s *absolute (cohomological) purity conjecture* [Gro77, I3.1.4] predicted that for a closed immersion $i : Y \rightarrow X$ between regular noetherian schemes of pure codimension c , and n an integer invertible on X , there is an isomorphism $i^!(Z/n) \simeq Z/n(-c)[-2c]$ in the derived category of torsion étale sheaves on Y . Though several cases were proved in the ’60s (see [Gro77, I5.1], [AGV73, XVIII 3.2.5]), proving this property in the generality conjectured turned out to be a highly non-trivial problem: it was not until the ’80s that a systematic approach for the general setting emerged, where a significant step was made by Thomason [Tho84], who proved the conjecture assuming that all prime divisors of n are sufficiently large, using the Atiyah-Hirzebruch spectral sequence for étale K -theory. The general case was eventually solved by Gabber [Fuj02], based on Thomason’s method and the rigidity property for algebraic K -theory. Further developments using a refinement of de Jong resolution of singularities, can be found in [ILO14, Exp. XVI].

Together with resolution of singularities, the absolute purity property lies at the very heart of Grothendieck’s profound insight on the cohomological study of algebraic varieties. It has played a crucial role in the following applications:

- (1) The proof that the six functors on the derived category of torsion étale sheaves preserve constructible objects.
- (2) The proof of what is now called Grothendieck–Verdier local duality. Most notably, if S is a noetherian scheme equipped with a dimension function, there exists a *dualizing complex* K_S over S such that $D_S = \underline{\mathrm{Hom}}(-, K_S)$ is an auto-equivalence of categories. The functor D_S then transforms f_* (resp. f^*) into $f_!$ (resp. $f^!$).
- (3) The construction of Gysin morphisms and cycle classes, and the cohomological study of intersection theory.
- (4) The study of the coniveau spectral sequence.

The attempt to answer these questions has inspired a number of important works based on new methods. Deligne managed to prove the first point unconditionally by avoiding the use of resolution of singularities [Del77, Th. Finitude]. The second point was known in dimension at most one in [Del77, Dualité 1.4], and proved by Gabber in general [ILO14, XVII]; furthermore, it implies the existence of the (self-dual) perverse t -structure over suitable base schemes, extending the fundamental work of [BBD82]. The third point already appeared in [AGV73, XVIII], and has since then been much worked on in the literature (see [DJK21]). The last point has also been discussed by a number of authors, such as [BO74] and [CTHK97].

With the growing development of the theory of *triangulated mixed motives* since the ’90s, absolute purity naturally finds its place in this framework. In fact, this property

already appears implicitly in Beilinson’s axiomatization modeled on the étale setting. Morel and Voevodsky’s motivic homotopy theory came later [MV99], together with a formalism of six functors in a similar fashion [Ayo07], in which absolute purity can be formulated for more general motivic categories, as highlighted in [CD16, A.2] and [DJK21, 4.3]. Here are the first cases where absolute purity is known:

- For mixed motives with rational coefficients, it was first formulated and proved by Cisinski–Déglise in [CD19, Th. 14.4.1].
- For modules over the algebraic K -theory spectrum, it is proved in [CD19, Th. 13.6.3]. Note that this serves as an ingredient in the proof of the previous case.
- For étale motives, it is proved in [Ayo14] and [CD15].

These examples provide some evidence that such a property should hold in greater generality, and should philosophically be an addition to the six functors formalism: it was conjectured in [Dég18b] that absolute purity should also hold for the motivic homotopy category SH itself.

From the case of rational motivic cohomology, it follows from [CD19, Th. 16.2.13] that absolute purity also holds for $\mathrm{SH}_{\mathbb{Q},+}$. In this note, we show that it holds for SH_- , and hence also for $\mathrm{SH}_{\mathbb{Q}}$ (Theorem 3.6). These properties are known under the assumption (Resol) in [BD17, 2.4.1], but the difficulty is that the category $\mathrm{SH}_{\mathbb{Q}}$ is not *separated* in the sense of [CD19, Def. 2.1.7]. The basis for our proof is the following simple but powerful observation (see Lemma 2.2 and Example 2.3(1)):

KEY LEMMA. — *For every scheme X , there is a canonical equivalence of symmetric monoidal stable ∞ -categories*

$$\mathrm{SH}(X)_- \simeq \mathrm{SH}(X_{\mathbb{Q}})_-,$$

where $X_{\mathbb{Q}} = X \times_{\mathrm{Spec} \mathbb{Z}} \mathrm{Spec} \mathbb{Q}$, which commutes with the six operations.

This boils down to the calculation that $\mathrm{SH}(\mathrm{Spec} k)_- = 0$ for any field k of positive characteristic. The same holds for $D_{A^1,-}(-, \Lambda)$, for any commutative ring Λ , as well as the ρ -inverted variants (see Example 2.3).

The Key Lemma reduces the question of absolute purity to the equicharacteristic case, which is well-known (we include a proof in Appendix C). We thus deduce absolute purity for the rational, minus, and ρ -inverted parts of the motivic sphere spectrum.

The finiteness and duality results (properties (III) and (IV)) are similarly deduced using the Key Lemma from the equicharacteristic case, which is known by work of Bondarko–Déglise [BD17] (see also [EK20a, §3.1]). See Propositions 3.3 and 3.4.

Finally, let us make the following remarks. First, if $f : X \rightarrow S$ is a smoothable morphism of regular schemes, and $\tilde{\omega}_{X/S}$ is the graded determinant of its relative virtual tangent bundle, then by (II) and (IV), the object

$$K_X = f^!(\mathbb{1}_S) \simeq \mathrm{Tw}_X(\tilde{\omega}_{X/S})$$

is a dualizing object in $\mathrm{SH}(X)_{\mathbb{Q}}$. Note the striking resemblance to Grothendieck–Serre duality for coherent cohomology [Har66, Def. III.2, Prop. 7.2].

(V) **COMPARISON.** — Let $\mathbf{H}_M\mathbb{Q}_S$ denote the rational motivic cohomology spectrum over a scheme S . This can be defined either as the Beilinson motivic cohomology spectrum [CD19, Def. 14.1.2] or equivalently as the inverse image along the structural map $S \rightarrow \mathrm{Spec} Z$ of the rationalization of Spitzweck’s integral motivic cohomology spectrum (by [Spi18, Th. 7.14]). Modules over $\mathbf{H}_M\mathbb{Q}$ provide a good triangulated category of rational mixed motives, equivalent to Voevodsky’s construction at least over excellent schemes [CD19, §16.1]. The comparison equivalence (Theorem A(V))

$$\mathrm{SH}(S)_{\mathbb{Q},+} \simeq \mathbf{H}_M\mathbb{Q}\text{-mod}(S)$$

was first observed by Morel over a field [Mor06] and generalized by Cisinski–Déglise [CD19, §16.2] to general bases.

For $\mathrm{SH}_{\mathbb{Q},-}$, an analogous statement was proved by Ananyevskiy–Levine–Panin over a field [ALP17, §4]. Our comparison generalizes this to arbitrary base schemes S and to a Λ -linear statement for any commutative ring Λ . Over any base scheme S , we define a motivic spectrum $\mathbf{H}_W\Lambda_S$, the Λ -linear *homotopical Witt spectrum* (see Definition 5.18). For S regular, $\mathbf{H}_W\Lambda_S$ is defined by the periodic \mathbb{G}_m -spectrum

$$(\underline{W}_S^\Lambda, \underline{W}_S^\Lambda, \dots)$$

where \underline{W}_S is the Zariski sheaf associated to the Witt presheaf

$$S \mapsto W(S_{\mathbb{Q}}) \otimes_Z \Lambda$$

on the site of smooth S -schemes of finite type (see Definition 5.2 and Paragraph 5.8).

We then prove $D_{A^1}(S, \Lambda)_- \simeq \mathbf{H}_W\Lambda\text{-mod}(S)$, hence in particular

$$\mathrm{SH}(S)_{\mathbb{Q},-} \simeq \mathbf{H}_W\mathbb{Q}\text{-mod}(S)$$

over every scheme S (see Corollary 5.16 and Paragraph 5.19). Following [ALP17], we can call the right-hand side the stable ∞ -category of *Witt motives* over S . For S regular, we deduce a computation of the minus part of the stable A^1 -derived cohomology of S :

$$H_{A^1}^{n,i}(S, \Lambda)_- = \mathrm{Hom}_{D_{A^1}(S, \Lambda)}(\mathbb{1}_S, \mathbb{1}_S(i)[n])_- \simeq H_{\mathrm{Zar}}^{n-i}(S_{\mathbb{Q}}, \underline{W}) \otimes_Z \Lambda.$$

Finally we put the plus and minus parts together to get a model for $\mathrm{SH}_{\mathbb{Q}}$. Extending the original definition of [BCD⁺20, Chap. 6], we define the rational *Milnor–Witt motivic cohomology* spectrum over any scheme S by

$$\mathbf{H}_{MW}\mathbb{Q}_S = \mathbf{H}_M\mathbb{Q}_S \oplus \mathbf{H}_W\mathbb{Q}_S,$$

where $\mathbf{H}_M\mathbb{Q}_S$ is the rational motivic cohomology ring spectrum over S (see Definition 6.1). We then deduce the comparison (Corollary 6.3)

$$\mathrm{SH}(S)_{\mathbb{Q}} \simeq \mathbf{H}_{MW}^{\mathbb{Q}}\text{-mod}(S).$$

Thus we may view the right-hand side as the stable ∞ -category of *Milnor–Witt motives* over S . For S the spectrum of an infinite perfect field of characteristic unequal to 2, this result was recently proved by Garkusha [Gar19].

In particular, for a regular scheme S , we get the following computation of the rational stable homotopy classes of endomorphisms of the sphere spectrum (see Corollary 6.2):

$$[\mathbb{1}_S, \mathbb{1}_S(i)[n]] \otimes_Z \mathbb{Q} \simeq K_{2i-n}^{(n)}(S)_{\mathbb{Q}} \oplus H_{\text{Zar}}^{n-i}(S_{\mathbb{Q}}, \mathbb{W})_{\mathbb{Q}}.$$

This result generalizes [ALP17, Rem. 4.5]. In the case where S is the spectrum of a field (more generally when S is a regular semi-local scheme over a field, see [BH21, Th.10.12]), this recovers Morel’s computation of the stable stems of A^1 -homotopy sheaves ([Mor12, Th. 6.40]).

We also apply these results to the dimensional homotopy t -structures of Bondarko–Déglise [BD17] on SH_- . If S is equipped with a dimension function δ , then we consider the δ -homotopy t -structure on $\text{SH}(S)_-$ and $\text{SH}(S)_{\mathbb{Q}}$ (Paragraph 4.1). The Key Lemma allows us to generalize the positivity/negativity criterion (Theorem 3 of op.cit.) to this setting (Proposition 4.2). Using this criterion we show that for a regular scheme S , the monoidal unit of $\text{D}_{A^1}(S, \Lambda)_-$ lies in the heart of the δ -homotopy t -structure, when $\delta = -\text{codim}_S$ (Corollary 5.14). (In contrast, this property does not hold for $\text{SH}_{\mathbb{Q},+}$.)

(VI) BIVARIANT THEORY. — In any motivic ∞ -category \mathcal{T} , the unit object represents a *bivariant theory* by the formula

$$H_{m,n}^{\mathcal{T}}(X/S) := [\mathbb{1}_X(n)[m], f^!(\mathbb{1}_S)]_{\mathcal{T}(X)}$$

for $f : X \rightarrow S$ a separated morphism of finite type (see [Dég18a, Def. 1.2.2]). (Note that for $f = \text{Id}_X$, we recover the cohomology theory represented by the unit, with opposite grading.) The last part of our main result is concerned with identifying these groups explicitly when \mathcal{T} is SH_- or $\text{SH}_{\mathbb{Q}}$, and S is a regular base scheme equipped with the dimension function $\delta = -\text{codim}_S$.

For $\text{SH}_{\mathbb{Q},+}$ this was achieved by Bondarko–Déglise [BD17, Th. 2], using absolute purity and the niveau spectral sequence. We follow the same ideas to get an analogous computation for SH_- and $\text{SH}_{\mathbb{Q}}$ (see Theorem 8.7). Let us make this computation explicit in the particular case $X = S$, a regular scheme. Then we get an isomorphism of cohomology theories:

$$H_{A^1}^{2n,n}(X, \mathbb{Q}) = [\mathbb{1}_X, \mathbb{1}_X\langle n \rangle]_{\mathbb{Q}} \simeq \widetilde{\text{CH}}^n(X) \otimes_Z \mathbb{Q}$$

where the right-hand side is the group of “rational equivalence classes of unramified quadratic cycles” of codimension n in X , i.e., the Chow–Witt group (see [Fas08, FS09]). More explicitly, this is the middle cohomology group of a complex of the form:

$$\bigoplus_{\eta \in X^{(n-1)}} \mathbf{K}_1^{\text{MW}}(\kappa_{\eta}, \omega_{\eta})_{\mathbb{Q}} \longrightarrow \bigoplus_{x \in X^{(n)}} \text{GW}(\kappa_x, \omega_x)_{\mathbb{Q}} \longrightarrow \bigoplus_{s \in X^{(n+1)}} W(\kappa_s, \omega_s)_{\mathbb{Q}}$$

where for $x \in X$, κ_x is the residue field and ω_x the determinant of the cotangent complex of the immersion of x in X , and \mathbf{K}_1^{MW} , GW , W are respectively the twisted form of the first Milnor–Witt, Grothendieck–Witt, and Witt group of the given field (see paragraph 8.6 for more details). Note that when 2 is invertible on S , this coincides with Theorem 4 of [FS09]. The above definition makes sense even without this

assumption: the Key Lemma tells us that the Witt part vanishes over the characteristic 2 part of S ! Note moreover that we give a formulation that also takes into account the presence of twists.

Using this computation we rule out the existence of Bondarko’s Chow weight structure on $\mathrm{SH}(S)_{\mathbb{Q}}$ for a large class of schemes S (see Corollary 8.14). Moreover, the computation is used in [DF20] to obtain the so-called Borel character over arbitrary regular bases, which provides a rational isomorphism between hermitian K-theory and Milnor–Witt motivic cohomology extending the Chern character.

ORGANIZATION OF THE PAPER. — The paper is divided into 8 sections and contains 5 appendices.

The Key Lemma is proved in Section 2. In Section 3, we prove properties (II), (III), and (IV) of Theorem B and Corollary C. Section 4 deals with dimensional homotopy t -structures. We also obtain a useful description of the generators of SH_{-} and $\mathrm{SH}_{\mathbb{Q}}$ (Proposition 4.4). In Section 5, we give the construction of the *homological Witt spectrum* and prove property (V) for SH_{-} . Section 6 deals with the same property (V) for $\mathrm{SH}_{\mathbb{Q}}$. In Section 7, we reformulate Panin–Walter’s theory of SL-orientations in terms of motivic categories, and explain its application to fundamental classes and Gysin morphisms. We also prove property (I), i.e., that $\mathrm{SH}_{\mathbb{Q}}$ and SH_{-} are SL-oriented. In Section 8, we explain the computation of the δ -niveau spectral sequence (a slight variation on the classical niveau spectral sequence using a dimension function δ as introduced in [BD17]) with respect to rational A^1 -bivariant theory (Definition 8.2), thus proving property (VI).

In Appendix A, we review the property of continuity in the language of motivic ∞ -categories. We use this property in Appendix B to extend the exceptional inverse image operation $f^!$ to essentially of finite presentation morphisms. In Appendix C we give a detailed proof of the absolute purity theorem for schemes defined over a field. (This result has already been noted in the literature, but without a detailed proof.) In Appendix D, we prove absolute purity for hermitian K-theory, on the model of the case of Quillen K-theory [CD19, Th. 13.6.3]. This result can be used to give another proof of Corollary C(II) (at least for schemes over $\mathbb{Z}[1/2]$, see [DFJK19]) and we include it for completeness. Finally in Appendix E, we prove the analogue of the Key Lemma for the 2-localized higher Witt groups of Balmer. We note that this readily implies the Gersten–Witt conjecture for arbitrary regular $\mathbb{Z}[1/2]$ -schemes, incidentally extending a result of Jacobson (see Remark E.5).

CONVENTIONS. — All schemes are assumed quasi-compact and quasi-separated. Regular schemes are assumed noetherian. Separated morphisms are assumed to be of finite type. We will say that a morphism $f : X \rightarrow S$ of schemes is *pro-étale* (resp. *pro-smooth*) if X is the projective limit of a cofiltered system of étale (resp. smooth) S -schemes with affine transition morphisms. Given a property \mathcal{P} of morphisms (e.g. étale, smooth, or separated), we will say that a morphism $f : X \rightarrow S$ of schemes *essentially* satisfies \mathcal{P} (or is essentially étale, smooth, or separated) if X can be

written as a projective limit of S -schemes satisfying \mathcal{P} with affine and étale transition morphisms. See Appendix B for details.

We will freely use the language of *motivic ∞ -categories*, although the reader can mostly replace these with motivic triangulated categories [CD19]. We will be more precisely concerned with the motivic stable homotopy category SH and the Λ -linear A^1 -derived category $D_{A^1}(-, \Lambda)$ for a commutative ring of coefficients Λ . Recall that, if T is any of these motivic ∞ -categories, the motivic ∞ -category $T[1/2]$ obtained by inverting 2 (or when $2 \in \Lambda^\times$) admits following Morel a canonical orthogonal decomposition:

$$T[1/2] = T_+ \times T_-$$

where T_+ (resp. T_-) is the (categorical) kernel of the projector $e_+ = (1 - \varepsilon)/2$ (resp. $e_- = (1 + \varepsilon)/2$) — here $\varepsilon = -\langle -1 \rangle$ in Morel’s notation. We also note that T_- can be identified with $T[1/2, \eta^{-1}]$, where η is the algebraic Hopf map (see e.g. [ALP17, Rem. 3.6]).

For a motivic ∞ -category T and a scheme S , the full subcategory $T_c(S)$ of *constructible* objects is the thick stable subcategory generated by objects of the form $f_! f^!(1_S)(n)$ for f smooth separated of finite type and $n \in \mathbb{Z}$. For the examples of T we consider, this will coincide with the subcategory of compact objects (see [CD19, Prop. 1.4.11]).

Acknowledgments. — F. Déglise would like to thank Denis-Charles Cisinski whose influence via years of collaborations in motivic homotopy theory has been invaluable for the results obtained in this work. F. Jin would like to thank Heng Xie for discussions about the paper [Jac17]. A. Khan would like to thank Tom Bachmann, Marc Hoyois, and Charanya Ravi for helpful discussions related to this paper. We would like to thank the referee for carefully reading the paper and for the comments which helped improving the quality of the paper.

2. THE KEY LEMMA

2.1. — For a scheme X , denote by $X_{\mathbb{Q}} = X \times_{\text{Spec}(\mathbb{Z})} \text{Spec}(\mathbb{Q})$ the characteristic zero fiber. We denote the inclusion by

$$\nu = \nu_X : X_{\mathbb{Q}} \longrightarrow X.$$

The following observation is the *key lemma* of this paper:

LEMMA 2.2. — *Let T be a continuous motivic ∞ -category. Suppose that for every field k of positive characteristic, $T(k) = 0$. Then for every scheme X , the functor*

$$\nu_X^* : T(X) \longrightarrow T(X_{\mathbb{Q}})$$

is an equivalence.

Proof. — If we write X as a cofiltered limit of schemes X_α of finite type over $\text{Spec } \mathbb{Z}$ (with affine transition maps), then by continuity (Definition A.2) ν_X^* will be the colimit of the functors $\nu_{X_\alpha}^*$. Thus we may assume X is noetherian. By Lemma B.10,

the co-unit $\nu^*\nu_* \rightarrow \text{Id}$ is invertible since ν is a pro-open immersion. It remains to show that the unit $\text{Id} \rightarrow \nu_*\nu^*$ is also invertible. By Proposition B.20, it will suffice to show that

$$i_x^! \longrightarrow i_x^! \nu_* \nu^*$$

is invertible for every point $x \in X$, where i_x is the inclusion of the residue field at x and where $i_x^!$ is defined as in Paragraph B.13. By Propositions B.18 and B.19 we may thus assume that X is the spectrum of a field k . If k is of characteristic 0, then ν is an isomorphism so the claim is obvious. If k is of positive characteristic, then both $T(X)$ and $T(X_\square)$ are zero by assumption, and the result follows.

EXAMPLE 2.3. — The condition of Lemma 2.2 is satisfied for the following T :

(1) $T = \text{SH}_-$, the minus part of the stable homotopy category. Indeed, recall that for any field k there is a canonical isomorphism

$$\text{Hom}_{\text{SH}(k)}(\mathbb{1}_k, \mathbb{1}_k) \simeq \text{GW}(k),$$

see [Mor12, Cor. 1.25],⁽⁵⁾ where $\text{GW}(k)$ is the Grothendieck–Witt ring of k . The decomposition $\text{SH}[1/2] \simeq \text{SH}_+ \times \text{SH}_-$ corresponds to the decomposition

$$\text{GW}(k)[1/2] \simeq Z[1/2] \times W(k)[1/2],$$

where $W(k)$ is the Witt ring (see [Mor12, Eq. (3.1)]). Thus there is a canonical isomorphism

$$\text{Hom}_{\text{SH}(k)_-}(\mathbb{1}_{k,-}, \mathbb{1}_{k,-}) \simeq W(k)[1/2].$$

But if k is of positive characteristic and hence not formally real, then $W(k)[1/2]$ vanishes (see e.g. [EKM08, §31.A] or [Lam05, Chap. VIII, Lem. 3.4]). It follows that $\mathbb{1}_{k,-} \simeq 0$ and hence $\mathbb{E}_- = \mathbb{E} \otimes \mathbb{1}_{k,-} \simeq 0$ for every $\mathbb{E} \in \text{SH}(k)$.

(2) $T = \text{D}_{\mathbb{A}^1}(-, \Lambda)_-$, where Λ is any commutative ring with $2 \in \Lambda^\times$. We may argue as in the first example, since the canonical map $\mathbb{1}_k \rightarrow \text{H}_{\mathbb{A}^1} Z_k$ induces an isomorphism on zeroth homotopy sheaves. Indeed, since k is pro-smooth over its prime subfield, this follows by base change from the case where k is perfect. In that case the claim is [Mor12, Th. 6.37].

(3) $T = \text{SH}(-)[1/\rho]$ or $T = \text{D}_{\mathbb{A}^1}(-)[1/\rho]$, where $\rho = -[-1]$ as in [Bac18, Cor. 19]. This follows from the fact that, when k is a field of positive characteristic, ρ is nilpotent in $\text{SH}(k)$ and $\text{D}_{\mathbb{A}^1}(k)$, see the proof of [Bac18, Cor. 32].

Moreover, each of these examples is continuous (see Example A.4).

REMARK 2.4

(1) Under the assumptions of Lemma 2.2, we moreover have canonical identifications $\nu^* \simeq \nu^!$, where $\nu^!$ is defined as in Paragraph B.13 (see Example B.15). As this functor is an equivalence, its right adjoint ν_* is also a left adjoint. Thus the functor $\nu^!$ admits a left adjoint $\nu_!$ (compare Warning B.17). Moreover, it immediately

⁽⁵⁾where the perfectness hypothesis is not necessary, see [BH21, Th. 10.12]

follows that both of the functors $\nu^* \simeq \nu^!$ and $\nu_* \simeq \nu_!$ commute with the six operations. Similarly, symmetric monoidality of ν^* implies that $\nu_* \simeq \nu_!$ is also symmetric monoidal.⁽⁶⁾

(2) Here is an alternative argument to show that the functor ν_X^* in Lemma 2.2 is an equivalence for $\mathcal{T} = \mathrm{SH}(-)[1/\rho]$ (respectively $\mathcal{T} = \mathrm{D}_{\mathbb{A}^1}(-)[1/\rho]$): by [Bac18, Th. 35] there is a canonical equivalence $\mathrm{SH}(X)[1/\rho] \simeq \mathrm{SH}(X_r)$, where X_r is the site associated to the *real scheme* of X , which is a topological space whose points are pairs (x, σ) where $x \in X$ is a point and σ is an ordering of the residue field of x ([Sch94, 0.4.1]); since no field of positive characteristic is orderable, the morphism ν_X induces a homeomorphism (and therefore an equivalence of sites) $(X_{\mathbb{Q}})_r \simeq X_r$, which induces an equivalence $\mathrm{SH}(X)[1/\rho] \simeq \mathrm{SH}(X_{\mathbb{Q}})[1/\rho]$.

3. DUALITY AND ABSOLUTE PURITY

In this section, we deduce from Lemma 2.2 several structural results for various motivic categories, corresponding to properties (II)–(IV) from the introduction.

3.1. — Let \mathcal{T} be one of the following motivic ∞ -categories:

- $\mathcal{T} = \mathrm{SH}_{\mathbb{Q}} \simeq \mathrm{D}_{\mathbb{A}^1}(-, \mathbb{Q})$;
- $\mathcal{T} = \mathrm{SH}_-$ or $\mathcal{T} = \mathrm{D}_{\mathbb{A}^1}(-, \Lambda)_-$, where Λ is any commutative ring of coefficients;
- $\mathcal{T} = \mathrm{SH}[1/\rho]$ or $\mathcal{T} = \mathrm{D}_{\mathbb{A}^1}[1/\rho]$, where ρ is as in Example 2.33.

Note that in the last two cases \mathcal{T} satisfies the condition in Lemma 2.2. In the first case we may use the equivalence $\mathrm{SH}_{\mathbb{Q}} \simeq \mathrm{SH}_{\mathbb{Q},+} \times \mathrm{SH}_{\mathbb{Q},-}$ to reduce to the plus and minus parts respectively. By [CD19, Th. 16.2.13] the plus part is identified with Beilinson motives $\mathrm{SH}_{\mathbb{Q},+} \simeq \mathrm{DM}_{\mathbb{B},\mathbb{Q}}$, for which the properties have already been established in op. cit.

3.2. — The first application is the fact that the six functors preserve constructibility and what is called *Grothendieck–Verdier local duality*, generalizing [BD17, Th. 2.4.9]:

PROPOSITION 3.3 (Constructibility). — *Let \mathcal{T} be as in 3.1. Given a morphism of schemes $f : X \rightarrow S$, the subcategory of constructible objects is stable under the following operations:*

- $f^*, f_!$ for f separated of finite type, \otimes_S ;
- f_* for f of finite type, $f^!$ for f separated of finite type, $\underline{\mathrm{Hom}}_S$, assuming in addition that X and S are quasi-excellent.

Proof. — If \mathcal{T} satisfies the condition in Lemma 2.2, then by Remark 2.4 one reduces to the case of characteristic zero schemes, for which the result follows from [BD17, Th. 2.4.9]. For Beilinson motives the result is proved in [CD16, Th. 6.3.15] (modulo the comparison of [CD19, Th. 16.1.2]).

⁽⁶⁾By [Lur18, Prop. 2.5.5.1] (resp. its dual), ν_* is lax symmetric monoidal (resp. $\nu_!$ is colax symmetric monoidal). That is, $\nu_* \simeq \nu_!$ is both lax and colax symmetric monoidal, hence it is strict symmetric monoidal.

PROPOSITION 3.4 (Grothendieck–Verdier duality). — *Let \mathcal{T} be as in 3.1. Let S be a regular quasi-excellent scheme and $K_S \in \mathcal{T}(S)$ a \otimes -invertible object. Then for any separated morphism $f : X \rightarrow S$ of finite type, the constructible object*

$$K_X = f^!(K_S) \in \mathcal{T}(X)$$

is dualizing. That is, for any constructible object $M \in \mathcal{T}(X)$, the canonical map

$$(3.4.a) \quad M \longrightarrow D_X(D_X(M))$$

is invertible, where D_X is the contravariant endofunctor of $\mathcal{T}(X)$ given by

$$D_X(M) = \underline{\mathrm{Hom}}(M, K_X).$$

Proof. — If \mathcal{T} satisfies the condition in Lemma 2.2, it suffices to show that the map (3.4.a) is an isomorphism after applying the equivalence $\nu^* : \mathcal{T}(X) \simeq \mathcal{T}(X_{\mathbb{Q}})$. Since this equivalence commutes with the six operations (Remark 2.4), we are reduced to the equicharacteristic case, which is proved in [BD17, Th. 2.4.9]. For Beilinson motives, the result is proved in [Cis19, Th. 2.3.2].

3.5. — The most important consequence we obtain is the absolute purity property. Let \mathcal{T} be a motivic ∞ -category. Recall that an *absolute object* E in \mathcal{T} is a cartesian section, i.e., the data of an object $E_X \in \mathcal{T}(X)$ for every scheme X together with natural isomorphisms $f^*E_Y \simeq E_X$ for every morphism $f : X \rightarrow Y$ (compatible up to coherent homotopy). An absolute object E is said to satisfy *absolute purity* if for any smoothable morphism $f : X \rightarrow Y$ between regular schemes with virtual tangent bundle T_f , the map

$$E_X \otimes \mathrm{Th}(T_f) \longrightarrow f^!E_Y$$

induced by the fundamental class of f (see [DJK21, Def. 4.3.7]), is an isomorphism.

THEOREM 3.6 (Absolute purity)

- (1) *If \mathcal{T} satisfies the condition in Lemma 2.2, then any absolute object in \mathcal{T} satisfies absolute purity.*
- (2) *The minus part $\mathbb{1}_-$ and the rational part $\mathbb{1}_{\mathbb{Q}}$ of the motivic sphere spectrum satisfy absolute purity in SH.*
- (3) *Given any commutative ring Λ with $2 \in \Lambda^\times$, the minus part of the A^1 -derived Λ -linear ring spectrum Λ_- satisfies absolute purity in $D_{A^1}(-, \Lambda)$.*

Proof. — If \mathcal{T} satisfies the condition in Lemma 2.2, then by Remark 2.4 we are reduced to the characteristic zero case, where the result follows from Theorem C.1. Consider the second claim. For the ring spectrum $\mathbb{1}_-$, one applies the previous claim with $\mathcal{T} = \mathrm{SH}_-$. The case of the rational sphere spectrum $\mathbb{1}_{\mathbb{Q}} \simeq \mathbb{1}_{\mathbb{Q},+} \oplus \mathbb{1}_{\mathbb{Q},-}$ follows from the case just treated and absolute purity for Beilinson motives ([CD19, Th. 14.4.1]). The third claim follows from the first one with $\mathcal{T} = D_{A^1}(S, \Lambda)$.

4. THE DIMENSIONAL HOMOTOPY t -STRUCTURE

We next turn to applications about the dimensional homotopy t -structure [BD17].

4.1. — Recall the following definitions:

(1) Let \mathcal{C} be a compactly generated stable ∞ -category which has small colimits. For \mathcal{F} a family of objects in \mathcal{C} , there is a t -structure on \mathcal{C} , called the t -structure generated by \mathcal{F} , where the subcategory of non-negative objects is the closure of \mathcal{F} under extensions, positive shifts and small coproducts ([BD17, Th. 1.2.6], [Ayo07, Prop. 2.1.70]).

(2) Let \mathcal{T} be a motivic ∞ -category. If $f : X \rightarrow S$ is a morphism of finite type, we denote $M^{\text{BM}}(X/S) = f_! \mathbb{1}_X \in \mathcal{T}(S)$ the *Borel-Moore motive*. If $E \in \mathcal{T}(S)$, we denote

$$E_{p,q}(X/S) = [M^{\text{BM}}(X/S)(q)[p], E]_{\mathcal{T}(S)}$$

the *bivariant group*.

(3) Let \mathcal{T} be a motivic ∞ -category and let S be a scheme equipped with a dimension function δ . The δ -homotopy (perverse) t -structure over S is the t -structure t_δ on $\mathcal{T}(S)$ generated by the family of elements of the form $M^{\text{BM}}(X/S)(n)[n + \delta(X)]$ for X a separated scheme of finite type over S and $n \in \mathbb{Z}$ [BD17, Def. 2.1.1].

(4) Let S be a scheme and let $x : \text{Spec } \kappa(x) \rightarrow S$ be a topological point of S . An S -model of x is an affine regular S -scheme $X = \text{Spec } A$ of finite type, where A is a subring of $\kappa(x)$ whose fraction field is $\kappa(x)$, such that x coincides with the composition $\text{Spec } \kappa(x) \rightarrow X \rightarrow S$. A morphism of S -models is an S -morphism of the underlying schemes compatible with the structure of subring of $\kappa(x)$. The category of S -models of x is denoted $\mathcal{M}(x)$ [BD17, Def. 3.2.1]. If S is excellent, $\mathcal{M}(x)$ is non-empty.

(5) Let E be an object in $\mathcal{T}(S)$. For x a topological point of S and $n, p \in \mathbb{Z}$, the fiber δ -homology of E in degree p is defined as

$$\widehat{H}_p^\delta(E)(x, n) = \varinjlim_{X \in \mathcal{M}(x)^{\text{op}}} E_{2\delta(x)+p+n, \delta(x)+n}(X/S),$$

see [BD17, Def. 3.2.3].⁽⁷⁾

Then we obtain the following generalization of [BD17, Th. 3.3.1]:

PROPOSITION 4.2. — *Let \mathcal{T} be as in 3.1 and let S be an excellent noetherian scheme equipped with a dimension function δ . Then for any object $E \in \mathcal{T}(S)$ the following conditions are equivalent:*

(1) E is t_δ -positive (resp. E is t_δ -negative), that is, $E \in \mathcal{T}(S)_{t_\delta > 1}$ (resp. $E \in \mathcal{T}(S)_{t_\delta \leq -1}$).

⁽⁷⁾We remark that, according to the extension of the six functors formalism done in Appendix B, we can write:

$$\widehat{H}_p^\delta(E)(x, n) = E_{2\delta(x)+p+n, \delta(x)+n}(\kappa(x)/S) := \text{Hom}_{\mathcal{T}(x)}(\mathbb{1}_x(\delta(x) + n)[2\delta(x) + p + n], \nu_x^! E).$$

with $\nu_x : \text{Spec}(\kappa(x)) \rightarrow S$ the canonical immersion (Example B.3). In particular, we could avoid the use of S -models from [BD17].

(2) For any topological point x of S and integers $n \in \mathbb{Z}$ and $p \neq 0$ (resp. $p > 0$), $\widehat{H}_p^\delta(E)(x, n) = 0$.

Proof. — Suppose first that \mathcal{T} satisfies the condition in Lemma 2.2, and let ν^* denote the equivalence of loc. cit. Denote by $\delta_\mathbb{Q}$ the dimension function over $S_\mathbb{Q}$ induced by δ . By [BD17, Prop. 2.1.6], the functor ν^* is t_δ -exact, so we find that E is t_δ -positive if and only if ν^*E is $t_{\delta_\mathbb{Q}}$ -positive. On the other hand, by [BD17, Rem. 3.2.5] we know that $\widehat{H}_p^\delta(E)(x, n) = \widehat{H}_p^{\delta_\mathbb{Q}}(\nu^*E)(x, n)$ if x is a point of $S_\mathbb{Q}$, and if x is not a point of $S_\mathbb{Q}$ we have $\widehat{H}_p^\delta(E)(x, n) = 0$ by the condition in Lemma 2.2. Therefore we are reduced to prove the same statement for $\nu^*E \in \mathcal{T}(S_\mathbb{Q})$, which follows from [BD17, Ex. 3.2.14 & Th. 3.3.1].

For Beilinson motives the result already follows from [BD17, Th. 3.3.1], since $\mathrm{DM}_{\mathbb{B}, \mathbb{Q}}$ is separated and therefore satisfies the condition (Resol) in [BD17, 2.4.1]. Hence we get the claim also for $\mathcal{T} = \mathrm{SH}_\mathbb{Q}$.

We deduce the analogues of the properties that are consequences of [BD17, Th. 3.3.1], as stated in [BD17, Cor. 3.3.5, 3.3.6, 3.3.7, 3.3.9, 3.3.10]. In particular, the t -structure t_δ is non-degenerate.

4.3. — Over the ring of integers in a number field, we also obtain the following description of generators analogous to [BD17, Cor. 2.4.8]:

PROPOSITION 4.4 (Generation). — *Let \mathcal{T} be as in 3.1, and let S be one of the following schemes:*

- (1) S is the spectrum of a commutative ring which is either
 - the localization of the ring of integers of a number field at a set of primes,
 - or the ring of integers in a local field.
- (2) S is a scheme such that -1 is a sum of squares in every residue field.

Then the stable ∞ -category $\mathcal{T}(S)$ coincides with the localizing subcategory generated by objects of the form $f_\mathbb{1}_X(n)$, where $f : X \rightarrow S$ is a projective morphism with X a regular scheme and $n \in \mathbb{Z}$.*

Proof. — For Beilinson motives the result is proved in [CD19, Th. 15.2.3]. Thus if -1 is a sum of squares in every residue field of S , then by [CD19, Cor. 16.2.14] we have $\mathrm{SH}(S, \mathbb{Q}) \simeq \mathrm{DM}_{\mathbb{B}}(S, \mathbb{Q})$, and the result follows.

Thus we assume $S = \mathrm{Spec}(R)$ is as in (1). If R has equal characteristic, the result is already proved in [BD17, Cor. 2.4.8]. Otherwise, we have $S_\mathbb{Q} = \mathrm{Spec} K$ where K is the fraction field of R . If \mathcal{T} satisfies the condition in Lemma 2.2, we only need to show that $\mathcal{T}(\mathrm{Spec} K)$ is generated as a localizing subcategory by objects of the form $\nu^*f_*\mathbb{1}_X(n) = \nu^*M^{\mathrm{BM}}(X/S)(n)$, where $f : X \rightarrow S$ is a projective morphism with X regular and $n \in \mathbb{Z}$. By [BD17, Cor. 2.4.8] and noetherian induction, it suffices to show that for any irreducible smooth projective K -scheme X , there exists a non-empty subscheme $U \subseteq X$ such that $M^{\mathrm{BM}}(U/K) \in \nu^*(\mathcal{T}')$. By de Jong's theory of alterations (see [Sch00, 1.2.2]), there exists a projective generically finite surjection $X' \rightarrow X$ such

that X' has a projective regular model over S . By the proof of [BD17, Cor. 2.4.8], there exists a non-empty open subscheme $U \subseteq X$ such that $M^{\text{BM}}(U/K)$ is a direct summand of an element in $\nu^*(T')$, and the result follows.

Applying Proposition 4.4, we have the following analogue of [BD17, Th. 2.4.3]:

PROPOSITION 4.5. — *Let T be as in 3.1 and let S be as in Proposition 4.4. Then the homotopy t -structure on T agrees with the t -structure generated by elements of the form $M^{\text{BM}}(X/S)(n)[n + \delta(X)]$ for X a regular scheme which is projective over S and $n \in \mathbb{Z}$.*

Note that via Lemma 2.2, we have avoided using Proposition 4.5 in the proof of Proposition 4.2, therefore removing the restriction on S .

4.6. — From Remark 2.4 it follows that the functor ν^* preserves the δ -effective subcategory ([BD17, Def. 2.2.1]). Consequently the analogues of Propositions 4.2 and 4.5 for the effective category also hold, by a similar argument.

5. THE MINUS A^1 -DERIVED CATEGORY AND WITT MOTIVES

5.1. — We recall the classical definition of the Witt ring $W(S)$ of an arbitrary scheme S , following Knebusch [Kne77, I, §5]. Given a scheme S , a *symmetric bundle* (called “space” in loc. cit.) over S is the data (V, ϕ) of a vector bundle V over S together with a non-degenerate symmetric bilinear form $\phi : V \otimes V \rightarrow A_S^1$; i.e., ϕ is a morphism of vector bundles such that $\phi \circ \varepsilon = \phi$, where ε is the isomorphism exchanging factors, and the adjoint map $\phi' : V \rightarrow \underline{\text{Hom}}(V, A_S^1)$ is an isomorphism. The notion of morphism of symmetric bundles is clear, and an isomorphism is called an *isometry*.

Then $W(S)$ is the monoid of isometry classes of symmetric bundles, modulo the sub-monoid consisting of classes of metabolic symmetric bundles (see p. 133 and p. 145 in [Kne77]). Tensor product of symmetric bundles induces a product on the abelian group $W(S)$, which turns it into a commutative ring. It is clear from the definition that $W(S)$ is contravariantly functorial in S , with respect to arbitrary morphisms of schemes, so that W defines a presheaf of commutative rings on the category of all schemes.

DEFINITION 5.2. — For any scheme S , we define the *Witt sheaf* \underline{W}_S on the site of smooth S -schemes as the Zariski sheaf associated with the presheaf $X \mapsto W(X)$. Given a subring Λ of \mathbb{Q} , we set $\underline{W}_S^\Lambda = \underline{W}_S \otimes_{\mathbb{Z}} \Lambda$.

REMARK 5.3. — According to the Gersten conjecture for the Witt ring (known in the equicharacteristic case by [BGPW02]), it is well-known that for a regular scheme S over a field, the Witt sheaf \underline{W}_S agrees with the unramified Witt sheaf, i.e., the sheaf of unramified elements in the Witt ring of the fraction field (in other words, the 0-th cohomology of the Gersten–Witt complex). In fact, according to Corollary E.4, we

further obtain that for *any* regular scheme S , the Λ -linear Witt sheaf \underline{W}_S^Λ agrees with the Λ -linear unramified Witt sheaf, as long as $2 \in \Lambda^\times$.

REMARK 5.4. — Note that \underline{W}_S can also be described as the restriction to Sm_S of the sheaf \underline{W} associated with W on the big Zariski site of schemes of finite presentation over S . In particular, for any morphism of schemes $f : X \rightarrow S$ one gets a morphism of sheaves:

$$(5.4.a) \quad \underline{W}_S \longrightarrow f_*(\underline{W}_X),$$

or equivalently by adjunction:

$$(5.4.b) \quad f^*(\underline{W}_S) \longrightarrow \underline{W}_X.$$

REMARK 5.5. — The above definition will be meaningful for us only when S is regular. As will be clear from the next results, this is explained by the failure of the Gersten–Witt conjecture over singular schemes.

We will use the following basic lemma on Witt groups, for which we could not find a reference in the literature.

LEMMA 5.6. — *Let X be the limit of a cofiltered system of schemes $(X_i)_{i \in I}$ with a ne transition maps. Then the canonical maps:*

$$\varinjlim_{i \in I} W(X_i) \longrightarrow W(X)$$

are isomorphisms. Moreover, for any integer $n > 0$, the canonical maps

$$\varinjlim_{i \in I} H^n(X_i, \underline{W}_{X_i}) \longrightarrow H^n(X, \underline{W}_X)$$

are isomorphisms.

Proof. — Recall that, under these assumptions, the category of vector bundles over X is equivalent to the 2-colimit of the category of vector bundles over X_i ([Gro64, 8.5.3–8.5.5]). This implies the same result for symmetric bundles, by definition of the latter (Paragraph 5.1). Moreover, if V is a symmetric bundle over X , inverse image of a symmetric bundle V_i over X_i for some $i \in I$, then it is metabolic if and only if V_j is metabolic for some $j > i$. Thus the claim follows by definition of the Witt group. The second claim then follows using the fact that the Zariski topology is coherent.

THEOREM 5.7. — *Let S be a regular scheme over a prime field k of characteristic not 2. Then the following assertions hold:*

(1) *The Zariski sheaf \underline{W}_S is a Nisnevich sheaf and for any smooth S -scheme X , the following comparison map is an isomorphism:*

$$(5.7.a) \quad H_{\mathrm{Zar}}^*(X, \underline{W}_S) \simeq H_{\mathrm{Nis}}^*(X, \underline{W}_S).$$

Moreover, \underline{W}_S is strictly A^1 -local.⁽⁸⁾

⁽⁸⁾meaning that its Nisnevich cohomology presheaves over Sm_S are A^1 -homotopy invariant.

(2) The algebraic Hopf map $\eta : (G_{m,S}, 1) \rightarrow S_+$ induces an isomorphism:

$$(5.7.b) \quad \eta_* : (\underline{W}_S)_{-1} = \underline{\mathbf{Hom}}(\Lambda(G_m, 1), \underline{W}_S) \simeq \underline{W}_S,$$

where the internal Hom is taken in the effective A^1 -derived category.

Proof. — Consider Assertion (1). When S is smooth over k , the claim follows from [Pan10]. The case of an arbitrary regular scheme S can be obtained using Popescu’s theorem (Theorem C.2), the preceding lemma, and the fact that the smooth Zariski and Nisnevich toposes over S are coherent ([AGV73, Exp. VI, 2.3]). Alternatively, the reader can check that the arguments of [Pan10] actually work under our more general assumptions.

Then, the assertion (2) is a consequence of the localization exact sequence for triangular Witt groups, given the isomorphism $\Lambda(G_{m,S}, 1) \simeq \Lambda(\mathrm{Th}(A_S^1))[-1]$ in $D_{A^1}^{\mathrm{eff}}(S, \Lambda)$.⁽⁹⁾

5.8. — Let us now consider a regular \mathbb{Q} -scheme S and $\Lambda \subseteq \mathbb{Q}$ a subring such that $2 \in \Lambda^\times$. Recall that $D_{A^1}^{\mathrm{eff}}(S, \Lambda)$ can be seen as the full subcategory of the derived ∞ -category of Λ -linear Nisnevich sheaves over Sm_S spanned by A^1 -local complexes. In particular, under the assumption of the previous theorem, the Witt sheaf \underline{W}_S^Λ defines such a complex (concentrated in degree 0). Moreover, according to Theorem 5.7 (2), it is an infinite loop space with respect to $\mathbb{1}_S(1)[1]$. More precisely, there is an object $\mathbf{HW}_S^\Lambda \in D_{A^1}(S, \Lambda)$ defined by the G_m -spectrum given by the collection

$$(\underline{W}_S^\Lambda, \underline{W}_S^\Lambda, \dots)$$

together with the isomorphisms (5.7.b). We call \mathbf{HW}_S^Λ the Λ -linear homotopical Witt spectrum over S (see Corollary 5.14 for the justification of the adjective “homotopical”). By construction, this spectrum is η -periodic and satisfies:

$$\mathrm{Hom}_{D_{A^1}(S, \Lambda)}(\Lambda_S(X), \mathbf{HW}_S^\Lambda(i)[n]) = H_{\mathrm{Zar}}^{n-i}(X, \underline{W}_S) \otimes_{\mathbb{Z}} \Lambda.$$

The right-hand side is also the $(n - i)$ -th cohomology group of the Λ -linearized Gersten–Witt complex. The commutative monoid structure on \underline{W}_S^Λ extends to a commutative ring structure on the spectrum \mathbf{HW}_S^Λ , in such a way that the above isomorphism is compatible with products.

Note that it follows from Theorem 5.7 (2) that $\mathbf{HW}_{S,+}^\Lambda = 0$ or equivalently, that $\mathbf{HW}_{S,-}^\Lambda \simeq \mathbf{HW}_S^\Lambda$ is an object of $D_{A^1}(S, \Lambda)_-$.

The Key Lemma 2.2 allows us to define \mathbf{HW}_S^Λ in the mixed characteristic case:

DEFINITION 5.9. — Let S be an arbitrary regular scheme and $\Lambda \subseteq \mathbb{Q}$ a subring such that $2 \in \Lambda^\times$. We define the Λ -linear homotopical Witt ring spectrum over S as

$$\mathbf{HW}_S^\Lambda := \nu_* (\mathbf{HW}_{S_0}^\Lambda),$$

where $\nu : S_{\mathbb{Q}} \rightarrow S$ is the inclusion of the characteristic zero fiber as in Paragraph 2.1.

⁽⁹⁾We refer the interested reader to the arguments of [CF14, Lem. 3.8] for details.

REMARK 5.10. — While this definition may seem a bit artificial, the main point is that it allows us to bypass a missing part of the theory of higher Witt groups, which at the moment is not well-behaved over \mathbb{F}_2 -schemes (for example, the localization exact sequence is not known for such schemes). Let us first mention that a recent work, [CDH⁺20a, CDH⁺20b, CDH⁺20c], may correct the situation. Moreover, Proposition E.2 shows that one can extend the definition of \mathbf{HW}_S^Λ in Paragraph 5.8 to the case of S a regular $\mathbb{Z}[1/2]$ -scheme (with Λ still a subring of \mathbb{Q} such that $2 \in \Lambda^\times$). In particular, in this latter case, the ring spectrum \mathbf{HW}_S^Λ represents the cohomology of the Λ -linear Gersten–Witt complex (see Corollary E.4).

5.11. — Before moving to our main theorem, let us prove an intermediate lemma (which will be strengthened in Remark 5.20). Let $f : X \rightarrow S$ be an arbitrary morphism of regular schemes. Then there is a canonical morphism of ring spectra

$$\Theta_f : f^*(\mathbf{HW}_S^\Lambda) \longrightarrow \mathbf{HW}_X^\Lambda$$

in $\mathcal{D}_{\mathbb{A}^1}(X, \Lambda)$, induced by applying ν_* to the canonical map of \mathbb{G}_m -spectra over $X_{\mathbb{Q}}$

$$(f^*\underline{\mathbf{W}}_{S_{\mathbb{Q}}}^\Lambda, f^*\underline{\mathbf{W}}_{S_{\mathbb{Q}}}^\Lambda, \dots) \longrightarrow (\underline{\mathbf{W}}_{X_{\mathbb{Q}}}^\Lambda, \underline{\mathbf{W}}_{X_{\mathbb{Q}}}^\Lambda, \dots),$$

which in turn comes from the functoriality (5.4.b).

LEMMA 5.12. — *With notation as above, assume moreover that X is pro-smooth over S . Then Θ_f is an isomorphism.*

Proof. — By definition, we may assume that S is defined over \mathbb{Q} . Since f is pro-smooth, the functor f^* preserves \mathbb{A}^1 -locality of complexes of Λ -linear Nisnevich sheaves (this follows using continuity, see Appendix A, from the smooth case). Thus it suffices to note that the map (5.4.b) is an isomorphism of Nisnevich sheaves over Sm_X .

THEOREM 5.13. — *Let S be a regular scheme and $\Lambda \subseteq \mathbb{Q}$ with $2 \in \Lambda^\times$. Then the unit map of the ring spectrum \mathbf{HW}_S induces an isomorphism of ring spectra*

$$\Lambda_{S,-} \longrightarrow \mathbf{HW}_S^\Lambda$$

in $\mathcal{D}_{\mathbb{A}^1}(S, \Lambda)$.

Proof. — Again using the Key Lemma 2.2, and the fact that $\nu_*(\Lambda_{S_{\mathbb{Q}},-}) \simeq \Lambda_{S,-}$ (by monoidality, see Remark 2.4), we are reduced to the case where S is a regular \mathbb{Q} -scheme. By Zariski descent [CD19, 2.3.8] and Lemma 5.12, we may also assume that S is affine. By Popescu’s desingularization theorem (Theorem C.2), any such S is pro-smooth over \mathbb{Q} , so by Lemma 5.12 we reduce to $S = \mathrm{Spec} \mathbb{Q}$. Then the result follows from [Bac18, Prop. 41].

COROLLARY 5.14. — *Let S be a regular scheme and $\Lambda \subseteq \mathbb{Q}$ with $2 \in \Lambda^\times$. Then we have:*

(1) *There are canonical isomorphisms*

$$H_{\mathbb{A}^1}^{n,m}(S, \Lambda)_- \simeq H_{\text{Zar}}^{n-m}(S_{\mathbb{Q}}, \underline{\mathbb{W}}) \otimes_{\mathbb{Z}} \Lambda$$

for any integers $(n, m) \in \mathbb{Z}^2$, compatible with pullbacks and products.

(2) *The ring spectrum $\Lambda_{S,-} \simeq \mathbf{HW}_S^\Lambda$ is in the heart of the δ -homotopy t -structure on $\mathbf{D}_{\mathbb{A}^1}(S, \Lambda)$ for $\delta = -\text{codim}_S$ (see Paragraph 4.1).*

(3) *The canonical map $\mathbf{KW}_S^\Lambda \rightarrow \mathbf{HW}_S^\Lambda$ induces an isomorphism*

$$H_0^\delta(\mathbf{KW}_S^\Lambda) \longrightarrow \mathbf{HW}_S^\Lambda$$

where \mathbf{KW}_S^Λ is the spectrum representing the Λ -linearized higher Witt groups.

Proof. — The first isomorphism is immediate from Theorem 5.13. To get the second assertion, one uses the following computation given a point $x \in S$:

$$\widehat{H}_p^\delta(\Lambda_{S,-})(x, n) \simeq H_{\mathbb{A}^1}^{n-p,n}(\kappa(x), \Lambda)_- \simeq H_{\text{Zar}}^{-p}(\kappa(x), \underline{\mathbb{W}}) \otimes_{\mathbb{Z}} \Lambda$$

which is obtained using absolute purity for $\mathbb{1}_-$ (Theorem 3.6), as in [BD17, Ex. 3.2.8].⁽¹⁰⁾ Then Proposition 4.2 gives the assertion. Finally, the last assertion can be reduced to the case of (characteristic zero) fields. Then the δ -homotopy t -structure is just Morel’s homotopy t -structure ([BD17, Ex. 2.3.5(2)]), and the computation follows from [ALP17, Th. 1, Rem. 1].

REMARK 5.15. — Note further that one obtains:

$$\widehat{H}_0^\delta(\Lambda_{S,-})(x, n) \simeq W(\kappa(x))_\Lambda.$$

This isomorphism can be made canonical after tensoring with $\det(N_x(X_{(x)}))^\times$.

COROLLARY 5.16. — *Let S be a regular scheme and $\Lambda \subseteq \mathbb{Q}$ with $2 \in \Lambda^\times$. The canonical functor*

$$\mathbf{D}_{\mathbb{A}^1}(S, \Lambda)_- \longrightarrow \mathbf{D}_{\mathbb{A}^1}(S, \Lambda)$$

induces an equivalence of symmetric monoidal ∞ -categories

$$\mathbf{D}_{\mathbb{A}^1}(S, \Lambda)_- \simeq \mathbf{HW}_S^\Lambda\text{-mod}$$

where the right hand-side is the ∞ -category of modules over \mathbf{HW}_S^Λ , seen as a commutative algebra in the monoidal ∞ -category $\mathbf{D}_{\mathbb{A}^1}(S, \Lambda)$.

Moreover, the δ -homotopy t -structure on $\mathbf{D}_{\mathbb{A}^1}(S, \Lambda)_-$ (Paragraph 4.1) is non-degenerate and the unit lies in the heart (for $\delta = -\text{codim}_S$).

REMARK 5.17. — The objects of the category $\mathbf{HW}_S^\Lambda\text{-mod}$ are called *Witt motives* in [ALP17, §4]. One can obtain a somewhat more concrete description of this category along the lines of [ALP17]. Consider the symmetric monoidal Grothendieck abelian category $\text{Sh}_{\underline{\mathbb{W}}}(S)$ of modules over the sheaf $\underline{\mathbb{W}}_S^\Lambda$ in the category of Nisnevich sheaves. Then define the Λ -linear ∞ -category of *Witt motives* $\text{DM}_W(S, \Lambda)$ as the

⁽¹⁰⁾Note the isomorphism depends on a choice of a trivialization of the normal bundle of x in $X_{(x)}$.

A^1 -localization and \mathbb{P}^1 -stabilization of the derived ∞ -category $D(\mathrm{Sh}_{\underline{W}}(S))$. The proof of the identification $\mathrm{DM}_W(S, \Lambda) \simeq \mathbf{HW}_S^\Lambda\text{-mod}$ is analogous to the ones in [RØ08], [CD15], [BF18], [EHK⁺20] and [EK20b], where the details are left to the reader.⁽¹¹⁾

One can formally extend the preceding identification to any base.

DEFINITION 5.18. — Let S be an arbitrary scheme. Let $\Lambda \subseteq \mathbb{Q}$ be a subring with $2 \in \Lambda^\times$. We define the Λ -linear homotopical Witt spectrum $\mathbf{H}_W\Lambda_S \in D_{A^1}(S, \Lambda)$ as the inverse image $p^*(\mathbf{H}_W\Lambda_Z)$ along the projection $p : S \rightarrow \mathrm{Spec}(Z)$.

5.19. — By construction, $\mathbf{H}_W\Lambda_S$ defines an absolute ring spectrum (in the sense of Paragraph 3.5) as S varies. Note that Theorem 5.13 implies that the unit map of $\mathbf{H}_W\Lambda_S$ induces a canonical isomorphism

$$\mathbf{H}_W\Lambda_S \simeq \Lambda_{S,-},$$

and that there is a canonical equivalence

$$D_{A^1}(S, \Lambda)_- \simeq \mathbf{H}_W\Lambda_S\text{-mod},$$

over every base S . In particular, whenever S is regular, we get by Theorem 5.13 the canonical isomorphism

$$(5.19.a) \quad \mathbf{H}_W\Lambda_S \simeq \Lambda_{S,-} \simeq \mathbf{HW}_S^\Lambda,$$

thus recovering Definition 5.9.

REMARK 5.20. — For S regular, the identification of Definitions 5.9 and 5.18 through the isomorphism (5.19.a) means in particular that the motivic spectrum $\mathbf{H}_W\Lambda_S$ is induced by a \mathbb{G}_m -periodic and strictly A^1 -local sheaf of abelian groups. One cannot expect this to hold over singular schemes S , due to the failure of the Gersten–Witt conjecture (see Remark 5.5). In particular, over a singular base S with dimension function δ , it is no longer true that $\mathbf{H}_W\Lambda_S$ is concentrated in only one degree for the δ -homotopy t -structure.

6. RATIONAL STABLE HOMOTOPY AND MILNOR–WITT MOTIVES

Based on the initial definition of [BCD⁺20, Chap. 3 & 6] and the comparison result of [Gar19, 4.2], we introduce the following definition:

DEFINITION 6.1. — Let S be a scheme. One defines the *Milnor–Witt rational motivic cohomology spectrum* over S , denoted $\mathbf{H}_{\mathrm{MW}\mathbb{Q}_S} \in \mathrm{SH}(S)_{\mathbb{Q}}$, by the formula:

$$\mathbf{H}_{\mathrm{MW}\mathbb{Q}_S} = \mathbf{H}_{\mathrm{M}\mathbb{Q}_S} \oplus \mathbf{H}_{\mathrm{W}\mathbb{Q}_S},$$

where $\mathbf{H}_{\mathrm{M}\mathbb{Q}_S}$ is the Beilinson motivic cohomology spectrum (cf. [CD19, 14.1.2]) and $\mathbf{H}_{\mathrm{W}\mathbb{Q}_S}$ is the homotopical Witt spectrum over S (Definition 5.18). This is a motivic ring spectrum, which forms an absolute ring spectrum as S varies.

⁽¹¹⁾Hint: use the description of the spectrum associated with \underline{W}_S^Λ .

Recall that $\mathrm{SH}(S)_{\mathbb{Q}}$ is canonically identified with $\mathrm{D}_{\mathbb{A}^1}(S, \mathbb{Q})$, as symmetric monoidal ∞ -categories [CD19, 5.3.35]. An immediate corollary of Theorem 5.13 and [CD19, 14.2.14] is:

COROLLARY 6.2. — *For any scheme S , the unit map of the ring spectrum $\mathbf{H}_{\mathrm{MW}}\mathbb{Q}_S$ induces an isomorphism of ring spectra in $\mathrm{SH}(S)_{\mathbb{Q}}$*

$$\mathbb{1}_S \otimes \mathbb{Q} \simeq \mathbf{H}_{\mathrm{MW}}\mathbb{Q}_S$$

compatible with base change. In particular, for any regular scheme S , one obtains canonical isomorphisms compatible with pullbacks and products:

$$\mathrm{H}_{\mathbb{A}^1}^{n,i}(S, \mathbb{Q}) := \mathrm{Hom}_{\mathrm{SH}(S)_{\mathbb{Q}}}(\mathbb{1}_S, \mathbb{1}_S(i)[n]) \simeq K_{2i-n}^{(i)}(S) \oplus H_{\mathrm{Zar}}^{n-i}(S_{\mathbb{Q}}, \mathbb{W}) \otimes_{\mathbb{Z}} \mathbb{Q}$$

for all $(n, i) \in \mathbb{Z}^2$.

Note that this formula was conjectured by Morel (see discussion after Theorem 5.2.2 in [Mor04]). We can deduce formally the following extension of a result due to Garkusha (see [Gar19]).

COROLLARY 6.3. — *For any scheme S , there is a canonical equivalence of symmetric monoidal ∞ -categories:*

$$\mathrm{SH}(S)_{\mathbb{Q}} \simeq \mathbf{H}_{\mathrm{MW}}\mathbb{Q}_S\text{-mod.}$$

Moreover, the collection of these equivalences for various schemes S induces an equivalence of motivic ∞ -categories:

$$\mathrm{SH}_{\mathbb{Q}} \simeq \mathbf{H}_{\mathrm{MW}}\mathbb{Q}\text{-mod,}$$

which is compatible with the six operations.

7. SL-ORIENTATIONS

Recall that Panin and Walter have defined several notions that weaken the classical notion of orientation in motivic homotopy theory. We refer the reader to [Ana19] or [DF20, Def. 2.1.3]. In this section, we adapt their definition of SL-orientation to motivic categories, on the model of the usual notion of orientation (see [CD19, 2.4.38, 2.4.40]).

7.1. — We let \mathcal{T} be a motivic ∞ -category over a category of schemes \mathcal{S} . Let S be a scheme in \mathcal{S} . Recall that the functor $E \mapsto \mathrm{Th}_S(E)$, sending a vector bundle E to its Thom space in $\mathcal{T}(S)$, can be extended to a monoidal functor:

$$\mathrm{Th}_S : \underline{\mathbf{K}}(S) \longrightarrow \mathrm{Pic}(\mathcal{T}(S)), \quad v \longmapsto \mathrm{Th}_S(v).$$

from the Picard groupoid $\mathrm{Pic}(\mathcal{T}(S)) = \mathrm{Pic}(\mathcal{T}(S), \otimes)$ of virtual vector bundles over S [Del87] to that of \otimes -invertible objects of $\mathcal{T}(S)$. Moreover, this assignment is functorial with respect to pullbacks, giving a natural transformation $\mathrm{Th} : \underline{\mathbf{K}} \rightarrow \mathrm{Pic}(\mathcal{T})$.

7.2. — An SL-orientation of a virtual vector bundle v over S is an isomorphism

$$\lambda : \det(v) \simeq A_S^1;$$

we refer to the pair (v, λ) as an SL-oriented vector bundle. The sum of SL-oriented bundles is given by

$$(v, \lambda) + (v', \lambda') = (v + v', \lambda \otimes \lambda').$$

The notion of isomorphism of SL-oriented virtual bundles being clear, one gets a Picard groupoid $\underline{\mathbf{K}}^{\text{SL}}(S)$ together with a (symmetric) monoidal forgetful functor $\psi_S : \underline{\mathbf{K}}^{\text{SL}}(S) \rightarrow \underline{\mathbf{K}}(S)$.⁽¹²⁾ This is obviously functorial and defines a natural transformation $\psi : \underline{\mathbf{K}}^{\text{SL}} \rightarrow \underline{\mathbf{K}}$.

Consider the canonical functor:

$$\langle - \rangle_S : \underline{\mathbf{Z}}(S) \longrightarrow \text{Pic}(\mathcal{T}(S)), \quad r \longmapsto \text{Th}_S(A_S^r) = \mathbb{1}_S \langle r \rangle,$$

where $\underline{\mathbf{Z}}(S)$ is the Picard groupoid of locally constant integral functions on S . As S varies this defines a natural transformation $\underline{\mathbf{Z}} \rightarrow \text{Pic}(\mathcal{T})$.

DEFINITION 7.3. — An SL-orientation of the motivic ∞ -category \mathcal{T} is a natural transformation \mathfrak{t} of presheaves over \mathcal{S} in monoidal categories, pictured as a 2-arrow in the following diagram:

$$\begin{array}{ccc} \underline{\mathbf{K}}^{\text{SL}} & \xrightarrow{\text{Th} \circ f} & \overline{\text{Pic}}(\mathcal{T}) \\ \text{rk} \Big\downarrow & \parallel \mathfrak{t} & \\ \underline{\mathbf{Z}} & \xrightarrow{\langle - \rangle} & \end{array}$$

such that for any scheme S , $\mathfrak{t}(A_S^1)$ is the identity of $\text{Th}(A_S^1)$ (where A^1 is endowed with its tautological orientation). We refer to the pair $(\mathcal{T}, \mathfrak{t})$ as an SL-oriented motivic ∞ -category.

REMARK 7.4. — Explicitly, an SL-orientation on \mathcal{T} amounts to the data of, for any scheme S and any SL-oriented virtual bundle (v, λ) over S , a natural isomorphism:

$$\mathfrak{t}(v, \lambda) : \text{Th}_S(v) \longrightarrow \mathbb{1}_S \langle \text{rk } v \rangle.$$

These isomorphisms are compatible with pullbacks and products.

EXAMPLE 7.5. — Clearly a classical orientation of \mathcal{T} induces by restriction an SL-orientation. In fact, a classical orientation can be formulated just as in Definition 7.3, except that the upper horizontal arrow is replaced by $\text{Th} : \underline{\mathbf{K}} \rightarrow \text{Pic}(\mathcal{T})$.

REMARK 7.6. — Recall that the map $\underline{\mathbf{K}} \rightarrow \text{Pic}(\mathcal{T})$ lifts to an \mathcal{E}_∞ -map from the Thomason–Trobaugh K-theory space \mathbf{K} to the Picard ∞ -groupoid of \otimes -invertible objects in \mathcal{T} (see e.g. [BH21, §16.2]). Let \mathbf{K}^{SL} denote the SL-oriented (higher) K-theory presheaf as in [BH21, Ex. 16.20]. There is again a forgetful map $\mathbf{K}^{\text{SL}} \rightarrow \mathbf{K}$, which we can use to formulate a stronger version of Definition 7.3. This amounts to the existence of a homotopy coherent system of compatibilities between the isomorphisms in

⁽¹²⁾One can also build this category from the exact category of SL-torsors.

Remark 7.3. We do not discuss this notion further, since we do not know whether the orientation of Theorem 7.8 is homotopy coherent. We note however that the orientation on the motivic ∞ -category of **MSL**-modules is homotopy coherent, as follows from the motivic Thom spectrum description of **MSL** given in [BH21, §16].

7.7. — We compare our definition with the original one of Panin and Walter. By universality of SH among motivic ∞ -categories [Rob15], there is an (essentially unique) adjunction of motivic ∞ -categories

$$\varphi^* : \mathrm{SH} \xrightarrow{\mathbb{O}} \mathcal{T} : \varphi_*$$

Let $H^{n,i}(S, \mathcal{T}) = \mathrm{Hom}_{\mathcal{T}(S)}(\mathbb{1}_S, \mathbb{1}_S(i)[n])$ be the *fundamental cohomology* associated with \mathcal{T} . For any scheme S in \mathcal{S} , $H^{n,i}(-, \mathcal{T})$ is represented over S by the motivic ring spectrum $\mathbf{H}_S^{\mathcal{T}} = \varphi_*(\mathbb{1}_S) \in \mathrm{SH}(S)$. The collection $(\mathbf{H}_S^{\mathcal{T}})_{S \in \mathcal{S}}$ form a section of SH over \mathcal{S} .⁽¹³⁾

Then it is easy to check that an SL-orientation \mathfrak{t} of \mathcal{T} is equivalent to a family of SL-orientations \mathfrak{t}_S^{PW} of the ring spectra $\mathbf{H}_S^{\mathcal{T}}$ in the sense of Panin–Walter, for all schemes S in \mathcal{S} , such that for any morphism of schemes $f : T \rightarrow S$, the canonical morphism:

$$f^*(\mathbf{H}_S^{\mathcal{T}}) \longrightarrow \mathbf{H}_T^{\mathcal{T}}$$

is compatible with the respective SL-orientations. Moreover, if \mathcal{S} admits a final object Σ , then an SL-orientation of \mathcal{T} is uniquely determined by an SL-orientation of $\mathbf{H}_{\Sigma}^{\mathcal{T}}$ (by functoriality of \mathfrak{t}).

THEOREM 7.8. — *The motivic ∞ -category $\mathrm{D}_{\mathbb{A}^1}(-, \mathbb{Z})_-$ admits a unique SL-orientation.*

Proof. — According to the preceding paragraph it is sufficient to give an SL-orientation of the ring spectrum $\mathbb{1}_{S,-}$, when $S = \mathrm{Spec} \mathbb{Z}[1/2]$. Then the result follows from [Ana19, Cor. 5.4] and Corollary 5.14: indeed, according to the latter, $H_{\mathbb{A}^1}^{0,0}(-, \mathbb{Z}[1/2])_- \simeq \underline{W}_S^{\mathbb{Z}[1/2]}$ is a Zariski sheaf over Sm_S .

As a corollary, we get the analog of the classical fact in topology that rational spectra are all orientable.

COROLLARY 7.9. — *The rational stable homotopy category $\mathrm{SH}_{\mathbb{Q}}$ admits a canonical SL-orientation.*

This follows from the decomposition $\mathrm{SH}_{\mathbb{Q}} = \mathrm{DM}_{\mathbb{B}} \oplus \mathrm{D}_{\mathbb{A}^1}(-, \mathbb{Q})_-$. Note that the orientation is unique if one asks in addition that its projection on the plus-part is induced by the additive orientation on Beilinson motivic cohomology $\mathbf{H}_{\mathbb{M}}\mathbb{Q}$.

⁽¹³⁾Note that this section is not cartesian in general. However, it will be in the two cases considered below.

7.10. — In the following discussion we consider an arbitrary SL-oriented motivic ∞ -category $(\mathcal{T}, \mathfrak{t})$ over \mathcal{S} , but the reader may focus on the case of $D_{\Lambda^1}(-, \Lambda)_-$ or $\mathrm{SH}_{\mathbb{Q}}$.

Recall that one can define following Deligne [Del87] the determinant functor

$$\widetilde{\mathrm{det}}_S : \underline{\mathbf{K}}(S) \longrightarrow \underline{\mathbf{Pic}}(S), \quad v \longmapsto (\mathrm{det}(v), \mathrm{rk}(v))$$

with values in the Picard groupoid of graded line bundles over S . This is again contravariantly functorial in S .

Motivated by the following definition, and by the theory of Chow–Witt groups, we consider the following twisting functor:

$$\begin{aligned} \mathrm{Tw}_S : \underline{\mathbf{Pic}}(S) &\longrightarrow \mathcal{T}(S) \\ (L, r) &\longmapsto \mathrm{Th}(L \oplus \mathbb{A}_S^{r-1}) = \mathrm{Th}(L)\langle r-1 \rangle. \end{aligned}$$

We will also write simply $\mathrm{Tw}_S(L) = \mathrm{Tw}_S(L, 0)$.

Let v be a virtual vector bundle of rank r over a scheme S in \mathcal{S} . Then the rank $r-1$ virtual bundle $v - \mathrm{det} v$ admits an obvious SL-orientation $\lambda : \mathrm{det}(v - \mathrm{det} v) \simeq \mathrm{det} v \otimes (\mathrm{det} v)^{-1} \simeq \mathbb{A}_S^1$.

DEFINITION 7.11. — Given a virtual vector bundle v over a scheme S in \mathcal{S} , we define the *twisted Thom class* associated with \mathfrak{t} as the following map $\mathfrak{t}(v) : \mathrm{Th}_S(v) \rightarrow \mathrm{Tw}_S(\mathrm{det} v)\langle \mathrm{rk}(v) \rangle$ defined as the composite:

$$\begin{aligned} \mathrm{Th}_S(v) \simeq \mathrm{Th}_S(\mathrm{det} v) \otimes \mathrm{Th}_S(v - \mathrm{det} v) &\xrightarrow{1 \otimes \mathfrak{t}(v - \mathrm{det} v, \lambda)} \mathrm{Tw}_S(\widetilde{\mathrm{det}} v) \\ &= \mathrm{Tw}_S(\mathrm{det} v, \mathrm{rk}(v)). \end{aligned}$$

REMARK 7.12. — Note in particular that to give an SL-orientation on \mathcal{T} in the sense of Definition 7.3 is equivalent to give a natural transformation as a 2-arrow in the following diagram:

$$\begin{array}{ccc} \underline{\mathbf{K}} & \xrightarrow{\quad \mathrm{Th} \quad} & \underline{\mathbf{Pic}}(\mathcal{T}) \\ \widetilde{\mathrm{det}} \Big| & \parallel \mathfrak{t} & \\ \underline{\mathbf{Pic}} & \xrightarrow{\quad \mathrm{Tw} \quad} & \end{array}$$

Note in particular that the twisted Thom class is multiplicative, roughly expressed by the relation $\mathfrak{t}(v+w) = \mathfrak{t}(v) \otimes \mathfrak{t}(w)$, using the fact Tw_S is monoidal (see below).

7.13. — Let \mathcal{T} be an SL-oriented motivic ∞ -category. Given line bundles L and M over S , there is an induced isomorphism:

$$\mathrm{Tw}(L) \otimes \mathrm{Tw}(M) = \mathrm{Th}(L \oplus M)\langle -2 \rangle \xrightarrow{\mathfrak{t}(L \oplus M)} \mathrm{Tw}(\mathrm{det}(L \oplus M)) \simeq \mathrm{Tw}(L \otimes M).$$

One deduces from this isomorphism that the twisting functor Tw_S has a canonical structure of a monoidal functor:

$$\mathrm{Tw}_S : \underline{\mathbf{Pic}}(S) \longrightarrow \underline{\mathbf{Pic}}(\mathcal{T}(S)).$$

According to a remark of Ananyevskiy (see [Ana19, Lem. 4.1]), one can go further. Indeed, for any line bundle L over S , one can define a morphism of S -schemes

$\phi_L : L^\times \rightarrow (L^\vee)^\times$ defined locally by sending a non-zero vector u to the unique function f such that $f(u) = 1$. The morphism of schemes ϕ_L is an isomorphism, so that, by definition of the Thom space, one gets a canonical isomorphism in $\mathcal{T}(S)$:

$$\mathrm{Th}(L) \simeq \mathrm{Th}(L^\vee).$$

Using the monoidal structure on Tw , one gets isomorphisms

$$\mathrm{Tw}(L) \longrightarrow \mathrm{Tw}(L^\vee) \simeq \mathrm{Tw}(L)^{\otimes, -1} \iff \mathrm{Tw}(L^{\otimes, 2}) \simeq \mathbb{1}_S.$$

In other words, the twisting functor Tw_S can be factorized and induces a functor from the Picard groupoid of S modulo squares.⁽¹⁴⁾

The following result is obtained by combining the construction of [DJK21], Theorem 3.6, Theorem 7.8 and the preceding construction.

THEOREM 7.14. — *Let \mathcal{T} be one of the motivic ∞ -categories $\mathrm{D}_{\mathbb{A}^1}(-, Z)_-$ or $\mathrm{SH}_{\mathbb{Q}}$. Then for any smoothable lci morphism $f : X \rightarrow S$, with graded determinant $\tilde{\omega}_f = \widetilde{\det}(L_f)$, there exists a fundamental class*

$$\eta_f : \mathrm{Tw}_X(\tilde{\omega}_f) \longrightarrow f^!(\mathbb{1}_S)$$

which is an isomorphism whenever X and S are regular, or f is smooth. Moreover, the classes η_f are compatible with composition, transversal pullbacks, and satisfy an excess intersection formula (see [DJK21] for the precise formulations).

REMARK 7.15. — As in [DJK21], Theorem 7.14 gives rise to trace maps and exceptional functoriality (Gysin or wrong-way maps) for the four homology/cohomology theories associated with any spectrum \mathbb{E} in $\mathrm{D}_{\mathbb{A}^1}_-$ or $\mathrm{SH}_{\mathbb{Q}}$. The only thing that changes is the new twists involved. We will illustrate this in the forthcoming section.

8. BIVARIANT \mathbb{A}^1 -THEORY AND CHOW–WITT GROUPS

8.1. — In this section, we will use our results to compute certain \mathbb{A}^1 -homology groups. We will use two kinds of ring spectra as coefficients, which naturally correspond to motivic ∞ -categories:

- $R = \mathrm{H}_{\mathbb{A}^1}Z_-$, corresponding to $\mathcal{T} = \mathrm{D}_{\mathbb{A}^1}(-, Z)_-$;
- $R = \mathrm{H}_{\mathbb{A}^1}\mathbb{Q}$, corresponding to $\mathcal{T} = \mathrm{D}_{\mathbb{A}^1}(-, \mathbb{Q}) \simeq \mathrm{SH}_{\mathbb{Q}}$.

DEFINITION 8.2. — Let R and \mathcal{T} be as above. Let X be a scheme, L a line bundle over X , and $(n, i) \in \mathbb{Z}^2$ a pair of integers. We define the L -th twisted cohomology of X in degree (n, i) and coefficients in R as:

$$\mathrm{H}_{\mathbb{A}^1}^{n,i}(X, R(L)) = \mathrm{Hom}_{\mathcal{T}(S)}(R_X, R_X \otimes \mathrm{Tw}(L)(i)[n]).$$

⁽¹⁴⁾This almost shows that the SL -orientation \mathfrak{t} corresponds to an SL^c -orientation in the terminology of Panin. See [Ana19, Rem. 4.4] for more discussion.

Given a morphism of $Z[1/2]$ -schemes $f : X \rightarrow S$ that is essentially of finite presentation (Definition B.2), we define the L -th twist of the bivariant theory of X/S in degree (n, i) and coefficients in R as:

$$H_{n,i}^{\mathbb{A}^1}(X/S, R(L)) = \text{Hom}_{\mathcal{T}(S)}(R_X \otimes \text{Tw}(L)(i)[n], f^!R_X),$$

where $f^!$ is as defined in Paragraph B.13.

Taking into account the continuity property of \mathcal{T} , one gets the following duality theorem as a corollary of Theorem 7.14.⁽¹⁵⁾

COROLLARY 8.3. — *Let R and \mathcal{T} be as above. Then for any smoothable morphism $f : X \rightarrow S$ between regular schemes, of relative dimension d and determinant ω_f , any $(n, i) \in \mathbb{Z}^2$ and any line bundle L/X , one has a canonical isomorphism:*

$$H_{n,i}^{\mathbb{A}^1}(X/S, R(L)) \simeq H_{\mathbb{A}^1}^{2d-n, d-i}(X, R(\omega_f - L)).$$

8.4. δ -niveau spectral sequences. — Now we study the *niveau spectral sequences* in motivic homotopy theory, along the lines of [BD17, §3.1]. Let us start by fixing our conventions for the next results:

- All schemes in this section will be assumed noetherian and of finite dimension.
- S is a regular noetherian scheme and $\delta = -\text{codim}_S$ is the canonical dimension function.
- X is an essentially of finite type over S . We denote the induced dimension function on X (see [BD17, 1.1.7]) again by δ . Since it is also assumed noetherian and finite-dimensional, δ is bounded.
- *Coefficients.* $R = H_{\mathbb{A}^1}\mathbb{Q}$ or $R = H_{\mathbb{A}^1}\mathbb{Z}$.
- *Twists.* L/X is a line bundle, $n \in \mathbb{Z}$.

The δ -niveau spectral sequence of X/S associated with R_S in $\mathcal{T}(S)$ has the following form:

$${}^\delta E_{p,q}^1 = \bigoplus_{x \in X_{(p)}} H_{p+q,n}^{\mathbb{A}^1}(\kappa(x)/S, R(L_x)) \implies H_{p+q,n}^{\mathbb{A}^1}(X/S, R(L)),$$

where $X_{(p)} = \{x \in X \mid \delta(x) = p\}$, for $x \in X$, $\kappa(x)$ is the residue field and L_x is the fiber of L at x . Given our assumptions, one can apply the previous corollary to the morphism of schemes $\text{Spec } \kappa(x) \rightarrow S$; note that it is of relative dimension $\delta(x)$, and denote by ω_x its determinant. This gives, for $x \in X_{(p)}$, a canonical isomorphism:

$$H_{p+q,n}^{\mathbb{A}^1}(\kappa(x)/S, R(L_x)) \simeq H_{\mathbb{A}^1}^{p-q, p-n}(\kappa(x), R(\omega_x - L_x))$$

Taking into account the computation of (5.14), and the known vanishing of Beilinson motivic cohomology of fields, one obtains that the previous spectral sequence is concentrated in one of the regions: $(p > n, q > n)$ or $(p \in [\delta_-(X), \delta_+(X)], q = n)$, where $\delta_-(X)$ (resp. $\delta_+(X)$) is the infimum (resp. supremum) of values of δ over X .

⁽¹⁵⁾This formula is obviously reminiscent to the duality of Grothendieck–Serre for coherent sheaves.

Moreover, using Morel’s computations of the graded stable homotopy groups of the zero sphere over fields as Milnor–Witt K-theory (cf. [Mor12]), one can compute the terms of the complex located on the line $q = n$ as (graded by the index p):

$$\dots \longrightarrow \bigoplus_{x \in X^{(p)}} M_{p-n}(x, \omega_x \otimes L_x^\vee) \xrightarrow{d_{p-1}} \bigoplus_{y \in X^{(p-1)}} M_{p-n-1}(y, \omega_y \otimes L_y^\vee) \dots$$

where we have put, without twists by a line bundle:

- if $R = \mathbb{H}_{\mathbb{A}^1} \mathbb{Q}$, $M_r(x) := \mathbf{K}_r^{\text{MW}}(\kappa(x))_{\mathbb{Q}}$;
- if $R = \mathbb{H}_{\mathbb{A}^1} \mathbb{Z}_-$, $M_r(x) := W(\kappa(x))[1/2]$;

and with twists:

$$M_r(x, \omega_x \otimes L_x^\vee) := M_r(x) \otimes_{Z[\kappa(x)^\times]} R[(\omega_x \otimes L_x^\vee)^\times].$$

DEFINITION 8.5. — We define the (L, n) -th twisted Gersten complex of X with coefficients in $M = \mathbf{K}_{*,\mathbb{Q}}^{\text{MW}}$ (resp. $M = W[1/2]$), with respect to the dimension function δ , as the complex just defined above for $R = \mathbb{H}_{\mathbb{A}^1} \mathbb{Q}$ (resp. $R = \mathbb{H}_{\mathbb{A}^1} \mathbb{Z}_-$). We denote it by $C_*^\delta(X, M(L, n))$, and by $A_p^\delta(X, M(L, n))$ its homology in degree p . We define the \mathbb{Q} -linear n -th Chow–Witt group of X twisted by L and graded by δ as:

$$\widetilde{\text{CH}}_{\delta=n}(X, L)_{\mathbb{Q}} := A_p^\delta(X, \mathbf{K}_*^{\text{MW}}(L, n)_{\mathbb{Q}}).$$

8.6. — Let us consider the case $R = \mathbb{H}_{\mathbb{A}^1} \mathbb{Q}$. By construction, the above Gersten complex in degrees around n is of the form:

$$\bigoplus_{\eta \in X_{(n+1)}} \mathbf{K}_1^{\text{MW}}(\eta, \omega_\eta \otimes L_\eta^\vee)_{\mathbb{Q}} \xrightarrow{d_n} \bigoplus_{x \in X_{(n)}} GW(x, \omega_x \otimes L_x^\vee)_{\mathbb{Q}} \longrightarrow \bigoplus_{s \in X_{(n-1)}} W(s, \omega_s \otimes L_s^\vee)_{\mathbb{Q}}$$

so that $\widetilde{\text{CH}}_{\delta=n}(X, L)_R$ is a sub-quotient of cycles on X with coefficients in quadratic forms and δ -dimension n . Using the theory of [Fel19], one can check that the differentials have the expected form, of the so-called *Rost-Schmid complex*,⁽¹⁶⁾ so that this definition extends the previously known one of [Fas08] when $X = S$ is regular. In the case where S is a smooth k -scheme, the reader can find the relevant result in [BCD⁺20, Chap. 6, Lem. 4.2.3].

Note also that in degree $p > n$, the Gersten complexes with coefficients in $(\mathbf{K}_*^{\text{MW}})_{\mathbb{Q}}$ and in $W_{\mathbb{Q}}$ agree. In the end, using the vanishing of the δ -niveau spectral sequence as explained in the paragraph preceding the definition, one gets:

THEOREM 8.7. — *Let S be a regular scheme, $\delta = -\text{codim}_S$ and X an essentially of finite type S -scheme. Then one gets canonical isomorphisms:*

$$\mathbb{H}_{r,n}^{\mathbb{A}^1}(X/S, \mathbb{Q}(L)) \simeq \begin{cases} \widetilde{\text{CH}}_{\delta=n}(X, L)_{\mathbb{Q}} & r = 2n, \\ A_{n-1}^\delta(X, \mathbf{K}^{\text{MW}}(L, n)_{\mathbb{Q}}) & r = 2n - 1, \\ A_i^\delta(X, \underline{W}(L)_{\mathbb{Q}}) & r = n + i, \ i > n. \end{cases}$$

⁽¹⁶⁾See [Fel19, §3.3] stated in the case where $X = S$ is a smooth scheme over a field. The general case works similarly.

REMARK 8.8. — The δ -niveau spectral filtration is well-known to be functorial in the S -scheme X , covariantly with respect to proper maps, contravariantly with respect to étale maps. With a bit more work, as in [Jin16], one can show that the above δ -niveau spectral sequence, and therefore the Gersten complex $C_*^\delta(X, \mathbf{K}^{\text{MW}}(L, n))$ satisfies the same functoriality. One then obtains that the above isomorphisms are natural with respect to that functoriality on the right hand-side and the usual functoriality of bivariant theories (as in [DJK21] in the motivic case).

With a little more efforts, one can get the contravariant functoriality in X/S of the Gersten complex with respect to smoothable lci maps as in Feld's theory of Milnor–Witt modules (see [Fel20]). The above isomorphisms are again compatible with this functoriality, where on the left hand-side one uses the Gysin maps of [DJK21].

It is interesting to rewrite the case $X = S$.

COROLLARY 8.9. — *Let X be a regular scheme. Then one has canonical isomorphisms:*

$$\begin{aligned} \text{Hom}_{\text{SH}(X)}(\mathbb{1}_X, \text{Tw}_X(L)(n)[2n]) \otimes_{\mathbb{Z}} \mathbb{Q} &\simeq \widetilde{\text{CH}}^n(X, L)_{\mathbb{Q}} \\ &\simeq \text{CH}^n(X)_{\mathbb{Q}} \oplus H_{\text{Zar}}^n(X_{\mathbb{Q}}, \underline{W}(L))_{\mathbb{Q}}, \end{aligned}$$

where $\widetilde{\text{CH}}^n(X, L)$ denotes Chow–Witt groups in codimension n with rational coefficients, and $\underline{W}(L)$ denotes the L -twisted Witt sheaf.

Note also that in the case $R = H_{\mathbb{A}^1} Z_-$ the spectral sequence degenerate at E_2 and gives the following:

THEOREM 8.10. — *Let S be a regular scheme, $\delta = -\text{codim}_S$ and X an essentially of finite type S -scheme. Then one gets a canonical isomorphism:*

$$\text{H}_{r,n}^{\mathbb{A}^1}(X/S, Z_-(L)) \simeq A_{r-n}^{\delta}(X_{\mathbb{Q}}, W(L, n)[1/2]).$$

REMARK 8.11. — The Gersten complex of $X_{\mathbb{Q}}$ with coefficients in $W(L, n)[1/2]$ can be identified with Gille's version of the Gersten–Witt complex associated with the dualizing sheaf $f^{-1}(\omega_S) \otimes L$ as defined in [Gil07].

8.12. — We give here an application to the study of weight structures on motivic categories. Let \mathcal{T} be a motivic ∞ -category and let S be a scheme. Recall that the *Chow weight structure* on $\mathcal{T}(S)$, if it exists, is the unique weight structure on the underlying motivic triangulated category whose heart agrees with the idempotent completion of the localizing subcategory generated by objects of the form $f_* \mathbb{1}_X(r)[2r]$, where $r \in \mathbb{Z}$ and $f : X \rightarrow S$ is a proper morphism with X regular (see [Bon14, Th. 2.1]).

We consider the question of the existence of the Chow weight structure on $\mathcal{T} = \text{SH}_{\mathbb{Q}}$. Using de Jong's alterations, if S is separated of finite type over an excellent scheme of dimension ≤ 2 , and -1 is a sum of squares in any residue field of S , then $\text{SH}(S)_{\mathbb{Q}} \simeq \text{DM}(S)_{\mathbb{Q}}$ has a Chow weight structure by [Héb11] and [Bon14]. In what follows we show that the converse is almost true, provided that a general version of resolution of singularities is available.

LEMMA 8.13. — *Let $X_0 \rightarrow S_1$ be a proper, dominant, generically finite morphism between integral schemes of characteristic 0, with X_0 regular. Assume that the function field κ of X_0 is formally real. Then the group $H_{0,0}^{A^1}((X_0 \times_{S_1} X_0)/X_0, \mathbb{Q}_-)$ is non-vanishing.*

Proof. — Denote $T = X_0 \times_{S_1} X_0$. By Theorem 8.10, the group $H_{0,0}^{A^1}(T/X_0, \mathbb{Q}_-)$ agrees with the kernel of the first differential in the Gersten-Witt complex

$$H_{0,0}^{A^1}(T/X_0, \mathbb{Q}_-) = \ker \left(\bigoplus_{x \in T^{(0)}} W(\kappa(x))_{\mathbb{Q}} \longrightarrow \bigoplus_{x \in T^{(1)}} W(\kappa(x), \omega_x)_{\mathbb{Q}} \right).$$

The residue fields of the generic points of T are all isomorphic to κ , and the group $W(\kappa)_{\mathbb{Q}}$ is non-vanishing by [Lam05, VIII 6.15]. This implies that the group $H_{0,0}^{A^1}(T/X_0, \mathbb{Q}_-)$ is non-vanishing: indeed, let η be a generic point of T , then the structure map

$$\mathbb{Q} \simeq W(\mathbb{Q})_{\mathbb{Q}} \longrightarrow W(\kappa(\eta))_{\mathbb{Q}}$$

factors through the kernel $H_{0,0}^{A^1}(T/X_0, \mathbb{Q}_-)$, and for an ordering σ of $\kappa(\eta)$ the composition with the signature map on $\kappa(\eta)$

$$\mathbb{Q} \simeq W(\mathbb{Q})_{\mathbb{Q}} \longrightarrow H_{0,0}^{A^1}(T/X_0, \mathbb{Q}_-) \longrightarrow \mathbb{Q}$$

agrees with the signature on \mathbb{Q} induced by σ , which shows that $H_{0,0}^{A^1}(T/X_0, \mathbb{Q}_-)$ is non-zero.

COROLLARY 8.14. — *Let S be a scheme and let s be a point of S . Assume that there exists a proper morphism $f : X \rightarrow S$ with X regular, such that*

- *the induced map between the characteristic 0 fibers $X_0 \rightarrow S_0$ factors through S_1 , the reduced closure of s in S_0 ;*
- *the morphism $X_0 \rightarrow S_1$ satisfies the condition in Lemma 8.13.⁽¹⁷⁾*

Then there cannot exist a Chow weight structure on $\mathrm{SH}(S)_{\mathbb{Q}}$.

Proof. — If the Chow weight structure exists, then the objects $f_* \mathbb{1}_X(r)[2r]$ are negative in the sense of [Bon14, Def. 1.5 VII]. In particular this means that the groups $H_{2r-n,r}^{A^1}((X \times_S X)/X, \mathbb{Q}_-) \simeq H_{2r-n,r}^{A^1}((X_0 \times_{S_1} X_0)/X_0, \mathbb{Q}_-)$ vanish for all $n > 0$. Since the spectrum $H_{A^1} Z_-$ is \mathbb{G}_m -periodic by Theorem 5.7, we have in particular $H_{0,0}^{A^1}((X \times_S X)/X, \mathbb{Q}_-) = 0$, which contradicts Lemma 8.13.

REMARK 8.15

(1) The criterion of Corollary 8.14 is to be understood as follows: if s is a formally real point of S , let S_0 be the characteristic 0 fiber of S and let S_1 be the closure of s in S_0 . Then Hironaka’s resolution of singularities says that there exists a proper birational morphism $X_0 \rightarrow S_1$ with X_0 regular. Corollary 8.14 states that if X_0 has a regular model over the whole S , then there is no Chow weight structure on $\mathrm{SH}(S)_{\mathbb{Q}}$.

⁽¹⁷⁾Note that these assumptions imply that the residue field of s is formally real.

(2) The condition in Corollary 8.14 holds when S has equal characteristic, or when S is the spectrum of a Dedekind domain (see [Sch00, 1.2.2]). The condition is almost a de Jong alteration, except that we require the morphism to be generically finite of *odd* degree, since a finite field extension of odd degree of a formally real field remains formally real.

APPENDIX A. CONTINUITY IN MOTIVIC ∞ -CATEGORIES

SETUP A.1. — Suppose given a cofiltered system $(S_\alpha)_\alpha$ of schemes with affine transition maps. We write S for its limit, $u_\alpha : S \rightarrow S_\alpha$ for the projection map, and $u_{\alpha,\beta} : S_\beta \rightarrow S_\alpha$ for the transition map induced by a morphism $\alpha \rightarrow \beta$ in the indexing category. The maps are labeled as follows:

$$\begin{array}{ccc} S & \xrightarrow{u_\beta} & S_\beta \\ & \searrow u_\alpha & \downarrow u_{\alpha,\beta} \\ & & S_\alpha \end{array}$$

DEFINITION A.2. — Let \mathcal{T} be a motivic ∞ -category. For every cofiltered system $(S_\alpha)_\alpha$ as in Setup A.1, consider the canonical functor

$$\varinjlim_\alpha \mathcal{T}(S_\alpha) \longrightarrow \mathcal{T}(S),$$

where the colimit is taken in the ∞ -category of presentable ∞ -categories and left-adjoint functors [Lur09, Def. 5.5.3.1], and the transition maps are the inverse image functors $u_{\alpha,\beta}^*$. We say that \mathcal{T} is *continuous* if this functor is an equivalence for all such systems $(S_\alpha)_\alpha$.

A.3. — By [Lur09, Cor. 5.5.3.4], the condition in Definition A.2 is equivalent to the assertion that the canonical functor

$$\mathcal{T}(S) \longrightarrow \varprojlim_\alpha \mathcal{T}(S_\alpha)$$

is an equivalence, where the limit is taken in the ∞ -category of presentable ∞ -categories and *right*-adjoint functors, and the transition maps are the direct image functors $(u_{\alpha,\beta})_*$. By [Lur09, Th. 5.5.3.18], this limit can be taken equivalently in the ∞ -category of (large) ∞ -categories.

In particular, the space of maps between any two objects $E, F \in \mathcal{T}(S)$ can be computed by the limit

$$\mathrm{Maps}_{\mathrm{SH}(S)}(E, F) \simeq \varprojlim_\alpha \mathrm{Maps}_{\mathrm{SH}(S_\alpha)}((u_\alpha)_*(E), (u_\alpha)_*(F)).$$

EXAMPLE A.4. — The motivic ∞ -category SH is continuous [Hoy14, Prop. C.12(4)]. We immediately deduce the same for $\mathrm{SH}[1/2]$, SH_+ , SH_- , $\mathrm{SH}_\mathbb{Q}$, $\mathrm{D}_{\mathrm{A}^1}(-, \Lambda)$ (for any commutative ring Λ), etc.

DEFINITION A.5. — We say that \mathcal{T} is *compactly generated* if the stable ∞ -category $\mathcal{T}(S)$ is compactly generated for every scheme S [Lur09, Def. 5.5.7.1], and for any morphism $f : S \rightarrow T$, the inverse image functor $f^* : \mathcal{T}(T) \rightarrow \mathcal{T}(S)$ preserves compact

objects. We write $\mathcal{T}(S)^\omega$ for the full subcategory of compact objects in $\mathcal{T}(S)$ for any scheme S . This is a thick stable subcategory, and we have $\text{Ind}(\mathcal{T}(S)^\omega) \simeq \mathcal{T}(S)$.

REMARK A.6. — Suppose that \mathcal{T} is compactly generated. Then by adjunction, the direct image functor $f_* : \mathcal{T}(S) \rightarrow \mathcal{T}(T)$ commutes with filtered colimits for any morphism $f : S \rightarrow T$. Since it is an exact functor between stable ∞ -categories, it therefore commutes with arbitrary small colimits.

DEFINITION A.7. — Recall from [CD19, Def. 4.2.1] the full subcategory \mathcal{T}_c of *constructible* objects. We say that a motivic ∞ -category \mathcal{T} is *constructibly generated* if it is compactly generated and the constructible objects are compact. Then we have $\mathcal{T}_c(S) = \mathcal{T}(S)^\omega$ for every scheme S , i.e., the compact and constructible objects coincide [CD19, Prop. 1.4.11].

REMARK A.8. — Suppose that \mathcal{T} is constructibly generated. In this case, the exceptional inverse image functor $f^!$ commutes with colimits for any morphism f of finite type. This follows by adjunction from the fact that $f_!$ preserves compact objects [CD19, Cor. 4.2.12] (where the separatedness and noetherian hypotheses are not necessary, see [Kha21, Cor. 2.62]).

EXAMPLE A.9. — The motivic ∞ -category SH is constructibly generated by [Hoy14, Prop. C.12(1-3)]. We deduce the same for $\text{SH}^{[1/2]}$, SH_+ , SH_- , SH_\square , $\text{D}_{\mathbb{A}^1}(-, \Lambda)$ (for any commutative ring Λ), etc.

A.10. — Let \mathcal{T} be a compactly generated motivic ∞ -category. It follows from Prop. 5.5.7.8 and Cor. 4.4.5.21 in [Lur09] that \mathcal{T} is continuous if and only if, for every cofiltered system $(S_\alpha)_\alpha$ as in Setup A.1, the canonical functor

$$\varinjlim_\alpha \mathcal{T}(S_\alpha)^\omega \longrightarrow \mathcal{T}(S)^\omega$$

is an equivalence, where the colimit is taken in the ∞ -category of (small) ∞ -categories and the transition arrows are the inverse image functors $u_{\alpha,\beta}^*$. Concretely, we have:

(1) For any index α_0 and objects $E_{\alpha_0}, F_{\alpha_0} \in \mathcal{T}(S_{\alpha_0})$ with E_{α_0} compact, we have canonical isomorphisms of spaces

$$\varinjlim_{\alpha > \alpha_0} \text{Maps}_{\mathcal{T}(S_\alpha)}(E_\alpha, F_\alpha) \longrightarrow \text{Maps}_{\mathcal{T}(S)}(E, F),$$

where $E_\alpha \in \mathcal{T}(S_\alpha)$ and $E \in \mathcal{T}(S)$ are the inverse images (and similarly for F_α and F). (Using the compactness of E_{α_0} , we can reduce to the case where F_{α_0} is also compact. Then the claim holds by the fully faithfulness of the functor above, since formation of mapping spaces in an ∞ -category commutes with filtered colimits.)

(2) For every compact object $E \in \mathcal{T}(S)^\omega$, there exists an index α , a compact object $F_\alpha \in \mathcal{T}(S_\alpha)^\omega$, and an isomorphism $E \simeq u_\alpha^*(F_\alpha)$.

We finish this appendix by formulating continuity in terms of the cohomology and bivariant theories represented by an object of a motivic ∞ -category.

NOTATION A.11. — Let \mathcal{T} be a motivic ∞ -category and E an absolute object of \mathcal{T} . Recall the following notation from [DJK21, Def. 2.2.1]: for any scheme X and virtual vector bundle v on X , the v -twisted cohomology space of X is

$$E(X, v) = \mathrm{Maps}_{\mathcal{T}(X)}(\mathbb{1}_X, E_X \otimes \mathrm{Th}_X(v)),$$

where Th_X is the Thom space functor in \mathcal{T} (see Paragraph 7.2). For any finite type morphism $f : X \rightarrow S$, we define the v -twisted bivariant theory space as

$$E(X/S, v) = \mathrm{Maps}_{\mathcal{T}(X)}(\mathrm{Th}_X(v), f^!E_S) \simeq \mathrm{Maps}_{\mathcal{T}(S)}(f_! \mathrm{Th}_X(v), E_S).$$

A.12. — Let \mathcal{T} be a constructibly generated, continuous motivic ∞ -category. Let $(S_\alpha)_\alpha$ be a cofiltered system of schemes as in Setup A.1, with limit S . By continuity for algebraic K-theory [TT90, Th. 7.2], any virtual vector bundle v on S is the inverse image of a virtual vector bundle v_{α_0} on S_{α_0} , for some sufficiently large index α_0 . For any absolute object E of \mathcal{T} , Paragraph A.10(1) yields that the canonical maps

$$\varinjlim_{\alpha > \alpha_0} E(S_\alpha, v_\alpha) \longrightarrow E(S, v)$$

are invertible (since the unit $\mathbb{1}_{S_{\alpha_0}} \in \mathcal{T}(S_{\alpha_0})$ is constructible and hence compact).

A.13. — Let \mathcal{T} be a constructibly generated, continuous motivic ∞ -category. Let $(S_\alpha)_\alpha$ be a cofiltered system of schemes as in Setup A.1, with limit S . Let X_{α_0} be a finite type S_{α_0} -scheme for some index α_0 , let $X_\alpha = X_{\alpha_0} \times_{S_{\alpha_0}} S_\alpha$ for every $\alpha > \alpha_0$, and let $X \simeq X_{\alpha_0} \times_{S_{\alpha_0}} S$ be the limit of $(X_\alpha)_{\alpha > \alpha_0}$. Denote by $f_\alpha : X_\alpha \rightarrow S_\alpha$ and $f : X \rightarrow S$ the structural morphisms. Let v be a vector bundle over X , which as in Paragraph A.12 we may assume (by modifying α_0 if necessary) is the inverse image of some virtual vector bundle v_{α_0} on X_{α_0} . Then the canonical maps

$$\varinjlim_{\alpha > \alpha_0} E(X_\alpha/S_\alpha, v_\alpha) \longrightarrow E(X/S, v)$$

are invertible. Indeed, the constructible generation assumption on \mathcal{T} implies (see Remark A.8) that $(f_\alpha)_!(\mathrm{Th}_{X_\alpha}(v_\alpha))$ is compact for every α . Moreover, by the base change formula, $f_! \mathrm{Th}_X(v)$ is the inverse image of $(f_\alpha)_!(\mathrm{Th}_{X_\alpha}(v_\alpha))$ for every α . Hence the claim follows from Paragraph A.10(1).

APPENDIX B. ESSENTIALLY OF FINITE PRESENTATION MORPHISMS

B.1. ESSENTIALLY OF FINITE PRESENTATION AND SMOOTH MORPHISMS. — The material in this subsection is well-known, but does not seem to be recorded in the literature yet. We recall our global conventions that all schemes are implicitly assumed to be quasi-compact and quasi-separated.

DEFINITION B.1. — A morphism of schemes $f : X \rightarrow Y$ is *pro-finitely presented* if it can be written as the limit of a cofiltered system of finitely presented morphisms $f_\alpha : X_\alpha \rightarrow Y$, with affine transition maps. We define *pro-smooth* and *pro-étale* morphisms, and *pro-open immersions* analogously. Note that the transition maps are automatically étale in the latter two cases.

DEFINITION B.2. — Let $f : X \rightarrow Y$ be a morphism of schemes. We say that f is *essentially of finite presentation* (resp. *essentially of finite type*) if it factors through a pro-étale morphism $p : X \rightarrow X'$ and a morphism of finite presentation (resp. of finite type) $g : X' \rightarrow Y$. We define *essentially smooth* and *essentially étale* morphisms analogously.

EXAMPLE B.3. — If X is a scheme and $x \in X$ is a (not necessarily closed) point, then the inclusion of the residue field $i_x : \text{Spec } \kappa(x) \rightarrow X$ is essentially of finite type via the canonical factorization

$$i_x : \{x\} \xrightarrow{j_x} \overline{\{x\}} \xrightarrow{k_x} X.$$

If X is noetherian, then i_x is essentially of finite presentation.

PROPOSITION B.4. — *The class of essentially of finite presentation (resp. essentially smooth, essentially étale) morphisms is stable under base change and composition.*

PROPOSITION B.5. — *A morphism $f : X \rightarrow Y$ is essentially of finite presentation (resp. essentially smooth, essentially étale) if and only if it can be written as the limit of a cofiltered system of finitely presented morphisms $f_\alpha : X_\alpha \rightarrow Y$ (resp. smooth morphisms, étale morphisms), with a pro-étale and étale transition maps.*

In particular, Proposition B.5 says that “essentially étale” is equivalent to “pro-étale”. Both Propositions B.4 and B.5 are simple consequences of the following lemma:

LEMMA B.6. — *Let \mathcal{P} be the class of finitely presented morphisms (resp. smooth morphisms, étale morphisms, open immersions). Let \mathcal{Q} be the class of morphisms $f : X \rightarrow Y$ that can be written as the limit of a cofiltered system of morphisms $f_\alpha : X_\alpha \rightarrow Y$ belonging to \mathcal{P} , with a pro-étale and étale transition maps. Then the class \mathcal{Q} is stable under base change and composition.*

Proof. — Stability under base change is clear. Let $f : X \rightarrow Y$ and $g : Y \rightarrow Z$ be morphisms in \mathcal{Q} . Write them as limits of cofiltered systems $(f_\alpha : X_\alpha \rightarrow Y)_\alpha$ and $(g_\beta : Y_\beta \rightarrow Z)_\beta$, respectively. Then for every index α , there exists by [Gro64, Th. 8.8.2(ii)] an index β , a finitely presented morphism $f_{\alpha,\beta} : X_{\alpha,\beta} \rightarrow Y_\beta$ and a Y -isomorphism $X_{\alpha,\beta} \times_{Y_\beta} Y \simeq X_\alpha$. Then $g \circ f : X \rightarrow Z$ is the limit of the finitely presented morphisms $X_{\beta,\alpha} \rightarrow Y_\beta \rightarrow Z$. In the respective cases, $f_{\alpha,\beta}$ is smooth, étale, or an open immersion by [Gro64, Prop. 17.7.8 or Prop. 8.6.3].

Proof of Proposition B.5. — The case of étale morphisms follows immediately from the fact that pro-étale morphisms are stable under composition (Lemma B.6). Let \mathcal{P} be the class of smooth or finitely presented morphisms and let \mathcal{Q} be as in Lemma B.6. If $f : X \rightarrow Y$ is essentially \mathcal{P} , then it is by definition the composite of two morphisms in \mathcal{Q} , and hence itself belongs to \mathcal{Q} by Lemma B.6. Conversely suppose that $f : X \rightarrow Y$ belongs to \mathcal{Q} . Write f as the limit of a cofiltered system of morphisms

$f_\alpha : X_\alpha \rightarrow Y$ in \mathcal{P} , with affine and étale transition maps. Then for every α , the projection $u_\alpha : X \rightarrow X_\alpha$ is pro-étale, so the factorization

$$f : X \xrightarrow{u_\alpha} X_\alpha \xrightarrow{f_\alpha} Y$$

exhibits f as essentially \mathcal{P} .

Proof of Proposition B.4. — Combine Proposition B.5 and Lemma B.6.

LEMMA B.7. — *Let $f : X \rightarrow Y$ be a pro-étale morphism. If f is of finite presentation, then it is étale.*

Proof. — It will suffice by definition to show that f is formally étale. If f is the limit of a cofiltered system of étale morphisms $f_\alpha : X_\alpha \rightarrow Y$, then the relative cotangent complex $L_{X/Y}$ is the colimit over α of $u_\alpha^* L_{X_\alpha/Y}$, where $u_\alpha : X \rightarrow X_\alpha$ is the projection (see e.g. [Sta21, Tags 08S9, 08R2]). Hence $L_{X/Y} \simeq 0$, i.e., f is formally étale.

LEMMA B.8. — *Let $f : X \rightarrow Y$ be a morphism of S -schemes. If X and Y are pro-étale over S , then f is pro-étale.*

Proof. — Write X and Y as limits of cofiltered systems $(X_\alpha)_\alpha$ and $(Y_\beta)_\beta$ of étale S -schemes, respectively. For every index β , there exists by [Gro64, Prop. 8.14.2] an index α and an S -morphism $f_\beta : X_\alpha \rightarrow Y_\beta$ through which the composite $X \rightarrow Y \rightarrow Y_\beta$ factors. Now f_β is étale (as a morphism between étale S -schemes), and $f : X \rightarrow Y$ is the limit of the étale maps $X_\alpha \times_{Y_\beta} Y \rightarrow Y$ (base changes of f_β).

LEMMA B.9. — *Let $f : X \rightarrow Y$ be a morphism of S -schemes. If X is pro-étale over S and Y is of finite presentation over S , then f is essentially of finite presentation.*

Proof. — Write X as the limit of a cofiltered system of étale S -schemes X_α (with affine transition maps). By [Gro64, Prop. 8.14.2], $f : X \rightarrow Y$ factors through an S -morphism $f_\alpha : X_\alpha \rightarrow Y$ for some index α . Then f is the limit of the finitely presented morphisms $f_\beta : X_\beta \rightarrow X_\alpha \rightarrow Y$ for all $\beta > \alpha$. Hence it is essentially of finite presentation by Proposition B.5.

B.2. f^* FOR PRO-ÉTALE MORPHISMS

LEMMA B.10. — *Let \mathcal{T} be a continuous motivic ∞ -category. Then for every pro-open immersion $j : U \rightarrow X$, the functor $j_* : \mathcal{T}(U) \rightarrow \mathcal{T}(X)$ is fully faithful. Equivalently, the co-unit $j^* j_* \rightarrow \text{Id}$ is invertible.*

Proof. — Write j as the limit of a cofiltered system of open immersions $(j_\alpha : U_\alpha \rightarrow X)_\alpha$. Let E and F be objects of $\mathcal{T}(U)$. By continuity (Paragraph A.3), the canonical map

$$\text{Maps}_{\mathcal{T}(U)}(E, F) \longrightarrow \text{Maps}_{\mathcal{T}(X)}(j_*(E), j_*(F))$$

is identified with the map

$$\varprojlim_{\alpha} \text{Maps}_{\mathcal{T}(U_\alpha)}((u_\alpha)_*(E), (u_\alpha)_*(F)) \longrightarrow \text{Maps}_{\mathcal{T}(X)}(j_*(E), j_*(F))$$

induced by the functors $(j_\alpha)_*$. Since each $(j_\alpha)_*$ is fully faithful by smooth base change [CD19, 1.1.20], this map is an isomorphism.

If \mathcal{T} is compactly generated, then Lemma B.10 follows alternatively from the following base change formula:

PROPOSITION B.11. — *Let \mathcal{T} be a continuous motivic ∞ -category. Assume that for every morphism of schemes f , the direct image f_* commutes with colimits (by Remark A.6, this is automatic if \mathcal{T} is compactly generated). Suppose given a cartesian square of schemes*

$$\begin{array}{ccc} X' & \xrightarrow{g} & Y' \\ p \downarrow & & \downarrow q \\ X & \xrightarrow{f} & Y \end{array}$$

with f and g pro-étale. Then there is a canonical isomorphism of functors

$$f^*q_* \simeq p_*g^*.$$

Proof. — The natural transformation is defined as the composite

$$f^*q_* \xrightarrow{\text{unit}} p_*p^*f^*q_* \simeq p_*g^*q^*q_* \xrightarrow{\text{counit}} p_*g^*.$$

Write f as the limit of a cofiltered system of étale maps $f_\alpha : X_\alpha \rightarrow X$, and write $u_\alpha : X \rightarrow X_\alpha$ for the projections. Denote by $g_\alpha : X'_\alpha \rightarrow Y$ and $v_\alpha : X' \rightarrow X'_\alpha$ the respective base changes of f_α and u_α along $q : Y' \rightarrow Y$. By continuity (Paragraph A.3), it will suffice to show that the induced natural transformation

$$(u_\alpha)_*f^*q_* \longrightarrow (u_\alpha)_*p_*g^*$$

is invertible for all α . Since $f = f_\alpha \circ u_\alpha$ and $g = g_\alpha \circ v_\alpha$, this is identified with

$$(u_\alpha)_*u_\alpha^*(p_\alpha)_*g_\alpha^* \simeq (u_\alpha)_*u_\alpha^*f_\alpha^*q_* \longrightarrow (u_\alpha)_*p_*v_\alpha^*g_\alpha^*,$$

where we have used the base change isomorphism $f_\alpha^*q_* \simeq (p_\alpha)_*g_\alpha^*$ (since f_α is étale). It remains therefore to demonstrate the base change formula $u_\alpha^*(p_\alpha)_* \simeq p_*v_\alpha^*$. For every morphism $\alpha \rightarrow \beta$ in the indexing category, we have the base change isomorphism $u_{\alpha,\beta}^*(p_\beta)_* \simeq (p_\alpha)_*v_{\alpha,\beta}^*$ (since the transition maps $u_{\alpha,\beta} : X_\alpha \rightarrow X_\beta$ are étale), via which the square

$$\begin{array}{ccc} \mathcal{T}(X'_\beta) & \xrightarrow{v_{\alpha,\beta}^*} & \mathcal{T}(X'_\alpha) \\ (p_\beta)_* \downarrow & & \downarrow (p_\alpha)_* \\ \mathcal{T}(X_\beta) & \xrightarrow{u_{\alpha,\beta}^*} & \mathcal{T}(X_\alpha) \end{array}$$

commutes. Since $(p_\alpha)_*$ is a left adjoint for each α (as \mathcal{T} is compactly generated, see Remark A.6), this can be regarded as a diagram in the ∞ -category of presentable

∞ -categories and left-adjoint functors. Thus by continuity, taking colimits in both rows results in a commutative square

$$\begin{array}{ccc} \mathcal{T}(X'_\alpha) & \xrightarrow{v_\alpha^*} & \mathcal{T}(X') \\ (p_\alpha)_* \Big\downarrow & & \Big\downarrow p_* \\ \mathcal{T}(X_\alpha) & \xrightarrow{u_\alpha^*} & \mathcal{T}(X) \end{array}$$

providing the desired base change isomorphism $u_\alpha^*(p_\alpha)_* \simeq p_*v_\alpha^*$.

PROPOSITION B.12. — *Let \mathcal{T} be a continuous motivic ∞ -category. Assume that for every morphism of schemes f (resp. of finite presentation), the direct image functor f_* (resp. exceptional inverse image functor $f^!$) commutes with colimits. (By Remark A.8, this is automatic if \mathcal{T} is constructibly generated.) Then for every commutative square of schemes*

$$\begin{array}{ccc} X' & \xrightarrow{g} & Y' \\ p \Big\downarrow & & \Big\downarrow q \\ X & \xrightarrow{f} & Y \end{array}$$

with f and g of finite presentation, there is a canonical morphism of functors

$$Ex^{*!} : p^*f^! \longrightarrow g^!q^*$$

which is invertible if q is pro-étale.

Proof. — The natural transformation is defined as the composite

$$p^*f^! \xrightarrow{\text{unit}} p^*f^!q_*q^* \simeq p^*p_*g^!q^* \xrightarrow{\text{counit}} g^!q^*,$$

where the middle isomorphism is Proposition B.11.

Note that the canonical map $a : X' \rightarrow X \times_Y Y'$ is pro-étale by Lemma B.8, since the source and target are both pro-étale over X . Since the source and target are of finite presentation over Y' , it follows that a is also of finite presentation and hence étale by Lemma B.7. In particular, we have a canonical isomorphism $a^* \simeq a^!$. Thus we reduce to the case where the square is cartesian.

Write q as a cofiltered limit of étale maps $q_\alpha : Y'_\alpha \rightarrow Y$, and denote by $v_\alpha : Y' \rightarrow Y'_\alpha$ the projections. Consider the cartesian square

$$\begin{array}{ccc} X' & \xrightarrow{g} & Y' \\ u_\alpha \Big\downarrow & & \Big\downarrow v_\alpha \\ X'_\alpha & \xrightarrow{g_\alpha} & Y'_\alpha \\ p_\alpha \Big\downarrow & & \Big\downarrow q_\alpha \\ X & \xrightarrow{f} & Y \end{array}$$

Regarding the left-hand triangle as a commutative square by inserting the identity $X \rightarrow X$, Proposition B.12 applies to give a canonical isomorphism $p^*a^! \simeq q^*$. This then yields the desired isomorphism

$$p^*(f')^! \simeq p^*a^!(f'')^! \simeq q^*(f'')^!.$$

EXAMPLE B.15. — If $f : X \rightarrow Y$ is pro-étale, then there is an equivalence $f^! \simeq f^*$.

Under mild assumptions on \mathcal{T} , we will be able to prove:

PROPOSITION B.16. — *Suppose that \mathcal{T} is constructibly generated (Definition A.7). Let $f : X \rightarrow Y$ and $g : Y \rightarrow Z$ be essentially of finite presentation morphisms. Then there is a canonical isomorphism of functors $(g \circ f)^! \simeq f^!g^!$.*

In fact, it suffices to assume that for every morphism of schemes f (resp. of finite presentation), the direct image functor f_* (resp. exceptional inverse image functor $f^!$) commutes with colimits (see Remark A.8).

WARNING B.17. — Below we will show that much of the usual behaviour of the operation $f^!$ extends to the essentially of finite presentation case. However, we caution the reader that we cannot typically expect $f^!$ to admit a left adjoint $f_!$ when f is not of finite type.

PROPOSITION B.18 (Base change). — *Assume that for every morphism of schemes f , the direct image f_* commutes with colimits. Suppose given a cartesian square of schemes*

$$\begin{array}{ccc} X' & \xrightarrow{g} & Y' \\ p \downarrow & & \downarrow q \\ X & \xrightarrow{f} & Y \end{array}$$

with f and g essentially of finite presentation. Then there is a canonical isomorphism of functors

$$p_*g^! \simeq f^!q_*.$$

Proof. — Choose a factorization $f = f' \circ a$ with f' of finite presentation and a pro-étale. Then the claim follows from the base change formula for finite type morphisms, the identification $a^! \simeq a^*$ (Example B.15), and Proposition B.11.

PROPOSITION B.19 (*!-exchange). — *Assume that for every morphism of schemes f (resp. of finite presentation), the direct image functor f_* (resp. exceptional inverse image functor $f^!$) commutes with colimits. Suppose given a commutative square of schemes*

$$\begin{array}{ccc} X' & \xrightarrow{g} & Y' \\ p \downarrow & & \downarrow q \\ X & \xrightarrow{f} & Y \end{array}$$

with f and g essentially of finite presentation. Then there is a canonical morphism of functors

$$Ex^{*!} : p^* f^! \longrightarrow g^! q^*$$

which is invertible if q is pro-étale.

Proof. — The natural transformation is defined as the composite

$$p^* f^! \xrightarrow{\text{unit}} p^* f^! q_* q^* \simeq p^* p_* g^! q^* \xrightarrow{\text{counit}} g^! q^*,$$

where the middle isomorphism is Proposition B.18. By factoring f through a pro-étale morphism and a finite presentation morphism, we immediately reduce to the case where f is of finite presentation, which is Proposition B.12.

Proof of Proposition B.16. — Follows directly from Proposition B.19.

We have the following !-analogue of [CD19, Cor. 4.3.17] (with almost the same proof):

PROPOSITION B.20. — *Let X be a noetherian scheme of finite dimension. As x ranges over the points of X , the family of functors*

$$i_x^! : T(X) \longrightarrow T(\text{Spec } \kappa(x))$$

is jointly conservative.

Proof. — Argue by induction on the dimension d . For $d \leq 0$ the claim is clear. Assume $d > 0$. The localization of X at a point x is a pro-open immersion $j_x : \text{Spec } \mathcal{O}_{X,x} \rightarrow X$. Since $j_x^! \simeq j_x^*$ (Example B.15), we may use [CD19, Prop. 4.3.9] (and Proposition B.16) to reduce to the case where X is local. Then the claim follows easily using the localization triangle

$$(i_{x_0})_* i_{x_0}^! \longrightarrow \text{Id} \longrightarrow j_* j^!,$$

where i_{x_0} is the inclusion of the closed point (which is essentially of finite presentation, see Example B.3), and j is the inclusion of the complement (which is of dimension $< d$).

APPENDIX C. ABSOLUTE PURITY IN EQUAL CHARACTERISTIC

Let \mathcal{T} be a motivic ∞ -category fibered over a category of schemes \mathcal{S} , and E an absolute spectrum in \mathcal{T} . As recalled in 3.5, for $f : X \rightarrow Y$ a smoothable morphism of regular schemes with virtual tangent bundle T_f , there is a map

$$(C.0.a) \quad \eta_f : E_X \otimes \text{Th}(T_f) \longrightarrow f^! E_Y$$

induced by the fundamental class of f [DJK21, Def. 4.3.7]. We say that E is f -pure if the map (C.0.a) is an isomorphism, and we say that E satisfies *absolute purity* if it is f -pure for every smoothable morphism f between regular schemes in \mathcal{S} . In fact, it suffices to check this property for closed immersions [DJK21, Rem. 4.3.12(i)].

This appendix is dedicated to the following result:

THEOREM C.1. — *Let k be a perfect field and let \mathcal{T} be a continuous, constructibly generated motivic ∞ -category (Definitions A.2 and A.7) fibered over k -schemes. Then any absolute spectrum in \mathcal{T} satisfies absolute purity.*

This statement was first asserted in [Dég18b, Ex.1.3.4(2)] (see also [CD15, Prop.6.2]). It is proved, as suggested in loc.cit., by using Popescu’s theorem to reduce to relative purity (the case of a closed immersion between smooth schemes). In this appendix we record the details of two variants of this proof.

Recall that a homomorphism of commutative rings $u : A \rightarrow B$ is *regular* if it is flat and for every residue field $A \rightarrow \kappa$, the ring $B \otimes_A \kappa$ is geometrically regular. Then Popescu’s theorem asserts (see [Spi99, Th. 1.1]):

THEOREM C.2 (Popescu). — *Let A and B be noetherian commutative rings. A ring homomorphism $u : A \rightarrow B$ is regular if and only if it is ind-smooth, i.e., it exhibits B as a filtered colimit of smooth A -algebras.*

Before proceeding to the proofs, we begin with a few generalities.

DEFINITION C.3. — A *regular closed pair* (X, Z) is a regular closed immersion $i : Z \rightarrow X$ between schemes. A *transverse morphism* of regular closed pairs $f : (Y, W) \rightarrow (X, Z)$ is a Tor-independent cartesian square

$$\begin{array}{ccc} W & \longrightarrow & Y \\ \downarrow & & \downarrow \\ Z & \longrightarrow & X \end{array}$$

LEMMA C.4. — *Let \mathcal{T} be a continuous, constructibly generated motivic ∞ -category. Let E be an absolute object in \mathcal{T} . Let (X, Z) be a regular closed pair which is the limit of a cofiltered system $(X_\alpha, Z_\alpha)_{\alpha \in \Lambda}$ of regular closed pairs with affine and transverse transition morphisms. If E is i_α -pure for every $\alpha \in \Lambda$, then E is i -pure.*

Proof of Lemma C.4. — The assumptions on \mathcal{T} imply (see Paragraph A.13) that the canonical map of spaces

$$\varinjlim_{\alpha \in \Lambda} E(Z_\alpha/X_\alpha, \langle -N_{i_\alpha} \rangle) \longrightarrow E(Z/X, \langle -N_i \rangle)$$

is invertible (see Notation A.11). By the transverse base change formula [DJK21, Th. 3.2.21], each base change morphism $E(Z_\alpha/X_\alpha, \langle -N_{i_\alpha} \rangle) \rightarrow E(Z/X, \langle -N_i \rangle)$ sends the fundamental class η_{i_α} to the fundamental class η_i . Since each of the classes η_{i_α} is invertible by assumption, the claim follows.

LEMMA C.5. — *Let \mathcal{T} be a continuous, constructibly generated motivic ∞ -category. Let E be an absolute object in \mathcal{T} . If i is a closed immersion between schemes that are essentially smooth over a scheme S , then E is i -pure.*

Proof. — The case where both schemes are smooth over S follows from relative purity as in [DJK21, Prop. 4.3.10(i)]. In general, write X and Z as the limits of projective systems of smooth S -schemes X_α and Z_α , respectively, with affine étale transition morphisms. Then by [Gro64, Th. 8.8.2(ii), Prop. 8.6.3], the closed immersion i descends to a closed immersion $i_\lambda : Z_\lambda \rightarrow X_\lambda$ for some sufficiently large λ . In particular, the pair (X, Z) can be written as the limit of a projective system of closed pairs of smooth S -schemes, with affine étale transition morphisms. Thus the claim follows from Lemma C.4 (note that étale morphisms of pairs are transverse).

C.6. — We now give the first proof. We will require the following lemma:

LEMMA C.7. — *Let $A \rightarrow B$ be a local homomorphism of regular local rings. Let $I \subset A$ be a proper ideal such that B/IB is regular, and set $n = \dim(B) - \dim(B/IB)$. If there exist n elements $f_1, \dots, f_n \in A$ generating I , then the f_i form a subset of a regular system of parameters for A , and the ring A/I is regular.*

Proof. — Let \mathfrak{m} and \mathfrak{n} be the respective maximal ideals of A and B . By assumption, the ideal IB is generated by the images $g_i \in B$ of f_i , and $\dim(B/IB) = \dim(B) - n$. By [Gro64, 0, Prop. 16.3.7], we deduce that the g_i form a subset of a system of parameters for B . Since B/IB is regular, this implies that they are B/\mathfrak{n} -linearly independent as elements of $\mathfrak{n}/\mathfrak{n}^2$ [Gro64, 0, Prop. 17.1.7]. This implies that the f_i are A/\mathfrak{m} -linearly independent in $\mathfrak{m}/\mathfrak{m}^2$. The claim now follows by another application of [Gro64, 0, Prop. 17.1.7].

PROPOSITION C.8. — *Let k be a perfect field and let $i : Z \rightarrow X$ be a closed immersion between a n -regular k -schemes. Then the closed pair (X, Z) is a projective limit of closed pairs of smooth k -schemes with transverse a n -transition morphisms.*

Proof. — Let $X = \text{Spec } A$ and let I be the ideal defining Z . By Theorem C.2, we may write A as a colimit of smooth k -algebras A_α for $\alpha \in \Lambda$. Truncating Λ if necessary, we may assume that for every $\alpha \in \Lambda$, there is an ideal I_α of A_α such that $I = I_\alpha A$. Let $X_\alpha = \text{Spec } A_\alpha$ and $Z_\alpha = \text{Spec } A_\alpha/I_\alpha$.

Let n be the codimension of Z in X , considered as a Zariski-locally constant function on Z . Truncating Λ again, we may also assume that for all $\alpha \in \Lambda$, the ideal I_α is locally generated by at most n elements. Indeed, the locus T_α where this is not true is closed in Z_α . As Z has codimension n in X , the projective limit of T_α is empty, so we have $T_\alpha = \emptyset$ for all sufficiently large α .

Now let X'_α be the open subscheme of X_α given by the complement of the singular locus of Z_α , and let $Z'_\alpha = X'_\alpha \cap Z_\alpha$. It follows from Lemma C.7 that

– X'_α contains the image of X in X_α , and consequently the map $X \rightarrow X_\alpha$ factors through X'_α .

– The induced map $X \rightarrow X'_\alpha$ is a transverse morphism of regular closed pairs $(X, Z) \rightarrow (X'_\alpha, Z'_\alpha)$, since both pairs are defined by regular closed immersions of the same codimension. The transition morphisms of the projective system $(X'_\alpha, Z'_\alpha)_{\alpha \in \Lambda}$ are also transverse for the same reason.

Since k is perfect, (X'_α, Z'_α) is a closed pair of smooth k -schemes. Furthermore, since (X, Z) is the projective limit of $(X_\alpha, Z_\alpha)_{\alpha \in \Lambda}$, it is also the projective limit of $(X'_\alpha, Z'_\alpha)_{\alpha \in \Lambda}$, and the result follows.

First proof of Theorem C.1. — Let \mathcal{T} be a motivic ∞ -category as in the statement. By Zariski descent [CD19, Prop. 3.3.4], it suffices to show that \bar{E} is i -pure for any closed immersion $i : Z \rightarrow X$ between affine regular k -schemes. This follows by combining Lemma C.4, Lemma C.5 and Proposition C.8.

C.9. — The second proof is similar but uses an argument of Gabber to first reduce to the case of punctual purity [ILO14, XVI.3.2].

LEMMA C.10. — *Let $A \rightarrow B$ be a local homomorphism of regular local rings. If there exist n elements $f_1, \dots, f_n \in A$ whose images $g_1, \dots, g_n \in B$ form a regular sequence of parameters for B , then (f_1, \dots, f_n) is a regular system of parameters for A . In particular, we have $\dim(A) = \dim(B) = n$.*

Proof. — This is a special case of Lemma C.7, but it is simpler to argue directly as follows. Since g_1, \dots, g_n form a minimal set of generators for the maximal ideal $\mathfrak{n} \subset B$, they induce a basis of the B/\mathfrak{n} -vector space $\mathfrak{n}/\mathfrak{n}^2$. It follows that the f_i induce a basis of the A/\mathfrak{m} -vector space $\mathfrak{m}/\mathfrak{m}^2$. By [Gro64, 0, Prop. 17.1.7] this means that the f_i form a regular system of parameters for A , as claimed.

We use the following variant of Proposition C.8:

PROPOSITION C.11. — *Let A be a regular local k -algebra with maximal ideal \mathfrak{m} . Then the regular closed pair $(\mathrm{Spec} A, \mathrm{Spec} A/\mathfrak{m})$ is a projective limit of closed pairs of essentially smooth n -schemes with transverse transition morphisms.*

Proof. — By Theorem C.2, we may write A as the colimit of an inductive system $(A_\gamma^0)_{\gamma \in \Gamma}$ of smooth k -algebras. If A_α are the localizations of the A_γ^0 's at prime ideals, then A is also the colimit of the inductive system $(A_\alpha)_{\alpha \in \Lambda}$ of essentially smooth local k -algebras. Let $f_1, \dots, f_n \in \mathfrak{m}$ be a regular system of parameters for A , where $n = \dim(A)$. For some sufficiently large $\lambda \in \Lambda$, the f_i belong to A_λ and form a regular system of parameters of length n , by Lemma C.7. In particular, $\mathfrak{m} = \mathfrak{m}_\alpha A$ for each $\alpha > \lambda$, where \mathfrak{m}_α is the maximal ideal of A_α , and the closed pair $(\mathrm{Spec} A, \mathrm{Spec} A/\mathfrak{m})$ is the limit of the projective system

$$(\mathrm{Spec} A_\alpha, \mathrm{Spec} A_\alpha/\mathfrak{m}_\alpha)_{\alpha > \lambda}.$$

All pairs under consideration are given by regular closed immersions of codimension n , so it follows that the transition morphisms are transverse.

Second proof of Theorem C.1. — Let $i : Z \rightarrow X$ be a closed immersion between regular k -schemes. Let z be a point of Z , and let $Z_z = \mathrm{Spec} \mathcal{O}_{Z,z}$ and $X_z = \mathrm{Spec} \mathcal{O}_{X,i(z)}$.

Denote the canonical closed immersions

$$\begin{array}{ccc}
 z & \xrightarrow{i_Z} & Z_z \\
 & \searrow i_X & \downarrow i' \\
 & & X_z
 \end{array}$$

by i_Z and i_X respectively. By [CD19, Lem. 4.3.17] and [CD19, Prop. 4.3.9], we only need to show that $i_Z^* \eta_{i'}$ is an isomorphism. By Lemma C.4, Lemma C.5 and Proposition C.11, we know that both η_{i_Z} and η_{i_X} are isomorphisms. By [DJK21, Th. 3.3.2] the fundamental classes are compatible with composition, and we know that $i_Z^! \eta_{i'}$ is an isomorphism.

Now use induction on the dimension of Z_z . Let $j_Z : U_z \rightarrow Z_z$ be the open immersion given by the complement of z in Z_z . Then U_z has dimension lower than Z_z , and the induction hypothesis implies that $j_Z^* \eta_{i'}$ is an isomorphism. From the canonical localization triangle

$$i_Z^! \eta_{i'} \longrightarrow i_Z^* \eta_{i'} \longrightarrow i_Z^* j_{Z*} j_Z^* \eta_{i'} \longrightarrow i_Z^! \eta_{i'}[1]$$

and the five lemma, we know that $i_Z^* \eta_{i'}$ is an isomorphism, which finishes the proof.

APPENDIX D. ABSOLUTE PURITY IN HERMITIAN K-THEORY

In this appendix, we prove absolute purity for hermitian K-theory. Recall that Panin and Walter have constructed in [PW19] for any regular $Z[1/2]$ -scheme S a ring spectrum \mathbf{KQ}_S which represents Schlichting’s higher Grothendieck–Witt groups (i.e., hermitian K-theory) in $\mathrm{SH}(S)$. According to [ST15], this ring spectrum has a geometric model (given by orthogonal grassmannians, see op. cit., Introduction, Th. 1). This implies the \mathbf{KQ}_S for various regular $Z[1/2]$ -schemes S are compatible with base change. Therefore, one can define \mathbf{KQ}_S for general $Z[1/2]$ -scheme S just by pulling back $\mathbf{KQ}_{Z[1/2]}$ along the canonical map $S \rightarrow \mathrm{Spec}(Z[1/2])$. Then \mathbf{KQ} is an absolute ring spectrum with respect to SH over the category of $Z[1/2]$ -schemes.

THEOREM D.1. — *The absolute ring spectrum \mathbf{KQ} satisfies absolute purity over $Z[1/2]$ -schemes (in the sense recalled in Paragraph 3.5).*

The proof follows the same lines as the analogous result for algebraic K-theory [CD19, Th. 13.6.3], and boils down to dévissage in Grothendieck–Witt theory.

Proof of Theorem D.1. — Let $i : Z \rightarrow X$ be a closed immersion of pure codimension c between regular schemes. The claim is that the morphism

$$\eta_i : \mathbf{KQ}_Z \otimes \mathrm{Th}_Z(-N_i) \longrightarrow i^!(\mathbf{KQ}_X)$$

defined by the fundamental class is invertible. Since i_* is conservative and the stable ∞ -category $\mathrm{SH}(X)$ is generated by objects of the form $\Sigma_X^\infty(T)_+(n)[2n]$, where T varies

among smooth X -schemes and $n \in \mathbb{Z}$, it will suffice to show that the induced map

$$[\Sigma_{\mathbb{Z}}^{\infty}(T_Z)_+(n)[2n], \mathbf{KQ}_Z \otimes \mathrm{Th}_Z(-N_i)] \longrightarrow [\Sigma_{\mathbb{Z}}^{\infty}(T_Z)_+(n)[2n], i^!(\mathbf{KQ}_X)]$$

is invertible, where $T_Z = T \times_X Z$. In the notation of [DJK21], this is the map

$$\mathbf{KQ}^0(T_Z, -n - \langle N_i \rangle) \longrightarrow \mathbf{KQ}_{T_Z}^0(T, -n)$$

induced by pairing with the fundamental class

$$\eta_{T_Z/Z} \in \mathbf{KQ}_0(T_Z/T, -\langle N_i \rangle).$$

By [PW19, Cor. 7.3] and since \mathbf{KQ} is SL^c -oriented, we have canonical isomorphisms

$$\begin{aligned} \mathbf{KQ}^0(T_Z, -n - \langle N_i \rangle) &\simeq \mathrm{GW}_0^{[-n-c]}(T_Z, \det(-N_i)) \\ \mathbf{KQ}_{T_Z}^0(T, -n) &\simeq \mathrm{GW}_0^{[-n]}(T \text{ on } T_Z) \end{aligned}$$

under which the map in question is identified with the Gysin map in Grothendieck–Witt theory induced by direct image of coherent sheaves (see [Sch17, (9.9)]). The latter is invertible by dévissage in Grothendieck–Witt theory [Sch17, Th. 9.5, 9.18 & 9.19].

APPENDIX E. 2-PRIMARY UNRAMIFIED WITT RING

This appendix brings some complements to the theory of Witt and unramified Witt sheaves (see Definition 5.2) of independent interest.

E.1. — The theory of Witt groups has been extended to higher Witt groups $W^i(S)$ by Balmer (see e.g. [Bal05]). Recall that these groups are defined by considering the derived category $\mathcal{T} = D^b(\mathrm{VB}_S)$ of the exact category of vector bundles over S equipped with the $(-1)^i$ -duality $D_i = \underline{\mathrm{Hom}}(-, \mathcal{O}_S)[i]$ (see [Bal05, 1.4.12]). It is then possible to mimic the classical definition of symmetric bilinear forms/symmetric objects, with respect to the duality D_i , and metabolic ones (called “neutral” in loc. cit.). One defines $W^i(X)$ as the monoid of isomorphisms classes of D_i -symmetric objects of \mathcal{T} modulo the submonoid of D_i -metabolic ones (see loc. cit. 1.4.3).

The following lemma is the Witt-version of our *key lemma* (Lemma 2.2).

PROPOSITION E.2. — *Let S be a $\mathbb{Z}[1/2]$ -scheme and $\Lambda \subseteq \mathbb{Q}$ a subring with $2 \in \Lambda^\times$. Let $\nu : S_{\mathbb{Q}} \rightarrow S$ be the pro-open immersion of the characteristic 0 part of S as in Paragraph 2.1.*

Then the map $\underline{W}_S^\Lambda \rightarrow \nu_(\underline{W}_{S_{\mathbb{Q}}}^\Lambda)$ from (5.4.a) is an isomorphism.*

Proof. — It will be sufficient to prove that for any regular scheme S , the following map on Witt groups is an isomorphism:

$$W(S)[2^{-1}] \xrightarrow{\nu^*} W(S_{\mathbb{Q}})[2^{-1}].$$

We will use Balmer–Witt groups with support, as defined in [Bal01, §1]. Given a closed pair (X, Z) , we write $W^i(X, Z)$ for the i -th Witt group of X with support in Z .

Consider a closed subscheme Z of $\text{Spec}(Z)$, with open complement U . Write S_Z, S_U for the pullbacks of $S/\text{Spec}(Z)$ to Z, U . Using the long exact sequence of Witt groups with support [Bal01, 1.7], we get an exact sequence:

$$W^0(S, S_Z) \longrightarrow W^0(S) \longrightarrow W^0(S_U) \longrightarrow W^1(S, S_Z).$$

Taking the filtered inductive limit over the category of closed subschemes of S , and using additivity in S_Z , we get an exact sequence:

$$\bigoplus_{p \neq 2} W^0(S, S_p) \longrightarrow W^0(S) \xrightarrow{\nu^*} W^0(S_{\mathbb{Q}}) \longrightarrow \bigoplus_{p \neq 2} W^1(S, S_p),$$

where p runs over the odd primes.

Therefore, to get the lemma, it is sufficient to prove that for any odd prime p , the triangular Witt group $W^i(S, S_p)[2^{-1}]$ is zero.

Let δ be the dimension function on S such that $\delta = -\text{codim}_S$ (see e.g. [BD17, §1.1]). From the long exact sequence of Witt groups with support (op.cit., Th. 1.6), one can build out of the δ -niveau filtration on S_p , as in [BD17, §3.1], and deduce a converging spectral sequence⁽¹⁸⁾ of the form:

$$E_{i,j}^1 = \bigoplus_{x \in S_{p,(i)}} W^{-i-j}(S_{(x)}, \{x\}) \implies W^{-i-j}(S, S_p),$$

where $S_{p,(i)}$ denotes the set of points of x such that $\delta(x) = i$. Since S is regular and defined over $\mathbb{Z}[1/2]$, purity (dévissage) for the Witt groups [BW02, Th. 6.1] yields:

$$W^{-i-j}(S_{(x)}, \{x\}) = \begin{cases} W(\kappa(x)) & i + j \equiv 0 \pmod{4} \\ 0 & \text{otherwise.} \end{cases}$$

This group is 2-primary torsion as $\kappa(x)$ has positive characteristic ([Lam05, VIII.3.2 & 6.15]). In particular $E_{i,j}^1[1/2]$ vanishes and one can conclude.

REMARK E.3. — Note that the Witt ring $W(k)$ of a characteristic 2 field is always a 2-torsion ring. In particular, the only obstruction to the above proof to work for an arbitrary regular scheme S is that Balmer’s theory needs 2 being invertible on \mathcal{O}_S .

Let us draw the following corollary:

COROLLARY E.4. — *Let S be a regular $\mathbb{Z}[1/2]$ -scheme and $\Lambda \subseteq \mathbb{Q}$ such that $2 \in \Lambda^\times$.*

- (1) *If S is in addition semi-local, the Gersten–Witt complex of S (see [BW02]) is a resolution of $W(S)$.*
- (2) *Points (1) and (2) of Theorem 5.7 hold for S (i.e., without assuming the existence of a base field).*

This immediately follows from the previous proposition, as it allows one to reduce to the case of a \mathbb{Q} -scheme and then applies [Bal01, 4.3] for the first point and Theorem 5.7 for the second one.

⁽¹⁸⁾This construction is nothing else than the version with support of the Gersten–Witt spectral sequence, whose first version can be found in [BW02].

REMARK E.5. — The Gersten conjecture for the 2-primary Witt ring of excellent $\mathbb{Z}[1/2]$ -schemes was already proved by Jacobson (see [Jac17, 5.3]). Our method gives an alternative (somewhat trivial!) proof and also extends the result as we do not need S to be excellent.

REFERENCES

- [Ana19] A. ANANYEVSKIY – “SL-oriented cohomology theories”, 2019, arXiv: 1901.01597.
- [ALP17] A. ANANYEVSKIY, M. LEVINE & I. PANIN – “Witt sheaves and the η -inverted sphere spectrum”, *J. Topology* **10** (2017), no. 2, p. 370–385.
- [AGV73] M. ARTIN, A. GROTHENDIECK & J.-L. VERDIER – *Théorie des topos et cohomologie étale des schémas*, Lect. Notes in Math., vol. 269, 270, 305, Springer-Verlag, 1972–1973, Séminaire de Géométrie Algébrique du Bois-Marie 1963–64 (SGA 4).
- [Ayo07] J. AYOUB – *Les six opérations de Grothendieck et le formalisme des cycles évanescents dans le monde motivique*, Astérisque, vol. 314-315, Société Mathématique de France, Paris, 2007.
- [Ayo14] ———, “La réalisation étale et les opérations de Grothendieck”, *Ann. Sci. École Norm. Sup. (4)* **47** (2014), no. 1, p. 1–145.
- [Bac18] T. BACHMANN – “Motivic and real étale stable homotopy theory”, *Compositio Math.* **154** (2018), no. 5, p. 883–917.
- [BCD⁺20] T. BACHMANN, B. CALMÈS, F. DÉGLISE, J. FASEL & P. A. ØSTVÆR – “Milnor-Witt motives”, 2020, arXiv: 2004.06634v1.
- [BF18] T. BACHMANN & J. FASEL – “On the effectivity of spectra representing motivic cohomology theories”, 2018, arXiv: 1710.00594.
- [BH21] T. BACHMANN & M. HOYOIS – *Norms in motivic homotopy theory*, Astérisque, Société Mathématique de France, Paris, 2021, to appear.
- [Bal01] P. BALMER – “Witt cohomology, Mayer-Vietoris, homotopy invariance and the Gersten conjecture”, *K-Theory* **23** (2001), no. 1, p. 15–30.
- [Bal05] ———, “Witt groups”, in *Handbook of K-theory. Vol. 1, 2*, Springer, Berlin, 2005, p. 539–576.
- [BW02] P. BALMER & C. WALTER – “A Gersten-Witt spectral sequence for regular schemes”, *Ann. Sci. École Norm. Sup. (4)* **35** (2002), no. 1, p. 127–152.
- [BGPW02] P. BALMER, S. GILLE, I. PANIN & C. WALTER – “The Gersten conjecture for Witt groups in the equicharacteristic case”, *Doc. Math.* **7** (2002), p. 203–217.
- [BB82] A. A. BEĪLINSKIĬ, J. BERNSTEIN & P. DELIGNE – “Faisceaux pervers”, in *Analysis and topology on singular spaces, I (Luminy, 1981)*, Astérisque, vol. 100, Société Mathématique de France, Paris, 1982, p. 5–171.
- [BO74] S. BLOCH & A. OGUS – “Gersten’s conjecture and the homology of schemes”, *Ann. Sci. École Norm. Sup. (4)* **7** (1974), no. 4, p. 181–201.
- [Bon14] M. BONDARKO – “Weights for relative motives: relation with mixed complexes of sheaves”, *Internat. Math. Res. Notices* (2014), no. 17, p. 4715–4767.
- [BD17] M. BONDARKO & F. DÉGLISE – “Dimensional homotopy t-structures in motivic homotopy theory”, *Adv. Math.* **311** (2017), p. 91–189.
- [CDH⁺20a] B. CALMÈS, E. DOTTO, J. HARPAZ, F. HEBESTREIT, M. LAND, K. MOI, D. NARDIN, T. NIKOLAUS & W. STEIMLE – “Hermitian K -theory for stable ∞ -categories I: Foundations”, 2020, arXiv: 2009.07223.
- [CDH⁺20b] ———, “Hermitian K -theory for stable ∞ -categories II: Cobordism categories and additivity”, 2020, arXiv: 2009.07224.
- [CDH⁺20c] ———, “Hermitian K -theory for stable ∞ -categories III: Grothendieck-Witt groups of rings”, 2020, arXiv: 2009.07225.
- [CF14] B. CALMÈS & J. FASEL – “Finite Chow-Witt correspondences”, 2014, arXiv: 1412.2989.
- [Cis19] D.-C. CISINSKI – “Cohomological methods in intersection theory”, arXiv: 1905.03478, 2019.
- [CD15] D.-C. CISINSKI & F. DÉGLISE – “Integral mixed motives in equal characteristics”, *Doc. Math.* (2015), p. 145–194, Extra volume: Alexander S. Merkurjev’s sixtieth birthday.

- [CD16] ———, “Étale motives”, *Compositio Math.* **152** (2016), no. 3, p. 556–666.
- [CD19] ———, *Triangulated categories of mixed motives*, Springer Monographs in Math., Springer, Cham, 2019.
- [CTHK97] J.-L. COLLIOT-THÉLÈNE, R. HOOBLER & B. KAHN – “The Bloch-Ogus-Gabber theorem”, in *Algebraic K-theory (Toronto, ON, 1996)*, Fields Inst. Commun., vol. 16, American Mathematical Society, Providence, RI, 1997, p. 31–94.
- [Dég18a] F. DÉGLISE – “Bivariant theories in motivic stable homotopy”, *Doc. Math.* **23** (2018), p. 997–1076.
- [Dég18b] ———, “Orientation theory in arithmetic geometry”, in *K-Theory—Proceedings of the International Colloquium (Mumbai, 2016)*, Hindustan Book Agency, New Delhi, 2018, p. 239–347.
- [DF20] F. DÉGLISE & J. FASEL – “The Borel character”, 2020, arXiv: 1903.11679.
- [DFJK19] F. DÉGLISE, J. FASEL, F. JIN & A. A. KHAN – “Borel isomorphism and absolute purity”, 2019, arXiv: 1902.02055.
- [DJK21] F. DÉGLISE, F. JIN & A. A. KHAN – “Fundamental classes in motivic homotopy theory”, *J. Eur. Math. Soc. (JEMS)* (2021), to appear.
- [Del77] P. DELIGNE – *Cohomologie étale*, Lect. Notes in Math., vol. 569, Springer-Verlag, 1977, Séminaire de Géométrie Algébrique du Bois-Marie SGA 4 $\frac{1}{2}$.
- [Del87] ———, “Le déterminant de la cohomologie”, in *Current trends in arithmetical algebraic geometry (Arcata, Calif., 1985)*, Contemp. Math., vol. 67, American Mathematical Society, Providence, RI, 1987, p. 93–177.
- [EKM08] R. ELMAN, N. KARPENKO & A. MERKURJEV – *The algebraic and geometric theory of quadratic forms*, AMS Colloquium Publications, vol. 56, American Mathematical Society, Providence, RI, 2008.
- [EHK+20] E. ELMANTO, M. HOYOIS, A. A. KHAN, V. SOSNILO & M. YAKERSON – “Modules over algebraic cobordism”, *Forum Math. Pi* **8** (2020), article no. e14 (44 pages).
- [EK20a] E. ELMANTO & A. A. KHAN – “Perfection in motivic homotopy theory”, *Proc. London Math. Soc. (3)* **120** (2020), no. 1, p. 28–38.
- [EK20b] E. ELMANTO & H. KOLDERUP – “On modules over motivic ring spectra”, *Ann. K-Theory* **5** (2020), no. 2, p. 327–355.
- [Fas08] J. FASEL – *Groupes de Chow-Witt*, Mém. Soc. Math. France (N.S.), vol. 113, Société Mathématique de France, Paris, 2008.
- [FS09] J. FASEL & V. SRINIVAS – “Chow-Witt groups and Grothendieck-Witt groups of regular schemes”, *Adv. Math.* **221** (2009), no. 1, p. 302–329.
- [Fel19] N. FELD – “Morel homotopy modules and Milnor-Witt cycle modules”, 2019, arXiv: 1912.12680.
- [Fel20] ———, “Milnor-Witt cycle modules”, *J. Pure Appl. Algebra* **224** (2020), no. 7, p. 41.
- [Fuj02] K. FUJIWARA – “A proof of the absolute purity conjecture (after Gabber)”, in *Algebraic geometry 2000, Azumino (Hotaka)*, Adv. Stud. Pure Math., vol. 36, Math. Soc. Japan, Tokyo, 2002, p. 153–183.
- [Ful98] W. FULTON – *Intersection theory*, 2nd ed., Ergeb. Math. Grenzgeb. (3), vol. 2, Springer-Verlag, Berlin, 1998.
- [Gar19] G. GARKUSHA – “Reconstructing rational stable motivic homotopy theory”, *Compositio Math.* **155** (2019), no. 7, p. 1424–1443.
- [Gil07] S. GILLE – “A graded Gersten-Witt complex for schemes with a dualizing complex and the Chow group”, *J. Pure Appl. Algebra* **208** (2007), no. 2, p. 391–419.
- [Gro64] A. GROTHENDIECK – “Éléments de géométrie algébrique. IV. étude locale des schémas et des morphismes de schémas. I”, *Publ. Math. Inst. Hautes Études Sci.* **20** (1964), p. 5–259.
- [Gro77] ———, *Cohomologie ℓ -adique et fonctions L*, Lect. Notes in Math., vol. 589, Springer-Verlag, 1977, Séminaire de Géométrie Algébrique du Bois-Marie 1965–66 (SGA 5).
- [Har66] R. HARTSHORNE – *Residues and duality*, Lect. Notes in Math., vol. 20, Springer-Verlag, Berlin-New York, 1966.
- [Héb11] D. HÉBERT – “Structure de poids à la Bondarko sur les motifs de Beilinson”, *Compositio Math.* **147** (2011), no. 5, p. 1447–1462.

- [Hoy14] M. HOYOIS – “A quadratic refinement of the Grothendieck-Lefschetz-Verdier trace formula”, *Algebraic Geom. Topol.* **14** (2014), no. 6, p. 3603–3658.
- [ILO14] L. ILLUSIE, Y. LASZLO & F. ORGOGOZO (eds.) – *Travaux de Gabber sur l’uniformisation locale et la cohomologie étale des schémas quasi-excellents*, Astérisque, vol. 363-364, Société Mathématique de France, Paris, 2014.
- [Jac17] J. JACOBSON – “Real cohomology and the powers of the fundamental ideal in the Witt ring”, *Ann. K-Theory* **2** (2017), no. 3, p. 357–385.
- [Jin16] F. JIN – “Borel-Moore motivic homology and weight structure on mixed motives”, *Math. Z.* **283** (2016), no. 3, p. 1149–1183.
- [Kha16] A. A. KHAN – “Motivic homotopy theory in derived algebraic geometry”, PhD Thesis, Universität Duisburg-Essen, 2016.
- [Kha19] ———, “Virtual fundamental classes of derived stacks I”, 2019, arXiv: 1909.01332.
- [Kha21] ———, “Voevodsky’s criterion for constructible categories of coefficients”, Preprint, available at <https://www.preschema.com/papers/six.pdf>, 2021.
- [Kne77] M. KNEBUSCH – “Symmetric bilinear forms over algebraic varieties”, in *Conference on Quadratic Forms—1976 (Kingston, Ont., 1976)*, Queen’s Papers in Pure and Appl. Math., vol. 46, 1977, p. 103–283.
- [Lam05] T. Y. LAM – *Introduction to quadratic forms over fields*, Graduate Studies in Math., vol. 67, American Mathematical Society, Providence, RI, 2005.
- [Lur09] J. LURIE – *Higher topos theory*, Annals of Math. Studies, vol. 170, Princeton University Press, Princeton, NJ, 2009.
- [Lur12] ———, “Higher algebra”, Preprint, available at <https://www.math.ias.edu/~lurie/papers/HigherAlgebra.pdf>, 2012.
- [Lur18] ———, “Spectral algebraic geometry”, Preprint, available at <https://www.math.ias.edu/~lurie/papers/SAG-rootfile.pdf>, 2018.
- [Mor04] F. MOREL – “On the motivic π_0 of the sphere spectrum”, in *Axiomatic, enriched and motivic homotopy theory*, NATO Sci. Ser. II Math. Phys. Chem., vol. 131, Kluwer Acad. Publ., 2004, p. 219–260.
- [Mor06] ———, “Rational stable splitting of Grassmannians and rational motivic sphere spectrum”, 2006.
- [Mor12] ———, *A^1 -algebraic topology over a field*, Lect. Notes in Math., vol. 2052, Springer, Heidelberg, 2012.
- [MV99] F. MOREL & V. VOEVODSKY – “ A^1 -homotopy theory of schemes”, *Publ. Math. Inst. Hautes Études Sci.* (1999), no. 90, p. 45–143.
- [Pan10] I. PANIN – “Homotopy invariance of the sheaf W_{Nis} and of its cohomology”, in *Quadratic forms, linear algebraic groups, and cohomology*, Dev. Math., vol. 18, Springer, New York, 2010, p. 325–335.
- [PW19] I. PANIN & C. WALTER – “On the motivic commutative ring spectrum BO ”, *St. Petersburg Math. J.* **30** (2019), no. 6, p. 933–972.
- [Rob15] M. ROBALO – “K-theory and the bridge from motives to noncommutative motives”, *Adv. Math.* **269** (2015), p. 399–550.
- [RØ08] O. RÖNDIGS & P. A. ØSTVÆR – “On modules over motivic ring spectra”, *Adv. Math.* **219** (2008), no. 2, p. 689–727.
- [Sch94] C. SCHEIDERER – *Real and étale cohomology*, Lect. Notes in Math., vol. 1588, Springer-Verlag, Berlin, 1994.
- [Sch17] M. SCHLICHTING – “Hermitian K-theory, derived equivalences and Karoubi’s fundamental theorem”, *J. Pure Appl. Algebra* **221** (2017), no. 7, p. 1729–1844.
- [ST15] M. SCHLICHTING & G. S. TRIPATHI – “Geometric models for higher Grothendieck-Witt groups in A^1 -homotopy theory”, *Math. Ann.* **362** (2015), no. 3-4, p. 1143–1167.
- [Sch00] A. SCHOLL – “Integral elements in K-theory and products of modular curves”, in *The arithmetic and geometry of algebraic cycles (Banff, AB, 1998)*, NATO Sci. Ser. C Math. Phys. Sci., vol. 548, Kluwer Acad. Publ., 2000, p. 467–489.
- [Spi18] M. SPITZWECK – *A commutative P^1 -spectrum representing motivic cohomology over Dedekind domains*, Mém. Soc. Math. France (N.S.), vol. 157, Société Mathématique de France, Paris, 2018.

- [Spi99] M. SPIVAKOVSKY – “A new proof of D. Popescu’s theorem on smoothing of ring homomorphisms”, *J. Amer. Math. Soc.* **12** (1999), no. 2, p. 381–444.
- [Sta21] STACKS PROJECT AUTHORS – “The stacks project”, <https://stacks.math.columbia.edu>, 2021.
- [Tho84] R. W. THOMASON – “Absolute cohomological purity”, *Bull. Soc. math. France* **112** (1984), no. 3, p. 397–406.
- [TT90] R. W. THOMASON & T. TROBAUGH – “Higher algebraic K -theory of schemes and of derived categories”, in *The Grothendieck Festschrift, Vol. III*, Progress in Math., vol. 88, Birkhäuser Boston, Boston, MA, 1990, p. 247–435.

Manuscript received 14th July 2020
accepted 18th January 2021

FRÉDÉRIC DÉGLISE, ENS de Lyon, UMPA, UMR 5669
46 allée d’Italie, 69364 Lyon Cedex 07, France
E-mail : frederic.deglise@ens-lyon.fr
Url : <http://perso.ens-lyon.fr/frederic.deglise/>

JEAN FASEL, Institut Fourier - UMR 5582, Université Grenoble-Alpes
CS 40700, 38058 Grenoble Cedex 9, France
E-mail : jean.fasel@gmail.com
Url : <https://www-fourier.univ-grenoble-alpes.fr/~fasel/j/>

FANGZHOU JIN, School of Mathematical Sciences, Tongji University
Siping Road 1239, 200092 Shanghai, China
E-mail : fangzhoujin@tongji.edu.cn
Url : <https://fangzhoujin.gitlab.io/>

ADEEL A. KHAN, Institut des Hautes Études Scientifiques
35 route de Chartres, 91440 Bures-sur-Yvette, France
and
Institute of Mathematics, Academia Sinica
Taipei 10617, Taiwan
E-mail : khan@ihes.fr
Url : <https://www.preschema.com/>