



Journal de l'École polytechnique

Mathématiques

Jérémi DARDÉ & Julien ROYER

Critical time for the observability of Kolmogorov-type equations

Tome 8 (2021), p. 859-894.

http://jep.centre-mersenne.org/item/JEP_2021__8__859_0

© Les auteurs, 2021.

Certains droits réservés.



Cet article est mis à disposition selon les termes de la licence
LICENCE INTERNATIONALE D'ATTRIBUTION CREATIVE COMMONS BY 4.0.
<https://creativecommons.org/licenses/by/4.0/>

L'accès aux articles de la revue « Journal de l'École polytechnique — Mathématiques » (<http://jep.centre-mersenne.org/>), implique l'accord avec les conditions générales d'utilisation (<http://jep.centre-mersenne.org/legal/>).

Publié avec le soutien
du Centre National de la Recherche Scientifique



Publication membre du
Centre Mersenne pour l'édition scientifique ouverte
www.centre-mersenne.org

CRITICAL TIME FOR THE OBSERVABILITY OF KOLMOGOROV-TYPE EQUATIONS

BY JÉRÉMI DARDÉ & JULIEN ROYER

ABSTRACT . — This paper is devoted to the observability of a class of two-dimensional Kolmogorov-type equations presenting a quadratic degeneracy. We give lower and upper bounds for the critical time. These bounds coincide in symmetric settings, giving a sharp result in these cases. The proof is based on Carleman estimates and on the spectral properties of a family of non-selfadjoint Schrödinger operators, in particular the localization of the first eigenvalue and Agmon type estimates for the corresponding eigenfunctions.

RÉSUMÉ (Temps critique pour l'observabilité d'équations de type Kolmogorov)

Nous nous intéressons à l'observabilité d'équations de type Kolmogorov bi-dimensionnelles présentant une dégénérescence quadratique. Nous donnons un majorant et un minorant du temps critique. Dans une configuration symétrique, ces bornes coïncident et donnent alors précisément le temps critique d'observabilité. La preuve est basée sur des estimées de Carleman et sur l'étude des propriétés spectrales d'une famille d'opérateurs de Schrödinger non auto-adjoints, en particulier la localisation de la première valeur propre et des estimées de type Agmon pour les fonctions propres correspondantes.

CONTENTS

1. Introduction.....	859
2. Strategy of the proof.....	863
3. Spectral properties of the Kolmogorov equation.....	867
4. The observability estimate in small time.....	885
References.....	893

1. INTRODUCTION

We study the observability properties of two-dimensional Kolmogorov-type equations with a quadratic degeneracy. Let $\ell_+, \ell_- > 0$. We set $I =]-\ell_-, \ell_+[$ and $\Omega = \mathbb{T} \times I$, where \mathbb{T} is the one-dimensional torus $\mathbb{R}/(2\pi\mathbb{Z})$. All along the paper, a generic point in Ω will be denoted by (x, y) , with $x \in \mathbb{T}$ and $y \in I$.

MATHEMATICAL SUBJECT CLASSIFICATION (2020). — 35K65, 93B07, 47B28, 47A10, 47D06.

KEYWORDS. — Observability, Kolmogorov equations, Carleman estimates, non-selfadjoint operators, resolvent estimates.

This work has been supported by the CIMI Labex, Toulouse, France, under grant ANR-11-LABX-0040-CIMI..

We consider $q \in C^3(\bar{I}, \mathbb{R})$ such that

$$q(0) = 0 \quad \text{and} \quad \min_{y \in I} q'(y) > 0.$$

In particular, $q(y) \neq 0$ for $y \neq 0$. The model case is $q(y) = y$.

Then, for $T > 0$, we consider on Ω the Kolmogorov-type equation

$$(1.1) \quad \begin{cases} \partial_t u + q(y)^2 \partial_x u - \partial_{yy} u = 0, & \text{on }]0, T[\times \Omega, \\ u(t, \cdot) = 0, & \text{on } \partial\Omega, \text{ for all } t \in]0, T[, \\ u|_{t=0} \in L^2(\Omega). \end{cases}$$

We are interested in the observability properties of the problem (1.1):

DEFINITION 1.1

(i) We say that (1.1) is observable in time T through an open subset ω of Ω if there exists $C > 0$ such that for any solution u of (1.1) we have

$$(1.2) \quad \|u(T)\|_{L^2(\Omega)}^2 \leq C \int_0^T \|u(t)\|_{L^2(\omega)}^2 dt.$$

(ii) We say that (1.1) is observable in time T through an open subset Γ of the boundary $\mathbb{T} \times \{-\ell_-, \ell_+\}$ of Ω if there exists $C > 0$ such that for any solution u of (1.1) we have

$$(1.3) \quad \|u(T)\|_{L^2(\Omega)}^2 \leq C \int_0^T \|\partial_\nu u(t)\|_{L^2(\Gamma)}^2 dt.$$

Null-controllability and observability properties of non-degenerate parabolic equations have been investigated for several decades now, since the pioneering works [Ego63] and [FR71] which proved independently the null-controllability of the one-dimensional heat-equation. Then [LR95] and [FI96] independently generalized this result in any dimension, showing that the heat equation is observable through any (interior or boundary) observation set, in any positive time, in any geometrical setting.

This is not the case for degenerate parabolic equations, which are a more recent subject of study. These equations may or may not be observable, depending on the location and the strength of the degeneracy, the geometrical setting, and the time horizon T . The case of a degeneracy of the equation at the boundary of the domain is now fairly well-understood (see [CMV16] and the references therein). In general, this type of degenerate equations are observable for weak degeneracy, and are not when the degeneracy becomes too strong.

In the case of interior degeneracy, there is no general theory, and equations are for the moment studied one after another. Interestingly, the known results show that, for precise strength of the degeneracy, a minimal time appears, under which observability is lost.

Among parabolic equations with interior degeneracy, the Grushin equation is so far the best understood: the two-dimensional case is now almost completely understood, and some partial results have been obtained in multi-dimensional settings

[BCG14, BMM15, Koe17, Lis19, BDE20, DK20, ABM20]. Other equations have also been studied, such as the heat equation on the Heisenberg group [BC17].

Finally, we highlight that a minimal time condition for observability might also appear for systems of parabolic equations, degenerate or not (see, among others, [AKBGBdT16, Dup17, BBM20]), for degenerate Schrödinger equations [BS19, LS20], and appears naturally for the wave equation (see [RT74, BLR92]). We point out that the closely related problem of approximate controllability is investigated in [LL17] for a large class of degenerate parabolic and hyperbolic equations.

Regarding the Kolmogorov equation (1.1), observability properties have already been investigated in the case $q(y) = y$, that is for the system

$$(1.4) \quad \begin{cases} \partial_t u + y^2 \partial_x u - \partial_{yy} u = 0, & \text{on }]0, T[\times \Omega, \\ u(t, \cdot) = 0, & \text{on } \partial\Omega, \text{ for all } t \in]0, T[, \\ u|_{t=0} \in L^2(\Omega). \end{cases}$$

It is proved in [Bea14] that a critical time \mathcal{T} appears for the observability through an open set of the form $\omega = \mathbb{T} \times]a, b[$ if $0 \notin]a, b[$:

THEOREM 1.2 ([Bea14]). — *Let $\omega = \mathbb{T} \times]a, b[$ with $-\ell_- < a < b < \ell_+$.*

(i) *If $a < 0 < b$, then the problem (1.4) is observable through ω in time T for any $T > 0$.*

(ii) *If $a > 0$ there exists $\mathcal{T} \geq a^2/2$ such that*

- *if $T > \mathcal{T}$ then (1.4) is observable through ω ,*
- *if $T < \mathcal{T}$ then (1.4) is not observable through ω .*

The model studied in [Bea14] also includes the equation

$$\partial_t u + y^\gamma \partial_x u - \partial_{yy} u = 0$$

with $\gamma = 1$. In that case, it is proved that the problem is observable through any open set ω , for any $T > 0$, generalizing the previous study [BZ09] where the sets of observation were horizontal strips. Theorem 1.2 corresponds to the case $\gamma = 2$. The case $\gamma = 3$ is studied in [BHHR15]. It is proved that if $0 < a < b < \ell_+$ then the problem is not observable through $\mathbb{T} \times (a, b)$ in any time $T > 0$.

The fact that the observation domain ω is a horizontal strip of Ω may seem quite restrictive. However, the recent study [Koe20] shows that it is a quasi-necessary condition for (1.4) to be observable.

THEOREM 1.3 ([Koe20]). — *Let $\omega = \omega_x \times I$, where ω_x is a strict open set of \mathbb{T} . Then (1.4) is not observable through ω in any time $T > 0$.*

Furthermore, it is shown that a minimal time is needed for the system to be possibly observable for most of observation sets ω .

THEOREM 1.4 ([Koe20]). — *Let ω be an open subset of $\mathbb{T} \times I$. Suppose that there exists $\tilde{x} \in \mathbb{T}$ and $a > 0$ such that*

$$\{(\tilde{x}, y) : y \in (-a, a)\} \cap \bar{\omega} = \emptyset.$$

Then the system (1.4) is not observable through ω in any time $T < a^2/2$.

In the present paper, we investigate the observability properties of (1.1) with a more general coefficient $q(y)^2$, when the domain of observation is the boundary

$$\Gamma = \partial\Omega = \mathbb{T} \times \{-\ell_-, \ell_+\}.$$

We could similarly consider observation through an open subset ω given by horizontal strips of Ω . Our main result is the following:

THEOREM 1.5. — *We set*

$$T_{\min} = \frac{1}{q'(0)} \min\left(\int_0^{\ell_+} q(s) ds, \int_{-\ell_-}^0 |q(s)| ds\right),$$

and

$$T_{\max} = \frac{1}{q'(0)} \max\left(\int_0^{\ell_+} q(s) ds, \int_{-\ell_-}^0 |q(s)| ds\right).$$

There exists $\bar{T} \in [T_{\min}, T_{\max}]$ such that

- (i) *if $T > \bar{T}$, the problem (1.1) is observable through Γ ,*
- (ii) *if $T < \bar{T}$, the problem (1.1) is not observable through Γ .*

In particular, in any configuration for which $T_{\max} = T_{\min}$, we obtain the critical time needed for observability of equation (1.1) to hold. This is in particular the case for symmetric configurations:

THEOREM 1.6. — *Suppose $\ell_- = \ell_+$ and q is odd. Let*

$$\bar{T} = \frac{1}{q'(0)} \int_0^{\ell_+} q(s) ds.$$

Then

- (i) *if $T > \bar{T}$ the problem (1.1) is observable through Γ ,*
- (ii) *if $T < \bar{T}$ the problem (1.1) is not observable through Γ .*

Note that in the case $q(y) = y$, the critical time is $\bar{T} = \ell_+^2/2$. This is the analog for the observation from the boundary of the time $a^2/2$ which appears in Theorems 1.2 and 1.4. Theorem 1.6 is, up to our knowledge, the first result giving the precise value of the critical time for the observation of a two-dimensional Kolmogorov-type equation.

REMARK 1.7. — By a classical duality argument, Theorem 1.5 is equivalent to controllability properties for the adjoint equation, with a boundary Dirichlet control acting on Γ . We refer to [TW09] for details on this equivalence.

Outline of the paper. — The article is organized as follows. After this introduction, we give in Section 2 the main ideas for the proof of Theorem 1.5. The details are then given in the following two sections. In Section 3 we discuss the well-posedness of the problem (1.1) and we prove some spectral properties for the non-selfadjoint Schrödinger operator $K_n = -\partial_{yy} + inq(y)^2$ which naturally appears in the analysis. We prove Agmon-type estimates for the first eigenfunction, which gives the negative result for $T < T_{\min}$, and we estimate the decay of the corresponding semigroup. Finally, in Section 4, we prove a Carleman estimate and deduce an observability estimate in arbitrarily small time which depends on the frequency n with respect to x . Together with the decay properties of e^{-tK_n} , this will give the observability of (1.1) for $T > T_{\max}$.

Acknowledgements. — We express our gratitude to Karine Beauchard, for enriching discussions on this work.

2. STRATEGY OF THE PROOF

In this section we describe the strategy for the proof of Theorem 1.5. We only give the main ideas, and the details will be postponed to the following two sections.

2.1. WELL-POSEDNESS AND FOURIER TRANSFORM OF THE KOLMOGOROV EQUATION

Before discussing the properties of the solutions of (1.1), we check that this problem is well-posed.

PROPOSITION 2.1. — *Let $u_\circ \in L^2(\Omega)$. Then there exists a unique*

$$u \in C^0([0, T], L^2(\Omega)) \cap C^0(]0, T], H^2(\Omega) \cap H_0^1(\Omega)) \cap C^1(]0, T], L^2(\Omega))$$

which satisfies (1.1) with $u(0) = u_\circ$.

Notice in particular that the equation is regularizing, so we do not have to impose any regularity on the initial condition to get a solution in the strong sense.

Many argument in our analysis, including the proof of Proposition 2.1, will be based on a Fourier transform. All along the paper, the Fourier coefficients are taken with respect to the variable $x \in \mathbb{T}$. Given $u \in L^2(\Omega)$, we denote by $u_n \in \ell^2(\mathbb{Z}, L^2(I))$ the sequence of Fourier coefficients of u :

$$u(x, y) = \sum_{n \in \mathbb{Z}} u_n(y) e^{inx}, \quad u_n(y) = \frac{1}{2\pi} \int_{\mathbb{T}} e^{-inx} u(x, y) dx.$$

The same applies if u (and then the $u_n, n \in \mathbb{Z}$) are also functions of the time t .

For $n \in \mathbb{Z}$ we consider the problem

$$(2.1) \quad \begin{cases} \partial_t u_n - \partial_{yy} u_n + inq(y)^2 u_n = 0, & \text{on }]0, T[\times I, \\ u_n(t, -\ell_-) = u_n(t, \ell_+) = 0, & \text{for } t \in]0, T[, \\ u_n(0) \in L^2(I). \end{cases}$$

Then the Fourier coefficients of a solution of (1.1) are given by the solutions of (2.1).

PROPOSITION 2.2. — *Let u be a solution of (1.1) and let u_n , $n \in \mathbb{Z}$, be the corresponding Fourier coefficients. Then for all $n \in \mathbb{Z}$ we have*

$$u_n \in C^0([0, T], L^2(I)) \cap C^0(]0, T], H^2(I) \cap H_0^1(I)) \cap C^1(]0, T], L^2(I)),$$

and u_n is the unique solution of (2.1) with $u_n(0) = u_{\mathbf{o},n}$, where $u_{\mathbf{o},n}$ is the n -th Fourier coefficient of $u_{\mathbf{o}} = u(0)$.

An important property of the problem (2.1) is the following exponential time decay.

PROPOSITION 2.3. — *Let*

$$\gamma < \frac{q'(0)}{\sqrt{2}}.$$

There exists $C > 0$ such that for $n \in \mathbb{Z}$, a solution u_n of (2.1) and $\theta_1, \theta_2 \in [0, T]$ with $\theta_1 \leq \theta_2$, one has

$$\|u_n(\theta_2)\|_{L^2(I)}^2 \leq C \exp(-2\gamma \sqrt{|n|} (\theta_2 - \theta_1)) \|u_n(\theta_1)\|_{L^2(I)}^2.$$

The proofs of Propositions 2.1 and 2.2 will be given in Section 3.1. Proposition 2.3 will be discussed in Section 3.3.

2.2. POSITIVE RESULT: UPPER BOUND FOR THE CRITICAL TIME. — We begin the proof of Theorem 1.5 with the first statement and prove observability for (1.1) when $T > T_{\max}$.

With the trace theorems, the regularity of the solution ensures that the right-hand side of (1.3) makes sense, even if it could be equal to $+\infty$ if the initial condition is not regular enough. In fact, we are going to prove the following stronger result for observability (note that with τ_1 chosen positive, the right-hand side of (2.2) is finite).

PROPOSITION 2.4. — *Let $T > T_{\max}$ and $\tau_1 \in]0, T - T_{\max}[$. Let $\tau_2 \in]\tau_1, T]$. Then there exists $C > 0$ such that for any solution u of (1.1) we have*

$$(2.2) \quad \|u(T)\|_{L^2(\Omega)}^2 \leq C \int_{\tau_1}^{\tau_2} \|\partial_\nu u(t)\|_{L^2(\partial\Omega)}^2 dt.$$

Obviously, Proposition 2.4 implies (1.3). The fact that we observe during an arbitrarily small time $\tau_2 - \tau_1$ might seem contradictory with the minimal time condition. It is not the case, since only the state at time $T > T_{\max}$ is controlled by the observation on the time interval $[\tau_1, \tau_2]$. As we will see below, the dissipation effect of the Kolmogorov equation plays a key role in obtaining (2.2). Roughly speaking, we have to wait long enough for the dissipation to fully play its role, and inequality (2.2) to be true.

By Proposition 2.2 and the Parseval identity, Proposition 2.4 is equivalent to an observability estimate for (2.1) uniform with respect to the Fourier parameter n . In other words, it is equivalent to prove the following result.

PROPOSITION 2.5. — *Let T , τ_1 and τ_2 be as in Proposition 2.4. There exists $C > 0$ such that for any $n \in \mathbb{Z}$ and any solution u_n of (2.1) one has*

$$(2.3) \quad \|u_n(T)\|_{L^2(I)}^2 \leq C \int_{\tau_1}^{\tau_2} (|\partial_y u_n(t, -\ell_-)|^2 + |\partial_y u_n(t, \ell_+)|^2) dt.$$

Note that it is sufficient to prove (2.3) for $n \in \mathbb{N}$. The case $n \in \mathbb{Z}$ then follows by complex conjugation of (2.1).

The difficulty in Proposition 2.5 is the uniformity with respect to the parameter n . For n fixed, it is already known that the one-dimensional heat equation with a complex-valued potential is observable through the boundary in any positive time:

PROPOSITION 2.6. — *Let $T > 0$ and $n \in \mathbb{N}$. Let $\tau_1, \tau_2 \in]0, T]$ with $\tau_1 < \tau_2$. There exists $C_n > 0$ such that for any solution u_n of (2.1) we have*

$$(2.4) \quad \|u_n(T)\|_{L^2(I)}^2 \leq C_n \int_{\tau_1}^{\tau_2} (|\partial_y u_n(t, -\ell_-)|^2 + |\partial_y u_n(t, \ell_+)|^2) dt.$$

A proof of Proposition 2.6 will be given in Section 4.2. With this result, it is now enough to prove Proposition 2.5 for n large. To do so, we first obtain a precise estimate of the constant C_n in the asymptotic n large.

PROPOSITION 2.7. — *Let $\tau_1, \tau_2 \in]0, T]$ with $\tau_1 < \tau_2$ and*

$$\kappa > \max\left(\frac{1}{\sqrt{2}} \int_0^{\ell_+} q(s) ds, \frac{1}{\sqrt{2}} \int_{-\ell_-}^0 |q(s)| ds\right) = \frac{q'(0)}{\sqrt{2}} T_{\max}.$$

There exist $n_0 \in \mathbb{N}$ and $C > 0$ such that for $n \geq n_0$ and a solution u_n of (2.1) one has

$$\|u_n(\tau_2)\|_{L^2(I)}^2 \leq C \exp(2\kappa\sqrt{n}) \int_{\tau_1}^{\tau_2} (|\partial_y u_n(t, -\ell_-)|^2 + |\partial_y u_n(t, \ell_+)|^2) dt.$$

The proof of this proposition is based on carefully constructed Carleman estimates, in the spirit of [BDE20]. We refer to Section 4.4 for the details.

The observability estimate of Proposition 2.7 is valid for any non-trivial interval of time, but it is not uniform with respect to n . As said above, the dissipation effect has to be taken into account here. More precisely, the second ingredient for the proof of Proposition 2.5 is the estimate given by Proposition 2.3, which precisely counterbalances the loss observed in Proposition 2.7 if we wait long enough.

Proof of Proposition 2.5, assuming Propositions 2.3, 2.6 and 2.7. — Let $\delta \in]0, 1[$ be so small that

$$(1 + \delta)T_{\max} < (1 - \delta)^2(T - \tau_1).$$

Then we set

$$\kappa = (1 + \delta) \frac{q'(0)}{\sqrt{2}} T_{\max}, \quad \gamma = (1 - \delta) \frac{q'(0)}{\sqrt{2}}.$$

Proposition 2.3 applied with $\theta_2 = T$ and

$$\theta_1 = \min(\tau_1 + \delta(T - \tau_1), \tau_2)$$

gives a constant $C_1 > 0$ such that for all $n \in \mathbb{N}$ and u_n solution of (2.1) we have

$$\|u_n(T)\|_{L^2(I)}^2 \leq C_1 \exp(-2\gamma\sqrt{n}(1 - \delta)(T - \tau_1)) \|u_n(\theta_1)\|_{L^2(I)}^2.$$

By Propositions 2.6 and 2.7, there exists $C_2 > 0$ such that for all $n \in \mathbb{N}$ and u_n solution of (2.1) we have

$$\|u_n(\theta_1)\|_{L^2(I)}^2 \leq C_2 \exp(2\kappa\sqrt{n}) \int_{\tau_1}^{\theta_1} (|\partial_y u_n(t, -\ell_-)|^2 + |\partial_y u_n(t, \ell_+)|^2) ds.$$

Since

$$\kappa - \gamma(1 - \delta)(T - \tau_1) = \frac{q'(0)}{\sqrt{2}} ((1 + \delta)T_{\max} - (1 - \delta)^2(T - \tau_1)) < 0,$$

these two inequalities give

$$\|u_n(T)\|_{L^2(I)}^2 \leq C_1 C_2 \int_{\tau_1}^{\tau_2} (|\partial_y u_n(t, -\ell_-)|^2 + |\partial_y u_n(t, \ell_+)|^2) dt,$$

and the proposition is proved. □

We recall that Proposition 2.5 implies Proposition 2.4 and hence the first statement of Theorem 1.5. Thus, it is enough to prove Propositions 2.3, 2.6 and 2.7 to get the observability of (1.1) through Γ for $T > T_{\max}$. These proofs are postponed to Sections 3 and 4.

2.3. NEGATIVE RESULT: LOWER BOUND FOR THE CRITICAL TIME. — In this paragraph we discuss the second statement of Theorem 1.5 about the non-observability of (1.1) if $T < T_{\min}$. The proof relies on the construction of a particular family of solutions of (1.1) for which the estimate (1.3) cannot hold if $T < T_{\min}$. In Section 3, we will prove the following result.

PROPOSITION 2.8. — *For all $n \in \mathbb{N}$, there exist $\lambda_n \in \mathbb{C}$ and $\psi_n \in H^2(I) \cap H_0^1(I)$ such that $\|\psi_n\|_{L^2(I)} = 1$,*

$$(2.5) \quad \lambda_n = \sqrt{n} q'(0) e^{i\pi/4} + o_{n \rightarrow +\infty}(\sqrt{n}),$$

and

$$(-\partial_{yy} + inq(y)^2)\psi_n = \lambda_n \psi_n.$$

Moreover, for any $\varepsilon > 0$ there exists $C > 0$ such that, for all $n \in \mathbb{N}$,

$$(2.6) \quad |\psi'_n(-\ell_-)|^2 + |\psi'_n(\ell_+)|^2 \leq Cn \exp(-\sqrt{2n}(1 - \varepsilon)q'(0)T_{\min}).$$

With this proposition we now prove that we cannot have observability through Γ in time $T < T_{\min}$.

Proof of Theorem 1.5.(ii), assuming Proposition 2.8. — Assume that (1.3) holds. For $m \in \mathbb{N}$, $t \in [0, T]$, $x \in \mathbb{T}$ and $y \in \bar{I}$ we set

$$u_m(t, x, y) = e^{-\lambda_m t} e^{imx} \psi_m(y),$$

where λ_m and ψ_m are given by Proposition 2.8. This defines a solution u_m of (1.1). Then (1.3) gives

$$2 \operatorname{Re}(\lambda_m) \leq C(e^{2T \operatorname{Re}(\lambda_m)} - 1) \left(|\psi'_m(-\ell_-)|^2 + |\psi'_m(\ell_+)|^2 \right).$$

Let $\varepsilon > 0$. By Proposition 2.8 there exists $C_1 > 0$ such that

$$(\sqrt{2} q'(0) + o(1))\sqrt{m} \leq C_1 m \exp\left(\sqrt{2m} q'(0) [T - (1 - \varepsilon)T_{\min} + o(1)]\right).$$

This implies

$$T \geq (1 - \varepsilon)T_{\min}.$$

Since this holds for any $\varepsilon > 0$, this implies that $T \geq T_{\min}$, and the conclusion follows. \square

3. SPECTRAL PROPERTIES OF THE KOLMOGOROV EQUATION

In this section we prove Propositions 2.1, 2.2, 2.3 and 2.8.

3.1. WELL-POSEDNESS AND FOURIER TRANSFORM OF THE KOLMOGOROV EQUATION

We begin with the well-posedness of the problems (1.1) and (2.1) for all $n \in \mathbb{Z}$. We also show that if u is a solution of (1.1) then its Fourier coefficients u_n , $n \in \mathbb{Z}$, are solutions of (2.1). We set

$$H_{0,y}^1(\Omega) = \{u \in L^2(\Omega) : \partial_y u \in L^2(\Omega), u(x, \ell_{\pm}) = 0 \text{ for almost all } x \in \mathbb{T}\}.$$

By the Poincaré inequality, this is a Hilbert space for the norm defined by

$$\|u\|_{H_{0,y}^1(\Omega)}^2 = \|\partial_y u\|_{L^2(\Omega)}^2.$$

We consider on $L^2(\Omega)$ the operator K defined by

$$Ku = -\partial_{yy}u + q(y)^2 \partial_x u$$

on the domain

$$\text{Dom}(K) = \{u \in H_{0,y}^1(\Omega) : Ku \in L^2(\Omega)\},$$

where Ku is understood in the sense of distributions. Similarly, for $n \in \mathbb{Z}$ we consider on $L^2(I)$ the operator

$$(3.1) \quad K_n = -\partial_{yy} + inq(y)^2,$$

defined on the domain (independent of n)

$$(3.2) \quad \text{Dom}(K_n) = H^2(I) \cap H_0^1(I).$$

We notice that K_0 is just the usual Dirichlet Laplacian on I . In particular, it is selfadjoint and non-negative. However, the operators K and K_n for $n \neq 0$ are not symmetric. We will show that they are at least accretive. For K this means that

$$\forall u \in \text{Dom}(K), \quad \text{Re} \langle Ku, u \rangle_{L^2(\Omega)} \geq 0.$$

In fact, they are even maximal accretive. This means in particular that any $z \in \mathbb{C}$ with $\text{Re}(z) < 0$ belongs to the resolvent set of K .

PROPOSITION 3.1

- (i) *The operator K is maximal accretive on $L^2(\Omega)$.*
- (ii) *For all $n \in \mathbb{Z}$, the operator K_n is maximal accretive on $L^2(I)$.*

(iii) Let $u \in \text{Dom}(K)$ and let $(u_n)_{n \in \mathbb{Z}}$ be the Fourier coefficients of u . Then u_n belongs to $\text{Dom}(K_n)$ for all $n \in \mathbb{Z}$ and the Fourier coefficients of Ku are the $K_n u_n$, $n \in \mathbb{Z}$.

Proof

– We begin with the second statement. It is easy to see that for $n \in \mathbb{Z}$ and $u \in \text{Dom}(K_n)$ we have

$$(3.3) \quad \text{Re} \langle K_n u, u \rangle = \|u'\|_{L^2(I)}^2 \geq 0,$$

which means that K_n is accretive. Then K_n is an accretive and bounded perturbation of the selfadjoint operator K_0 , so it is maximal accretive.

– Now let $u \in \text{Dom}(K)$ and $v = Ku \in L^2(\Omega)$. We denote by $(u_n)_{n \in \mathbb{Z}}, (v_n)_{n \in \mathbb{Z}} \in \ell^2(\mathbb{Z}, L^2(I))$ the sequences of Fourier coefficients of u and v , respectively. Let $n \in \mathbb{Z}$, $\phi_n \in C_0^\infty(I)$ and $\phi : (x, y) \mapsto e^{inx} \phi_n(y)$. By the Parseval identity we have

$$\begin{aligned} \langle u_n, -\phi_n'' - inq(y)^2 \phi_n \rangle_{L^2(I)} &= \frac{1}{2\pi} \langle u, -\partial_{yy} \phi - q(y)^2 \partial_x \phi \rangle_{L^2(\Omega)} \\ &= \frac{1}{2\pi} \langle v, \phi \rangle_{L^2(\Omega)} = \langle v_n, \phi_n \rangle_{L^2(I)}. \end{aligned}$$

This implies that $u_n'' \in L^2(I)$ (hence $u_n \in H^2(I)$) and

$$-u_n'' + inq(y)^2 u_n = v_n.$$

On the other hand, it is clear from the definition of u_n that $u_n(-\ell_-) = u_n(\ell_+) = 0$, so $u_n \in \text{Dom}(K_n)$. Then we can write $K_n u_n = v_n$. This gives the last statement of the proposition.

– As above we can see that the Fourier coefficients of $\partial_y u$ are the u_n' , $n \in \mathbb{Z}$. Then, by (3.3) and the Parseval identity we get

$$(3.4) \quad \|\partial_y u\|_{L^2(\Omega)}^2 = 2\pi \sum_{n \in \mathbb{Z}} \|u_n'\|_{L^2(I)}^2 = 2\pi \text{Re} \sum_{n \in \mathbb{Z}} \langle v_n, u_n \rangle = \text{Re} \langle v, u \rangle = \text{Re} \langle Ku, u \rangle.$$

– We check that the operator K is closed. Let $(u_m)_{m \in \mathbb{N}}$ be a sequence in $\text{Dom}(K)$ such that $u_m \rightarrow u$ and $Ku_m \rightarrow v$ in $L^2(\Omega)$, for some $u, v \in L^2(\Omega)$. In the sense of distributions we have

$$(3.5) \quad -\partial_{yy} u + q(y)^2 \partial_x u = \lim_{m \rightarrow +\infty} (-\partial_{yy} u_m + q(y)^2 \partial_x u_m) = v \in L^2(\Omega).$$

For $m, p \in \mathbb{N}$ we have $u_m - u_p \in \text{Dom}(K)$, so by (3.4) we have

$$\|u_m - u_p\|_{H_{0,y}^1(\Omega)}^2 = \text{Re} \langle K(u_m - u_p), u_m - u_p \rangle_{L^2(\Omega)} \xrightarrow{m \rightarrow +\infty} 0.$$

This implies that the sequence $(u_m)_{m \in \mathbb{N}}$ has a limit in $H_{0,y}^1(\Omega)$, which is necessarily u . By the trace theorem, we see that u_m also goes to u in $L^2(\partial\Omega)$, so u vanishes on $\partial\Omega$. Finally, we have proved that u belongs to $\text{Dom}(K)$ and, by (3.5), $Ku = v$. This proves that K is closed.

– By (3.4) the operator K is accretive on $L^2(\Omega)$. Then, for $u \in \text{Dom}(K)$ we have

$$(3.6) \quad \|(K + 1)u\|_{L^2(\Omega)}^2 \geq \|Ku\|_{L^2(\Omega)}^2 + \|u\|_{L^2(\Omega)}^2,$$

so $(K + 1)$ is injective with closed range. Now let $v \in L^2(\Omega)$ be such that

$$\forall u \in \text{Dom}(K), \quad \langle (K + 1)u, v \rangle = 0.$$

Then, in the sense of distributions we have

$$-\partial_{yy}v - q(y)^2\partial_x v + v = 0.$$

As above we can check that the operator $\tilde{K} = -\partial_{yy} - q(y)^2\partial_x$, defined on the domain

$$\text{Dom}(\tilde{K}) = \{u \in H_{0,y}^1(\Omega) : \tilde{K}u \in L^2(\Omega)\},$$

is accretive. This implies that $v = 0$ (in fact, \tilde{K} is the adjoint of K). Thus $\text{Ran}(K + 1)^\perp = \{0\}$ and $(K + 1)$ is invertible. By (3.6), its inverse is bounded. This proves that -1 belongs to the resolvent set of K , and hence K is maximal accretive. \square

By the Lummer-Philipps theorem (see for instance [EN00]), the operator $(-K)$ generates a contractions semigroup $(e^{-tK})_{t \geq 0}$ on $L^2(\Omega)$. Given $u_o \in \text{Dom}(K)$, the function $u : t \mapsto e^{-tK}u$ belongs to $C^0(\mathbb{R}_+, \text{Dom}(K)) \cap C^1(\mathbb{R}_+, L^2(\Omega))$. This gives a strong solution of (1.1). More generally, for $u_o \in L^2(\Omega)$, the function $t \mapsto e^{-tK}u_o$ belongs to $C^0(\mathbb{R}_+, L^2(\Omega))$. This gives a weak solution of (1.1). The same applies on $L^2(I)$ to K_n and (2.1) for any $n \in \mathbb{Z}$. In particular, if u_n is a solution of (2.1) then for all $t_1, t_2 \in [0, T]$ such that $t_1 \leq t_2$ we have

$$(3.7) \quad \|u_n(t_2)\|_{L^2(I)}^2 \leq \|u_n(t_1)\|_{L^2(I)}^2.$$

Now, we show that the solutions of (2.1) for $n \in \mathbb{Z}$ give the Fourier coefficients of a solution of (1.1).

PROPOSITION 3.2. — *Let $u_o \in L^2(\Omega)$. For $t \geq 0$ we set $u(t) = e^{-tK}u_o \in L^2(\Omega)$. We denote by $(u_{o,n})_{n \in \mathbb{Z}}$ and $(u_n(t))_{n \in \mathbb{Z}}$ the Fourier coefficients of u_o and $u(t)$, $t \geq 0$, respectively. Then for all $n \in \mathbb{Z}$ and $t \geq 0$ we have*

$$u_n(t) = e^{-tK_n}u_{o,n}.$$

Proof. — First assume that $u_o \in \text{Dom}(K)$. Let $n \in \mathbb{Z}$ and $t \geq 0$. By differentiation under the integral sign and Proposition 3.1 we have in $L^2(I)$, for $h > 0$,

$$\begin{aligned} \frac{u_n(t+h) - u_n(t)}{h} &= \frac{1}{2\pi} \int_{\mathbb{T}} e^{-inx} \frac{u(t+h, x) - u(t, x)}{h} dx \\ &\xrightarrow{h \rightarrow 0} -\frac{1}{2\pi} \int_{\mathbb{T}} e^{-inx} K u(t, x) dx = -K_n u_n(t). \end{aligned}$$

The conclusion follows in this case.

In general, since $\text{Dom}(K)$ is dense in $L^2(\Omega)$, we can consider a sequence $(u_o^m)_{m \in \mathbb{N}}$ in $\text{Dom}(K)$ which converges to u_o in $L^2(\Omega)$. For $m \in \mathbb{N}$ we denote by $u_{o,n}^m$, $n \in \mathbb{Z}$,

the Fourier coefficients of u_o^m . Then $u_{o,n}^m$ goes to $u_{o,n}$ in $L^2(I)$ for all $n \in \mathbb{Z}$ and, by continuity of e^{-tK} and e^{-tK_n} in $L^2(\Omega)$ and $L^2(I)$ respectively,

$$\begin{aligned} u_n(t) &= \frac{1}{2\pi} \int_{\mathbb{T}} e^{-inx} (e^{-tK} u_o)(x) dx = \lim_{m \rightarrow +\infty} \frac{1}{2\pi} \int_{\mathbb{T}} e^{-inx} (e^{-tK} u_o^m)(x) dx \\ &= \lim_{m \rightarrow +\infty} e^{-tK_n} u_{o,n}^m = e^{-tK_n} u_{o,n}. \end{aligned}$$

The proposition is proved. □

Before we can state Theorem 1.5, we still have to check that the right-hand side of (1.3) makes sense (one would not have this difficulty with observability through an open subset of Ω). To do so, we investigate the regularizing effect of equation (1.1), and prove that even if the initial condition u_o merely belongs to $L^2(\Omega)$, the solution is smooth enough for the right-hand side of (1.3) to be well defined. The proof of this result relies on Proposition 2.3, which will be proved in Section 3.3 below.

PROPOSITION 3.3. — *For $u_o \in L^2(\Omega)$ and $\tau > 0$ we have*

$$e^{-\tau K} u_o \in H^2(\Omega) \cap H_0^1(\Omega) \subset \text{Dom}(K).$$

Proof

– For $t > 0$ we set $u(t) = e^{-tK} u_o$. We denote by $u_{o,n}$ and $u_n(t)$, $n \in \mathbb{Z}$, the Fourier coefficients of u_o and $u(t)$, respectively. For $n \in \mathbb{Z}$ and $t > 0$ we have $u_n(t) = e^{-tK_n} u_{o,n}$ by Proposition 3.2.

– By Proposition 2.3 there exists $c > 0$ such that for $\tau > 0$ and $k \in \mathbb{N}$ we have

$$(3.8) \quad n^k \|u_n(\tau)\|_{L^2(I)} \leq \frac{c}{\tau^{2k}} \|u_{o,n}\|_{L^2(I)}.$$

This implies in particular that $u(\tau) \in C^\infty(\mathbb{T}, L^2(I))$.

– Let $\tau > 0$. Assume that $u_o \in C_0^\infty(\Omega) \subset \text{Dom}(K^2)$. Then $u \in C^1([0, \tau], \text{Dom}(K))$ and $u_n \in C^1([0, \tau], \text{Dom}(K_n))$ for all $n \in \mathbb{Z}$. Let $n \in \mathbb{Z}$. Since $(\partial_t + K_n)u_n = 0$ we have

$$\begin{aligned} 0 &= \text{Re} \int_{\tau}^{2\tau} (t - \tau) \langle (\partial_t + K_n)u_n(t), \partial_t u_n(t) \rangle_{L^2(I)} dt \\ &= \int_{\tau}^{2\tau} (t - \tau) \|\partial_t u_n(t)\|_{L^2(I)}^2 dt + \int_{\tau}^{2\tau} (t - \tau) \text{Re} \langle K_n u_n(t), \partial_t u_n(t) \rangle_{L^2(I)} dt. \end{aligned}$$

Since

$$\begin{aligned} \text{Re} \langle K_n u_n(t), \partial_t u_n(t) \rangle &= \text{Re} \langle -\partial_{yy} u_n(t), \partial_t u_n(t) \rangle + \text{Re} \langle inq(y)^2 u_n(t), \partial_t u_n(t) \rangle \\ &\geq \frac{1}{2} \frac{d}{dt} \langle -\partial_{yy} u_n(t), u_n(t) \rangle - n \|q\|_\infty^2 \|u_n(t)\| \|\partial_t u_n(t)\| \\ &\geq \frac{1}{2} \frac{d}{dt} \text{Re} \langle K_n u_n(t), u_n(t) \rangle - \frac{n^2 \|q\|_\infty^4 \|u_n(t)\|^2}{2} - \frac{\|\partial_t u_n(t)\|^2}{2}, \end{aligned}$$

with (3.7) this gives

$$\int_{\tau}^{2\tau} (t - \tau) \frac{\|\partial_t u_n(t)\|^2}{2} dt \leq -\frac{1}{2} \operatorname{Re} \int_{\tau}^{2\tau} (t - \tau) \frac{d}{dt} \langle K_n u_n(t), u_n(t) \rangle dt + \frac{n^2 \tau^2 \|q\|_{\infty}^4 \|u_n(\tau)\|^2}{4}.$$

An integration by parts gives

$$\begin{aligned} -\operatorname{Re} \int_{\tau}^{2\tau} (t - \tau) \frac{d}{dt} \langle K_n u_n(t), u_n(t) \rangle dt &= -\tau \operatorname{Re} \langle K_n u_n(2\tau), u_n(2\tau) \rangle + \int_{\tau}^{2\tau} \operatorname{Re} \langle K_n u_n(t), u_n(t) \rangle dt \\ &\leq -\frac{1}{2} \int_{\tau}^{2\tau} \frac{d}{dt} \|u_n(t)\|^2 dt \leq \frac{\|u_n(\tau)\|^2}{2} - \frac{\|u_n(2\tau)\|^2}{2}. \end{aligned}$$

On the other hand, since the function $t \mapsto \partial_t u_n(t)$ is also a solution of (1.1), its norm is non-increasing, so

$$\frac{1}{2} \int_{\tau}^{2\tau} (t - \tau) \|\partial_t u_n(t)\|^2 dt \geq \frac{\tau^2 \|\partial_t u_n(2\tau)\|^2}{4}.$$

Finally, with (3.8) we get

$$\begin{aligned} \|\partial_t u_n(2\tau)\|^2 + \frac{2\|u_n(2\tau)\|^2}{\tau^2} &\leq \frac{2\|u_n(\tau)\|^2}{\tau^2} + n^2 \|q\|_{\infty}^4 \|u_n(\tau)\|^2 \\ &\leq \frac{2\|u_{o,n}\|^2}{\tau^2} + \frac{c^2 \|q\|_{\infty}^4 \|u_{o,n}\|^2}{\tau^4}. \end{aligned}$$

Hence, by the Parseval identity,

$$(3.9) \quad \|Ku(2\tau)\|^2 = \|\partial_t u(2\tau)\|_{L^2(\Omega)}^2 \leq \frac{2\|u_o\|^2}{\tau^2} + \frac{c^2 \|q\|_{\infty}^4 \|u_o\|^2}{\tau^4}.$$

– Let $u_o \in L^2(\Omega)$ and $(u_{o,m})_{m \in \mathbb{N}}$ be a sequence in $C_0^\infty(\Omega)$ which goes to u_o in $L^2(\Omega)$. For $\tau > 0$ we set $u(\tau) = e^{-\tau K} u_o$ and $u_m(\tau) = e^{-\tau K} u_{o,m}$, $m \in \mathbb{N}$. Let $\delta > 0$. $u_m(t)$ converges to $u(t)$ for any $t \geq 0$ and the function $t \mapsto u'_m(t)$ has a uniform limit on $[\delta, +\infty[$. This implies that the function u belongs to $C^1(]0, +\infty[, L^2(\Omega))$. Then, since $-K$ is the generator of the semigroup e^{-tK} , $u(t)$ belongs to $\operatorname{Dom}(K)$ for all $t > 0$ and $u'(t) = -Ku(t)$.

– Finally, for $u_o \in L^2(\Omega)$ and $\tau > 0$ we have $(-\partial_{yy} - q(y)^2 \partial_x)u(\tau) \in L^2(\Omega)$ and $\partial_x u(\tau) \in L^2(\Omega)$, so $-\partial_{yy} u(\tau) \in L^2(\Omega)$. Since we also have $\partial_{xx} u(\tau) \in L^2(\Omega)$, this proves that $u(\tau)$ belongs to $H^2(\Omega)$. The fact that $u(\tau)$ is also in $H_0^1(\Omega)$ is a consequence of the fact that it is in $\operatorname{Dom}(K) \subset H_{0,y}^1(\Omega)$, and the proof is complete. \square

3.2. GENERAL SPECTRAL PROPERTIES FOR NON-SELFADJOINT SCHRÖDINGER OPERATORS

In the rest of this section, we prove Propositions 2.3 and 2.8. They can both be rewritten in terms of the operator K_n defined by (3.1)-(3.2).

We have seen in Proposition 3.1 that K_n is a maximal accretive operator on $L^2(I)$. In particular, the resolvent set of K_n is not empty. And since $\operatorname{Dom}(K_n)$ is compactly

embedded in $L^2(I)$, the resolvent of K_n is compact. This implies that the spectrum of K_n consists of eigenvalues which have finite algebraic multiplicities.

We have already said that $(-K_n)$ generates a contractions semigroup on $L^2(I)$ (see (3.7)). However, this is not enough for Proposition 2.3. For $n = 0$, the operator K_0 is selfadjoint and the decay of the corresponding semigroup is given by the functional calculus. If we denote by λ_0 the first eigenvalue of K_0 , then λ_0 is positive and for all $t \geq 0$ we have

$$\|e^{-tK_0}\|_{L(L^2(I))} \leq e^{-t\lambda_0}.$$

For $n \neq 0$, the operator K_n is not selfadjoint, and the link between the exponential decay of e^{-tK_n} and the real parts of the eigenvalues of K_n is not that direct.

The purpose of the rest of this section is then to give some spectral properties for the non-selfadjoint operator K_n . We are interested in the location of the spectrum (and in particular the eigenvalue with the smallest real part), the size of the resolvent $(K_n - z)^{-1}$ for z outside this spectrum (for a non-selfadjoint operator, the resolvent can have a large norm even for z far from the spectrum) and then an estimate of the propagator e^{-tK_n} for $t \geq 0$.

The properties of the operator K_n will be deduced from analogous results for the classical complex harmonic oscillators and the complex Airy operators.

With the Agmon estimates (see Section 3.4 below), we will see that for large n the eigenvectors of K_n associated to “small” eigenvalues should be in some sense localized close to 0. And near 0 we have

$$inq(y)^2 \sim inq'(0)^2 y^2.$$

Thus, it is expected that, at least for a small spectral parameter, the spectral properties of K_n for large n should be close to those of the harmonic oscillator

$$(3.10) \quad H_n = -\partial_{yy} + inq'(0)^2 y^2,$$

defined on the domain

$$\text{Dom}(H_n) = \{u \in H^2(\mathbb{R}) : yu \in H^1(\mathbb{R}), y^2 u \in L^2(\mathbb{R})\}.$$

It is known (see for instance [Hel13, §14.4]) that H_n defines for all $n \in \mathbb{N}^*$ a maximal accretive operator on $L^2(\mathbb{R})$. Its spectrum consists of a sequence of (geometrically and algebraically) simple eigenvalues, given by

$$(2k - 1)\sqrt{n} q'(0) e^{i\pi/4}, \quad k \in \mathbb{N}^*,$$

and for each $k \in \mathbb{N}^*$, a corresponding eigenfunction is given by

$$(3.11) \quad y \mapsto P_k(e^{i\pi/8} \alpha y) e^{-((\alpha y)^2/2\sqrt{2}) - i(\alpha y)^2/2\sqrt{2}}, \quad \alpha = n^{1/4} q'(0)^{1/2},$$

where P_k is a polynomial of degree k . In particular,

$$\inf_{\sigma \in \text{Sp}(H_n)} \text{Re}(\sigma) = \frac{\sqrt{n} q'(0)}{\sqrt{2}}.$$

This is not enough to get a decay estimate for the propagator e^{-tH_n} , $t \geq 0$. However, it is also known that for $\gamma < q'(0)/\sqrt{2}$ there exists $c > 0$ such that

$$(3.12) \quad \sup_{\operatorname{Re}(z) \leq \gamma\sqrt{n}} \|(H_n - z)^{-1}\|_{L(L^2(\mathbb{R}))} \leq \frac{c}{\sqrt{n}}$$

(in fact we have more precise resolvent estimates [HSV13, KSTV15]). Then we deduce (see for instance [EN00, Th. V.1.11]) that there exists $C > 0$ such that for all $t \geq 0$ we have

$$(3.13) \quad \|e^{-tH_n}\|_{L(L^2(\mathbb{R}))} \leq Ce^{-t\gamma\sqrt{n}}.$$

Proposition 2.3 precisely says that we have a similar estimate for the propagator generated by K_n , while Proposition 2.8 shows that for large n the first eigenvalue of K_n is close to the first eigenvalue of H_n . The decay of the corresponding eigenfunction has the same form as in (3.11), but it depends on the values of q on the whole interval I , and not only on its behavior in a neighborhood of 0.

For the proofs, we will also compare K_n to some complex Airy operators. Near $y_0 \in I \setminus \{0\}$, the potential $inq(y)^2$ looks like $inq(y_0)^2 + 2inq(y_0)q'(y_0)(y - y_0)$, with $q(y_0)q'(y_0) \neq 0$. It is then useful to recall the properties of Schrödinger operators with linear purely imaginary potentials.

Given $\alpha \in \mathbb{R} \setminus \{0\}$, we consider on $L^2(\mathbb{R})$ the operator

$$(3.14) \quad A_\alpha u = -\partial_{yy}u + i\alpha yu,$$

defined on the domain

$$\operatorname{Dom}(A_\alpha) = \{u \in L^2(\mathbb{R}) : (-u'' + i\alpha yu) \in L^2(\mathbb{R})\}.$$

This complex Airy operator is now well understood, see for instance [Hel11, KS15] and references therein. We notice that for $\alpha > 0$ we have

$$\Theta_\alpha^{-1}(A_\alpha - z)^{-1}\Theta_\alpha = \frac{1}{\alpha^{2/3}}(A_1 - z\alpha^{-2/3})^{-1},$$

where Θ_α is the unitary operator defined on $L^2(\mathbb{R})$ by

$$(\Theta_\alpha u)(y) = \alpha^{1/6}u(\alpha^{1/3}y).$$

Moreover, $A_{-\alpha} = A_\alpha^*$. Then, from the properties of A_1 we deduce the following result.

PROPOSITION 3.4

- (i) *The spectrum of A_α is empty for any $\alpha \in \mathbb{R} \setminus \{0\}$.*
- (ii) *Let $\gamma \in \mathbb{R}$. Then there exists $c > 0$ such that for all $\alpha \in \mathbb{R} \setminus \{0\}$ we have*

$$\sup_{\operatorname{Re}(z) \leq \gamma|\alpha|^{2/3}} \|(A_\alpha - z)^{-1}\| \leq \frac{c}{|\alpha|^{2/3}}.$$

To understand the behavior of K_n near the boundary points $\pm\ell_\pm$, we introduce the complex Airy operator on \mathbb{R}_+ . For $\alpha \in \mathbb{R} \setminus \{0\}$ we consider on $L^2(\mathbb{R}_+)$ the operator defined by

$$A_\alpha^+ u = -\partial_{yy}u + i\alpha yu,$$

on the domain

$$\text{Dom}(A_\alpha^+) = \{u \in L^2(\mathbb{R}_+) : (-u'' + i\alpha y u) \in L^2(\mathbb{R}_+) \text{ and } u(0) = 0\}.$$

To prove the following proposition, we use in $L^2(\mathbb{R}_+)$ the same dilation Θ_a as above and we apply [Hel11, Lemma 5.1]. For the properties of the Airy function we refer for instance to [VS10].

PROPOSITION 3.5

(i) *Let $\alpha > 0$. The spectrum of A_α^+ consists of a sequence of simple eigenvalues. These eigenvalues are given by*

$$\lambda_k^+ = \alpha^{2/3} e^{i\pi/3} |\mu_k|, \quad k \in \mathbb{N}^*,$$

where $\dots < \mu_k < \dots < \mu_2 < \mu_1 < 0$ are the zeros of the Airy function.

(ii) *Let $\gamma < |\mu_1|/2$. There exists $C > 0$ such that for all $\alpha \in \mathbb{R} \setminus \{0\}$ we have*

$$\sup_{\text{Re}(z) \leq \gamma |\alpha|^{2/3}} \|(A_\alpha^+ - z)^{-1}\| \leq \frac{c}{|\alpha|^{2/3}}.$$

Of course, we have similar properties on $L^2(\mathbb{R}_-)$ for the operator

$$A_\alpha^- : u \mapsto -\partial_{yy} u + i\alpha y u,$$

defined on the domain

$$\text{Dom}(A_\alpha^-) = \{u \in L^2(\mathbb{R}_-) : (-u'' + i\alpha y u) \in L^2(\mathbb{R}_-) \text{ and } u(0) = 0\}.$$

3.3. RESOLVENT ESTIMATES. — In this paragraph, we prove Proposition 2.3 (see Proposition 3.7 below) and the first part of Proposition 2.8, about the eigenvalue λ_n (see Proposition 3.9). The estimate of a corresponding eigenfunction at the boundary will be given in the next paragraph.

We prove estimates for the resolvent $(K_n - z)^{-1}$ when z has real part smaller than $\gamma\sqrt{n}$, with γ as in Proposition 2.3. More precisely, we estimate the difference between $(K_n - z)^{-1}$ and the model resolvent $(H_n - z)^{-1}$, in a suitable sense. By the theory of semigroups, this will give Proposition 2.3. This will also give the existence of an eigenvalue λ_n which satisfies (2.5).

To compare $(K_n - z)^{-1}$ and $(H_n - z)^{-1}$, we follow the ideas of [Hen14]. Our one-dimensional setting is simpler than the general case considered therein so, for the reader convenience, we provide a complete proof adapted to our problem. Notice also that (2.5) is not contained in the results given in [Hen14], where the imaginary parts of the eigenvalues are not an issue.

We denote by $\mathbf{1}_I$ the operator which maps $u \in L^2(\mathbb{R})$ to its restriction on I : $\mathbf{1}_I u = u|_I \in L^2(I)$. Then $\mathbf{1}_I^*$ maps a function $v \in L^2(I)$ to its extension by 0 on \mathbb{R} .

PROPOSITION 3.6

(i) *Let $\gamma \in]0, q'(0)/\sqrt{2}[$. There exist $n_0 \in \mathbb{N}^*$ and $c > 0$ such that for $n \geq n_0$ and $z \in \mathbb{C}$ with $\text{Re}(z) \leq \gamma\sqrt{n}$ we have $z \in \rho(K_n)$ and*

$$\|(K_n - z)^{-1}\|_{L(L^2(I))} \leq \frac{c}{\sqrt{n}}.$$

(ii) We have

$$\|\mathbb{1}_I^* K_n^{-1} \mathbb{1}_I - H_n^{-1}\|_{L(L^2(\mathbb{R}))} = o_{n \rightarrow +\infty}(1/\sqrt{n}).$$

Proof. — The proof consists in using localized versions of the resolvents of the complex harmonic operator H_n and of Airy-type operators to construct an approximation $Q_n(z)$ of the resolvent $(K_n - z)^{-1}$. We first introduce suitable cut-off functions, then we define $Q_n(z)$ and finally we check that it is indeed an approximation of $(K_n - z)^{-1}$ up to a uniformly bounded operator. The proposition will then follow from estimates on $Q_n(z)$. For $n \in \mathbb{N}^*$ we set

$$\mathbb{C}_n^- = \{z \in \mathbb{C} : \operatorname{Re}(z) \leq \gamma\sqrt{n}\}.$$

– For $n \in \mathbb{N}^*$ and $z \in \mathbb{C}_n^-$ we set

$$R_n(z) = \mathbb{1}_I(H_n - z)^{-1} \mathbb{1}_I^*.$$

This defines a bounded operator on $L^2(I)$. Our purpose is to prove that $R_n(z)$ gives an approximate inverse of $(K_n - z)$ near 0, in the following sense. We consider

$$\rho \in]1/6, 1/4[,$$

a cut-off function $\chi \in C_0^\infty(\mathbb{R}, [0, 1])$ supported in I and equal to 1 on a neighborhood of 0, and for $n \in \mathbb{N}^*$ and $y \in \bar{I}$ we set

$$\chi_n(y) = \chi(n^\rho y).$$

Then we set

$$T_n(z) = R_n(z)\chi_n(K_n - z) - \chi_n.$$

We prove that $T_n(z)$ extends to a bounded operator on $L^2(I)$ and

$$(3.15) \quad \|T_n(z)\|_{L(L^2(I))} \xrightarrow{n \rightarrow +\infty} 0,$$

where the convergence is uniform with respect to $z \in \mathbb{C}_n^-$.

Let $u \in \operatorname{Dom}(K_n)$. For $n \in \mathbb{N}^*$ we have $\chi_n u \in \operatorname{Dom}(K_n)$ and $\mathbb{1}_I^* \chi_n u \in \operatorname{Dom}(H_n)$. For $y \in \bar{I}$ we set

$$r(y) = q(y)^2 - q'(0)^2 y^2.$$

Then for $z \in \mathbb{C}_n^-$ we have

$$R_n(z)(K_n - z)\chi_n u = \chi_n u + inR_n(z)r\chi_n u.$$

This gives

$$(3.16) \quad R_n(z)\chi_n(K_n - z)u = \chi_n u + inR_n(z)r\chi_n u - R_n(z)\chi_n'' u + 2R_n(z)(\chi_n' u)'$$

Since $|r(y)\chi_n(y)| \lesssim n^{-3\rho}$, we have by (3.12)

$$\|nR_n(z)r\chi_n\|_{L(L^2(I))} \lesssim n^{1-3\rho-1/2} \xrightarrow{n \rightarrow +\infty} 0.$$

We also have

$$\|R_n(z)\chi_n''\|_{L(L^2(I))} \lesssim n^{2\rho-1/2} \xrightarrow{n \rightarrow +\infty} 0.$$

For the last term we observe that for $v \in L^2(\mathbb{R})$ we have

$$\begin{aligned} \|\partial_y(H_n^* - \bar{z})^{-1}v\|_{L^2(\mathbb{R})}^2 &= \operatorname{Re} \langle v, (H_n^* - \bar{z})^{-1}v \rangle_{L^2(\mathbb{R})} + \operatorname{Re}(z) \|(H_n^* - \bar{z})^{-1}v\|_{L^2(\mathbb{R})}^2 \\ &\lesssim \frac{\|v\|_{L^2(\mathbb{R})}^2}{\sqrt{n}}. \end{aligned}$$

Taking the adjoint gives

$$\begin{aligned} \|R_n(z)\partial_y(\chi'_n u)\|_{L^2(I)} &\leq \|(H_n - z)^{-1}\partial_y(\mathbf{1}_I^* \chi'_n u)\|_{L^2(\mathbb{R})} \\ &\lesssim n^{-1/4} \|\mathbf{1}_I^* \chi'_n u\|_{L^2(\mathbb{R})} \lesssim n^{\rho-1/4} \|u\|_{L^2(I)}, \end{aligned}$$

and (3.15) follows.

– Then we consider

$$\tilde{\rho} \in](1 + 2\rho)/6, (1 - \rho)/3[.$$

In particular, $\tilde{\rho} > \rho$. For $n \in \mathbb{N}^*$ we denote by ν_n the integer part of $1 + (\ell_+ + \ell_-)n^{\tilde{\rho}}$, and for $j \in \{0, \dots, \nu_n\}$ we set

$$a_{j,n} = -\ell_- + j\delta_n, \quad \delta_n = \frac{\ell_+ + \ell_-}{\nu_n}.$$

We also consider $\theta \in C_0^\infty(\mathbb{R})$ supported in $] - 2/3, 2/3[$, equal to 1 on $[- 1/3, 1/3]$ and such that $\theta(-y) = 1 - \theta(1 - y)$ for $y \in [0, 1]$. Then for all $y \in \mathbb{R}$ we have

$$\sum_{m \in \mathbb{Z}} \theta(y - m) = 1.$$

For $n \in \mathbb{N}^*$, $j \in \{0, \dots, \nu_n\}$ and $y \in \bar{I}$ we set

$$\theta_{j,n}(y) = \theta((y - a_{j,n})/\delta_n)(1 - \chi_n)(y).$$

For $n \in \mathbb{N}^*$ large enough and $j \in \{1, \dots, \nu_n - 1\}$ such that $\theta_{j,n} \neq 0$ we have $|a_{j,n}| \gtrsim n^{-\rho}$, and in particular $a_{j,n} \neq 0$. We define $A_{j,n}$ by

$$A_{j,n}u = -\partial_{yy}u + inq(a_{j,n})^2u + 2inq(a_{j,n})q'(a_{j,n})(y - a_{j,n})u$$

on the domain

$$\operatorname{Dom}(A_{j,n}) = \{u \in H^2(\mathbb{R}) : yu \in L^2(\mathbb{R})\}.$$

With the notation (3.14) we have

$$A_{j,n} = \tau_{a_{j,n}} A_{2n(qq')(a_{j,n})} \tau_{-a_{j,n}} + inq(a_{j,n})^2,$$

where $\tau_{\pm a_{j,n}}$ is the usual translation operator: $(\tau_{\pm a_{j,n}} u)(y) = u(y \mp a_{j,n})$. Thus $A_{j,n}$ satisfies the properties of Proposition 3.4 with $\alpha = 2n(qq')(a_{j,n})$. We similarly set

$$A_{0,n} = \tau_{-\ell_-} A_{2n(qq')(-\ell_-)}^+ \tau_{\ell_-} + inq(-\ell_-)^2$$

and

$$A_{\nu_n,n} = \tau_{\ell_+} A_{2n(qq')(\ell_+)}^- \tau_{-\ell_+} + inq(\ell_+)^2.$$

Notice that $A_{0,n}$ and $A_{\nu_n,n}$ are operators on $L^2(-\ell_-, +\infty)$ and $L^2(-\infty, \ell_+)$, respectively. They satisfy the same properties as the model operators (see Proposition 3.5).

For $j \in \{1, \dots, \nu_n - 1\}$ we set $\mathbb{1}_j = \mathbb{1}_I$. We also denote by $\mathbb{1}_0$ the operator which maps $u \in L^2(-\ell_-, +\infty)$ to its restriction on I , and by $\mathbb{1}_{\nu_n}$ the operator which maps $u \in L^2(-\infty, \ell_+)$ to its restriction to I . For $z \in \mathbb{C}_n^-$ we set

$$R_{j,n}(z) = \mathbb{1}_j (A_{j,n} - z)^{-1} \mathbb{1}_j^*$$

and

$$T_{j,n}(z) = R_{j,n}(z)\theta_{j,n}(K_n - z) - \theta_{j,n}.$$

We proceed as above. For $j \in \{0, \dots, \nu_n\}$ we have

$$R_{j,n}(z)\theta_{j,n}(K_n - z)u = \theta_{j,n}u + inR_{j,n}(z)\theta_{j,n}r_{j,n}u - R_{j,n}(z)\theta_{j,n}''u + 2R_{j,n}(z)\partial_y(\theta_{j,n}'u),$$

where

$$r_{j,n}(y) = q^2(y) - q^2(a_{j,n}) - 2(y - a_{j,n})q'(a_{j,n})q(a_{j,n}).$$

Let $n \in \mathbb{N}^*$ and $j \in \{0, \dots, \nu_n\}$ such that $\theta_{j,n} \neq 0$. Then we have $2nq'(a_{j,n})|q(a_{j,n})| \gtrsim n^{1-\rho}$ and hence, for $z \in \mathbb{C}_n^-$ (in particular $\text{Re}(z) \leq \gamma n^{1/2} \ll n^{(2/3)(1-\rho)}$), Proposition 3.5 gives

$$(3.17) \quad \|R_{j,n}(z)\|_{L(L^2(I))} \lesssim n^{-(2/3)(1-\rho)}.$$

Then, as above we have $|\tilde{r}_{j,n}(y)\theta_{j,n}(y)| \lesssim n^{-2\tilde{\rho}}$ so

$$n\|R_{j,n}(z)\tilde{r}_{j,n}\theta_{j,n}\|_{L(L^2(I))} \lesssim n^{1-2\tilde{\rho}-(2/3)(1-\rho)} \xrightarrow{n \rightarrow +\infty} 0.$$

Moreover,

$$\begin{aligned} \|R_{j,n}(z)\theta_{j,n}''u\|_{L^2(I)} &\lesssim n^{2\tilde{\rho}-(2/3)(1-\rho)}\|u\|_{L^2(I)}, \\ \|R_{j,n}(z)(\theta_{j,n}'u)'\|_{L^2(I)} &\lesssim n^{\tilde{\rho}-(1/3)(1-\rho)}\|u\|_{L^2(I)}. \end{aligned}$$

All these estimates being uniform with respect to $j \in \{0, \dots, \nu_n\}$, we finally get

$$(3.18) \quad \sup_{z \in \mathbb{C}_n^-} \sup_{0 \leq j \leq \nu_n} \|T_{j,n}(z)\|_{L(L^2(I))} \xrightarrow{n \rightarrow +\infty} 0.$$

– For $u \in L^2(I)$ we write

$$u = \chi_n u + \sum_{j=0}^{\nu_n} \theta_{j,n} u,$$

We want to sum (3.15) and the estimates (3.18), for $j \in \{0, \dots, \nu_n\}$, to get an approximate inverse for $(K_n - z)$. We have seen that each contribution goes to 0, but the number of terms grows with n .

Let $\tilde{\theta} \in C_0^\infty(\mathbb{R}, [0, 1])$ be equal to 1 on $[-2/3, 2/3]$ and supported in $] -1, 1[$. Then for $n \in \mathbb{N}^*$, $j \in \{1, \dots, \nu_n\}$ and $y \in \bar{I}$ we set

$$\tilde{\theta}_{j,n}(y) = \tilde{\theta}((y - a_{j,n})/\delta_n),$$

and then

$$(3.19) \quad Q_n(z) = R_n(z)\chi_n + \sum_{j=0}^{\nu_n} \tilde{\theta}_{j,n} R_{j,n}(z)\theta_{j,n}.$$

For $u \in \text{Dom}(K_n)$ and $z \in \mathbb{C}_n^-$ we have $\theta_{j,n}u = \theta_{j,n}\tilde{\theta}_{j,n}u$ and

$$\theta_{j,n}(K_n - z)(1 - \tilde{\theta}_{j,n})u = 0,$$

so

$$\|Q_n(z)(K_n - z)u - u\|_{L^2(I)} \leq \|T_n(z)u\|_{L^2(I)} + \left\| \sum_{j=0}^{\nu_n} \tilde{\theta}_{j,n} T_{j,n}(z) \tilde{\theta}_{j,n} u \right\|_{L^2(I)}.$$

Moreover $\tilde{\theta}_{j,n} \tilde{\theta}_{k,n} = 0$ whenever $|j - k| \geq 2$, so by almost orthogonality (twice) we can write

$$\begin{aligned} \left\| \sum_{j=0}^{\nu_n} \tilde{\theta}_{j,n} T_{j,n}(z) \tilde{\theta}_{j,n} u \right\|_{L^2(I)}^2 &\lesssim \sum_{j=0}^{\nu_n} \|\tilde{\theta}_{j,n} T_{j,n}(z) \tilde{\theta}_{j,n} u\|_{L^2(I)}^2 \\ &\lesssim \sup_{0 \leq j \leq \nu_n} \|T_{j,n}(z)\|_{L^2(I)}^2 \sum_{j=0}^{\nu_n} \|\tilde{\theta}_{j,n} u\|_{L^2(I)}^2 \\ &\lesssim \sup_{0 \leq j \leq \nu_n} \|T_{j,n}(z)\|_{L^2(I)}^2 \|u\|_{L^2(I)}^2. \end{aligned}$$

This proves

$$(3.20) \quad \sup_{\substack{z \in \mathbb{C}_n^- \\ \|u\|_{L^2(I)}=1}} \sup_{u \in \text{Dom}(K_n)} \|Q_n(z)(K_n - z)u - u\|_{L^2(I)} \xrightarrow{n \rightarrow +\infty} 0.$$

Thus for n large enough the operator K_n has no eigenvalue and hence no spectrum in \mathbb{C}_n^- . Moreover for $z \in \mathbb{C}_n^-$ we have

$$(3.21) \quad (K_n - z)^{-1} = B_n(z)Q_n(z),$$

where

$$B_n(z) = \left(1 + (Q_n(z)(K_n - z) - 1)\right)^{-1}$$

is bounded on $L^2(I)$ uniformly in $z \in \mathbb{C}_n^-$ and n large enough.

– Let $u \in L^2(I)$ and $z \in \mathbb{C}_n^-$. By (3.17), and using again the almost orthogonality, we obtain

$$\left\| B_n(z) \sum_{j=0}^{\nu_n} \tilde{\theta}_{j,n} R_{j,n}(z) \theta_{j,n} u \right\|_{L^2(I)}^2 \lesssim \frac{\|u\|_{L^2(I)}^2}{n^{(4/3)(1-\rho)}},$$

so

$$(3.22) \quad \|(K_n - z)^{-1} - B_n(z)R_n(z)\chi_n\|_{L^2(I)} \lesssim \frac{1}{n^{(2/3)(1-\rho)}}.$$

With (3.12), this gives the first statement of the proposition.

– We now consider the case $z = 0$ to prove the second part of the proposition. By (3.20) we have

$$\|B_n(0) - 1\|_{L^2(I)} = \|(1 + (Q_n(0)K_n - 1))^{-1} - 1\|_{L^2(I)} \xrightarrow{n \rightarrow +\infty} 0,$$

so (3.22) and (3.12) give

$$(3.23) \quad \|K_n^{-1} - \mathbf{1}_I H_n^{-1} \mathbf{1}_I^* \chi_n\|_{L^2(I)} = o_{n \rightarrow +\infty}(1/\sqrt{n}).$$

On $\text{supp}(1 - \chi_n)$ we have $|y| \gtrsim n^{-\rho}$ and for $v \in \text{Dom}(H_n^*)$ we have

$$\langle ny^2 v, v \rangle = |\text{Im} \langle H_n^* v, v \rangle|,$$

so for $u \in L^2(I)$ we can write

$$\begin{aligned} \|(1 - \chi_n)\mathbb{1}_I(H_n^*)^{-1}\mathbb{1}_I^*u\|_{L^2(I)}^2 &\lesssim n^{2\rho}\langle y^2(H_n^*)^{-1}\mathbb{1}_I^*u, (H_n^*)^{-1}\mathbb{1}_I^*u \rangle_{L^2(\mathbb{R})} \\ &\lesssim n^{2\rho-1}|\operatorname{Im}\langle \mathbb{1}_I^*u, (H_n^*)^{-1}\mathbb{1}_I^*u \rangle_{L^2(\mathbb{R})}| \\ &\lesssim \frac{\|u\|_{L^2(I)}^2}{n^{(3/2)-2\rho}}. \end{aligned}$$

Taking the adjoint gives

$$\|\mathbb{1}_I H_n^{-1}\mathbb{1}_I^*(1 - \chi_n)\|_{L(L^2(I))} = \underset{n \rightarrow +\infty}{o}(1/\sqrt{n}).$$

With (3.23), the proof is complete. \square

Now we are in position to prove Proposition 2.3. It is a direct consequence of the following result.

PROPOSITION 3.7. — *Let $\gamma < q'(0)/\sqrt{2}$. There exist $n_0 \in \mathbb{N}^*$ and $C > 0$ such that for $n \geq n_0$ and $t \geq 0$ we have*

$$\|e^{-tK_n}\|_{L(L^2(I))} \leq Ce^{-t\gamma\sqrt{n}}.$$

Proof. — Let $n_0 \in \mathbb{N}^*$ and $c > 0$ by given by Proposition 3.6. For $n \geq n_0$ we set $\tilde{K}_n = -K_n + \gamma\sqrt{n}$. Then for $n \geq n_0$ and $z \in \mathbb{C}$ with $\operatorname{Re}(z) \geq 0$ we have $z \in \rho(\tilde{K}_n)$ and

$$\|(\tilde{K}_n - z)^{-1}\|_{L(L^2(I))} \leq \frac{c}{\sqrt{n}}.$$

Moreover for $t \geq 0$ we have

$$\|e^{t\tilde{K}_n}\| \leq e^{t\gamma\sqrt{n}}.$$

Then we apply [EN00, Th. V.1.11 p. 302] to the operator \tilde{K}_n . With the notation used in the proof therein, we have $\omega_0 \leq \gamma\sqrt{n}$, $M \leq c/\sqrt{n}$ and $L = 2\pi$. We obtain that the semigroup $(e^{t\tilde{K}_n})_{t \geq 0}$ is bounded uniformly in $t \geq 0$ and $n \geq n_0$, so there exists $C > 0$ such that for all $n \geq n_0$ and $t \geq 0$ we have

$$\|e^{-tK_n}\| = e^{-t\gamma\sqrt{n}}\|e^{t\tilde{K}_n}\| \leq Ce^{-t\gamma\sqrt{n}}.$$

We also refer to [HS10] to get bounds on a semigroup from bounds on the resolvent of the corresponding generator. \square

Now we turn to the proof of (2.5). A more general version of the following result is given in [Kat80, §IV.3.5].

PROPOSITION 3.8. — *Let T be a closed operator on a Hilbert space H . Let $\lambda \in \mathbb{C}$. Assume that λ is an isolated eigenvalue of T . Let $(B_m)_{m \in \mathbb{N}}$ be a sequence of bounded operators on H such that $\|B_m\|_{L(H)} \rightarrow 0$ as $m \rightarrow +\infty$. For $m \in \mathbb{N}$ we set $T_m = T + B_m$. Let $\varepsilon > 0$. Then for m large enough the operator T_m has an isolated eigenvalue λ_m such that $|\lambda_m - \lambda| \leq \varepsilon$.*

Proof. — We set $C = \{\zeta \in \mathbb{C} : |\zeta - \lambda| = \varepsilon\}$. Without loss of generality, we can assume that $\varepsilon > 0$ is so small that λ is the only point of $\text{Sp}(T)$ in the disk $D(\lambda, 2\varepsilon)$. We set $M = \sup_{\zeta \in C} \|(T - \zeta)^{-1}\|$. Since

$$T_m - \zeta = (T - \zeta)(1 + (T - \zeta)^{-1}B_m),$$

we see that $C \cap \text{Sp}(T_m) = \emptyset$ as soon as $M\|B_m\| < 1/2$. Moreover, in this case, we have for $\zeta \in C$,

$$\|(T_m - \zeta)^{-1}\| \leq 2M.$$

We set

$$P = \frac{1}{2i\pi} \int_C (T - \zeta)^{-1} d\zeta.$$

We similarly define P_m by replacing T by T_m . Then we have by the resolvent identity

$$\|P_m - P\| = \left\| \frac{1}{2i\pi} \int_C (T - z)^{-1} B_m (T_m - \zeta)^{-1} d\zeta \right\| \leq 2\varepsilon M^2 \|B_m\|.$$

Thus for m large enough we have $\|P_m - P\| < 1$. By [Kat80, §I.4.6] this implies that

$$\dim(\text{Ran}(P_m)) = \dim(\text{Ran}(P)) \in \mathbb{N}^*.$$

This proves that T_m has an eigenvalue λ_m such that $|\lambda - \lambda_m| < \varepsilon$. □

PROPOSITION 3.9. — *For $n \in \mathbb{N}^*$ large enough there exists an eigenvalue λ_n of K_n such that*

$$|\lambda_n - e^{i\pi/4} q'(0) \sqrt{n}| = o_{n \rightarrow +\infty}(\sqrt{n}).$$

Proof. — We consider on $L^2(\mathbb{R})$ the unitary operator Θ_n which maps u to

$$\Theta_n u : x \mapsto n^{1/8} u(n^{1/4} x).$$

Then we have $\Theta_n^{-1} H_n \Theta_n = \sqrt{n} H_1$. By Proposition 3.6,

$$\left\| \sqrt{n} \Theta_n^{-1} \mathbb{1}_I^* K_n^{-1} \mathbb{1}_I \Theta_n - H_1^{-1} \right\|_{L(L^2(\mathbb{R}))} \xrightarrow{n \rightarrow +\infty} 0.$$

We set $\lambda = e^{i\pi/4} q'(0)$. Then $\mu = \lambda^{-1}$ is an eigenvalue of H_1^{-1} . By Proposition 3.8, there exists an eigenvalue μ_n of $\sqrt{n} \Theta_n^{-1} \mathbb{1}_I^* K_n^{-1} \mathbb{1}_I \Theta_n$ such that μ_n goes to μ as n goes to $+\infty$. Then $n^{-1/2} \mu_n$ is an eigenvalue of $\mathbb{1}_I^* K_n^{-1} \mathbb{1}_I$, and hence an eigenvalue of K_n^{-1} . We conclude the proof by setting $\lambda_n = \sqrt{n} \mu_n^{-1}$. □

3.4. AGMON ESTIMATES. — To conclude the proof of Proposition 2.8, it remains to prove the estimate (2.6) for an eigenfunction ψ_n of K_n corresponding to the eigenvalue λ_n .

This estimate is given by an Agmon estimate. The Agmon estimates measure how the eigenfunctions corresponding to the smallest eigenvalues of a Schrödinger operator concentrate near the minimum of the potential. Exponential decay of eigenfunctions and precise Agmon estimates are classical results for real-valued potentials (see for instance [Agm85, Hel88]). We refer to [KRRS17] for Agmon estimates for a general non-selfadjoint Laplacian.

Here, it is expected that for large n an eigenfunction corresponding to the first eigenvalue λ_n of K_n will concentrate near 0, where the potential q^2 reaches its minimum. In particular, such an eigenfunction will be small at the boundary, so it is indeed a good candidate to break an observability estimate like (2.4) when $T < T_{\min}$.

PROPOSITION 3.10. — *Let $E > 0$ and $\varepsilon \in]0, 1[$. For $n \in \mathbb{N}$ and $y \in \bar{I}$ we set*

$$(3.24) \quad W_{n,\varepsilon}(y) = \frac{1-\varepsilon}{\sqrt{2}} \left| \int_0^y \sqrt{(nq(s)^2 - \sqrt{n}(E + \varepsilon))_+} ds \right|,$$

where for $\sigma \in \mathbb{R}$ we write σ_+ for $\max(0, \sigma)$. There exists $C > 0$ such that for $n \in \mathbb{N}$, $u \in \text{Dom}(K_n)$ and $\lambda \in \mathbb{C}$ with

$$(3.25) \quad |\text{Re}(\lambda)| + |\text{Im}(\lambda)| \leq E\sqrt{n},$$

we have

$$\|e^{W_{n,\varepsilon}}u'\|_{L^2(I)}^2 + \sqrt{n} \|e^{W_{n,\varepsilon}}u\|_{L^2(I)}^2 \leq C\sqrt{n} \|u\|_{L^2(I)}^2 + \frac{C}{\sqrt{n}} \|e^{W_{n,\varepsilon}}(K_n - \lambda)u\|_{L^2(I)}^2.$$

This result is proved with more generality in [KRRS17]. For the reader convenience we recall a proof in our 1-dimensional setting.

Proof. — We denote by Q_n the quadratic form corresponding to K_n . It is defined for $f, g \in H_0^1(I)$ by

$$Q_n(f, g) = \int_I f' \bar{g}' + in \int_I q^2 f \bar{g}.$$

– Let $u \in \text{Dom}(K_n)$. For $\zeta \in W^{1,\infty}(\bar{I}, \mathbb{R})$, we have

$$\langle u', (\zeta^2 u)' \rangle_{L^2(I)} = \langle \zeta u', 2\zeta' u + \zeta u' \rangle_{L^2(I)} = \langle (\zeta u)' - \zeta' u, (\zeta u)' + \zeta' u \rangle_{L^2(I)},$$

so

$$\text{Re} \langle u', (\zeta^2 u)' \rangle_{L^2(I)} = \|(\zeta u)'\|_{L^2(I)}^2 - \|\zeta' u\|_{L^2(I)}^2.$$

– Let $W \in W^{1,\infty}(\bar{I}, \mathbb{R})$. Applied with $\zeta = e^W$, this equality gives

$$\text{Re}(Q_n(u, e^{2W}u)) = \text{Re} \langle u', (e^{2W}u)' \rangle_{L^2(I)} = \|(e^W u)'\|_{L^2(I)}^2 - \|W' e^W u\|_{L^2(I)}^2.$$

On the other hand, a direct computation shows that

$$\text{Im}(Q_n(u, e^{2W}u)) = \text{Im} \langle u', 2W' e^{2W}u \rangle_{L^2(I)} + n \|qe^W u\|_{L^2(I)}^2.$$

Let $\alpha \in]0, 1[$. Since

$$\begin{aligned} |\text{Im} \langle u', 2W' e^{2W}u \rangle_{L^2(I)}| &= 2 |\text{Im} \langle (e^W u)', W' e^W u \rangle_{L^2(I)}| \\ &\leq \alpha \|(e^W u)'\|_{L^2(I)}^2 + \alpha^{-1} \|W' e^W u\|_{L^2(I)}^2, \end{aligned}$$

we have

$$\text{Im}(Q_n(u, e^{2W}u)) \geq n \|qe^W u\|_{L^2(I)}^2 - \alpha \|(e^W u)'\|_{L^2(I)}^2 - \alpha^{-1} \|W' e^W u\|_{L^2(I)}^2,$$

and hence

$$\begin{aligned} \operatorname{Re}(Q_n(u, e^{2W}u)) + \operatorname{Im}(Q_n(u, e^{2W}u)) \\ \geq (1 - \alpha) \|(e^W u)'\|_{L^2(I)}^2 + \int_I (nq^2 - (1 + \alpha^{-1})W'^2) |e^W u|^2. \end{aligned}$$

Finally,

$$\begin{aligned} \|(e^W u)'\|_{L^2(I)}^2 &\geq \|e^W u'\|_{L^2(I)}^2 + \|W' e^W u\|_{L^2(I)}^2 - 2\|e^W u'\|_{L^2(I)} \|W' e^W u\|_{L^2(I)} \\ &\geq \frac{1}{2} \|e^W u'\|_{L^2(I)}^2 - \|W' e^W u\|_{L^2(I)}^2, \end{aligned}$$

so if we set $\beta = 2 + \alpha^{-1} - \alpha$ and $\varepsilon_1 = (1 - \alpha)/2$, we get

$$(3.26) \quad \begin{aligned} \operatorname{Re}(Q_n(u, e^{2W}u)) + \operatorname{Im}(Q_n(u, e^{2W}u)) \\ \geq \varepsilon_1 \|e^W u'\|_{L^2(I)}^2 + \int_I (nq^2 - \beta W'^2) |e^W u|^2. \end{aligned}$$

– On the other hand, for $\lambda \in \mathbb{C}$ we have

$$Q_n(u, e^{2W}u) = \lambda \|e^W u\|_{L^2(I)}^2 + \langle (K_n - \lambda)u, e^{2W}u \rangle_{L^2(I)}.$$

We take the real and imaginary parts of this equality. With (3.26) this gives

$$(3.27) \quad \begin{aligned} \varepsilon_1 \|e^W u'\|_{L^2(I)}^2 + \int_I (nq^2 - \beta W'^2 - \operatorname{Re}(\lambda) - \operatorname{Im}(\lambda)) |e^W u|^2 \\ \leq 2 \|e^W (K_n - \lambda)u\|_{L^2(I)} \|e^W u\|_{L^2(I)}. \end{aligned}$$

– Now assume that (3.25) holds. Let $\delta_n^\pm \in]0, \ell_\pm]$ be such that

$$[-\delta_n^-, \delta_n^+] = \{y \in \bar{I} : nq(y)^2 \leq \sqrt{n}(E + \varepsilon)\}.$$

Let $W_{n,\varepsilon}$ be given by (3.24). We choose $\alpha \in]0, 1[$ in such a way that

$$\beta = \frac{2}{(1 - \varepsilon)^2}.$$

On $[-\delta_n^-, \delta_n^+]$, $W_{n,\varepsilon}$ and hence $W'_{n,\varepsilon}$ vanish, so

$$\beta W'_{n,\varepsilon}(y)^2 + \operatorname{Re}(\lambda) + \operatorname{Im}(\lambda) - nq(y)^2 \leq E\sqrt{n},$$

while on $I \setminus [-\delta_n^-, \delta_n^+]$ we have

$$\beta W'_{n,\varepsilon}(y)^2 = nq(y)^2 - \sqrt{n}(E + \varepsilon),$$

so

$$nq(y)^2 - \beta W'_{n,\varepsilon}(y)^2 - \operatorname{Re}(\lambda) - \operatorname{Im}(\lambda) \geq \varepsilon\sqrt{n}.$$

Then, by (3.27),

$$\begin{aligned} \varepsilon_1 \|e^{W_{n,\varepsilon}} u'\|_{L^2(I)}^2 + \varepsilon \sqrt{n} \int_{I \setminus [-\delta_n^-, \delta_n^+]} |e^{W_{n,\varepsilon}} u|^2 \\ \leq 2 \|e^{W_{n,\varepsilon}} (K_n - \lambda) u\|_{L^2(I)} \|e^{W_{n,\varepsilon}} u\|_{L^2(I)} + E \sqrt{n} \int_{-\delta_n^-}^{\delta_n^+} |u|^2 \\ \leq \frac{\varepsilon \sqrt{n}}{2} \|e^{W_{n,\varepsilon}} u\|_{L^2(I)}^2 + \frac{2}{\varepsilon \sqrt{n}} \|e^{W_{n,\varepsilon}} (K_n - \lambda) u\|_{L^2(I)}^2 + E \sqrt{n} \int_{-\delta_n^-}^{\delta_n^+} |u|^2, \end{aligned}$$

and finally,

$$\begin{aligned} \varepsilon_1 \|e^{W_{n,\varepsilon}} u'\|_{L^2(I)}^2 + \varepsilon \sqrt{n} \|e^{W_{n,\varepsilon}} u\|_{L^2(I)}^2 \\ \leq \frac{\varepsilon \sqrt{n}}{2} \|e^{W_{n,\varepsilon}} u\|_{L^2(I)}^2 + \frac{2}{\varepsilon \sqrt{n}} \|e^{W_{n,\varepsilon}} (K_n - \lambda) u\|_{L^2(I)}^2 + (E + \varepsilon) \sqrt{n} \int_{-\delta_n^-}^{\delta_n^+} |u|^2. \end{aligned}$$

The proposition is proved. □

For $\varepsilon \in]0, 1]$ and $y \in I$ we set

$$\kappa_\varepsilon(y) = \frac{(1 - \varepsilon)}{\sqrt{2}} \int_0^y q(s) ds.$$

PROPOSITION 3.11. — *Let $E > 0$ and $\varepsilon \in]0, 1]$. There exists $C_\varepsilon > 0$ such that for $n \in \mathbb{N}$ and $y \in I$ we have*

$$(3.28) \quad \sqrt{n} \kappa_\varepsilon(y) - C_\varepsilon \leq W_{n,\varepsilon/2}(y) \leq \sqrt{n} \kappa_{\varepsilon/2}(y).$$

Proof. — The second inequality is clear. It is enough to prove the first for n large. Let $\alpha \geq 1$ to be fixed large enough later. For n large enough we consider $\eta_n^\pm \in]0, \ell_\pm]$ such that

$$q(\pm \eta_n^\pm)^2 = \frac{\alpha}{\sqrt{n}} (E + \varepsilon/2).$$

We have

$$\eta_n^\pm = O_{n \rightarrow +\infty} (n^{-1/4}),$$

and hence

$$\sqrt{n} \kappa_\varepsilon(\pm \eta_n^\pm) = O_{n \rightarrow +\infty} (1).$$

In particular, for n large enough the first inequality in (3.28) holds for $y \in [-\eta_n^-, \eta_n^+]$ if C_ε is chosen large enough, since then the left-hand side is negative. On the other hand, for $y \geq \eta_n^+$ we have

$$\int_{\eta_n^+}^y \sqrt{nq(s)^2 - (E + \varepsilon/2)\sqrt{n}} ds \geq \sqrt{1 - \alpha^{-1}} \sqrt{n} \int_{\eta_n^+}^y q(s) ds.$$

Then

$$W_{n,\varepsilon/2}(y) \geq \frac{1 - \varepsilon/2}{1 - \varepsilon} \sqrt{1 - \alpha^{-1}} \sqrt{n} \kappa_\varepsilon(y) + O_{n \rightarrow +\infty} (1).$$

For α large enough this gives (3.28). We proceed similarly for $y \leq -\eta_n^-$. □

Combining Propositions 3.10 and 3.11 we obtain the following version of the Agmon estimates:

PROPOSITION 3.12. — *Let $E > 0$ and $\varepsilon \in]0, 1]$. There exists $C > 0$ such that for $n \in \mathbb{N}$, $u \in \text{Dom}(K_n)$ and $\lambda \in \mathbb{C}$ with $|\text{Re}(\lambda)| + |\text{Im}(\lambda)| \leq E\sqrt{n}$ we have*

$$\|e^{\sqrt{n}\kappa_\varepsilon} u'\|_{L^2(I)}^2 + \sqrt{n} \|e^{\sqrt{n}\kappa_\varepsilon} u\|_{L^2(I)}^2 \leq C\sqrt{n} \|u\|_{L^2(I)}^2 + \frac{C}{\sqrt{n}} \|e^{\sqrt{n}\kappa_\varepsilon/2} (K_n - \lambda)u\|_{L^2(I)}^2.$$

Proof. — If we denote by $\tilde{C} > 0$ the constant given by Proposition 3.10, then by Proposition 3.11 we obtain the estimate of Proposition 3.12 with $C = e^{C_\varepsilon \tilde{C}}$. \square

From Proposition 3.12 we deduce the pointwise estimate (2.6).

PROPOSITION 3.13. — *Let $E > 0$ and $\varepsilon \in]0, 1[$. There exists $C > 0$ such that for $n \in \mathbb{N}$, an eigenvalue μ_n of K_n with $\text{Re}(\mu_n) + \text{Im}(\mu_n) \leq E\sqrt{n}$ and $\psi_n \in \ker(K_n - \mu_n)$, we have*

$$\|e^{\sqrt{n}\kappa_\varepsilon} \psi_n'\|_{L^\infty(I)}^2 \leq Cn \|\psi_n\|_{L^2(I)}^2.$$

Proof. — By Proposition 3.12 we have

$$(3.29) \quad \|e^{\sqrt{n}\kappa_\varepsilon} \psi_n\|_{L^2(I)}^2 \lesssim \|\psi_n\|_{L^2(I)}^2, \quad \|e^{\sqrt{n}\kappa_\varepsilon} \psi_n'\|_{L^2(I)}^2 \lesssim \sqrt{n} \|\psi_n\|_{L^2(I)}^2.$$

– We prove

$$(3.30) \quad \|e^{\sqrt{n}\kappa_\varepsilon} \psi_n''\|_{L^2(I)}^2 \lesssim n^{3/2} \|\psi_n\|_{L^2(I)}^2.$$

We have $\psi_n'' = inq^2\psi_n - \mu_n\psi_n$. With (3.29) we get

$$\|e^{\sqrt{n}\kappa_\varepsilon} \mu_n \psi_n\|_{L^2(I)}^2 \lesssim |\mu_n|^2 \|\psi_n\|_{L^2(I)}^2 \lesssim n \|\psi_n\|_{L^2(I)}^2.$$

For the other term we have by an integration by parts

$$\begin{aligned} \sqrt{2}(1 - \varepsilon) \|e^{\sqrt{n}\kappa_\varepsilon} nq^2\psi_n\|_{L^2(I)}^2 &= \int_I 2\sqrt{n}\kappa'_\varepsilon e^{2\sqrt{n}\kappa_\varepsilon} n^{3/2}q^3 |\psi_n|^2 dy \\ &= - \int_I e^{2\sqrt{n}\kappa_\varepsilon} n^{3/2} (3q^2q' |\psi_n|^2 + 2q^3 \text{Re}(\overline{\psi_n}\psi_n')) dy. \end{aligned}$$

On the one hand we have

$$\left| \int_I e^{2\sqrt{n}\kappa_\varepsilon} n^{3/2} 3q^2q' |\psi_n|^2 dy \right| \lesssim n^{3/2} \|e^{\sqrt{n}\kappa_\varepsilon} \psi_n\|_{L^2(I)}^2 \lesssim n^{3/2} \|\psi_n\|_{L^2(I)}^2.$$

On the other hand,

$$\begin{aligned} \left| \int_I e^{2\sqrt{n}\kappa_\varepsilon} n^{3/2} 2q^3 \text{Re}(\overline{\psi_n}\psi_n') dy \right| &\leq 2 \|e^{\sqrt{n}\kappa_\varepsilon} nq^2\psi_n\|_{L^2(I)} \|qe^{\sqrt{n}\kappa_\varepsilon} \sqrt{n}\psi_n'\|_{L^2(I)} \\ &\leq (1 - \varepsilon) \|e^{\sqrt{n}\kappa_\varepsilon} nq^2\psi_n\|_{L^2(I)}^2 + \frac{\|q\|_\infty^2}{1 - \varepsilon} \|e^{\sqrt{n}\kappa_\varepsilon} \sqrt{n}\psi_n'\|_{L^2(I)}^2. \end{aligned}$$

This gives (3.30).

– For $u \in H^1(I)$ we have

$$\|u\|_{L^\infty(I)}^2 \leq \frac{1}{\ell_+ + \ell_-} \|u\|_{L^2(I)}^2 + 2\|u\|_{L^2(I)}\|u'\|_{L^2(I)}$$

so, by (3.29) and (3.30),

$$\|e^{\sqrt{n}\kappa_\varepsilon}\psi'_n\|_{L^\infty(I)}^2 \lesssim n\|\psi_n\|_{L^2(I)}^2.$$

This completes the proof. □

Notice that (3.30) is better than the naive estimate obtained from (3.29) and the expression of ψ''_n . In fact we do not have to be optimal here, since the power of n in the right-hand side of (2.6) is not important for the proof of the second part of Theorem 1.5.

4. THE OBSERVABILITY ESTIMATE IN SMALL TIME

In this section we prove Propositions 2.6 (see Section 4.2) and 2.7 (see Section 4.4). The proofs rely on some Carleman estimates and the construction of a suitable weight function.

In this section we will not use an index n for a solution u of (2.1). No confusion will be possible since we will never consider a solution of the initial x -dependent problem (1.1). Moreover, we use an index for the partial derivatives, so u_t stands for $\partial_t u$, u_{yy} for $\partial_{yy} u$, etc.

4.1. A GENERIC CARLEMAN ESTIMATE. — We begin our analysis with a generic Carleman estimate. In the following statement, ϕ is a Carleman weight function. It will be applied to $w = e^{-\phi}u$, where u is a solution of a problem of the form (2.1), possibly with a source term (see (4.12) below). We also impose that w vanishes at initial and final times.

PROPOSITION 4.1. — *Let $n \in \mathbb{N}$, $\tau_1, \tau_2 > 0$ with $\tau_1 < \tau_2$, $a, b \in \mathbb{R}$ with $a < b$, and $g \in L^2(] \tau_1, \tau_2[\times] a, b[)$. Let $\phi \in C^4(] \tau_1, \tau_2[\times] a, b[, \mathbb{R}_+)$. We consider*

$$w \in C^0(] \tau_1, \tau_2], H^2(a, b)) \cap C^1(] \tau_1, \tau_2], L^2(a, b))$$

such that

$$(4.1) \quad w_t - w_{yy} + inq(y)^2 w + \phi_t w - 2\phi_y w_y - \phi_y^2 w - \phi_{yy} w = g.$$

We assume that w also satisfies the Dirichlet boundary condition

$$(4.2) \quad \forall t \in] \tau_1, \tau_2[, \quad w(t, a) = w(t, b) = 0,$$

and the initial and final conditions

$$(4.3) \quad \forall y \in] a, b[, \quad w(\tau_1, y) = w(\tau_2, y) = 0, \quad w_y(\tau_1, y) = w_y(\tau_2, y) = 0.$$

Then we have

$$\int_{\tau_1}^{\tau_2} \int_a^b (\Phi_0 |w|^2 + \Phi_1 |w_y|^2) dy dt \leq - \int_{\tau_1}^{\tau_2} [\phi_y |w_y|^2]_a^b dt + \frac{1}{2} \int_{\tau_1}^{\tau_2} \int_a^b |g|^2 dy dt,$$

where

$$(4.4) \quad \Phi_0 = -2\phi_y^2\phi_{yy} - \frac{\phi_{tt}}{2} + \frac{\phi_{yyyy}}{2} + 2\phi_{ty}\phi_y - \frac{n^{3/2}q^2q'}{\sqrt{2}}$$

and

$$(4.5) \quad \Phi_1 = -2\phi_{yy} - \sqrt{2n}q'.$$

Proof. — We can rewrite (4.1) as

$$(-w_{yy} + \Phi w) + (w_t - 2\phi_y w_y - \phi_{yy} w + inq^2 w) = g,$$

where $\Phi = \phi_t - \phi_y^2$. The identity $2 \operatorname{Re}(\alpha\bar{\beta}) \leq |\alpha + \beta|^2$ then gives, after integration,

$$(4.6) \quad \operatorname{Re} \int_{\tau_1}^{\tau_2} \int_a^b (-w_{yy} + \Phi w)(\bar{w}_t - 2\phi_y \bar{w}_y - \phi_{yy} \bar{w} - inq^2 \bar{w}) dy dt \\ \leq \frac{1}{2} \int_{\tau_1}^{\tau_2} \int_a^b |g|^2 dy dt.$$

We estimate the left-hand side with integrations by parts, using (4.2) and (4.3). The terms involving \bar{w}_t give

$$\operatorname{Re} \int_{\tau_1}^{\tau_2} \int_a^b (-w_{yy}) \bar{w}_t dy dt = 0$$

and

$$\operatorname{Re} \int_{\tau_1}^{\tau_2} \int_a^b (\Phi w) \bar{w}_t dy dt = -\frac{1}{2} \int_{\tau_1}^{\tau_2} \int_a^b \Phi_t |w|^2 dy dt.$$

On the other hand, for all $t \in]\tau_1, \tau_2[$ we have

$$\operatorname{Re} \int_a^b (-w_{yy})(-2\phi_y \bar{w}_y) dy = [\phi_y |w_y|^2]_a^b - \int_a^b \phi_{yy} |w_y|^2 dy, \\ \operatorname{Re} \int_a^b (-w_{yy})(-\phi_{yy} \bar{w}) dy = -\int_a^b \phi_{yy} |w_y|^2 dy + \frac{1}{2} \int_a^b \phi_{yyyy} |w|^2 dy, \\ \operatorname{Re} \int_a^b (-w_{yy})(-inq^2 \bar{w}) dy = 2n \int_a^b qq' \operatorname{Im}(w_y \bar{w}) dy,$$

and

$$\operatorname{Re} \int_a^b (\Phi w)(-2\phi_y \bar{w}_y - \phi_{yy} \bar{w}) dy = \int_a^b \Phi_y \phi_y |w|^2 dy, \\ \operatorname{Re} \int_a^b (\Phi w)(-inq^2 \bar{w}) dy = 0.$$

We integrate these five equalities over $t \in]\tau_1, \tau_2[$, and then (4.6) gives

$$\int_{\tau_1}^{\tau_2} [\phi_y |w_y|^2]_a^b dt + \int_{\tau_1}^{\tau_2} \int_a^b \left(-\frac{\Phi_t}{2} + \frac{\phi_{yyyy}}{2} + \Phi_y \phi_y \right) |w|^2 dy dt \\ - 2 \int_{\tau_1}^{\tau_2} \int_a^b \phi_{yy} |w_y|^2 dy dt + 2n \int_{\tau_1}^{\tau_2} \int_a^b qq' \operatorname{Im}(w_y \bar{w}) dy dt \leq \frac{1}{2} \int_{\tau_1}^{\tau_2} \int_a^b |g|^2 dy dt.$$

Since

$$2nqq' \operatorname{Im}(w_y \bar{w}) \geq -\sqrt{2}\sqrt{n}q'|w_y|^2 - \frac{n^{3/2}q^2q'|w|^2}{\sqrt{2}},$$

the conclusion follows. □

4.2. OBSERVABILITY INEQUALITY FOR A FIXED FOURIER PARAMETER. — In this paragraph we prove Proposition 2.6 about observability for a fixed Fourier parameter $n \in \mathbb{N}$. As already said, this is nothing but the observability inequality for a heat equation with a (complex) potential. Nevertheless, we recall a proof of this well-known result to enlighten the construction of the refined weight function in the next paragraphs.

The proof of Proposition 2.6 relies on Proposition 4.1. For the time dependence of the weight ϕ , we will use the function θ given in the following lemma.

LEMMA 4.2. — Let $\tau_1, \tau_2 > 0$ with $\tau_1 < \tau_2$. There exists θ in $C^\infty(] \tau_1, \tau_2[)$ such that

- (i) $\theta \geq 1$ on $] \tau_1, \tau_2[$, $\theta \equiv 1$ on $[(2\tau_1 + \tau_2)/3, (\tau_1 + 2\tau_2)/3]$,
- (ii) $\lim_{t \rightarrow \tau_1} \theta(t) = \lim_{t \rightarrow \tau_2} \theta(t) = +\infty$,
- (iii) there exists a constant $C > 0$ such that for all $t \in] \tau_1, \tau_2[$,

$$|\theta'(t)| \leq C\theta(t)^2, \quad |\theta''(t)| \leq C\theta(t)^3.$$

Proof. — Let $\chi \in C_0^\infty(] \tau_1, \tau_2[, [0, 1])$ be equal to 1 on $[(2\tau_1 + \tau_2)/3, (\tau_1 + 2\tau_2)/3]$. For $t \in] \tau_1, \tau_2[$ we set

$$\theta(t) = 1 + \frac{1 - \chi(t)}{(t - \tau_1)(\tau_2 - t)}.$$

Then θ satisfies all the required properties. □

Now we can prove Proposition 2.6:

Proof of Proposition 2.6. — For $y \in \bar{I}$ we set

$$\psi(y) = \psi_1\left(\frac{2y + \ell_- - \ell_+}{\ell_- + \ell_+}\right), \quad \text{where } \psi_1(\eta) = -\frac{\eta^2}{2} \pm 2\eta + 3, \quad \eta \in [-1, 1]$$

(the sign in front of 2η is not important here, but it has to be chosen carefully if we only observe from one side of the boundary, as will be the case in Proposition 4.4 below). In particular, for some $c_0 > 0$ we have on \bar{I}

$$(4.7) \quad \psi'' \leq -c_0, \quad |\psi'| \geq c_0, \quad \psi \geq c_0.$$

Let u be a solution of (2.1). Let $s > 1$ to be chosen large enough later. For $t \in] \tau_1, \tau_2[$ and $y \in \bar{I}$ we set

$$\phi(t, y) = s\theta(t)\psi(y),$$

where θ is given by Lemma 4.2, and

$$w(t, y) = u(t, y)e^{-\phi(t, y)}.$$

Then w satisfies (4.1)-(4.3) with $a = -\ell_-$, $b = \ell_+$ and $g \equiv 0$. Therefore, Proposition 4.1 gives

$$\int_{\tau_1}^{\tau_2} \int_I (\Phi_0|w|^2 + \Phi_1|w_y|^2) dy dt \leq - \int_{\tau_1}^{\tau_2} [\phi_y|w_y|^2]_{-\ell_-}^{\ell_+} dt,$$

with

$$(4.8) \quad \Phi_0 = s^3 \left(-2\theta^3(\psi')^2\psi'' - \frac{\theta''\psi}{2s^2} + \frac{\theta\psi^{(4)}}{2s^2} + \frac{2\theta'\theta(\psi')^2}{s} - \frac{n^{3/2}q^2q'}{s^3\sqrt{2}} \right)$$

and

$$(4.9) \quad \Phi_1 = s \left(-2\theta\psi'' - \frac{\sqrt{2n}q'}{s} \right).$$

Thus, by Lemma 4.2 and (4.7) we can fix s so large that $\Phi_0 \geq 1$ and $\Phi_1 \geq 1$ on $]\tau_1, \tau_2[\times \bar{I}$. This gives

$$\int_{\tau_1}^{\tau_2} \int_I |w(t, y)|^2 dy dt \lesssim \int_{\tau_1}^{\tau_2} (|w_y(t, -\ell_-)|^2 + |w_y(t, \ell_+)|^2) dt,$$

and then, since $\theta \equiv 1$ on $[(2\tau_1 + \tau_2)/3, (\tau_1 + 2\tau_2)/3]$ and ψ is bounded away from 0,

$$\int_{(2\tau_1 + \tau_2)/3}^{(\tau_1 + 2\tau_2)/3} \int_I |u(t, y)|^2 dy dt \lesssim \int_{\tau_1}^{\tau_2} (|u_y(t, -\ell_-)|^2 + |u_y(t, \ell_+)|^2) dt.$$

We have $\|u(T)\|_{L^2(I)}^2 \leq \|u(t)\|_{L^2(I)}^2$ for all $t \in [(2\tau_1 + \tau_2)/3, (\tau_1 + 2\tau_2)/3]$. After integration this gives

$$\|u(T)\|_{L^2(I)}^2 \leq \frac{3}{\tau_2 - \tau_1} \int_{(2\tau_1 + \tau_2)/3}^{(\tau_1 + 2\tau_2)/3} \|u(t)\|_{L^2(I)}^2 dt \lesssim \int_{\tau_1}^{\tau_2} (|u_y(t, -\ell_-)|^2 + |u_y(t, \ell_+)|^2) dt,$$

which ends the proof. □

Notice that in this rough proof we have not tried to control the dependence of C_n with respect to n . It is the purpose of the next paragraph to get a precise estimate of the cost of observability for (2.1). The interest of Proposition 2.6 is that it is now enough to consider only large values of n .

To obtain estimates in the high frequency regime, we will use the same strategy, but we will choose more carefully the parameter s and the phase function ψ (both should be chosen as small as possible).

From (4.8), we see that s^3 has to be at least of order $n^{3/2}$, while in (4.9), s has to be of order \sqrt{n} . From these observations, we deduce that the correct scaling should be $s \sim \sqrt{n}$.

Finally, with $s = \sqrt{n}$, it is then the choice of ψ that will make Φ_0 and Φ_1 positive for n large enough. We see from (4.8)-(4.9) that ψ should satisfy

$$(4.10) \quad -2(\psi')^2\psi'' - \frac{q^2q'}{\sqrt{2}} > 0 \quad \text{and} \quad -2\psi'' - \sqrt{2}q' > 0.$$

This leads to the construction of the weight function given in the next paragraph.

REMARK 4.3. — The strategy we develop in the following sections shares certain similarities with the strategy developed in [Bak12] to obtain a sharp upper bound on the vanishing order of solutions to Schrödinger equations with smooth potentials. The correct scaling $s \sim \sqrt{n}$ is natural in view of the main result of this work.

4.3. A REFINED CARLEMAN ESTIMATE. — In this paragraph we prove a refined version of Proposition 4.1 for n large and a suitable choice for ψ . As discussed at the end of Section 4.2, we will choose ϕ proportional to \sqrt{n} . The choice of ψ satisfying (4.10) will be discussed in Proposition 4.5.

PROPOSITION 4.4. — Let $a, b \in \bar{I}$ with $a < b$ and $\psi \in C^4([a, b], \mathbb{R})$. We assume that for some $\varepsilon > 0$ we have on $[a, b]$:

$$\psi \geq \varepsilon, \quad -2(\psi')^2\psi'' - \frac{q^2q'}{\sqrt{2}} \geq \varepsilon, \quad -2\psi'' - \sqrt{2}q' \geq \varepsilon.$$

Let $\tau_1, \tau_2 \in]0, T]$ with $\tau_1 < \tau_2$. For $t \in]\tau_1, \tau_2[$ and $y \in [a, b]$ we set $\varphi(t, y) = \theta(t)\psi(y)$, where θ is given by Lemma 4.2. Let $n \in \mathbb{N}$ and u in

$$(4.11) \quad C^0([\tau_1, \tau_2], H^2(a, b) \cap H_0^1(a, b)) \cap C^1([\tau_1, \tau_2], L^2(a, b)).$$

We set

$$(4.12) \quad f = u_t - u_{yy} + inq(y)^2u,$$

and

$$w = ue^{-\sqrt{n}\varphi}, \quad g = fe^{-\sqrt{n}\varphi}.$$

Then there exist $N \in \mathbb{N}$ and $C > 0$ such that the following statements hold if $n \geq N$.

(i) If $\psi' > 0$,

$$\int_{\tau_1}^{\tau_2} \int_a^b (n^{3/2}\theta^3|w|^2 + \sqrt{n}\theta|w_y|^2) dy dt \leq C\sqrt{n} \int_{\tau_1}^{\tau_2} |w_y(t, a)|^2 dt + C \int_{\tau_1}^{\tau_2} \int_a^b |g|^2 dy dt.$$

(ii) If $\psi' < 0$,

$$\int_{\tau_1}^{\tau_2} \int_a^b (n^{3/2}\theta^3|w|^2 + \sqrt{n}\theta|w_y|^2) dy dt \leq C\sqrt{n} \int_{\tau_1}^{\tau_2} |w_y(t, b)|^2 dt + C \int_{\tau_1}^{\tau_2} \int_a^b |g|^2 dy dt.$$

Proof. — We observe that φ belongs to $C^4(] \tau_1, \tau_2[\times [a, b])$, the functions f and g are in $C^0([\tau_1, \tau_2], L^2(a, b))$, w extends to a function in (4.11) and we have

$$w_t - w_{yy} + inq(y)^2w + \sqrt{n}\varphi_t w - 2\sqrt{n}\varphi_y w_y - n\varphi_y^2 w - \sqrt{n}\varphi_{yy} w = g.$$

Moreover, w satisfies the boundary conditions (4.2) and the initial and final conditions (4.3). Then, by Proposition 4.1 applied with $\phi = \sqrt{n}\varphi$, we have

$$\int_{\tau_1}^{\tau_2} \int_a^b (n^{3/2}\Phi_0|w|^2 + \sqrt{n}\Phi_1|w_y|^2) dy dt \leq -\sqrt{n} \int_{\tau_1}^{\tau_2} [\varphi_y|w_y|^2]_a^b dt + \frac{1}{2} \int_{\tau_1}^{\tau_2} \int_a^b |g|^2 dy dt,$$

where

$$\Phi_0 = -2\varphi_y^2\varphi_{yy} - \frac{q^2q'}{\sqrt{2}} - \frac{\varphi_{tt}}{2n} + \frac{\varphi_{yyyy}}{2n} + \frac{2\varphi_{ty}\varphi_y}{\sqrt{n}}$$

and

$$\Phi_1 = -2\varphi_{yy} - \sqrt{2}q'.$$

The properties of θ and the boundedness of the derivatives of ψ give, for n large enough,

$$\Phi_0(t, y) \geq \frac{\varepsilon\theta^3}{2} \quad \text{and} \quad \Phi_1(t, y) \geq \varepsilon\theta.$$

Thus,

$$\frac{\varepsilon}{2} \int_{\tau_1}^{\tau_2} \int_a^b (n^{3/2} \theta^3 |w|^2 + \sqrt{n} \theta |w_y|^2) dy dt \leq -\sqrt{n} \int_{\tau_1}^{\tau_2} [\varphi_y |w_y|^2]_a^b dt + \frac{1}{2} \int_{\tau_1}^{\tau_2} \int_a^b |g|^2 dy dt.$$

Notice that the assumptions on ψ imply that ψ' does not vanish. If ψ' takes positive values then we have

$$-\sqrt{n} \int_{\tau_1}^{\tau_2} \varphi_y(t, b) |w_y(t, b)|^2 dt \leq 0,$$

which gives the first inequality. Otherwise $\psi' < 0$ and we similarly get the second estimate. □

4.4. PRECISE ESTIMATE OF THE COST OF OBSERVATION IN SMALL TIME FOR n LARGE

In this paragraph we finish the proof of Proposition 2.7. We could apply directly Proposition 4.4 and observe from one side of I only. However, we can reduce the cost of observability if we observe from both sides.

More precisely, the part of u in $[0, \ell_+]$ will be controled by the values of u_y at ℓ_+ , and the part of u in $[-\ell_-, 0]$ will be controled by the values of u_y at $-\ell_-$. Thus, with the notation of the previous paragraph, we have to choose ψ such that $\psi' < 0$ on the right and $\psi' > 0$ on the left. Since ψ' does not vanish, we have to apply Proposition 4.4 separately on the right and on the left.

PROPOSITION 4.5. — *Let τ_1, τ_2 and κ be as given by Proposition 2.7. There exist $N \in \mathbb{N}^*, \varphi \in C^0([\tau_1, \tau_2] \times \bar{I}, \mathbb{R})$ and $C > 0$ such that*

$$(4.13) \quad \forall t \in [(2\tau_1 + \tau_2)/3, (\tau_1 + 2\tau_2)/3], \forall y \in I, \quad 0 \leq \varphi(t, y) \leq \kappa,$$

and for any $n \geq N$ and any solution u of (2.1) we have

$$\int_{\tau_1}^{\tau_2} \int_I (n^{3/2} |u|^2 + \sqrt{n} |u_y|^2) e^{-2\sqrt{n}\varphi} dy dt \leq C\sqrt{n} \int_{\tau_1}^{\tau_2} (|u_y(t, -\ell_-)|^2 + |u_y(t, \ell_+)|^2) dt.$$

Proof

– Let $\beta > 1/\sqrt{2}$ and $\varepsilon_0 > 0$ be such that

$$(4.14) \quad \varepsilon_0 + \beta \max\left(\int_0^{\ell_+} (q(s) + 3\varepsilon_0) ds, \int_{-\ell_-}^0 (|q(s)| + 3\varepsilon_0) ds\right) < \kappa.$$

Let $\delta \in]0, \min(\ell_-, \ell_+)]$ be such that $\max(|q(-\delta)|, q(\delta)) \leq \varepsilon_0$. For $y \in [-\delta, \ell_+]$ we set

$$\psi_+(y) = \varepsilon_0 + \beta \int_y^{\ell_+} (q(s) + 3\varepsilon_0) ds + c_+,$$

with $c_+ \geq 0$ to be chosen later. Then we have

$$\psi_+ \geq \varepsilon_0, \quad \psi'_+ = -\beta(q + 3\varepsilon_0) \leq -2\beta\varepsilon_0, \quad \psi''_+ = -\beta q',$$

so

$$-2\psi_+{}^2 \psi''_+ - \frac{q^2 q'}{\sqrt{2}} = 2\beta^3 (q + 3\varepsilon_0)^2 q' - \frac{q^2 q'}{\sqrt{2}} \geq \frac{q'}{\sqrt{2}} ((q + 3\varepsilon_0)^2 - q^2) \geq \varepsilon_0^2 \min(q')$$

and

$$-2\psi_+'' - \sqrt{2}q' \geq 2\left(\beta - \frac{1}{\sqrt{2}}\right)q' \geq 2\left(\beta - \frac{1}{\sqrt{2}}\right)\min(q').$$

Thus ψ_+ satisfies the assumptions of Proposition 4.4 on $[-\delta, \ell_+]$. Then, for $t \in]\tau_1, \tau_2[$ we set

$$(4.15) \quad \varphi_+(t, y) = \theta(t)\psi_+(y),$$

where θ is given by Lemma 4.2.

– We consider $\chi_+ \in C^\infty(\bar{I}, [0, 1])$ such that $\chi_+ = 1$ on $[0, \ell_+]$ and $\chi_+ = 0$ on $[-\ell_-, -\delta]$. Then we set $u_+ = \chi_+u$. It satisfies

$$\forall t \in]\tau_1, \tau_2[, \quad u_+(t, -\delta) = u_+(t, \ell_+) = 0$$

and

$$\forall t \in]\tau_1, \tau_2[, \forall y \in [-\delta, \ell_+], \quad (\partial_t - \partial_{yy} + inq^2)u_+(t, y) = f_+(t, y),$$

where

$$f_+ = -\chi_+''u - 2\chi_+'u_y.$$

In particular, $f_+(t, \cdot)$ is supported in $[-\delta, 0]$. We set

$$w_+ = u_+e^{-\sqrt{n}\varphi_+} \quad \text{and} \quad g_+ = f_+e^{-\sqrt{n}\varphi_+}.$$

We have

$$\sqrt{n}|\partial_y u_+|^2 e^{-2\sqrt{n}\varphi_+} \lesssim \sqrt{n}|\partial_y w_+|^2 + n^{3/2}|w_+|^2 \theta(t)^2.$$

Then, by the second case in Proposition 4.4, we obtain

$$(4.16) \quad \begin{aligned} & \int_{\tau_1}^{\tau_2} \int_0^{\ell_+} (n^{3/2}|u_+|^2 + \sqrt{n}|\partial_y u_+|^2) e^{-2\sqrt{n}\varphi_+} dy dt \\ & \lesssim \int_{\tau_1}^{\tau_2} \int_0^{\ell_+} (n^{3/2}\theta^2|w_+|^2 + \sqrt{n}|\partial_y w_+|^2) dy dt \\ & \lesssim \sqrt{n} \int_{\tau_1}^{\tau_2} |\partial_y w_+(t, \ell_+)|^2 dt + \int_{\tau_1}^{\tau_2} \int_{-\delta}^0 |g_+|^2 dy dt \\ & \lesssim \sqrt{n} \int_{\tau_1}^{\tau_2} |\partial_y u_+(t, \ell_+)|^2 dt + \int_{\tau_1}^{\tau_2} \int_{-\delta}^0 |f_+|^2 e^{-2\sqrt{n}\varphi_+} dy dt. \end{aligned}$$

– For $y \in [-\ell_-, \delta]$ we set

$$\psi_-(y) = \varepsilon_0 + \beta \int_{-\ell_-}^y (|q(s)| + 3\varepsilon_0) ds + c_-,$$

with $c_- \geq 0$ to be chosen later, and for $t \in]0, T[$,

$$\varphi_-(t, y) = \theta(t)\psi_-(y).$$

Let $\chi_- \in C^\infty([-\ell_-, \ell_+], [0, 1])$ such that $\chi_- = 1$ on $[-\ell_-, 0]$ and $\chi_- = 0$ on $y \in [\delta, \ell_+]$. We set $u_- = \chi_-u$ and $f_- = -\chi_-''u - 2\chi_- 'u_y$. Then, as above, but using the first

statement in Proposition 4.4, we obtain

$$(4.17) \quad \int_{\tau_1}^{\tau_2} \int_{-\ell_-}^0 (n^{3/2}|u_-|^2 + \sqrt{n}|\partial_y u_-|^2) e^{-2\sqrt{n}\varphi_-} dy dt$$

$$\lesssim \sqrt{n} \int_{\tau_1}^{\tau_2} |\partial_y u_-(t, -\ell_-)|^2 dt + \int_{\tau_1}^{\tau_2} \int_0^\delta |f_-|^2 e^{-2\sqrt{n}\varphi_-} dy dt.$$

– We set $c_+ = \max(0, c)$ and $c_- = \max(0, -c)$ where

$$c = \beta \left(\int_{-\ell_-}^0 (|q(s)| + 3\varepsilon_0) ds - \int_0^{\ell_+} (q(s) + 3\varepsilon_0) ds \right),$$

so that $\psi_+(0) = \psi_-(0)$. Then for $t \in]\tau_1, \tau_2[$ and $y \in \bar{I}$ we set

$$\varphi(t, y) = \begin{cases} \varphi_-(t, y) & \text{if } y \leq 0, \\ \varphi_+(t, y) & \text{if } y \geq 0. \end{cases}$$

In particular, by construction, φ is continuous on $] \tau_1, \tau_2[\times \bar{I}$ and satisfies (4.13). Moreover, $\varphi_+ \geq \varphi$ on $[-\delta, 0]$, $\varphi_- \geq \varphi$ on $[0, \delta]$ and, on $[-\delta, \delta]$,

$$|f_+| + |f_-| \lesssim |u| + |u_y|.$$

Then, by summing (4.16) and (4.17),

$$\int_{\tau_1}^{\tau_2} \int_I (n^{3/2}|u|^2 + \sqrt{n}|u_y|^2) e^{-2\sqrt{n}\varphi} dy dt$$

$$\lesssim \sqrt{n} \int_{\tau_1}^{\tau_2} (|u_y(t, -\ell_-)|^2 + |u_y(t, \ell_+)|^2) dt + \int_{\tau_1}^{\tau_2} \int_{-\delta}^\delta (|u|^2 + |u_y|^2) e^{-2\sqrt{n}\varphi} dy dt.$$

For n large enough, the last term is smaller than one half of the left-hand side, and the conclusion follows. □

We can now prove Proposition 2.7.

Proof of Proposition 2.7. — Let N be given by Proposition 4.5 and $n \geq N$. Let u be a solution of (2.1). Let φ be given by Proposition 4.5. By (4.13) we have in particular

$$\int_{(2\tau_1 + \tau_2)/3}^{(\tau_1 + 2\tau_2)/3} \int_I |u|^2 dy dt \leq \frac{C e^{2\kappa\sqrt{n}}}{n} \int_{\tau_1}^{\tau_2} (|u_y(t, -\ell_-)|^2 + |u_y(t, \ell_+)|^2) dt.$$

By (3.7) we have $\|u(\tau_2)\|_{L^2(I)}^2 \leq \|u(t)\|_{L^2(I)}^2$ for all $t \in](2\tau_1 + \tau_2)/3, (\tau_1 + 2\tau_2)/3[$, so

$$\|u(\tau_2)\|_{L^2(I)}^2 \leq \frac{3 C e^{2\kappa\sqrt{n}}}{(\tau_2 - \tau_1)n} \int_{\tau_1}^{\tau_2} (|u_y(t, -\ell_-)|^2 + |u_y(t, \ell_+)|^2) dt,$$

and Proposition 2.7 is proved. □

REFERENCES

- [Agm85] S. AGMON – “Bounds on exponential decay of eigenfunctions of Schrödinger operators”, in *Schrödinger operators (Como, 1984)*, Lect. Notes in Math., vol. 1159, Springer, Berlin, 1985, p. 1–38.
- [ABM20] D. ALLONSIUS, F. BOYER & M. MORANCEY – “Analysis of the null-controllability of degenerate parabolic systems of Grushin type via the moments method”, 2020, to appear in *J. Evol. Equ.*, HAL:hal-02471592.
- [AKBGBdT16] F. AMMAR KHODJA, A. BENABDALLAH, M. GONZÁLEZ-BURGOS & L. DE TERESA – “New phenomena for the null controllability of parabolic systems: minimal time and geometrical dependence”, *J. Math. Anal. Appl.* **444** (2016), no. 2, p. 1071–1113.
- [Bak12] L. BAKRI – “Quantitative uniqueness for Schrödinger operator”, *Indiana Univ. Math. J.* **61** (2012), no. 4, p. 1565–1580.
- [BLR92] C. BARDOS, G. LEBEAU & J. RAUCH – “Sharp sufficient conditions for the observation, control, and stabilization of waves from the boundary”, *SIAM J. Control Optim.* **30** (1992), no. 5, p. 1024–1065.
- [Bea14] K. BEAUCHARD – “Null controllability of Kolmogorov-type equations”, *Math. Control Signals Systems* **26** (2014), no. 1, p. 145–176.
- [BC17] K. BEAUCHARD & P. CANNARSA – “Heat equation on the Heisenberg group: observability and applications”, *J. Differential Equations* **262** (2017), no. 8, p. 4475–4521.
- [BCG14] K. BEAUCHARD, P. CANNARSA & R. GUGLIELMI – “Null controllability of Grushin-type operators in dimension two”, *J. Eur. Math. Soc. (JEMS)* **16** (2014), no. 1, p. 67–101.
- [BDE20] K. BEAUCHARD, J. DARDÉ & S. ERVEDOZA – “Minimal time issues for the observability of Grushin-type equations”, *Ann. Inst. Fourier (Grenoble)* **70** (2020), no. 1, p. 247–312.
- [BHHR15] K. BEAUCHARD, B. HELFFER, R. HENRY & L. ROBBIANO – “Degenerate parabolic operators of Kolmogorov type with a geometric control condition”, *ESAIM Control Optim. Calc. Var.* **21** (2015), no. 2, p. 487–512.
- [BMM15] K. BEAUCHARD, L. MILLER & M. MORANCEY – “2D Grushin-type equations: minimal time and null controllable data”, *J. Differential Equations* **259** (2015), no. 11, p. 5813–5845.
- [BZ09] K. BEAUCHARD & E. ZUAZUA – “Some controllability results for the 2D Kolmogorov equation”, *Ann. Inst. H. Poincaré Anal. Non Linéaire* **26** (2009), no. 5, p. 1793–1815.
- [BBM20] A. BENABDALLAH, F. BOYER & M. MORANCEY – “A block moment method to handle spectral condensation phenomenon in parabolic control problems”, *Ann. H. Lebesgue* **3** (2020), p. 717–793.
- [BS19] N. BURQ & C. SUN – “Time optimal observability for Grushin Schrödinger equation”, 2019, HAL:hal-02309020.
- [CMV16] P. CANNARSA, P. MARTINEZ & J. VANCOSTENOBLE – *Global Carleman estimates for degenerate parabolic operators with applications*, Mem. Amer. Math. Soc., vol. 239, no. 1133, American Mathematical Society, Providence, RI, 2016.
- [Dup17] M. DUPREZ – “Controllability of a 2×2 parabolic system by one force with space-dependent coupling term of order one”, *ESAIM Control Optim. Calc. Var.* **23** (2017), no. 4, p. 1473–1498.
- [DK20] M. DUPREZ & A. KOENIG – “Control of the Grushin equation: non-rectangular control region and minimal time”, *ESAIM Control Optim. Calc. Var.* **26** (2020), article no. 3 (18 pages).
- [Ego63] J. V. EGOROV – “Some problems in the theory of optimal control”, *Ž. Vyčisl. Mat. i Mat. Fiz.* **3** (1963), p. 887–904.
- [EN00] K.-J. ENGEL & R. NAGEL – *One-parameter semigroups for linear evolution equations*, Graduate Texts in Math., vol. 194, Springer-Verlag, New York, 2000.
- [FR71] H. O. FATTORINI & D. L. RUSSELL – “Exact controllability theorems for linear parabolic equations in one space dimension”, *Arch. Rational Mech. Anal.* **43** (1971), p. 272–292.
- [FI96] A. V. FURSIKOV & O. Y. IMANUVILOV – *Controllability of evolution equations*, Lect. Notes Series, vol. 34, Seoul National University, Seoul, 1996.

- [Hel88] B. HELFFER – *Semi-classical analysis for the Schrödinger operator and applications*, Lect. Notes in Math., vol. 1336, Springer-Verlag, Berlin, 1988.
- [Hel11] ———, “On pseudo-spectral problems related to a time-dependent model in superconductivity with electric current”, *Confluentes Math.* **3** (2011), no. 2, p. 237–251.
- [Hel13] ———, *Spectral theory and its applications*, Cambridge Studies in Advanced Math., vol. 139, Cambridge University Press, Cambridge, 2013.
- [HS10] B. HELFFER & J. SJÖSTRAND – “From resolvent bounds to semi-group bounds”, 2010, Actes du colloque d’Évian 2009, [arXiv:1001.4171](https://arxiv.org/abs/1001.4171).
- [Hen14] R. HENRY – “On the semi-classical analysis of Schrödinger operators with purely imaginary electric potentials in a bounded domain”, 2014, [arXiv:1405.6183](https://arxiv.org/abs/1405.6183).
- [HSV13] M. HITRIK, J. SJÖSTRAND & J. VIOLA – “Resolvent estimates for elliptic quadratic differential operators”, *Anal. PDE* **6** (2013), no. 1, p. 181–196.
- [Kat80] T. KATO – *Perturbation theory for linear operators*, Classics in Mathematics, Springer-Verlag, Berlin, 1980.
- [Koe17] A. KOENIG – “Non-null-controllability of the Grushin operator in 2D”, *Comptes Rendus Mathématique* **355** (2017), no. 12, p. 1215–1235.
- [Koe20] ———, “Lack of null-controllability for the fractional heat equation and related equations”, *SIAM J. Control Optim.* **58** (2020), no. 6, p. 3130–3160.
- [KRRRS17] D. KREJČÍŘÍK, N. RAYMOND, J. ROYER & P. SIEGL – “Non-accretive Schrödinger operators and exponential decay of their eigenfunctions”, *Israel J. Math.* **221** (2017), no. 2, p. 779–802.
- [KSTV15] D. KREJČÍŘÍK, P. SIEGL, M. TATER & J. VIOLA – “Pseudospectra in non-Hermitian quantum mechanics”, *J. Math. Phys.* **56** (2015), no. 10, article no. 103513 (32 pages).
- [KS15] D. KREJČÍŘÍK & P. SIEGL – “Elements of spectral theory without the spectral theorem”, in *Non-selfadjoint operators in quantum physics*, Wiley, Hoboken, NJ, 2015, p. 241–291.
- [LL17] C. LAURENT & M. LÉAUTAUD – “Tunneling estimates and approximate controllability for hypoelliptic equations”, 2017, to appear in Mem. Amer. Math. Soc., [HAL:hal-01717587](https://hal.archives-ouvertes.fr/hal-01717587).
- [LR95] G. LEBEAU & L. ROBBIANO – “Contrôle exact de l’équation de la chaleur”, *Comm. Partial Differential Equations* **20** (1995), no. 1-2, p. 335–356.
- [LS20] C. LETROUT & C. SUN – “Observability of Baouendi-Grushin-type equations through resolvent estimates”, 2020, [HAL:hal-02964024](https://hal.archives-ouvertes.fr/hal-02964024).
- [Lis19] P. LISSY – “A non-controllability result for the half-heat equation on the whole line based on the prolate spheroidal wave functions and its application to the Grushin equation”, 2019, [HAL:hal-02420212v1](https://hal.archives-ouvertes.fr/hal-02420212v1).
- [RT74] J. RAUCH & M. TAYLOR – “Exponential decay of solutions to hyperbolic equations in bounded domains”, *Indiana Univ. Math. J.* **24** (1974), p. 79–86.
- [TW09] M. TUCSNAK & G. WEISS – *Observation and control for operator semigroups*, Birkhäuser Advanced Texts, Birkhäuser Verlag, Basel, 2009.
- [VS10] O. VALLÉE & M. SOARES – *Airy functions and applications to physics*, 2nd ed., Imperial College Press, London, 2010.

Manuscript received 17th June 2020

accepted 2nd March 2021

JÉRÉMI DARDÉ, Institut de Mathématiques de Toulouse, UMR5219, Univ. de Toulouse, CNRS, UPS F-31062 Toulouse Cedex 9, France

E-mail : jeremi.darde@math.univ-toulouse.fr

Url : <http://www.jeremidarde.toile-libre.org/>

JULIEN ROYER, Institut de Mathématiques de Toulouse, UMR5219, Univ. de Toulouse, CNRS, UPS F-31062 Toulouse Cedex 9, France

E-mail : julien.royer@math.univ-toulouse.fr

Url : <https://www.math.univ-toulouse.fr/~jroyer/>