



Journal de l'École polytechnique

Mathématiques

Camille LAURENT-GENGOUX & Leonid RYVKIN

The neighborhood of a singular leaf

Tome 8 (2021), p. 1037-1064.

http://jep.centre-mersenne.org/item/JEP_2021__8__1037_0

© Les auteurs, 2021.

Certains droits réservés.



Cet article est mis à disposition selon les termes de la licence
LICENCE INTERNATIONALE D'ATTRIBUTION CREATIVE COMMONS BY 4.0.
<https://creativecommons.org/licenses/by/4.0/>

L'accès aux articles de la revue « Journal de l'École polytechnique — Mathématiques » (<http://jep.centre-mersenne.org/>), implique l'accord avec les conditions générales d'utilisation (<http://jep.centre-mersenne.org/legal/>).

Publié avec le soutien
du Centre National de la Recherche Scientifique



Publication membre du
Centre Mersenne pour l'édition scientifique ouverte
www.centre-mersenne.org

THE NEIGHBORHOOD OF A SINGULAR LEAF

BY CAMILLE LAURENT-GENGOUX & LEONID RYVKIN

In memory of Kirill Mackenzie.

ABSTRACT . — An important result for regular foliations is their formal semi-local triviality near simply connected leaves. We extend this result to singular foliations for all 2-connected leaves and a wide class of 1-connected leaves by proving a semi-local Levi-Malcev theorem for the semi-simple part of their holonomy Lie algebroid.

RÉSUMÉ (Le voisinage d'une feuille singulière) . — Un résultat important concernant les feuilletages réguliers est leur trivialité semi-locale formelle au voisinage des feuilles simplement connexes. Nous étendons ce résultat aux feuilletages singuliers pour toute feuille 2-connexe et pour une classe importante de feuilles 1-connexes en démontrant un théorème de Levi-Malcev semi-local pour la partie semi-simple de leur algèbroïde de Lie d'holonomie.

CONTENTS

Introduction.....	1037
1. Holonomy and connections.....	1039
2. The formal neighborhood of a simply connected (singular) leaf.....	1048
3. Local and semi-local structure of a singular foliation.....	1059
References.....	1063

INTRODUCTION

Although much less studied than the well-understood regular foliations, singular foliations appear more frequently in differential geometry: orbits of Lie groups actions, symplectic leaves of a Poisson structure, vector fields tangent to an affine variety or annihilating given functions are all instances where the dimension of the leaves may not be constant. All these instances fall into the following category:

MATHEMATICAL SUBJECT CLASSIFICATION (2020). — 53C12, 57R30, 93B18.

KEYWORDS. — Singular foliations, singular leaves, linearizations in control theory, Levi-Malcev theorems, Lie algebroids.

Both authors are supported by the interdisciplinary CNRS projet *Granum* 80Prime. L. R. is supported by the PRIME Program of the German Academic Exchange Service with funds from the German Federal Ministry of Education and Research.

DEFINITION. — A singular foliation on a manifold M is a sub-sheaf F of the sheaf of C^∞ -modules $\mathfrak{X}(M)$ of vector fields, which is (i) stable under Lie bracket and (ii) locally finitely generated.⁽¹⁾

This definition permits to partition M into submanifolds called *leaves* ([Her63]). There is an open subset of M where F is a regular foliation. In particular, at least formally, in a neighborhood of any such a leaf L , the foliation F is entirely described by a group morphism from the fundamental group $\pi_1(L)$ to the group of formal diffeomorphisms of a transversal. In particular, regular foliations are (formally) trivial near simply-connected leaves [Ree52].

For singular leaves, there have been recent advances in understanding the semi-local structure. Androulidakis and Zambon [AZ13, AZ14] have shown that the holonomy groupoid of F (defined previously by Androulidakis and Skandalis [AS09]) acts on the normal bundle of the leaf. When the singular foliation is linearizable, this describes the whole semi-local structure. In this article, we mainly focus on the case where L is simply-connected. To our great surprise, we were able to prove that, despite having possibly extremely rich transverse structures, singular foliations remain (formally) trivial near simply-connected locally closed leaves, when the transverse singular foliation is made of vector fields vanishing at order at least 2 (Theorems 3.8 and 3.11). When the transverse linear part is not trivial but L is 2-connected, we still have a Levi-Malcev type theorem decomposing F as a semi-direct product of a semi-simple linearizable Lie groupoid action on some transverse singular foliation, at least on a formal level. The same conclusion holds for simply-connected leaves provided a Levi-Malcev decomposition exists for the linear holonomy Lie algebroid (Theorems 2.8 and 2.22).

The paper is organized as follows: In Section 1, we review the notion of holonomy Lie algebroid A_L of a leaf L . Using the Artin-Rees theorem, we show that the sub-algebroid of A_L coming from vector fields in F that vanish at least quadratically along L form a nilpotent Lie algebra bundle. This allows us to describe the semi-simple quotient $A_L^{\mathbb{S}}$ of A_L as a quotient of the linear part A_L^{lin} of A_L . Using the method of Euler-like vector fields developed in [BLM19], we show that singular foliations that contain a transverse Euler vector field admit homogeneous generators (see Theorem 1.32). In Section 2, we state our most central result (Theorem 2.8) and give a geometric reformulation of it (Theorem 2.22).

Section 3 applies these results to leaves of dimension 0, recovering some results of Cerveau [Cer79] and deriving consequences for the NMRLA class of [LGLS20]. Finally, for arbitrary locally closed leaves, we show formal semi-local triviality of transversally quadratic leaves (Theorem 3.8) and linearly trivial leaves (Theorem 3.11), a phenomenon that is a distinctive feature of singular foliations, with no analogue in the Lie algebroid or Poisson manifold categories (Remark 3.9).

⁽¹⁾In most of this paper, we will deal with *locally real analytic* singular foliations F (i.e., M is covered by coordinate neighborhoods in which F admits real analytic generators—the change of coordinates does not need to be real analytic), see [LGLS20].

Acknowledgements. — We thank Marco Zambon for a crucial discussion at an early stage of the project, and several precious comments on the final version. We acknowledge valuable comments by Iakovos Androulidakis at relevant places in the text. Karandeep Singh also suggested several improvements. We would like to express our sincere gratitude to the anonymous referees for their helpful comments that helped to improve the quality of the manuscript.

1. HOLONOMY AND CONNECTIONS

1.1. **THE LINEAR HOLONOMY LIE ALGEBROID OF A LEAF.** — Let $F \subset \mathfrak{X}(M)$ be a singular foliation on a manifold M . The singular foliation F induces a “singular distribution” defined for every $p \in M$ by:

$$T_p F := \{X(p) \mid X \in F\}.$$

A fundamental Lemma about singular foliations, originating from Cerveau [Cer79], then proved in this context by Dazord [Daz85], and rediscovered by Androulidakis and Skandalis [AS09], says that singular foliations satisfy a local splitting property, in the following sense:

LEMMA 1.1 ([Cer79, Daz85, AS09]). — *Let $F \subset \mathfrak{X}(M)$ be a singular foliation on a manifold M of dimension n . Every point $p \in M$ admits a neighborhood U on which F is isomorphic to the direct product of the following two singular foliations:*

- (1) *all vector fields on an open ball of dimension d , where $d = \dim(T_p F)$, and*
- (2) *a singular foliation \top on an open ball of dimension $n - d$ made of vector fields that vanish at the origin, and called transverse singular foliation.*

The germ of the transverse singular foliation does not depend on any choice: any two local isomorphisms as above lead to transverse singular foliations which are locally isomorphic in a neighborhood of the origin.

This splitting lemma is crucial for proving the following results:

(1) By [Her63], M has a unique decomposition into submanifolds called *leaves* which are the maximal integral subsets for F . Moreover, the tangent space of the leaf through $p \in M$ is $T_p F$.

(2) Any two points p_1, p_2 on the the same leaf admit neighborhoods U_1, U_2 on which the singular foliations $F|_{U_1}$ and $F|_{U_2}$ are isomorphic ([Daz85]). In particular, their germs of transverse singular foliations are isomorphic. It makes sense, therefore, to speak of *the* transverse singular foliation of a given leaf.

In this subsection, we will define several Lie algebroids describing the behaviour of F near a chosen leaf L .

DEFINITION 1.2 ([AS09, AZ13]). — Let F be a singular foliation and L a locally closed leaf. Let $I_L \subset C^\infty(M)$ be the ideal of functions vanishing along L . The *holonomy Lie algebroid* $A_L \rightarrow L$ is defined implicitly by the equality $\Gamma(A_L) = F/I_L F$.

To verify that this yields a well-defined Lie algebroid, one shows that $F/I_L F$ is a Lie Rinehart-algebra and a projective $C^\infty(L)$ -module ([AZ13]). It is therefore a Lie algebroid. By construction, this Lie algebroid is transitive. It is therefore locally trivial. We denote by $\mathfrak{g}_L = \ker(\rho: A_L \rightarrow TL)$ its isotropy bundle of Lie algebras. At a given point $p \in L$, \mathfrak{g}_p is by construction the Androulidakis-Skandalis isotropy Lie algebra (see [AS09]) of F at p .

The holonomy Lie algebroid A_L acts on the normal bundle $\nu = TM|_L/TL$ of L in M , see [AZ13]. Algebraically, this action can be seen as follows: The space $\Gamma(\nu)$ is isomorphic to $\mathfrak{X}(M)/\mathfrak{X}_L(M)$, where $\mathfrak{X}_L(M) \subset \mathfrak{X}(M)$ are the vector fields tangent to L , i.e., the vector fields X satisfying $X(I_L) \subset I_L$. Now, $F \subset \mathfrak{X}_L(M)$ acts on this quotient and $I_L F$ acts trivially. The induced action of $F/I_L F$ on $\mathfrak{X}(M)/\mathfrak{X}_L(M)$ is a Lie algebroid action of A_L on the normal vector bundle ν .

This action is equivalently given by a homomorphism of transitive Lie algebroids $A_L \rightarrow \text{CDO}(\nu)$, where $\text{CDO}(\nu)$ is the Lie algebroid of *covariant differential operators* of $\nu \rightarrow L$ introduced by Mackenzie ([Mac87]). Recall that this Lie algebroid fits into the exact sequence

$$0 \longrightarrow \mathfrak{gl}(\nu) \longrightarrow \text{CDO}(\nu) \longrightarrow TL \longrightarrow 0$$

and that its sections can be interpreted as fiberwise linear vector fields on ν .

EXAMPLE 1.3. — For L a regular leaf, $A_L = TL$ and the TL -action on the normal bundle is the Bott-connection.

DEFINITION 1.4. — We call *linear holonomy Lie algebroid of L* the image Lie algebroid of $A_L \rightarrow \text{CDO}(\nu)$. We denote this Lie algebroid by A_L^{lin} .

EXAMPLE 1.5. — Let the leaf $L = \{p\}$ be a point. Then $A_L = F/I_p F = \mathfrak{g}_p$ is a Lie algebra and $\nu = T_p M$. The linear holonomy Lie algebroid is the Lie subalgebra of $\mathfrak{gl}(T_p M)$ obtained by linearizing all the vector fields in F :

$$\begin{array}{ccc} F & \xrightarrow{\text{lin}(X)} & \mathfrak{gl}(T_p M) \\ | & \searrow & \\ \mathfrak{g}_p & & \end{array}$$

The dotted arrow is well-defined, as vector fields in $I_p F$ vanish quadratically at p . Its image is the Lie algebra considered in [Cer79].

By definition of A_L^{lin} , there is a surjective Lie algebroid morphism $A_L \rightarrow A_L^{\text{lin}}$. Let us understand its kernel, which is a locally trivial bundle of Lie algebras by transitivity of A_L . Since all vector fields in F are tangent to the leaf L , derivation with respect to $X \in F$ preserves the filtration:

$$C^\infty(M) \supset I_L \supset I_L^2 \supset \dots$$

This implies that for all $i, j > 0$:

$$[F \cap I_L^i \mathfrak{X}(M), F \cap I_L^j \mathfrak{X}(M)] \subset F \cap I_L^{i+j-1} \mathfrak{X}(M).$$

This induces a natural filtration on $\Gamma(A_L)$ by

$$\Gamma(A_L)^i := \frac{F \cap I_L^i \mathfrak{X}(M)}{I_L F \cap I_L^i \mathfrak{X}(M)}.$$

In words, $\Gamma(A_L)^i$ is “the space of sections in A_L that can be represented by a vector field in F that vanishes at order i along L ”. This filtration obviously satisfies $[\Gamma(A_L)^i, \Gamma(A_L)^j] \subset \Gamma(A_L)^{i+j-1}$ and $[\Gamma(A_L)^i, \Gamma(A_L)] \subset \Gamma(A_L)^i$. By construction, we have:

LEMMA 1.6. — *There exists a vector bundle filtration of A_L :*

$$A_L = A_L^0 \supset A_L^1 \supset A_L^2 \supset \dots$$

such that $\Gamma(A_L^i) = \Gamma(A_L)^i$. Moreover, for every $i > 0$, A_L^i is a Lie algebra bundle, $A_L^1 = \mathfrak{g}_L$ and $A_L^2 = \ker(A_L \rightarrow A_L^{\text{lin}})$. In particular $A_L^{\text{lin}} = A_L/A_L^2$.

REMARK 1.7. — In general, the above filtration does not need to terminate, i.e., $\bigcap_{i>0} A_L^i$ might not be the zero vector bundle. For instance, consider the foliation on \mathbb{R} defined by the vector field $e^{-1/x^2} \partial/\partial x$. Then $\{0\}$ is a leaf for which $A_{\{0\}}^i = A_{\{0\}} = \mathbb{R}$ for all $i \in \mathbb{N}$, hence $\bigcap_i A_{\{0\}}^i = \mathbb{R}$. We will see, in the next subsection, that this pathology cannot happen for locally real analytic singular foliations.

1.2. NILPOTENCE AND THE SEMI-SIMPLE HOLONOMY. — For a locally real analytic singular foliation F , upon restriction to a neighborhood U of a point $p \in M$, we may assume that $F|_U$ has real analytic generators $(X_i)_{i=1}^r$ in some local coordinates, and consider the module F^{ra} over real analytic functions it generates, and define the real analytic holonomy Lie algebroid of the leaf through p (in U) by $F^{\text{ra}}/I F^{\text{ra}}$ (where I are real analytic functions vanishing on $L \cap U$ —which is easily seen to be a real analytic subvariety).

LEMMA 1.8. — *In the above setting, the (smooth) holonomy Lie algebroid and the real analytic holonomy Lie algebroid are isomorphic (as filtered Lie algebras). In equation: $\Gamma(A_L|_{U \cap L}) = F^{\text{ra}}/I F^{\text{ra}}$.*

Proof. — We have to show that the natural filtered Lie algebra morphism

$$F^{\text{ra}}/I F^{\text{ra}} \longrightarrow F/I_L F$$

induced by the inclusion is an isomorphism. This is a direct consequence of the statement that smooth functions form a faithfully flat module over real analytic functions ([Mal67, Cor. VI.1.12]).

1.2.1. The Artin-Rees Lemma and nilpotence. — In this section, we show that for locally real analytic foliations, the kernel of the linearization homomorphism $A_L \rightarrow A_L^{\text{lin}}$ is a bundle of nilpotent Lie algebras. The proof is based on the following statement of commutative algebra:

THEOREM 1.9 (Artin-Rees [AM69]). — *Let X be a finitely generated module over a Noetherian ring C , I be an ideal of C and $F \subset X$ a submodule. Then there is a positive integer c such that*

$$(1) \quad I^n X \cap F = I^{n-c} (I^c X \cap F) \quad \text{for all } n > c.$$

The classical formulation of the Artin-Rees lemma is more general, but we stated the form which is most directly applicable to our situation. In fact, we need the following immediate consequence of (1), applied with $n = c + 1$:

$$(2) \quad I^{c+1} X \cap F \subset I F.$$

For O, I, F, X as in Theorem 1.9, we call *Artin-Rees bound* of F in X at I the smallest integer that satisfies Condition (1).

THEOREM 1.10. — *Let F be a locally real analytic singular foliation and L a leaf. Then $A_L^{c+1} = 0$ where c is the Artin-Rees bound of F in $\mathfrak{X}(M)$ at I_L . In particular, the Lie algebra bundle A_L^2 is nilpotent.*

Proof. — Let $p \in L$. Upon restriction to a neighborhood U of $p \in M$, we may assume that $F|_U$ has real analytic generators $(X_i)_{i=1}^r$ in some local coordinates. By Lemma 1.8, we may use the real analytic holonomy Lie algebroid. According to Artin-Rees Theorem 1.9 applied in the following context:

- C is the algebra of real analytic functions on U (which is Noetherian by [Mal67, Th. III.3.8]),
- I is the ideal of real analytic functions on U vanishing along L ,
- X is the C -module of real analytic vector fields on U ,
- $F^{\text{ra}} \subset X$ is the sub-module generated by the vector fields $(X_i)_{i=1}^r$,

there exists $c \in \mathbb{N}$ that satisfies (1) and therefore (2). Geometrically, (2) applied with $n = c + 1$ implies that a vector field in F^{ra} that vanishes at order $c + 1$ along L belongs to $I F^{\text{ra}}$. In terms of the filtration in Lemma 1.6, it means that $A_L^{c+1} = 0$. In particular A_L^2 is a bundle of nilpotent Lie algebras of depth less than or equal to $c + 1$.

The following example (inspired by Grabowska and Grabowski [GG20]) illustrates that the Artin-Rees bound can be arbitrarily large in our situation, i.e., that the filtration on A_L may have arbitrarily many non-zero terms.

EXAMPLE 1.11. — On $M = \mathbb{R}^n$, let us give to the coordinates (x_1, \dots, x_n) the weights $(1, \dots, n)$. Real analytic functions on M then become a filtered algebra. Real analytic vector fields that preserve this filtration form a module F stable under Lie bracket and generated by the finite family

$$\{x_1^{i_1} \dots x_n^{i_n} \partial / \partial x_k \mid k \in [1 : n], i_1, \dots, i_n \in [0 : n] \text{ and } i_1 + 2i_2 + \dots + ni_n > k\}.$$

Since all vector fields in F vanish at the origin 0 , $L = \{0\}$ is a leaf. The vector field $x_1^n \partial / \partial x_n$ is an element in $F \cap I_L^n \mathfrak{X}(M)$, but does not belong to $I_L F$. This implies

that the Artin-Rees bound c is greater than or equal to n . Since $I_L^i \mathfrak{X}(M) \subset F$ for $i > n$, we have $I_L^{n+1} \cap \mathfrak{X}(M) \subset F = I_L(I_L^n \cap \mathfrak{X}(M))$, so that $c = n$.

The Artin-Rees bound also bounds the possible degrees of generators of a singular foliation which is preserved by some Euler vector field, as stated below.

PROPOSITION 1.12. — *Let F be a real analytic singular foliation in a neighborhood of 0 in \mathbb{R}^n made of vector fields that vanish at 0 . If F is preserved by the Euler vector field $E = \sum_{i=1}^n x_i \partial/\partial x_i$, then:*

- (1) *Every homogeneous component of a vector field in F belongs to F .*
- (2) *F admits homogeneous generators whose degrees are less than or equal to the Artin-Rees bound of F at 0 .*

Notice that we will extend Proposition 1.12 to neighborhood of leaves (see Theorem 1.32 below). To prove Proposition 1.12, we start with a lemma:

LEMMA 1.13. — *For $k > 1$, the operator*

$$P^k : \mathfrak{X}(\mathbb{R}^n) \longrightarrow \mathfrak{X}(\mathbb{R}^n)$$

$$X \longmapsto ([E, X] - (k-1) \cdot X)$$

restricts to an invertible isomorphism of $F \cap I^{k+1} \mathfrak{X}(\mathbb{R}^n)$, where $I = I_{\{0\}}$ is the ideal of functions vanishing at 0 .

Proof. — The operator P^k restricts to an endomorphism of $F \cap I^{k+1} \mathfrak{X}(\mathbb{R}^n)$, since the identity map and the Lie bracket with E preserve both F and $I^{k+1} \mathfrak{X}(\mathbb{R}^n)$. Furthermore, the restriction $P^k|_{I^{k+1} \mathfrak{X}(\mathbb{R}^n)}$ of P^k to $I^{k+1} \mathfrak{X}(\mathbb{R}^n)$ is injective as the kernel of P^k is given by homogeneous vector fields of degree k , a space in trivial intersection with $I^{k+1} \mathfrak{X}(\mathbb{R}^n)$. Hence, the restriction $P^k|_{F \cap I^{k+1} \mathfrak{X}(\mathbb{R}^n)}$ is also injective.

We claim that an explicit inverse to $P^k|_{I^{k+1} \mathfrak{X}(\mathbb{R}^n)}$ is given by

$$Q^k : I^{k+1} \mathfrak{X}(\mathbb{R}^n) \longrightarrow I^{k+1} \mathfrak{X}(\mathbb{R}^n)$$

$$(3) \quad Z = \sum_{i=1}^n z_i(x) \frac{\partial}{\partial x_i} \longmapsto Q^k(Z) = \int_{t=0}^1 \left(\frac{1}{t^{k+1}} \sum_{i=1}^n z_i(tx) \frac{\partial}{\partial x_i} \right) dt.$$

The convergence of the integral is guaranteed by the fact that all functions $t \mapsto z_i(tx)$ vanish at order $k + 1$ at 0 . A simple integration by part gives $P^k \circ Q^k = \text{id}_{I^{k+1} \mathfrak{X}(\mathbb{R}^n)}$. It is also clear that each one of the functions $\int_{t=0}^1 (1/t^{k+1}) z_i(tx) dt$ belongs to I^{k+1} , so that the image of Q^k is indeed in $I^{k+1} \mathfrak{X}(\mathbb{R}^n)$.

To conclude the proof, we need to show that Q^k preserves the subspace

$$F \cap I^{k+1} \mathfrak{X}(\mathbb{R}^n).$$

First, let us interpret Equation (3) as:

$$(4) \quad Q^k(Z) = \int_{t=0}^1 \frac{1}{t^k} \mu_*^t(Z) dt,$$

where μ^t is the homothety $x \mapsto x/t$. Since μ^t is the the flow at time $-\ln(t)$ of E , it preserves F by Proposition 1.6 in [AS09]. In particular, if $Z \in F$, then $\mu_x^t(Z) \in F$ for all $t \in]0, 1]$. Now, since F admits real analytic generators, F is closed with respect to the Fréchet topology (see [Tou68, Th. 2]—the result is attributed to Malgrange). In particular, F is stable under the integration (4), so that if $Z \in F \cap I^{k+1}\mathfrak{X}(\mathbb{R}^n)$, then $Q^k(Z) \in F$. This proves the lemma.

Proof of Proposition 1.12. — Let us decompose $X \in F \cap I^k\mathfrak{X}(\mathbb{R}^n)$ as $X = X^{(k)} + R$, with $X^{(k)}$ homogeneous of degree k , and $R \in I^{k+1}\mathfrak{X}(\mathbb{R}^n)$. As $X^{(k)}$ is in the kernel of P^k , we have

$$P^k(R) = P^k(X) \in F \cap I^{k+1}\mathfrak{X}(\mathbb{R}^n).$$

By Lemma 1.13, this implies that $R \in F \cap I^{k+1}\mathfrak{X}(\mathbb{R}^n)$, so that $X^{(k)} = X - R \in F$. This proves that the lowest component of an element in F is in F . The first item of the proposition follows by an immediate finite induction.

For (e_1, \dots, e_b) a local trivialization of A_L^k/A_L^{k+1} , let us choose (X_1, \dots, X_b) a b -tuple of elements in F that represent it. The b -tuple $(X_1^{(k)}, \dots, X_b^{(k)})$ of their homogeneous components of degree k is again made of element of F by the first item, and still represents (e_1, \dots, e_b) . Applying this procedure for all $k = 0, \dots, r$, we obtain a basis of $\mathfrak{g} = A_L = A_L^0$ which are all represented by homogeneous vector fields in F of degree less than or equal to the Artin-Rees bound. In view of [AS09, Prop. 1.5 (a)], these vector fields are generators of F . This proves the second item.

REMARK 1.14. — In the case $\mathfrak{g}_0 = \mathfrak{g}_0^{\text{lin}}$, Proposition 1.12 reduces to [Cer79, Th. 8.1]—a result extended to a neighborhood of a leaf by Marco Zambon [Zam19].

1.2.2. *The Levi exact sequence and the semi-simple holonomy Lie algebroid.* — For a Lie algebra \mathfrak{g} , the Levi-Malcev decomposition theorem goes as follows:

- (i) \mathfrak{g} has a unique maximal solvable ideal $\mathfrak{rad}(\mathfrak{g})$,
- (ii) the quotient $\mathfrak{g}/\mathfrak{rad}(\mathfrak{g})$ is a semi-simple Lie algebra \mathfrak{g}^{S} , and
- (iii) there is a section $\mathfrak{g}^{\text{S}} \hookrightarrow \mathfrak{g}$ which is a Lie algebra homomorphism.

The Levi-Malcev theorem does not easily generalize to transitive Lie algebroid. Remark 1.17 gives a counter-example to step (iii), but steps (i) and (ii) admit generalizations that we now describe. For $A \rightarrow L$ a transitive Lie algebroid with anchor ρ the isotropy Lie algebra bundle $\mathfrak{g} := \ker(\rho)$ is locally trivial ([Mac87, Th. 8.2.1]), so that the *fiberwise radical*, i.e., the disjoint union $\coprod_{l \in L} \mathfrak{rad}(\mathfrak{g}_l)$ is indeed a Lie algebra bundle over L . We denote it by $\mathfrak{rad}(A)$. Since A is, near every point $m \in L$, a direct product of $TL \rightarrow L$ with its isotropy Lie algebra at m (see, e.g. [Zun03, Th. 1.2]), sections of $\mathfrak{rad}(A)$ form an ideal of $\Gamma(A)$. This proves the following proposition:

PROPOSITION 1.15. — *For every transitive Lie algebroid A over L , the quotient $A/\mathfrak{rad}(A)$ is a transitive Lie algebroid over L , with semi-simple isotropies, and*

$$(5) \quad \mathfrak{rad}(A) \longrightarrow A \longrightarrow A/\mathfrak{rad}(A)$$

is a short exact sequence of Lie algebroids.

These general considerations lead to the following definition.

DEFINITION 1.16. — Let L be a leaf of a singular foliation F . We call the quotient $A_L/\mathfrak{rad}(\mathfrak{g}_L)$ the *semi-simple holonomy Lie algebroid* of L and denote it by $A_L^{\mathbb{S}}$.

REMARK 1.17. — It is not true in general that the short exact sequence (5) admits a Lie algebroid section, even for holonomy Lie algebroid A_L of a leaf of a singular foliation. For instance, for $A = TM \oplus \mathbb{R}$ equipped with the Lie algebroid structure associated to a closed 2-form $\omega \in \Omega^2(M)$, the semi-simple holonomy Lie algebroid $A/\mathfrak{rad}(A)$ is the tangent Lie algebroid TM , but a Lie algebroid section $TM \hookrightarrow A$ exists if and only if ω is exact.

PROPOSITION 1.18. — Let L be a leaf of a locally real analytic singular foliation F . There is a natural Lie algebroid morphism $A_L^{\text{lin}} \twoheadrightarrow A_L^{\mathbb{S}}$ that makes the following diagram commutative:

$$\begin{array}{ccc} A_L & \xrightarrow{\quad} & A_L^{\text{lin}} \\ & \searrow & \vdots \\ & & A_L^{\mathbb{S}} \end{array}$$

Proof. — By Theorem 1.10, the kernel of $A_L \rightarrow A_L^{\text{lin}}$ is a bundle of nilpotent Lie algebras. It is therefore contained in $\mathfrak{rad}(A_L)$, i.e., in the kernel of the natural projection $A_L \rightarrow A_L^{\mathbb{S}}$. The result follows.

1.3. CONNECTION THEORY

1.3.1. Ehresmann or Levi F -connections and flatness. — According to Proposition 1.18, for every leaf L of a locally singular foliation F , we have the following sequence of surjective morphisms of transitive Lie algebroids over L :

$$F \dashrightarrow A_L \longrightarrow A_L^{\text{lin}} \longrightarrow A_L^{\mathbb{S}} \longrightarrow TL,$$

where the leftmost arrow is dashed, as it is not a morphism of Lie algebroids. The main purpose of this article is to describe the behaviour of F in a neighborhood of L , using the semi-simple holonomy Lie algebroid $A_L^{\mathbb{S}}$.

To start, we will consider as in [LGR19] neighborhoods U of L in M which are small enough in the following sense: they have to admit a projection $\pi: U \rightarrow L$ such that $T_x F + \ker(T_x \pi) = T_x M$ for all $x \in U$. These pairs (U, π) shall be called *F -neighborhoods* and satisfy several important properties, in particular, by Proposition 2.21 in [LGR19]:

LEMMA 1.19. — Every locally closed leaf L of a singular foliation F admits an F -neighborhood (U, π) .

As we are only interested in the behaviour of F near L , for the rest of the section we assume that $M = U$ is an F -neighborhood equipped with some projection $\pi: M \rightarrow L$. The $C^\infty(L)$ -modules of π -vertical (resp. π -projectable) vector fields will be denoted by

$\mathfrak{X}^v \subset \mathfrak{X}^{\text{proj}} \subset \mathfrak{X}(U)$. We also write F^v and F^{proj} for the π -vertical and π -projectable vector fields in F .

LEMMA 1.20. — *The Lie algebra F^{proj} of π -projectable vector fields in F form a Lie-Rinehart algebra over $C^\infty(L)$. Moreover, there is a sequence of surjective Lie-Rinehart algebra morphisms:*

$$(6) \quad F^{\text{proj}} \longrightarrow \Gamma(A_L) \longrightarrow \Gamma(A_L^{\text{lin}}) \longrightarrow \Gamma(A_L^s) \longrightarrow \mathfrak{X}(L).$$

We introduce several types of connection adapted to this context:

DEFINITION 1.21 ([And17, LGR19]). — Let L be a leaf of a foliation F on $(M = U, \pi)$.

– An *Ehresmann F -connection* is a $C^\infty(L)$ -linear section $\mathfrak{X}(L) \rightarrow F^{\text{proj}}$ of the surjection $F^{\text{proj}} \rightarrow \mathfrak{X}(L)$ in (6).

– A *Levi F -connection* is a $C^\infty(L)$ -linear section $s : \Gamma(A_L^S) \rightarrow F^{\text{proj}}$ of the surjection $F^{\text{proj}} \rightarrow \Gamma(A_L^S)$ in (6).

Existence of an Ehresmann F -connection was already established in [And17, LGR19]. We now extend this result:

PROPOSITION 1.22. — *Let L be a leaf of a foliation F on $(M = U, \pi)$. Then, possibly on a sub- F -neighborhood of L , Levi F -connections and Ehresmann F -connections exist.*

The proposition follows from the following lemma:

LEMMA 1.23. — *Let L be a leaf of a foliation F on $(M = U, \pi)$. Then, possibly on a sub- F -neighborhood of L the surjection $F^{\text{proj}} \rightarrow \Gamma(A_L)$ admits a $C^\infty(L)$ -linear section.*

Proof. — When L is a point, A_L is a Lie algebra, and such a section s exists. In particular, there exists for all $p \in L$ a linear section $s_p : \mathfrak{g}_p \rightarrow F_p^v$. In view of the splitting Lemma 1.1, every point p admits a neighborhood U in L which admits a neighborhood V on M such that

$$A_L|_U \simeq TL|_U \oplus \mathfrak{g}_p \text{ and } F^{\text{proj}}|_V \simeq \Gamma(TL) \oplus F^v.$$

Under this identification, $(\text{id} \times s_p)$ is a section on $F^{\text{proj}}|_V \rightarrow \Gamma(A_L)|_U$.

Let $(U_i)_{i \in I}$ be an open cover of L , such that each U_i comes equipped with a $C^\infty(U_i)$ -linear section s_i of $p : F^{\text{proj}}|_{V_i} \rightarrow \Gamma(A_L)|_{U_i}$ for some open subset $V_i \subset p^{-1}(U_i)$. Without any loss of generality, one can assume that the open cover $(U_i)_{i \in I}$ is locally finite and comes with a partition of unit $(\chi_i)_{i \in I}$. Then:

$$V := \bigcup_{x \in L} \left(\bigcap_{i \in I | x \in U_i} V_i \right)$$

is an open neighborhood of L , and $s := \sum_{i \in I} \chi_i s_i$ is a well-defined $C^\infty(L)$ -linear section of $p : F^{\text{proj}}|_V \rightarrow \Gamma(A_L)$

Proof of Proposition 1.22. — The composition of any linear section $TL \rightarrow A_L$ (resp. $A_L^{\mathcal{S}} \rightarrow A_L$) with a section as in Lemma 1.23 yields an Ehresmann F -connection (resp. a Levi F -connection). This proves the statement.

DEFINITION 1.24. — An Ehresmann/Levi F -connection is called *flat* if it is bracket-preserving (i.e., a morphism of Lie-Rinehart algebras).

Let us give the geometric interpretation of the existence of flat Ehresmann/Levi connections:

PROPOSITION 1.25. — *Let L be a leaf of a singular foliation F .*

– *A flat Ehresmann F -connection exists if and only if near L there exists a regular foliation included into F admitting L as a leaf.*

– *A flat Levi F -connection exists if and only if there exists a Lie algebroid action of $A_L^{\mathcal{S}}$ on $\pi: M \rightarrow L$ made of vector fields in F .*

REMARK 1.26. — Let L be a locally closed leaf. A flat Ehresmann connection induces a flat section of the anchor map $A_L^{\mathcal{S}} \rightarrow TL$. Also, if L is Ehresmann-flat, then its normal bundle ν is a flat bundle. This gives clear obstructions to the existence of flat Ehresmann connections. In contrast, flat Levi F -connection can be assured to exist under relatively mild topological conditions, as we will see later.

1.3.2. Linear Ehresmann or Levi F -connections. — An additional desirable property for a Ehresmann or Levi F -connection is (transverse) linearity. For this purpose, we need to notion of fiberwise linearity. This is completed through the following definition adapted from [BLM19]:

DEFINITION 1.27. — Consider an F -neighborhood (U, π) of a locally closed leaf L . A vector field $E \in U$ which is:

- (i) tangent to the fibers of $\pi: U \rightarrow L$,
- (ii) vanishes along L ,
- (iii) whose linearization is the Euler vector field on the normal bundle ν , and
- (iv) that is complete

is said to be an Euler-like vector field on (U, π) .

Upon rescaling the vector field and shrinking the tubular neighborhood (U, π) the completeness condition (iv) can always be assumed for vector field satisfying (i)–(iii). The following lemma is the adaptation of [BLM19] to the case where π is given and E is assumed to be tangent to it.

LEMMA 1.28. — *Euler-like vector fields on an F -neighborhood (U, π) are in one-to-one correspondence with vector bundle structures on the fiber bundle $\pi: U \rightarrow L$.*

REMARK 1.29. — Every such a vector bundle is isomorphic to the normal bundle $\nu = TM|_L/TL$ of L in M .

DEFINITION 1.30. — Consider an F -neighborhood (U, π) of a locally closed leaf L . We say that a Levi (resp. Ehresmann) F -connection $s: \Gamma(A_L^S) \hookrightarrow F_U^{\text{proj}}$ (resp. $s: \Gamma(TL) \hookrightarrow F_U^{\text{proj}}$) is *linear* with respect to an Euler-like vector field E if there exists a neighborhood of the zero section on which every vector field in the image of s commutes with E .

REMARK 1.31. — Upon identifying U with a vector bundle as in Lemma 1.28, vector fields commuting with E are simply fiberwise linear vector fields.

We say that a vector field X is homogeneous of degree k with respect to E if $[E, X] = (k - 1)X$. (Linear vector fields are then homogeneous of degree 1). Upon choosing adapted coordinates (x, y) where $E = \sum_{i=1}^d x_i \partial / \partial x_i$, homogeneous vector fields of degree k are vector fields of the form:

$$\sum_i f_i(x, y) \frac{\partial}{\partial x_i} + \sum_j g_j(x, y) \frac{\partial}{\partial y_j},$$

where $x \mapsto f_i(x, y)$ and $x \mapsto g_j(x, y)$ are homogeneous polynomials of degree k and $k - 1$ respectively for every value y .

THEOREM 1.32. — *Let F be a locally real analytic singular foliation with leaf L . If F is preserved by an Euler-like vector field E along L , then near L :*

- (1) *Any homogeneous component of a vector field in F is in F .*
- (2) *The foliation F is generated by homogeneous vector fields (of degree less than or equal than the Artin-Rees bound of the transverse foliation).*
- (3) *There exists a $C^\infty(L)$ -linear section $s: \Gamma(A_L^{\text{lin}}) \rightarrow F^{\text{proj}}$ preserving the Lie bracket.*

Proof. — Fixing a tubular neighborhood adapted to E , the expressions for P^k and Q^k from Lemma 1.13 still make sense, and satisfy the same relations on vector fields tangent to L . Hence, we can proceed identically to Proposition 1.12. This proves the first two items. A bracket-preserving isomorphism from $\Gamma(A_L^{\text{lin}})$ to linear vector fields in F is obtained by mapping $X \in \Gamma(A_L^{\text{lin}})$ to the linear component of any of its inverse images in F .

For foliations as in Theorem 1.32, the existence of a flat section $s: \Gamma(A_L^S) \rightarrow F^{\text{proj}}$ is therefore equivalent to the existence of a Lie algebroid section $s: A_L^S \rightarrow A_L^{\text{lin}}$.

COROLLARY 1.33. — *Let F be a locally real analytic singular foliation with leaf L . If F is preserved by an Euler-like vector field E along L , then a flat Levi F -connection exists if and only if a Lie algebroid section $A_L^S \rightarrow A_L^{\text{lin}}$ exists.*

2. THE FORMAL NEIGHBORHOOD OF A SIMPLY CONNECTED (SINGULAR) LEAF

2.1. LIE ALGEBROID COHOMOLOGY OF SMALL DEGREES. — In order to prove our central Theorem 2.8, we will need the following statements about Lie algebroid cohomology.

The Whitehead lemmas I and II admit generalizations for Lie algebroids that we now state, using several results of Kirill Mackenzie (in [Mac05]):

LEMMA 2.1 (Whitehead Lemma I for Lie algebroids). — *Let $A \rightarrow L$ be a transitive Lie algebroid with semi-simple isotropies $\mathfrak{g}_L = \ker(\rho)$. If $\pi_1(L) = 0$, then the Lie algebroid cohomology group $H^1(A, E)$ is trivial for any flat finite-dimensional A -module $E \rightarrow L$.*

Proof. — Theorem 7.4.5 in Kirill Mackenzie [Mac05] asserts that there is a spectral sequence whose first page is $H^t(L, H_{\text{CE}}^s(\mathfrak{g}_L, E))$ converging to $H^\bullet(A, E)$. The only two terms contributing to $H^1(A, E)$ are $H^0(L, H_{\text{CE}}^1(\mathfrak{g}_L, E))$ and $H^1(L, H_{\text{CE}}^0(\mathfrak{g}_L, E))$. The former space is trivial, due to Whitehead’s lemma for Lie algebras and the semi-simplicity of \mathfrak{g}_L . The second one is trivial, because L is simply connected.

LEMMA 2.2 (Whitehead Lemma II for Lie algebroids). — *Let $A \rightarrow L$ be a transitive Lie algebroid with semi-simple isotropies $\mathfrak{g}_L = \ker(\rho)$. If $\pi_1(L) = \pi_2(L) = 0$, then the Lie algebroid cohomology group $H^2(A, E)$ is trivial for any flat finite-dimensional A -module $E \rightarrow L$.*

Proof. — Let us use again Mackenzie’s spectral sequence used the proof of Lemma 2.1. Here, the components of the first page required to be trivial are $H^0(L, H_{\text{CE}}^2(\mathfrak{g}_L, E))$, $H^1(L, H_{\text{CE}}^1(\mathfrak{g}_L, E))$, and $H^2(L, H_{\text{CE}}^0(\mathfrak{g}_L, E))$. The former two are again trivial by the Whitehead lemmas for Lie algebras. As L is simply connected the flat vector bundle $H_{\text{CE}}^0(\mathfrak{g}_L, E) = \Gamma(E^{\mathfrak{g}_L})$ is in fact a trivial vector bundle, so that $H^2(L, H_{\text{CE}}^0(\mathfrak{g}_L, E)) = H^2(L) \otimes H_{\text{CE}}^0(\mathfrak{g}_L, E)$. By the Hurewicz theorem, $\pi_1(L) = \pi_2(L) = 0$ implies that $H^2(L) = 0$. The third component is therefore zero and the result follows.

We prove the following generalization of the Levi-Malcev theorem, which is also a prototype for our central Theorem 2.8.

PROPOSITION 2.3. — *Let A be a transitive Lie algebroid over a 2-connected base L (i.e., $\pi_1(L) = \pi_2(L) = 0$) and A^S its semi-simple quotient. Then there exists a Lie algebroid section $s : A^S \rightarrow A$ of the projection $\pi : A \rightarrow A^S$.*

Proof. — The Lie algebra bundle $\mathfrak{rad}(A) = \ker(\pi)$, defined in Section 1.2.2 comes with a terminating natural filtration by Lie algebra bundles

$$(7) \quad \mathfrak{r}^0 = \mathfrak{rad}(A) \supset \mathfrak{r}^1 = [\mathfrak{r}^0, \mathfrak{r}^0] \supset \mathfrak{r}^2 = [\mathfrak{r}^1, \mathfrak{r}^1] \supset \dots \supset \mathfrak{r}^N = 0$$

such that the subquotients $\mathfrak{r}^i/\mathfrak{r}^{i+1}$ are Abelian. We construct s by induction. Let $s^0 : A^S \rightarrow A$ be any linear section: Its curvature is valued in the radical $\mathfrak{r}^0 = \mathfrak{rad}(A)$. Assume that there exists a section $s^i : A^S \rightarrow A$ whose curvature c^i is \mathfrak{r}^i -valued, then:

- (1) the quotient space $\mathfrak{r}^i/\mathfrak{r}^{i+1}$ is an A^S -module for: $(\xi, \bar{b}) \mapsto [\overline{s^i(\xi)}, \bar{b}]$, with $\xi \in \Gamma(A^S)$, $b \in \Gamma(\mathfrak{r}^i)$,
- (2) the skew-symmetric bilinear map:

$$(\xi, \zeta) \longmapsto c^i(\xi, \zeta) \pmod{\mathfrak{r}^{i+1}}$$

is a Lie algebroid 2-cocycle of A^S , valued in $\mathfrak{r}^i/\mathfrak{r}^{i+1}$.

By the second Whitehead lemma 2.2, this cocycle is a coboundary, so that there exists $\sigma_i : A \rightarrow \mathfrak{r}^i$ such that:

$$c_i(\xi, \zeta) = [s^i(\xi), \sigma^i(\zeta)] + [\sigma^i(\xi), s^i(\zeta)] - \sigma^i([\xi, \zeta]) \pmod{\mathfrak{r}^{i+1}}.$$

This means that the curvature of $s^{i+1} = s^i + \sigma^i$ is \mathfrak{r}^{i+1} -valued. Since the filtration (7) terminates at degree N , s^N is a Lie algebroid section.

REMARK 2.4. — Proposition 2.3 should not be confused from the Zung and the Monnier-Zung's Levi theorems for Lie algebroids [Zun03, MZ04], which is valid for any Lie algebroid, not only transitive ones, but is a local result.

Let us state an immediate consequence of Proposition 2.3 and Theorem 1.10:

COROLLARY 2.5. — *Let F be a singular foliation and L a locally closed leaf. If $\pi_1(L) = \pi_2(L) = 0$, then there exists a Lie algebroid section $A_L^{\mathbb{S}} \rightarrow A_L^{\text{lin}}$. In case F is locally real analytic, then there even exists a Lie algebroid section $A_L^{\mathbb{S}} \rightarrow A_L$.*

The local analyticity assumption on F is probably not necessary: to suppress it, we would need to extend Theorem 1.10 to all smooth singular foliations.

2.2. FORMAL SINGULAR FOLIATIONS. — In this section, we define the formal counterparts of several notions that have been studied in this article.

For L be a submanifold of M , we denote by \widehat{C} the algebra of formal functions along L , i.e., (by Borel's theorem) the quotient of $C^\infty(M)$ by the ideal of functions vanishing with all their derivatives along L . We call *formal vector fields along L* derivations of \widehat{C} .

We call *formal singular foliation along L* locally finitely generated \widehat{C} -submodules of formal vector fields along L which are closed under Lie bracket. For every singular foliation F , the tensor product $\widehat{F} := \widehat{C} \otimes_{C^\infty(M)} F$ is a formal singular foliation along L called the *formal jet of F along L* . We will only consider formal singular foliations for which L is a leaf, i.e., such that the restriction to L is onto $\mathfrak{X}(L)$.

As for the non-formal case, we call *holonomy Lie algebroid* of a formal singular foliation \widehat{F} along L the Lie algebroid whose space of sections is the quotient $\widehat{F}/\widehat{I}_L\widehat{F}$, with $\widehat{I}_L = \widehat{C} \otimes_{C^\infty(M)} I_L \subset \widehat{C}$ the ideal of formal functions vanishing along L . As in Lemma 1.8, since formal functions are a faithfully flat module over real analytic functions (cf. [Mal67, Th. III.4.9]), we have:

LEMMA 2.6. — *If a singular foliation F is locally real analytic, then for every leaf L , the holonomy Lie algebroids of F and of its formal jet \widehat{F} along L are isomorphic. In equation: $F/I_L F \simeq \widehat{F}/\widehat{I}_L\widehat{F}$.*

The notions of Levi F -connections and Ehresmann F -connections have formal equivalents for a formal singular foliation \widehat{F} near a leaf L . First, let us choose a tubular neighborhood $\pi : U \rightarrow L$. A formal vector field X is said to be projectable if it preserves $\pi^*C^\infty(L) \subset \widehat{C}$. Denote formal projectable vector fields in \widehat{F} by $\widehat{F}^{\text{proj}}$.

By construction, $\widehat{F}^{\text{proj}}$ is a Lie-Rinehart algebra over $C^\infty(L)$. There are natural Lie-Rinehart algebra morphisms from $\widehat{F}^{\text{proj}}$ to $\Gamma(A_L^\mathbb{S})$ and $\mathfrak{X}(L)$ respectively as in (6). We call *formal Levi F -connection* (resp. *formal Ehresmann F -connection*) a $C^\infty(L)$ -linear section $\Gamma(A_L^\mathbb{S}) \rightarrow \widehat{F}^{\text{proj}}$ (resp. $\mathfrak{X}(L) \rightarrow \widehat{F}^{\text{proj}}$) of these natural surjections.

REMARK 2.7. — For future use, notice that the time t -flow of a formal vector field X tangent to L is a well-defined algebra isomorphism of \widehat{C} , as long as the time t -flow of its restriction to L is defined. If X is π -projectable for some $\pi: M \rightarrow L$, this formal diffeomorphism maps π -fibers to π -fibers.

2.3. EXISTENCE OF FLAT LEVI F -CONNECTIONS. — The main goal of the section is to prove the following theorem:

THEOREM 2.8. — *Let L be a locally closed leaf of the locally real analytic foliation F . If $\pi_1(L) = 0$, and there exists a Lie algebroid section z from the semi-simple holonomy $A_L^\mathbb{S}$ to the linear holonomy A_L^{lin} , then there exist*

- an F -neighborhood (U, π) ,
- a formal Levi F -connection $s^\infty: \Gamma(A_L^\mathbb{S}) \rightarrow \widehat{F}^{\text{proj}}$,
- and a π -vertical formal Euler-like vector field E^∞ ,

such that

- s^∞ is linear with respect to E^∞ , i.e., $[s^\infty(\xi), E^\infty] = 0$ for all $\xi \in \Gamma(A_L^\mathbb{S})$,
- s^∞ is flat, i.e., $[s^\infty(\xi), s^\infty(\zeta)] = s^\infty([\xi, \zeta])$ for all $\xi, \zeta \in \Gamma(A_L^\mathbb{S})$.

Here is an immediate consequence of this theorem and Corollary 2.5.

COROLLARY 2.9. — *Let L be a locally closed leaf of the locally real analytic foliation F . If $\pi_1(L) = \pi_2(L) = 0$, then the conclusions of Theorem 2.8 hold.*

Theorem 2.8 will be proved by induction. The initial case is given by Lemma 2.11. Proposition 2.12 gives the induction step. Both result depend on the technical Lemma 2.10.

We denote by \mathfrak{X}^ν vertical vector fields for $\pi: U \rightarrow L$ so that, for every $k \in \mathbb{N}$, $I_L^k \mathfrak{X}^\nu$ stands for vertical vector fields vanishing at order k along L .

LEMMA 2.10. — *Let E be a π -vertical Euler-like vector field on (U, π) . For every π -vertical vector field $X \in \mathfrak{X}^\nu$ and every $k > 2$:*

- (1) if $X \in I_L^k \mathfrak{X}^\nu$, then $(1/(k-1))[E, X] - X \in I_L^{k+1} \mathfrak{X}^\nu$;
- (2) if the linearization of X along L is zero, and $[E, X] \in I_L^k \mathfrak{X}^\nu$, then $X \in I_L^k \mathfrak{X}^\nu$.

Proof. — It suffices to check both items in local adapted coordinates (x, y) where $y = (y_1, \dots, y_d)$ are local coordinates on L and $x = (x_1, \dots, x_c)$ are local coordinates on the fibers of $\pi: (x, y) \mapsto y$ such that $E = \sum_{i=1}^c x_i \partial / \partial x_i$. Since the ideal I_L is generated by x_1, \dots, x_c , the Taylor expansion implies that for every $X \in I_L^k \mathfrak{X}^\nu$ there

exists functions $f_{i;i_1,\dots,i_c}(y)$ such that

$$X = \sum_{i=1}^c \sum_{i_1+\dots+i_c=k} f_{i;i_1,\dots,i_c}(y) x_1^{i_1} \dots x_c^{i_c} \frac{\partial}{\partial x_i} \pmod{I_L^{k+1}\mathfrak{X}^v}.$$

Since $[E, I_L^{k+1}\mathfrak{X}^v] \subset I_L^{k+1}\mathfrak{X}^v$, the first item follows from the easily checked identity:

$$(8) \quad \left[E, x_1^{i_1} \dots x_c^{i_c} \frac{\partial}{\partial x_i} \right] = (k-1) x_1^{i_1} \dots x_c^{i_c} \frac{\partial}{\partial x_i}.$$

Let us prove the second item. By assumption, the linearization of $X \in \mathfrak{X}^v$ along L is zero, so that $X \in I_L^2\mathfrak{X}^v$. The conclusion then follows from considering the Taylor expansion of X , in view of Equation (8).

A pair (s, E) with s a Levi F -connection and E a π -vertical Euler-like vector field are said to be *flat and linear up to order k* if:

- $[s(\xi), E] = 0 \pmod{I_L^k\mathfrak{X}^v}$ for all $\xi \in \Gamma(A_L^\mathbb{S})$,
- $[s(\xi), s(\zeta)] = s([\xi, \zeta]) \pmod{I_L^k\mathfrak{X}^v}$ for all $\xi, \zeta \in \Gamma(A_L^\mathbb{S})$.

LEMMA 2.11. — *For every Euler-like vector field E and every Levi F -connection s :*

- $[E, s(\xi)] = 0 \pmod{I^2\mathfrak{X}^v}$,
- *If, in addition, s projects to a Lie algebroid morphism $z: A_L^\mathbb{S} \rightarrow A_L^{\text{lin}}$, then $[s(\xi), s(\zeta)] = s([\xi, \zeta]) \pmod{I_L^2\mathfrak{X}^v}$ for all $\xi, \zeta \in \Gamma(A_L^\mathbb{S})$.*

Proof. — Since E is an Euler-like vector field, a neighborhood U of L can be covered by local charts, equipped with coordinates $(x_1, \dots, x_c, y_1, \dots, y_d)$, such that:

- (1) L is locally given by $x = 0$, and π is locally given by $(x, y) \mapsto y$,
- (2) E coincides with the vector field $\sum_{i=1}^c x_i \partial / \partial x_i$.

The Taylor expansion of any π -projectable vector field X is given by

$$X = \sum_{i=0}^d f_i(y) \frac{\partial}{\partial y_i} + \sum_{i,j=1}^c a_{ij}(y) x_i \frac{\partial}{\partial x_j} + I_L^2\mathfrak{X}^v.$$

A simple calculation gives $[E, X] \in I_L^2\mathfrak{X}^v$, as $\sum_{i=0}^d f_i(y) \partial / \partial y_j + \sum_{i,j=1}^c a_{ij}(y) x_i \partial / \partial x_j$ commutes with E and $I_L^2\mathfrak{X}^v$ is preserved by E by Lemma 2.10. Since $s(\xi)$ is π -projectable for all $\xi \in \Gamma(A_L^\mathbb{S})$, the first assertion follows. The second assertion is a consequence of the following facts:

- (1) the assumption on s means that for all $\xi, \zeta \in \Gamma(A_L^\mathbb{S})$, the π -vertical vector field $[s(\xi), s(\zeta)] - s([\xi, \zeta])$ is contained in the kernel of $F^{\text{proj}} \rightarrow \Gamma(A_L^{\text{lin}})$,
- (2) the kernel of $F^{\text{proj}} \rightarrow \Gamma(A_L^{\text{lin}})$ is contained in $I_L^2\mathfrak{X}^v = \mathfrak{X}^v \cap I_L^2\mathfrak{X}^{\text{proj}}$ by definition of A_L^{lin} .

This completes the proof of the second argument.

PROPOSITION 2.12. — *Let (s^k, E^k) be a Levi F -connection and an Euler-like vector field respectively, which are flat and linear up to order k , with $k > 2$. Then there exists (s^{k+1}, E^{k+1}) , a Levi F -connection and an Euler-like vector field respectively, such that*

- the pair (s^{k+1}, E^{k+1}) is flat and linear up to order $k + 1$,
- the vector fields E^k and E^{k+1} coincide up to order k ,
- the vector fields $s^k(\xi)$ and $s^{k+1}(\xi)$ coincide up to order k for all $\xi \in \Gamma(A_L^S)$.

Proof. — By assumption, the pair (s^k, E^k) satisfies two conditions: “Linearity up to order k ” and “Flatness up to order k ”, i.e., for all $\xi, \zeta \in \Gamma(A_L^S)$

$$(\text{Lin}^k) \quad [s^k(\xi), E^k] = 0 \quad \text{mod } I_L^k \mathfrak{X}^v$$

$$(\text{Flat}^k) \quad [s^k(\xi), s^k(\zeta)] = s^k([\xi, \zeta]) \quad \text{mod } I_L^k \mathfrak{X}^v.$$

We will prove the proposition in three steps:

Step 1. — We construct a class that measures the failure of (Lin^k) to hold at order $k + 1$.

– The $C^\infty(L)$ module $I_L^k \mathfrak{X}^v / (F \cap I_L^k \mathfrak{X}^v + I^{k+1} \mathfrak{X}^v)$ is projective, i.e., isomorphic to the section space $\Gamma(V^k)$ of a vector bundle $V^k \rightarrow L$. Indeed, the quotient $I_L^k \mathfrak{X}^v / I_L^{k+1} \mathfrak{X}^v$ is given by sections of some vector bundle over L : It is a direct consequence of the splitting Lemma 1.1 that V^k is a quotient of that bundle.

– A Lie algebroid action of A_L^S on V^k is defined by $\nabla_\xi(\bar{\sigma}) := \overline{[s^k(\xi), \sigma]}$, with $\sigma \in I_L^k \mathfrak{X}^v$. The action is well-defined, because $[s^k(\xi), \cdot]$ preserves \mathfrak{X}^v, I_L and F , hence it preserves the numerator and denominator of $I_L^k \mathfrak{X}^v / (F \cap I_L^k \mathfrak{X}^v + I_L^{k+1} \mathfrak{X}^v)$. As $k > 2$, Equation (Flat^k) implies that the action ∇ is flat.

– By (Lin^k) , there is a well-defined $C^\infty(L)$ -linear map $\text{def} : \Gamma(A_L^S) \rightarrow \Gamma(V^k)$ given by $\xi \mapsto \overline{[s^k(\xi), E^k]}$ describing the defect of linearity up to order $k + 1$. The curvature

$$(9) \quad c^k(\xi, \zeta) = [s^k(\xi), s^k(\zeta)] - s^k([\xi, \zeta])$$

is valued in F and in $I_L^k \mathfrak{X}^v$, by assumption (Flat^k) . The first item in Lemma 2.10 implies that applying $[\cdot, E^k]$ to a vector field in $F \cap I_L^k \mathfrak{X}^v$ yields an element of $F \cap I_L^k \mathfrak{X}^v + I^{k+1} \mathfrak{X}^v$: Therefore upon applying $[\cdot, E^k]$ to Equation (9), we obtain

$$[s^k(\xi), [s^k(\zeta), E^k]] - [s^k(\zeta), [s^k(\xi), E^k]] = [s^k([\xi, \zeta]), E^k] \quad \text{mod } F \cap I_L^k \mathfrak{X}^v + I^{k+1} \mathfrak{X}^v,$$

which is exactly the cocycle condition: $\nabla_\xi \text{def}(\zeta) - \nabla_\zeta \text{def}(\xi) - \text{def}([\xi, \zeta])$.

Step 2. — We construct (E^{k+1}, s^{k+1}) satisfying (Lin^{k+1}) .

– Since L is simply connected, Lemma 2.1 implies that the class $[\text{def}] \in H^1(A_L^S, V^k)$ is zero. Choose $\varepsilon^k \in I_L^k \mathfrak{X}^v$, such that $\overline{\varepsilon^k}$ is a primitive of def :

$$\text{def}(\xi) = \nabla_\xi \varepsilon^k \quad \text{i.e., } [s^k(\xi), E^k] = [s^k(\xi), \varepsilon^k] \quad \text{mod } F \cap I_L^k \mathfrak{X}^v + I_L^{k+1} \mathfrak{X}^v.$$

We define the new Euler-like vector field by $E^{k+1} = E^k - \varepsilon^k$.

– By construction of E^{k+1} , for every given $\xi \in \Gamma(A_L^S)$, there exists a vector field $\sigma^k(\xi)$ in $F \cap I_L^k \mathfrak{X}^v$ such that

$$[s^k(\xi), E^{k+1}] = \sigma^k(\xi) \quad \text{mod } I_L^{k+1} \mathfrak{X}^v.$$

Using local trivializations and partitions of unity on L , the map $\xi \mapsto \sigma^k(\xi)$ can be achieved to be $C^\infty(L)$ -linear. We now define $s^{k+1} = s^k - \sigma^k / (k - 1)$. By construction, s^{k+1} is still a Levi F -connection.

– Let us verify (Lin^{k+1}) :

$$\begin{aligned} [s^{k+1}(\xi), E^{k+1}] &= \left[s^k(\xi) - \frac{\sigma^k(\xi)}{k-1}, E^{k+1} \right] \\ &= \sigma^k(\xi) - \frac{1}{k-1} [\sigma^k(\xi), E^{k+1}] = 0 \pmod{I_L^{k+1} \mathfrak{X}^v}. \end{aligned}$$

The last equality holds by the first item of Lemma 2.10.

Step 3. — Consider the curvature

$$c^{k+1}(\xi, \zeta) = [s^{k+1}(\xi), s^{k+1}(\zeta)] - s^{k+1}([\xi, \zeta]).$$

We show that (Lin^{k+1}) implies (Flat^{k+1}) , i.e., $c^{k+1} = 0$ modulo $I_L^{k+1} \mathfrak{X}^v$.

– Since $s^{k+1} = s^k$ modulo $I_L^k \mathfrak{X}^v$, we know that

$$c^{k+1} = c^k = 0 \pmod{I_L^k \mathfrak{X}^v}.$$

– In view of (Lin^{k+1}) , all underbraced terms in the following expression are in $I_L^{k+1} \mathfrak{X}^v$.

$$\begin{aligned} [c^{k+1}(\xi, \zeta), E^{k+1}] &= [s^{k+1}(\xi), \underbrace{[s^{k+1}(\zeta), E^{k+1}]}] - [s^{k+1}(\zeta), \underbrace{[s^{k+1}(\xi), E^{k+1}]}] - \underbrace{[s^{k+1}([\xi, \zeta]), E^{k+1}]}]. \end{aligned}$$

Since $s^{k+1}(\xi)$ and $s^{k+1}(\zeta)$ are projectable vector fields tangent to L , their bracket with $I_L^{k+1} \mathfrak{X}^v$ takes values in $I_L^{k+1} \mathfrak{X}^v$. Hence, $[c^{k+1}, E^{k+1}] = 0 \pmod{I_L^{k+1} \mathfrak{X}^v}$.

– The second item of Lemma 2.10, implies that $c^{k+1} = 0 \pmod{I_L^{k+1} \mathfrak{X}^v}$, i.e., (Flat^{k+1}) holds.

This completes the proof.

2.4. EXAMPLES AND COUNTER-EXAMPLES. — Let us give counter examples to naive generalizations of Theorem 2.8. Let us explore the non-simply connected case.

EXAMPLE 2.13. — For L a leaf in a regular foliation, we have

$$A_L = A_L^{\text{lin}} = A_L^{\text{S}} = TL,$$

and every tubular neighborhood (U, π) induces a unique flat Levi F -connection: it suffices to lift a vector field in L to the unique π -projectable vector field in F . However, the transverse formal Euler-like vector field can only exist if the holonomy $\Phi(\gamma)$ is a formally linearizable diffeomorphism of the transversal for all $\gamma \in \pi_1(L)$. The regular foliations (with dimension 1 leaves) obtained by suspension of diffeomorphism $\phi: \mathbb{R}^n \rightarrow \mathbb{R}^n$ fixing 0 are instances of such foliations with $L \simeq S^1$ if ϕ is not formally linearizable at zero (e.g. $n = 2$ and $(x, y) \mapsto (x, y + x^2)$).

EXAMPLE 2.14. — Consider the “self-eating snake” singular foliation, as in Figure 1, realized as follows. Let S be the “foliation by concentric circles”, i.e., the singular foliation on \mathbb{R}^2 of all vector fields X such that $X[\phi] = 0$, with $\phi = \sum_{i=1}^2 x_i^2$. Then consider the direct product singular foliations on $\mathbb{R}^2 \times \mathbb{R}$ given by $F := S \times \mathfrak{X}(\mathbb{R})$. This

foliation goes to the quotient through the equivalence relation $(x, t) \sim (\frac{1}{2}x, t + 1)$, for all $(x, t) \in \mathbb{R}^2 \times \mathbb{R}$. The only singular leaf of the quotient singular foliation is $L = S^1$.

In this case, the normal bundle ν is trivial as a vector bundle, a flat Levi F -connection exists (which is also a flat Ehresmann connection since $A_L^S = TL$), but there is no Ehresmann F -connection that makes the normal bundle isomorphic to the trivial one (i.e., the first return map on ν induced by any Ehresmann F -connection is non-trivial).




FIGURE 1. Example 2.14

Here is an example of a leaf for which a section $A_L^S \rightarrow A_L^{lin}$ exists (because it is transversally quadratic and $A_L^S = A_L^{lin}$) but does not admit a flat Levi F -connection.

EXAMPLE 2.15. — Let L be a manifold, $\alpha, \beta \in \Omega^1(L)$ closed 1-forms such that the class of $\alpha \wedge \beta$ in $H^2(L)$ is not trivial. On $M := L \times \mathbb{R}$, we consider for all $a, b, c \in \mathbb{N}$ with $2 \leq a < b$ and $c = a + b - 1$ the $C^\infty(M)$ -submodule $F \subset \mathfrak{X}(M)$ generated by the vector fields:

$$t^c \frac{\partial}{\partial t} \quad \text{and} \quad \psi(u) := u + \alpha(u)t^a \frac{\partial}{\partial t} + \beta(u)t^b \frac{\partial}{\partial t} \quad \text{with } u \in \mathfrak{X}(L).$$

A direct computation shows that:

$$\begin{aligned} \left[\psi(u), t^c \frac{\partial}{\partial t} \right] &= ((c - a)\alpha(u)t^{a-1} + (c - b)\beta(u)t^{b-1}) t^c \frac{\partial}{\partial t} \\ \psi([u, v]) - [\psi(u), \psi(v)] &= (b - a) (\alpha(u)\beta(v) - \beta(u)\alpha(v)) t^c \frac{\partial}{\partial t}, \end{aligned}$$

so that F is a singular foliation.

By construction, $L \times \{0\}$ is a leaf of F and I_L is then the ideal generated by t . The computations above imply that the holonomy Lie algebroid is the vector bundle $A_L = TL \oplus \mathbb{R}$ equipped with the bracket given for all $u, v \in \mathfrak{X}(L), f, g \in C^\infty(L)$ by:

$$[(u, f), (v, g)] = ([u, v], u[g] - v[f] + \omega(u, v)),$$

where $\omega = (b - a) \alpha \wedge \beta \in \Omega^2(L)$. The projection on TL is the anchor map. Any Ehresmann F -connection is of the form:

$$u \longmapsto \psi(u) + \gamma_t(u)t^c \frac{\partial}{\partial t}$$

for some t -dependent 1-form γ on L . Its curvature is $(\omega + d_{\text{dR}}\gamma_0)t^c\partial/\partial t + O(t^{c+1})$. Since the 2-form ω is not exact, the leaf L does not admit a flat Ehresmann F -connection.

The following is an example of a possibly simply connected leaf, which does not admit a section $A_L^{\text{S}} \rightarrow A_L^{\text{lin}}$ and therefore does not admit a flat Levi F -connection.

EXAMPLE 2.16. — Let L be a manifold and let $\omega \in H^2(L, \mathbb{Z})$ be the Chern class of an S^1 -bundle $P \rightarrow L$ with connection θ . Consider the associated \mathbb{C} -bundle $\pi: E \rightarrow L$, and equip it with the linear Ehresmann connection associated to θ . The horizontal lifts \widehat{u} of vector fields u on L , together with the infinitesimal vector field R of the S^1 -action generate a singular foliation F on E . For this foliation F the zero section L is a leaf. In view of the relation:

$$[\widehat{u} + \pi^*(f) \cdot R, \widehat{v} + \pi^*(g) \cdot R] = [\widehat{u}, \widehat{v}] + \pi^*(u[g] - v[f] + \omega(u, v)) \cdot R$$

for all $u, v \in \mathfrak{X}(L), f, g \in C^\infty(L)$,

the holonomy Lie algebroid of L is $A_L = TL \oplus \mathbb{R}$, its anchor map is the projection onto TL , and its Lie bracket is given for all $u, v \in \mathfrak{X}(L), f, g \in C^\infty(L)$ by

$$[(u, f), (v, g)] = ([u, v], u[g] - v[f] + \omega(u, v)).$$

As ω is nonzero in cohomology, there can be no Lie algebroid section from $TL = A_L^{\text{S}}$ to $A_L = A_L^{\text{lin}}$.

Let us construct examples for which Theorem 2.8 holds. Let $\tau: \Gamma(A) \rightarrow \mathfrak{X}(V)$ be the action of a Lie algebroid $A \rightarrow L$ on the vector bundle $V \rightarrow L$ and $R \subset \mathfrak{X}(V)$ a singular foliation made of vertical vector fields. If

- (1) R is invariant under the Lie algebroid action,
- (2) $\tau(\Gamma(A))$ intersects R trivially,

then vector fields in R , together with the vector fields for the infinitesimal A -action on V , generate a singular foliation F on V .

The same construction can be completed when R is substituted by a formal singular foliation \widehat{R} along the zero section L . Let us fix notations:

DEFINITION 2.17. — Let $A \rightarrow L$ be a Lie algebroid acting faithfully on V . For every (maybe formal) singular foliation R satisfying the above conditions (1) and (2), then the above singular foliation F is called the *semi-direct product* of A with R and is denoted⁽²⁾ by

$$F := A \widehat{\cap} R.$$

In all cases we are interested in, $A \rightarrow L$ is a transitive Lie algebroid. Also, R will be made of vertical vector fields vanishing along the zero section L . Under these conditions, the zero section L is a leaf of F . Let us give two examples.

⁽²⁾Notice that it is not true that $F^{\text{proj}} \simeq \Gamma(A) \cap R$ as Lie algebras (there is only an inclusion).

EXAMPLE 2.18. — Consider a singular foliation \mathcal{R} on \mathbb{R}^n . The *direct product* of L with \mathcal{R} is obtained by choosing, in Definition 2.17, A to be the Lie algebroid $TL \rightarrow L$ and V to be the trivial TL -module $\mathbb{R}^n \times L$.

EXAMPLE 2.19. — Let $n > 3$. Since $SO(n + 1)$ acts on the sphere S^n , there is a natural action of the transformation Lie algebroid $A = \mathfrak{so}(n + 1) \times S^n \rightarrow S^n$ on $V = TS^n \rightarrow S^n$. Let \mathcal{R} be the singular foliation on TS^n generated by the Euler vector field. The assumptions in Definition 2.17 are easily satisfied. The semi-direct product $A \widehat{\cap} \mathcal{R}$ is a singular foliation on TS^n , admitting the zero section $L = S^n$ as a leaf, with $A_L^{\mathcal{S}} = A = \mathfrak{so}(n + 1) \times L \rightarrow L$. For this leaf, the Euler field and the Lie algebroid action of A on TS^n give the formal Euler-like field and the formal Levi F -connection whose existence is guaranteed by Theorem 2.8.

For $n = 2$, the construction of the singular foliation F still makes sense, and the zero section $L = S^2$ is still a leaf. However, since A_L^{lin} is the transitive transformation Lie algebroid $(\mathfrak{so}(3) \oplus \mathbb{R}) \times L \rightarrow L$, with \mathbb{R} acting trivially, and since the isotropy of the Lie algebroid A_L^{lin} at every point in S^2 is an Abelian two-dimensional Lie algebra, we now have $A_L^{\mathcal{S}} = TS^2 \rightarrow S^2$. Since there is no Lie algebroid section $TL \rightarrow (\mathfrak{so}(3) \oplus \mathbb{R} \times L) \rightarrow L$ for $L = S^2$, Theorem 2.8 does not apply.

2.5. GEOMETRIC REFORMULATION. — Let L be a locally closed leaf of F . Assume both conditions in Theorem 2.8 are satisfied: L is simply connected and a section $z : A_L^{\mathcal{S}} \rightarrow A_L^{\text{lin}}$ exists. Theorem 2.8 then provides:

- ♠ a formal Euler-like vector field E , tangent to the fibers of $\pi : U \rightarrow L$,
- ◇ a formal Levi F -connection $s : \Gamma(A_L^{\mathcal{S}}) \rightarrow \widehat{F}^{\text{proj}}$.

Moreover, the image of s is made of vector fields commuting with E . Let us spell out the content of this data:

♠ The formal Euler-like vector field E yields a formal isomorphism Φ between the fibers of normal bundle $\pi : \nu = TM|_L/TL \rightarrow L$ to the the fibers of $\pi : U \rightarrow L$ that identifies, by construction, E with the Euler vector field E_ν of the normal bundle. We use Φ to transport to ν the formal jet \widehat{F} of F .

◇ The composition $\Phi^{-1} \circ s$ now becomes a flat Levi $\Phi^{-1}(\widehat{F})$ -connection on the fibers of $\pi : \nu \rightarrow L$.

Moreover, the image of $\Phi^{-1} \circ s$ is made of vector fields commuting with E_ν , i.e., linear vector fields on ν , so that the flat Levi F -connection of item ◇ is now a Lie algebroid action of $A_L^{\mathcal{S}}$ on the normal bundle. This proves the following Lemma.

LEMMA 2.20. — *The image of $\Phi^{-1} \circ s$ is made of linear vector fields on $\nu \rightarrow L$. More precisely, for every $\xi \in \Gamma(A_L^{\mathcal{S}})$, the linear vector field on ν describing the natural Lie algebroid action of $z(\xi) \in \Gamma(A_L^{\text{lin}})$ on ν coincides with $\Phi^{-1} \circ s(\xi)$.*

We call *radical foliation of F* the subspace $R \subset F^\vee$ of all vector fields in F^\vee whose image through the linearization map along L is in the radical of A_L^{lin} . In view

of the definition of $A_L^{\mathcal{S}}$, it can be defined by:

$$R := \text{Ker}(F^{\text{proj}} \longrightarrow \Gamma(A_L^{\mathcal{S}})).$$

LEMMA 2.21. — *The space R is a singular foliation on U , included in F^{\vee} , and*

$$[s(\Gamma(A_L^{\mathcal{S}})), \widehat{R}] \subset \widehat{R} \quad \text{and} \quad s(\Gamma(A_L^{\mathcal{S}})) \oplus \widehat{R} = \widehat{F}^{\text{proj}}.$$

Proof. — The formal jet \widehat{R} of R along L is the kernel of $\widehat{F}^{\text{proj}} \rightarrow \Gamma(A_L^{\mathcal{S}})$, and s is a Lie algebra section of that projection.

It follows from Lemma 2.21 that $\widehat{F}^{\text{proj}}$ is, as a Lie algebra, isomorphic to the semi-direct product:

$$\widehat{F}^{\text{proj}} \simeq s(\Gamma(A_L^{\mathcal{S}})) \ltimes \widehat{R}.$$

Using the formal diffeomorphism Φ^{-1} , we see that \widehat{F} is indeed a singular foliation of the form described in Definition 2.17 applied to $A = A_L^{\mathcal{S}} \rightarrow L, \nu = V$ and $\widehat{R}_{\nu} = \Phi^{-1}(\widehat{R})$. Using this language Theorem 2.8 takes the following form:

THEOREM 2.22. — *Let L be a locally closed leaf of the locally real analytic foliation F on a manifold M . If $\pi_1(L) = 0$, and there exists a Lie algebroid section z from the semi-simple holonomy $A_L^{\mathcal{S}}$ to the linear holonomy A_L^{lin} , then:*

(1) *the normal bundle $\nu = TM|_L/TL \rightarrow L$ comes equipped with a flat $A_L^{\mathcal{S}}$ -connection,*

(2) *there is a formal diffeomorphism between M and ν (near L) that identifies \widehat{F} and a semi-direct product⁽³⁾ singular foliation on $\nu \rightarrow L$ of the form:*

$$\widehat{F} = A_L^{\mathcal{S}} \widehat{\ltimes} \widehat{R}_{\nu},$$

where \widehat{R}_{ν} is a vertical singular foliation on ν , tangent to the fibers of $\nu \rightarrow L$, invariant under the $A_L^{\mathcal{S}}$ -action on ν , isomorphic to the formal jet of the radical foliation R of F .

REMARK 2.23. — The decomposition $\widehat{F} = A_L^{\mathcal{S}} \widehat{\ltimes} \widehat{R}_{\nu}$ should not confuse the reader. Vector fields arising from the infinitesimal action of sections $\Gamma(A_L^{\mathcal{S}})$ of the Lie algebroid $A_L^{\mathcal{S}}$ on ν are in direct sum with \widehat{R}_{ν} . But the $C^{\infty}(\nu)$ -module generated by this infinitesimal action is a singular foliation that does in general intersect \widehat{R}_{ν} . The corollary below gives a context where this module contains \widehat{R}_{ν} .

COROLLARY 2.24. — *Let L be a locally closed leaf of the locally real analytic foliation F on a manifold M . If $\pi_1(L) = 0$, and $A_L^{\mathcal{S}} = A_L$, then there is a formal diffeomorphism between F and the singular foliation associated to the natural Lie algebroid action of the holonomy Lie algebroid A_L on the normal bundle.*

⁽³⁾For the notation $A_L^{\mathcal{S}} \widehat{\ltimes} \widehat{R}_{\nu}$, see Definition 2.17.

Proof. — According to [AS09, Prop. 1.5], in a neighborhood of $p \in L$, the singular foliation F is generated by any family X_1, \dots, X_d of vector fields in F whose image in $\Gamma(A_L)$ is a local trivialization of A_L . As a consequence, the image of s generates F . The result then follows from Theorem 2.22.

3. LOCAL AND SEMI-LOCAL STRUCTURE OF A SINGULAR FOLIATION

3.1. LOCAL STRUCTURE OF A SINGULAR FOLIATION: LEVI THEOREMS. — Let us explore the consequences of Theorem 2.8 in the neighborhood of a point p in a manifold M equipped with a singular foliation F . Splitting Lemma 1.1 allows to make the additional assumption that all vector fields vanish at p , upon replacing M with a small disk transversal to the leaf through m if necessary.

Throughout Section 3.1, F shall be a locally real analytic singular foliation made of vector fields that vanish at a point $p \in M$.

3.1.1. Relation with Cerveau's Levi theorems. — The requirements of Theorem 2.8 (namely “If $\pi_1(L) = 0$, and there exists a Lie algebroid section z from the semi-simple holonomy A_L^S to the linear holonomy A_L^{lin} ”) hold automatically:

- (1) Since L is reduced to the point $\{p\}$, it is simply connected.
- (2) The Lie algebroids $A_L, A_L^{\text{lin}}, A_L^S$ are finite dimensional Lie algebras:
 - (a) A_L is the isotropy Lie algebra \mathfrak{g}_p at p ,
 - (b) A_L^{lin} is the quotient $\mathfrak{g}_p/\mathfrak{g}_p^{>2}$,
 - (c) A_L^S is the semi-simple part \mathfrak{g}_p^S of the Lie algebra \mathfrak{g}_p .

Now, in view of the usual Levi-Malcev decomposition theorem for finite dimensional Lie algebras, a Lie algebra section $\mathfrak{g}_p^S \rightarrow \mathfrak{g}_p$ exists. Its composition with the natural projection $\mathfrak{g}_p \rightarrow \mathfrak{g}_p^{\text{lin}}$ is a Lie algebra section $\mathfrak{g}_p^S \rightarrow \mathfrak{g}_p^{\text{lin}}$.

Theorem 2.8 specializes therefore to yield the following corollary:

COROLLARY 3.1 (Cerveau). — *Let \mathfrak{g}^S be the semi-simple part of the isotropy Lie algebra of F at \mathfrak{g} . Then there exists a Lie algebra morphism $s : \mathfrak{g}^S \rightarrow \widehat{F}_p$ and a formal Euler-like vector field E with respect to which the image of s is made of formally linear vector fields.*

A comparison of this Corollary with [Cer79, Th. 2.1] shows that both statements are equivalent (although stated and proved quite differently here). Also, for $L = \{p\}$, Corollary 2.24 recovers the second part of [Cer79, Th. 2.2].

3.1.2. Levi theorem for projective foliations. — Let us assume that F is a projective $C^\infty(M)$ -module.⁽⁴⁾ In this case, there exists a Lie algebroid $(A, [\cdot, \cdot], \rho)$, such that the anchor map $\rho : A \rightarrow \coprod_{m \in M} T_m F$, although it is not an isomorphism at every point, is an isomorphism (of $C^\infty(M)$ -modules) at the level of sections:

$$\rho : \Gamma(A) \cong F .$$

⁽⁴⁾i.e., “a Debord foliation” in the terminology of [LGLS20], see [Deb01].

Since all vector fields in F vanish at p , we have that $\rho|_p = 0$, so that the fiber of A_p is a Lie algebra: it is easily shown to coincide with the isotropy Lie algebra \mathfrak{g}_p . Applying Corollary 3.1 to this situation yields the following result, where $\widehat{\Gamma}(A)$ stands for formal sections of a vector bundle A near p :

COROLLARY 3.2 ([Wei00, Duf01, Zun03]). — *Let A be the Lie algebroid associated to a projective singular foliation made of vector fields vanishing at $\{p\}$. Denote by A_p^S the semi-simple part of the isotropy Lie algebra A_p . Then there exists a Lie algebra morphism $s : A_p^S \rightarrow \widehat{\Gamma}(A)$ and a formal Euler-like vector field E with respect to which the image of $\rho \circ s$ is made of formally linear vector fields.*

This statement indeed holds true for any Lie algebroid, see [Wei00, Duf01, Zun03].

3.2. SECTIONS TO THE HOLONOMY LIE (∞) -ALGEBROID. — Let F be a locally real analytic singular foliation. For every leaf L such that $\pi_1(L) = \pi_2(L) = 0$, Corollary 2.5 ensures the existence of a Lie algebroid section $A_L^S \rightarrow A_L$. Using Theorem 2.8, we can loosen the 2-connectedness condition for M as follows:

PROPOSITION 3.3. — *Let F be a locally real analytic singular foliation and L a simply connected and locally closed leaf, such that there exists a Lie algebroid section $A_L^S \rightarrow A_L^{\text{lin}}$. Then there exists a Lie algebroid section $A_L^S \rightarrow A_L$.*

Proof. — Let c be the Artin-Rees bound for F at L . By “stopping early” in the iteration for Theorem 2.8, we obtain a section $s = s^{c+1} : \Gamma(A_L^S) \rightarrow F$ and an Euler-like vector field $E = E^{c+1}$ such that $[s(\xi), s(\zeta)] - s([\xi, \zeta]) \in I_L^{c+1} \mathfrak{X}^v \cap F \subset I_L F$ for all $\xi, \zeta \in \Gamma(A_L^S)$. Such a map s induces a section $A_L^S \rightarrow A_L$ which is a Lie algebroid section.

Proposition 3.3 can be generalized as follows. For the sake of simplicity, we will assume below that the formal A_L^S -action in Theorem 2.8 is convergent, and that the leaf L is compact, so that we may refer to existing results in [LGLS20] and [LGR19]. These additional assumptions are certainly not relevant for both propositions below, but avoiding them would require to extend to the formal setting the statements we will refer to. In [LGLS20], it is shown that every real analytic singular foliation F is, locally on a neighborhood U of a point, the image through the anchor map of a *universal Lie ∞ -algebroid*, i.e., a Lie ∞ -algebroid $U^F = (E_{-i}, [\cdot \cdot \cdot]_i, \rho)$ whose 1-ary bracket $d = [\cdot]_1$, together with its anchor map:

$$\dots \xrightarrow{d} \Gamma(E_{-2}) \xrightarrow{d} \Gamma(E_{-1}) \xrightarrow{\rho} F|_U$$

form a projective resolution of F . In [LGR19, Th. 2.26], the universal Lie ∞ -algebroid is shown to exist in a neighborhood of a compact leaf. The restriction of such a Lie ∞ -algebroid U^F to L yields a transitive Lie ∞ -algebroid over L denoted by $U^F|_L$. It admits a canonical Lie ∞ -morphism onto A_L . We call Π its composition with the projection $A_L \rightarrow A_L^S$.

PROPOSITION 3.4. — *Let F be a locally real analytic singular foliation and L a simply connected and compact leaf, such that there exists a Lie algebroid section $A_L^S \rightarrow A_L^{\text{lin}}$. We assume that the formal section $\Gamma(A_L^S) \rightarrow F$ whose existence is guaranteed by Theorem 2.8 can be chosen to converge in a neighborhood of L . Then Π admits a Lie ∞ -algebroid section $A_L^S \rightarrow \mathcal{U}^F|_L$.*

Proof. — The Lie algebroid action of A_L^S on U defines a sub-foliation A_L^S in F , namely the image through the anchor map of the transformation Lie algebroid of this action. In view of [LGLS20, Th. 2.9], there exists a Lie ∞ -algebroid morphism Φ from this transformation Lie algebroid to the universal Lie ∞ -algebroid \mathcal{U}^F . The desired morphism is the restriction of Φ to the leaf L .

An important question for a given singular foliation is to know whether or not it comes from a Lie algebroid action [AZ13]. When the leaf L is a point $\{p\}$, the rank of such a Lie algebroid has to be greater than or equal to the dimension of the isotropy Lie algebra \mathfrak{g}_p . Although the general problem remains open, \mathfrak{g}_p carries a Chevalley-Eilenberg cohomology 3-class, called the NMRLA-class that obstructs the possibility to have a Lie algebroid whose rank is minimal i.e., equal to $\dim(\mathfrak{g}_p)$ (see [LGLS20, Prop. 4.29]). Proposition 3.4 has strong implications for this class: it shows that it is effaçable.

Recall that for \mathfrak{g} a Lie algebra and V a finite dimensional \mathfrak{g} -module, a class ω in a Chevalley-Eilenberg cohomology group $H^k(\mathfrak{g}, V)$ is *effaçable* (or erasable) if there exists a finite dimensional \mathfrak{g} -module W containing V such that the image of ω in $H^k(\mathfrak{g}, W)$ is zero.

Let us briefly describe the NMRLA class assuming $L = \{p\}$ is a point leaf. In this case, $\mathcal{U}^F|_{\{p\}}$ is a Lie ∞ -algebra whose 1-ary bracket can be assumed to be zero. Then, its degree (-1) component is a Lie algebra isomorphic to \mathfrak{g}_p (see [LGLS20, Prop. 4.14]), its degree (-2) -component is a \mathfrak{g}_p -module V , and the restriction to \mathfrak{g}_p of the 3-ary bracket is a Chevalley-Eilenberg 3-cocycle valued in V (see [LGLS20, Prop. 4.27]), defining the NMRLA class.

PROPOSITION 3.5. — *Let F be a locally real analytic singular foliation and $\{p\}$ a point leaf such that the formal section $\mathfrak{g}_p^S \rightarrow F$ whose existence is guaranteed by Corollary 3.1 can be chosen to converge in a neighborhood of p . Then the NMRLA-class of F at p is effaçable.*

Proof. — In view of [Hoc54, Th. 1], a cohomology class is effaçable if and only if its restriction to a maximal semi-simple Lie subalgebra is zero. Let $\Phi: \mathfrak{g}_p^S \rightarrow \mathcal{U}^F|_p$ be a Lie ∞ -algebroid morphism as in Proposition 3.4. The Taylor coefficient $\Phi_1: \mathfrak{g}_p^S \rightarrow \mathfrak{g}_p^S$ of Φ is the identity map and the second Taylor coefficient $\Phi_2: \wedge^2 \mathfrak{g}_p^S \rightarrow V$ satisfies (see [LGLS20, Eq. (4.10)]) for all $a, b, c \in \mathfrak{g}_p^S$,

$$\{a, b, c\}_3 = \{a, \Phi_2(b, c)\}_2 - \Phi_2(\{a, b\}_2, c) + \text{abc}.$$

This means that the restriction of the 3-ary bracket to \mathfrak{g}_p^S is a Chevalley-Eilenberg coboundary. This concludes the proof.

3.3. TRANSVERSALLY QUADRATIC SIMPLY CONNECTED LEAVES. — Let F be a locally real analytic foliation and L a leaf. We say that a leaf L is *transversally quadratic* if its transverse singular foliation (see Lemma 1.1) is made of vector fields vanishing at least quadratically. There is an easy characterization in terms of the holonomy Lie algebroid of the leaf L :

PROPOSITION 3.6. — *A leaf L is transversally quadratic if and only if $A_L^{\text{lin}} = A_L^{\text{S}} = TL$. In particular, the normal bundle ν carries a natural flat connection ∇^ν .*

Proof. — By definition of A_L^{lin} , the first part of the proposition follows from the following intermediate characterization of transversally quadratic leaves. A leaf is transversally quadratic if $F^\vee \subset I_L^2 \mathfrak{X}^\vee$. The second part of the proposition follows from the existence, for every leaf L , of a natural A_L^{lin} -action on ν , see Section 1.1.

REMARK 3.7. — It follows immediately from Proposition 3.6 that a leaf $L \subset M$ whose normal bundle is not flat cannot be transversally quadratic, which is a very strong constraint. For instance, $S^2 \subset TS^2$ cannot be transversally quadratic.

For a regular foliation, it is well-known that in a neighborhood of a simply connected leaf L , the foliation is “trivial”, i.e., formally, it is isomorphic to the direct product of the leaf L with an open disk. The same phenomenon occurs for transversally quadratic leaves:

THEOREM 3.8. — *Every simply-connected, transversally quadratic and locally closed leaf L of a locally real analytic singular foliation F is formally trivial, i.e., the formal jet \widehat{F} along L is isomorphic to the direct product⁽⁵⁾ of L with the formal jet of the transverse foliation.*

Proof. — Both conditions in Theorem 2.22 are satisfied: L is simply connected by assumption and a section $A_L^{\text{S}} \rightarrow A_L^{\text{lin}}$ exists since both algebroids coincide with TL by Proposition 3.6. There is therefore a formal isomorphism between \widehat{F} and $TL\widehat{\cap}\widehat{R}$, with \widehat{R} being the radical foliation. In this case, however, there are several obvious identifications:

(1) The radical foliation \widehat{R} of \widehat{F} is simply the transverse singular foliation.

(2) By Proposition 3.6, the normal bundle ν is flat. Since L is simply connected, it is indeed a trivial vector bundle: $\nu \simeq L \times \nu_p$, with ν_p some given fiber.

The semi-direct product is then reduced to a direct product. This gives the desired formal isomorphism.

REMARK 3.9. — Theorem 3.8 is a purely singular foliation phenomenon: there is no such a result for Lie algebroids or Poisson structures. In fact, even for regular Poisson or Lie algebroid structures there is no such a result. For instance, choose of a volume form ω on the 2-sphere S^2 , let $\pi = \omega^{-1}$ be its inverse Poisson structure and consider the Poisson structure on $S^2 \times \mathbb{R}$ given by $e^t \pi \oplus 0$ with t the parameter on \mathbb{R} . The

⁽⁵⁾Direct products of L with a singular foliation are discussed in Example 2.18.

symplectic leaves are the fibers of the projection $S^2 \times \mathbb{R} \rightarrow \mathbb{R}$. They are therefore simply-connected and their transverse Poisson structure is zero (in particular, it is transversally quadratic: it vanishes at order at least 2). But since the volumes of all the symplectic leaves are different, this Poisson structure cannot be isomorphic to a direct product of $\pi = \omega^{-1}$ with the trivial Poisson structure on \mathbb{R} in a neighborhood of a given leaf (even formally).

Similarly, consider sections of the vector bundle $A = T(S^2 \oplus \mathbb{R})$ over the manifold $S^2 \times \mathbb{R}$ as pairs (X, f) or (Y, g) with X, Y being t -dependent vector fields tangent to S^2 and f, g t -dependent real-valued functions on S^2 (with t the parameter along \mathbb{R}). The bracket:

$$[(X, f), (Y, g)] = ([X, Y], X[g] - Y[f] + t\omega(X, Y))$$

is a Lie algebroid bracket on A . The leaves of A are 2-spheres: they are therefore simply connected. The transverse Lie algebroid $T\mathbb{R} \rightarrow \mathbb{R}$ has trivial anchor and trivial bracket. The restriction of the Lie algebroid A to any two leaves are isomorphic, except for the exceptional leaf $t = 0$. Hence the Lie algebroid A is not a direct product near the leaf $t = 0$.

By applying the “stopping early” strategy from Proposition 3.3 in the proof of the previous Theorem, we obtain the following result:

COROLLARY 3.10. — *The holonomy Lie algebroid A_L of a simply connected, transversally quadratic and locally closed leaf L is the direct sum of TL with the isotropy Lie algebra of its transverse foliation.*

Proof. — By Proposition 3.3, a Lie algebroid section $\underline{s} : \mathfrak{X}(L) \rightarrow F^\vee / I_L F^\vee \simeq \Gamma(A_L)$ exists. This section makes the isotropy Lie algebra bundle $\ker(\rho)$ of A_L a flat Lie algebra bundle. Since L is simply connected, it is a trivial Lie algebra bundle.

The proof of Theorem 3.8 in fact shows the following more general statement:

THEOREM 3.11. — *A simply-connected and locally closed leaf L of a locally real analytic singular foliation F is formally trivial (i.e., the formal jet \widehat{F} along L is isomorphic to the direct product of L with the formal jet of the transverse foliation) if and only if there exists a Lie algebroid section $TL \rightarrow A_L^{\text{lin}}$.*

REFERENCES

- [And17] I. ANDROULIDAKIS – Personal communication, 2017.
- [AS09] I. ANDROULIDAKIS & G. SKANDALIS – “The holonomy groupoid of a singular foliation”, *J. reine angew. Math.* **626** (2009), p. 1–37.
- [AZ13] I. ANDROULIDAKIS & M. ZAMBON – “Smoothness of holonomy covers for singular foliations and essential isotropy”, *Math. Z.* **275** (2013), no. 3-4, p. 921–951.
- [AZ14] ———, “Holonomy transformations for singular foliations”, *Adv. Math.* **256** (2014), p. 348–397.
- [AM69] M. F. ATIYAH & I. G. MACDONALD – *Introduction to commutative algebra*, Addison-Wesley Publishing Co., Reading, MA, 1969.
- [BLM19] H. BURSZTYN, H. LIMA & E. MEINRENKEN – “Splitting theorems for Poisson and related structures”, *J. reine angew. Math.* **754** (2019), p. 281–312.

- [Cer79] D. CERVEAU – “Distributions involutives singulières”, *Ann. Inst. Fourier (Grenoble)* **29** (1979), no. 3, p. 261–294.
- [Daz85] P. DAZORD – “Feuilletages à singularités”, *Indag. Math.* **47** (1985), p. 21–39.
- [Deb01] C. DEBORD – “Holonomy groupoids of singular foliations”, *J. Differential Geom.* **58** (2001), no. 3, p. 467–500.
- [Duf01] J.-P. DUFOUR – “Normal forms for Lie algebroids”, in *Lie algebroids and related topics in differential geometry (Warsaw, 2000)*, Banach Center Publ., vol. 54, Polish Acad. Sci. Inst. Math., Warsaw, 2001, p. 35–41.
- [GG20] K. GRABOWSKA & J. GRABOWSKI – “Solvable Lie algebras of vector fields and a Lie’s conjecture”, *SIGMA Symmetry Integrability Geom. Methods Appl.* **16** (2020), article no. 065 (14 pages).
- [Her63] R. HERMANN – “On the accessibility problem in control theory”, in *Internat. Sympos. Nonlinear Differential Equations and Nonlinear Mechanics*, Academic Press, New York, 1963, p. 325–332.
- [Hoc54] G. HOCHSCHILD – “Cohomology classes of finite type and finite dimensional kernels for Lie algebras”, *Amer. J. Math.* **76** (1954), p. 763–778.
- [LGLS20] C. LAURENT-GENGOUX, S. LAVAU & T. STROBL – “The universal Lie ∞ -algebroid of a singular foliation”, *Doc. Math.* **25** (2020), p. 1571–1652.
- [LGR19] C. LAURENT-GENGOUX & L. RYVKIN – “The holonomy of a singular leaf”, 2019, [arXiv: 1912.05286](https://arxiv.org/abs/1912.05286).
- [Mac87] K. MACKENZIE – *Lie groupoids and Lie algebroids in differential geometry*, London Math. Society Lect. Note Series, vol. 124, Cambridge University Press, Cambridge, 1987.
- [Mac05] ———, *General theory of Lie groupoids and Lie algebroids*, London Math. Society Lect. Note Series, vol. 213, Cambridge University Press, Cambridge, 2005.
- [Mal67] B. MALGRANGE – *Ideals of differentiable functions*, TIFR Studies in Math., vol. 3, Tata Institute of Fundamental Research/Oxford University Press, Bombay/London, 1967.
- [MZ04] P. MONNIER & N. T. ZUNG – “Levi decomposition for smooth Poisson structures”, *J. Differential Geom.* **68** (2004), no. 2, p. 347–395.
- [Ree52] G. REEB – *Sur certaines propriétés topologiques des variétés feuilletées*, Actualités Sci. Ind., vol. 1183, Hermann & Cie., Paris, 1952.
- [Tou68] J.-C. TOUGERON – “Idéaux de fonctions différentiables. I”, *Ann. Inst. Fourier (Grenoble)* **18** (1968), no. 1, p. 177–240.
- [Wei00] A. WEINSTEIN – “Linearization problems for Lie algebroids and Lie groupoids”, *Lett. Math. Phys.* **52** (2000), no. 1, p. 93–102.
- [Zam19] M. ZAMBON – Personal communication, 2019.
- [Zun03] N. T. ZUNG – “Levi decomposition of analytic Poisson structures and Lie algebroids”, *Topology* **42** (2003), no. 6, p. 1403–1420.

Manuscript received 27th June 2020

accepted 3rd May 2021

CAMILLE LAURENT-GENGOUX, Institut Élie Cartan de Lorraine, UMR 7502, Université de Lorraine Metz, France

E-mail : camille.laurent-gengoux@univ-lorraine.fr

Url : <https://iecl.univ-lorraine.fr/membre-iecl/laurent-gengoux-camille/>

LEONID RYVKIN, Institut Mathématiques de Jussieu, Université Paris Diderot Paris, France

Faculty of Mathematics, Universität Duisburg-Essen

Essen, Germany

Institut für Mathematik, Georg-August-Universität Göttingen

Göttingen, Germany

E-mail : Leonid.Ryvkin@mathematik.uni-goettingen.de

Url : <http://www.ryvkin.eu/>