



# Journal de l'École polytechnique

## *Mathématiques*

Ismael BAILLEUL & Masato HOSHINO

**Paracontrolled calculus and regularity structures II**

Tome 8 (2021), p. 1275-1328.

[http://jep.centre-mersenne.org/item/JEP\\_2021\\_\\_8\\_\\_1275\\_0](http://jep.centre-mersenne.org/item/JEP_2021__8__1275_0)

© Les auteurs, 2021.

*Certains droits réservés.*



Cet article est mis à disposition selon les termes de la licence

LICENCE INTERNATIONALE D'ATTRIBUTION CREATIVE COMMONS BY 4.0.

<https://creativecommons.org/licenses/by/4.0/>

L'accès aux articles de la revue « Journal de l'École polytechnique — Mathématiques » (<http://jep.centre-mersenne.org/>), implique l'accord avec les conditions générales d'utilisation (<http://jep.centre-mersenne.org/legal/>).

Publié avec le soutien  
du Centre National de la Recherche Scientifique



Publication membre du  
Centre Mersenne pour l'édition scientifique ouverte  
[www.centre-mersenne.org](http://www.centre-mersenne.org)

## PARACONTROLLED CALCULUS AND REGULARITY STRUCTURES II

BY ISMAEL BAILLEUL & MASATO HOSHINO

**ABSTRACT.** — We prove a general equivalence statement between the notions of models and modeled distributions over a regularity structure, and paracontrolled systems indexed by the regularity structure. This takes in particular the form of a parameterization of the set of models over a regularity structure by the set of reference functions used in the paracontrolled representation of these objects. A number of consequences are emphasized. The construction of a modeled distribution from a paracontrolled system is explicit, and takes a particularly simple form in the case of the regularity structures introduced by Bruned, Hairer and Zambotti for the study of singular stochastic partial differential equations.

**RÉSUMÉ** (Calcul paracontrôlé et structures de régularités (II)). — Nous démontrons un énoncé général d'équivalence entre les notions de modèles et de distributions modelées définis sur une structure de régularité et la notion de système paracontrôlé indexé par cette structure de régularité. Cet énoncé donne en particulier une paramétrisation de l'ensemble des modèles sur une structure donnée par l'ensemble des fonctions de référence utilisées dans la représentation paracontrôlée de ces objets. Un certain nombre de conséquences sont données. La construction d'une distribution modelée à partir d'un système paracontrôlé est explicite et prend une forme particulièrement simple dans le cadre des structures de régularités introduites par Bruned, Hairer et Zambotti pour l'étude des équations aux dérivées partielles stochastiques singulières.

### CONTENTS

1. Introduction.....	1276
2. Functional setting.....	1284
3. From regularity structures and models to paracontrolled systems.....	1287
4. From paracontrolled systems to models and modeled distributions.....	1296
Appendix A. Concrete regularity structures.....	1315
Appendix B. Technical estimates.....	1317
Appendix C. The slowly growing setting.....	1326
References.....	1327

**MATHEMATICAL SUBJECT CLASSIFICATION (2020).** — 60L30, 60L40.

**KEYWORDS.** — Regularity structures, models, modeled distributions/functions, paracontrolled calculus, parameterization.

## 1. INTRODUCTION

The set of singular stochastic partial differential equations (PDEs) is characterized by the appearance in each equation of this class of ill-defined products, typically the product of a distribution with a function that is not sufficiently regular. The parabolic Anderson model equation

$$(\partial_t - \Delta)u = u\zeta,$$

on the two dimensional torus is a typical example of singular PDE. The space white noise  $\zeta$  has almost surely parabolic Hölder regularity  $\alpha - 2$ , for any  $\alpha < 1$ , and  $u$  cannot be expected to have better regularity than being  $\alpha$ -Hölder. So the product  $u\zeta$  does not make sense, since  $\alpha + (\alpha - 2) < 0$ . Two different sets of tools for the study of singular stochastic PDEs have emerged recently, under the form of Hairer's theory of regularity structures [14, 7, 9, 6] and paracontrolled calculus [13, 4, 3], after Gubinelli, Imkeller and Perkowski' seminal work. Both of them implement the same mantra: Make sense of the equation in a restricted space of functions/distributions whose elements look like the linear combination of reference random quantities, for which the ill-defined terms that come from the analysis of the product problems can be defined using probabilistic tools. Within the setting of regularity structures, Taylor-like pointwise expansions and jet-like objects are used to make sense of what it means to look like a linear combination of reference quantities

$$f(\cdot) \sim \sum_{\tau} f_{\tau}(z)(\Pi_z^{\mathbf{g}}\tau)(\cdot), \quad \text{near } z, \text{ for all spacetime points } z.$$

In the paracontrolled approach, one uses paraproducts to implement this mantra

$$f \sim \sum_{\tau} P_{f_{\tau}}[\tau].$$

Each term  $P_a b$  is a function or a distribution. This approach is justified at an intuitive level by the fact that  $P_{f_{\tau}}[\tau]$  can be thought of as a modulation of the reference function/distribution  $[\tau]$ . The two options seem technically very different from one another.

While Hairer's theory has now reached the state of a ready-to-use black box for the study of singular stochastic PDEs, like Cauchy-Lipschitz well-posedness theorem for ordinary differential equations, the task of giving a self-contained treatment of renormalisation matters within paracontrolled calculus remains to be done. It happens nonetheless to be possible to compare the two languages, independently of their applications to the study of singular stochastic PDEs. This task was initiated in Gubinelli, Imkeller, Perkowski' seminal work [13] and Martin and Perkowski's work [22], and in our previous work [5], where we proved that the set of admissible models  $\mathbf{M} = (\mathbf{g}, \Pi)$  over a concrete regularity structure  $\mathcal{T} = ((T^+, \Delta^+), (T, \Delta))$  equipped with an abstract integration map is parameterized by a *paracontrolled representation* of  $\Pi$  on the set of elements  $\tau$  with non-positive homogeneity. (Admissible models play a crucial in the regularity structures approach to the study of singular stochastic PDEs.) Theorem 21 in [5] says indeed that given *any* family  $(\llbracket \tau \rrbracket \in \mathcal{C}^{|\tau|})_{|\tau| \in \mathbb{Z}_0^-}$ , with  $\tau$

in a linear basis of  $T$ , there exists a unique admissible model  $(g, \Pi)$  on  $T$  such that one has

$$(1.1) \quad \Pi\tau = \sum_{\sigma < \tau} P_{g(\tau/\sigma)}[\sigma] + \llbracket \tau \rrbracket,$$

for all  $\tau \in T$  in the basis, with non-positive homogeneity. (All notations and words are explained below.) This result provides a parameterization of the *nonlinear* set of admissible models by a *linear* space, providing for instance a natural notion of tangent space to the space of admissible models. The distribution  $\llbracket \tau \rrbracket$  appears in (1.1) as ‘the’ part of  $\Pi\tau$  of regularity  $|\tau|$  in this decomposition, while the paraproducts  $P_{g(\tau/\sigma)}[\sigma]$  have regularity  $|\sigma| < |\tau|$ , for  $\sigma < \tau$ .

To understand the practical relevance of this linear parameterization of the space of admissible models on  $T$ , assume  $T$  stands for the Bruned, Hairer, and Zambotti’s regularity structure [7] associated with a singular stochastic PDE and  $M^\varepsilon = (g^\varepsilon, \Pi^\varepsilon)$  stands for the naive interpretation model associated with a smoothed noise in the equation, with regularization parameter  $\varepsilon$ . The BPHZ renormalization procedure for the model involves a real-valued map  $k$  acting on a side space  $T^-$ , which also defines a homogeneity-preserving linear map  $\tilde{k}$  from  $T$  into itself. It follows from [5, Th. 21] that the bracket data associated with the renormalized model  ${}^kM^\varepsilon$  is simply given by the  $\llbracket \tilde{k}(\tau) \rrbracket$ , for  $\tau$  of negative homogeneity. The convergence of renormalized admissible models has thus a direct counterpart in terms of bracket data. This answers one of the problems mentioned at the end of Tapia and Zambotti’s work [24] on the parameterization problem for the set of branched rough paths, in the present general setting.

Here is another illustration of the use of the parameterization result of admissible models proved in [5] that will be developed in Section 4.1 and Section 4.4. Consider the elementary setting of branched rough paths; they are admissible models on particular examples of regularity structures. Theorem 21 in [5] gives a direct proof of Lyons’ extension theorem, saying that a branched Hölder  $p$ -rough path has a unique extension into a branched Hölder  $q$ -rough path, for any  $q > p$ . (Recall weak geometric rough paths are branched rough paths.) This result allows to define the signature of a branched rough path. Similarly, let  $T$  be a regularity structure built from integration operators, with elements of arbitrary large positive homogeneity. It follows from [5, Th. 21] that an admissible model defined on the quotient space of  $T$ , modulo elements of a given positive homogeneity  $\alpha$ , has a unique extension into an admissible model over the regularity structure  $T$  quotiented by the elements of homogeneity  $\beta$ , for any  $\beta > \alpha$ . This allows to define the signature of an admissible model.

Such statements are concerned with admissible models on regularity structures associated with singular stochastic PDEs. We step back in the present work and prove a general result giving a parameterization of the *nonlinear* space of arbitrary models  $M = (g, \Pi)$  on any reasonable concrete regularity structure, by a *linear* space, in terms of representations of the maps  $g$  and  $\Pi$  by paracontrolled systems, similar to identity (1.1). (The set of models on any given regularity structure is always nonempty, as it

contains the element  $M_0 = (g_0, \Pi_0)$ , with  $g_0$  the character on  $T^+$  that sends any basis element of  $T^+$  on 1, and  $\Pi_0$  the null map. The nonlinearity of the space of models can be seen from the analytical constraints that they need to satisfy, that involves nonlinear operations on  $g$ .) Being reasonable means here satisfying assumptions (A–C) from Section 3 and Section 4. We insist here on the fact that these assumptions are not related to any kind of singular stochastic PDE or any dynamics or structure that could be modeled with such a regularity structure. As we shall see, the regularity structures used for the study of singular stochastic PDEs enjoy these properties, so all our results hold for them.

The result takes the following form. Given a concrete regularity structure

$$\mathcal{T} = ((T^+, \Delta^+), (T, \Delta)),$$

denote by  $M_{\text{rap}}(\mathcal{T}, \mathbb{R}^d)$  the space of models on  $\mathbb{R}^d$  decreasing rapidly at infinity. Once again, all terms will be properly defined below.

**THEOREM 1.** — *Let  $\mathcal{T}$  be a concrete regularity structure satisfying assumptions (A–C). Then one can construct a locally Lipschitz continuous map*

$$(1.2) \quad \begin{aligned} M_{\text{rap}}(\mathcal{T}, \mathbb{R}^d) &\longrightarrow \prod_{\sigma \in \mathcal{B}^+ \cap \mathcal{B}_X^+} \mathcal{C}_{\text{rap}}^{|\sigma|}(\mathbb{R}^d) \times \prod_{\tau \in \mathcal{B} \cap \mathcal{B}_X} \mathcal{C}_{\text{rap}}^{|\tau|}(\mathbb{R}^d) \\ (\mathbf{g}, \Pi) &\longmapsto \left( \llbracket \sigma \rrbracket^{\mathbf{M}}, \llbracket \tau \rrbracket^{\mathbf{g}}; \sigma \in \mathcal{B}^+ \cap \mathcal{B}_X^+, \tau \in \mathcal{B} \cap \mathcal{B}_X \right) \end{aligned}$$

by giving paracontrolled representations of  $\mathbf{g}$  and  $\Pi$ , for  $(\mathbf{g}, \Pi) \in M_{\text{rap}}(\mathcal{T}, \mathbb{R}^d)$ . Furthermore,  $M_{\text{rap}}(\mathcal{T}, \mathbb{R}^d)$  is locally bi-Lipschitz homeomorphic to the product space

$$(1.3) \quad \prod_{\sigma \in \mathcal{G}_\circ^+} \mathcal{C}_{\text{rap}}^{|\sigma|}(\mathbb{R}^d) \times \prod_{\tau \in \mathcal{B}_\bullet, |\tau| < 0} \mathcal{C}_{\text{rap}}^{|\tau|}(\mathbb{R}^d).$$

The first claim in Theorem 1 is part of [5, Th. 21]; see formulas (3.5) and (3.6) below for an explicit description of the map (1.2). The sets  $\mathcal{B}$  and  $\mathcal{B}^+$  are fixed linear bases of the spaces  $T$  and  $T^+$ , respectively, consisting of homogeneous vectors. The set  $\mathcal{B}_\bullet$  in (1.3) parametrizes part of the basis  $\mathcal{B}$ , while the set  $\mathcal{G}_\circ^+$  parametrizes part of the basis  $\mathcal{B}^+$ . The letter  $\mathcal{G}$  stands for ‘generator’. In the present setting of a general concrete regularity structure, the space  $T^+$  is not related to  $T$ , unlike what happens with the special regularity structures used for the study of singular stochastic PDEs. It is thus not surprising that there is some freedom in the construction/parameterization of the map  $\mathbf{g}$ . The degrees of freedom are parametrized by the set  $\mathcal{G}_\circ^+$ , described in assumption (C). Assumption (A) is a harmless requirement on how polynomials sit within  $T$  and  $T^+$ . Assumption (B) is a very mild requirement on the splitting map  $\Delta : T \rightarrow T \otimes T^+$ , and assumption (C) is a structure requirement on  $T^+$  and  $\Delta^+$  that provides a fundamental induction structure. The three assumptions are met by all concrete regularity structures built for the study of singular stochastic PDEs.

This type of parameterization is not entirely new as Tapia and Zambotti described in [24] a free transitive action of a product of Hölder spaces on the space of branched rough paths, a particular example of model over a particular regularity structure. This

action was not proved to be continuous however. In relation with the renormalization problem of stochastic models, Theorem 1 describes precisely the freedom that we have to tweak a divergent family of models and turn it into a convergent family of models. The renormalization process needs to give converging bracket data  $[[\sigma]]^M, [[\tau]]^\mathfrak{g}$ , for all  $\sigma \in \mathcal{G}_\circ^+$  and  $\tau \in \mathcal{B}_\bullet$  with  $|\tau| < 0$ . See a forthcoming work.

We single out here two direct consequences of Theorem 1 about density and extension questions on the space of models.

– The set of models with rapid decrease is equipped with a family of norms  $M \mapsto \|M\|_a$ , indexed by positive exponents  $a$ . Smooth functions are known to be dense in any Hölder space  $\mathcal{C}_a^\beta(\mathbb{R}^d)$ , with growth exponent  $a$ , if one sees the latter as a subset of  $\mathcal{C}_a^{\beta-\varepsilon}(\mathbb{R}^d)$ , for any positive  $\varepsilon$ . Theorem 1 provides as a consequence a direct proof of the following density result, proved in Section 4.4 – see Singh and Teichmann’s work [23] for a similar result, proved therein from an explicit mollification procedure on models.

**COROLLARY 2.** — *Given any positive exponent  $\varepsilon$ , the set of smooth models with rapid decrease is dense in the set of models with finite  $\mathcal{C}_{\text{rap}}^{|\tau|}$ -norms, for the topology induced by the  $\mathcal{C}_{\text{rap}}^{|\tau|-\varepsilon}$ -norms.*

– Branched rough paths are models on a finite time interval  $[0, T]$ , over a particular example of concrete regularity structure of the form  $T^+ = ((T^+, \Delta^+), (T^+, \Delta^+))$ , that satisfies assumptions (A–C). These models are entirely determined by their  $\mathfrak{g}$ -maps, and the elements of  $T^+$  are planted rooted trees with decorations on the nodes in a finite set  $\{1, \dots, \ell\}$ ; edges are not decorated. Defining a branched rough path above an  $\ell$ -dimensional control  $h = (h^1, \dots, h^\ell)$  means defining a  $\mathfrak{g}$ -map over  $(T^+, \Delta^+)$  such that  $\mathfrak{g}$  assigns  $h^j$  to the tree with only one node with decoration  $j$ , for all  $j$  such that  $1 \leq j \leq \ell$ , and no edge. The first proof that this is possible for any choice of Hölder control  $h$  was found by Lyons and Victoir [20], for geometric rough paths, using the axiom of choice. This unexpected device stimulated further explorations of this questions, and different proofs not using the axiom of choice were given subsequently [25, 14, 24, 18]. Unterberger constructs in [25] a rough path above  $h$  using paraproduct-like tools. Hairer uses in [14] the reconstruction theorem for that purpose, while Liu, Prömel and Teichmann use in [18] a version of the reconstruction theorem for Sobolev models and the notion of Sobolev rough path to extend Lyons-Victoir extension in their setting. Tapia and Zambotti used in [24] an explicit form of the Baker-Campbell-Hausdorff formula to bypass the use of the axiom of choice in the construction of a lift, and gave a parameterization of the set of all branched rough paths above  $h$ . Theorem 1 provides a direct access to such an extension result, in so far as the family of trees with only one node with decoration  $j$ , for  $1 \leq j \leq \ell$ , is a subset of the generator set  $\mathcal{G}_\circ^+$  in that setting.

**COROLLARY 3** (Lyons-Victoir’s extension theorem). — *Given any  $\mathbb{R}^\ell$ -valued Hölder control  $h$  on the time interval  $[0, T]$ , there exists a branched rough path above  $h$ .*

The above branched rough path is said to be a lift of the control  $h$ . Like in [24], Theorem 1 actually gives a parameterization of the set of all branched rough paths above  $h$ . One can formulate the same extension problem for the models on the class of concrete regularity structures introduced for the study of singular stochastic PDEs by Bruned, Hairer and Zambotti in [7] – we talk of BHZ regularity structures. We refer the reader to Section 4.3 for basics on BHZ regularity structures, and simply mention here that as in the case of branched rough paths, the elements of BHZ regularity structures  $T$  are rooted decorated trees. Their roots may have decorations, and we denote by  $(\bullet_j)_{1 \leq j \leq \ell}$  the family of one node trees with no edges, and decoration  $j$ .

**COROLLARY 4** (Extension result for models on BHZ regularity structures)

*Given a multidimensional noise  $\zeta = (\zeta_1, \dots, \zeta_\ell)$ , with  $\zeta_i \in \mathcal{C}_{\text{rap}}^{\bullet_i}(\mathbb{R}^m)$ , for all  $i$  such that  $1 \leq i \leq \ell$ , there exists a model  $M = (\mathbf{g}, \Pi)$  on the BHZ regularity structure  $T$  such that  $\Pi(\bullet_j) = \zeta_j$ , for all  $1 \leq j \leq \ell$ .*

The above model is said to be a lift of the noise  $\zeta$ . The parameterization of admissible models proved in [5] shows that one can further impose to the extension that it is an admissible model. (Recall the notion of admissibility is related to a peculiar feature of the regularity structures used for the study of singular stochastic PDEs.) As for the case of branched rough paths, Theorem 1 actually gives a description of the set of all models above the  $\ell$ -dimensional noise  $\zeta$ . Corollary 3 and Corollary 4 are proved in Section 4.4. Corollary 3 and Corollary 4 are actually previously unnoticed consequences of [5, Th. 21]. The general extension result stated in Corollary 4 is outside the scope of [5, Th. 21].

Enough for the consequences of Theorem 1; we now turn to the problem of the parameterization of the space of modeled distributions associated with a given model.

Given a model  $M = (\mathbf{g}, \Pi)$  on a concrete regularity structure, natural regularity spaces are given by the Hölder-type spaces  $\mathcal{D}^\gamma(T, \mathbf{g})$ , with generic element

$$f = \sum_{\tau \in \mathcal{B}, |\tau| < \gamma} f_\tau \tau.$$

For  $M = (\mathbf{g}, \Pi) \in \mathcal{M}_{\text{rap}}(T, \mathbb{R}^d)$ , there is an associated notion of rapidly decreasing space of modeled distributions taking values in the vector space  $T$ , with regularity exponent  $\gamma$ , denoted by  $\mathcal{D}_{\text{rap}}^\gamma(T, \mathbf{g})$ . The parameterization of  $\mathcal{D}_{\text{rap}}^\gamma(T, \mathbf{g})$  by data in paracontrolled representations of elements of that space requires in general a *structure condition* on these data reminiscent of a similar condition introduced by Martin and Perkowski in [22]; it is stated in Theorem 22. This non-trivial structure condition has a clear meaning in terms of an extension problem for the map  $\mathbf{g}$  from the Hopf algebra  $T^+$  to a larger Hopf algebra; an interesting technical point on its own. The structure condition happens nonetheless to take a very simple form for special concrete regularity structures satisfying assumption (D).

**THEOREM 5.** — *Let a concrete regularity structure  $T$  satisfy assumptions (A–D). Pick  $\gamma \in \mathbb{R} \setminus \{0\}$  such that  $\gamma - |\tau| \notin \mathbb{N}$  for any basis element  $\tau$  of  $T$  with  $|\tau| < \gamma$ , and*

$M = (\mathbf{g}, \Pi) \in M_{\text{rap}}(T, \mathbb{R}^d)$ . Then one can construct a locally Lipschitz continuous map

$$\mathcal{D}_{\text{rap}}^\gamma(T, \mathbf{g}) \longrightarrow \prod_{\tau \in \mathcal{B}, |\tau| < \gamma} \mathcal{C}_{\text{rap}}^{\gamma-|\tau|}(\mathbb{R}^d)$$

by giving a paracontrolled representation of elements in  $\mathcal{D}_{\text{rap}}^\gamma(T, \mathbf{g})$ . Furthermore,  $\mathcal{D}_{\text{rap}}^\gamma(T, \mathbf{g})$  is locally bi-Lipschitz homeomorphic to the product space

$$\prod_{\tau \in \mathcal{B}_\bullet, |\tau| < \gamma} \mathcal{C}_{\text{rap}}^{\gamma-|\tau|}(\mathbb{R}^d).$$

See formula (3.7) for the paracontrolled representation of a modeled distribution in  $\mathcal{D}_{\text{rap}}^\gamma(T, \mathbf{g})$ . Emphasize that the set  $\mathcal{B}_\bullet$  contains an element  $\mathbf{1}$  in the setting of regularity structures built for the study of singular stochastic PDEs. The only non-zero component in the parameterization of a classically regular function  $f$  is in that case given by its  $\mathbf{1}$ -component, equal to  $f \in \mathcal{C}_{\text{rap}}^\gamma(\mathbb{R}^d)$  itself.

Similarly, we can see the further homeomorphism result

$$M_{\text{rap}} \cap \mathcal{D}_{\text{rap}}^\gamma \simeq \prod_{\sigma \in \mathcal{G}_\circ^+} \mathcal{C}_{\text{rap}}^{|\sigma|}(\mathbb{R}^d) \times \prod_{\tau \in \mathcal{B}_\bullet, |\tau| < 0} \mathcal{C}_{\text{rap}}^{|\tau|}(\mathbb{R}^d) \times \prod_{\tau \in \mathcal{B}_\bullet, |\tau| < \gamma} \mathcal{C}_{\text{rap}}^{\gamma-|\tau|}(\mathbb{R}^d),$$

where  $M_{\text{rap}} \cap \mathcal{D}_{\text{rap}}^\gamma$  is the space of all pairs  $((\mathbf{g}, \Pi), \mathbf{f})$  of models  $(\mathbf{g}, \Pi) \in M_{\text{rap}}(T, \mathbb{R}^d)$  and modeled distributions  $\mathbf{f} \in \mathcal{D}_{\text{rap}}^\gamma(T, \mathbf{g})$ . Following Corollary 2, say here that given a model  $M$  on a concrete regularity structure  $T$ , the set of modeled distributions with rapid decrease is equipped with a family of norms  $\mathbf{f} \mapsto \|\mathbf{f}\|_a$ , indexed by a positive growth exponent  $a$ . The following result is obtained as a direct consequence of Theorem 5 and the density of smooth functions in any Hölder space  $\mathcal{C}_a^\beta(\mathbb{R}^d)$ , equipped with the weaker  $\mathcal{C}_a^{\beta-\varepsilon}(\mathbb{R}^d)$ -topology, for any positive exponent  $\varepsilon$ . As pointed out in Section 2 of Singh and Teichmann’s work [23], one can use the reconstruction theorem to define a mollification operator on modeled distributions and obtain as a consequence a density statement for the set of smooth modeled distributions. Theorem 5 shows that any mollification operation on Hölder spaces induces a mollification operation on the space of modeled distributions; this result is independent of the reconstruction theorem. See Section 4.4 for a proof.

**COROLLARY 6.** — *Let a concrete regularity structure  $T$  satisfy assumptions (A–D). Fix a model on  $T$ . Given any exponents  $\gamma \in \mathbb{R}$  as in Theorem 5 and  $\varepsilon > 0$ , the set of smooth elements  $((\mathbf{g}, \Pi), \mathbf{f})$  in  $M_{\text{rap}} \cap \mathcal{D}_{\text{rap}}^\gamma$  is dense in the same space but with the topology induced by the  $\mathcal{C}_{\text{rap}}^{|\tau|-\varepsilon}$ -norms and the  $\mathcal{D}_{\text{rap}}^{\gamma-\varepsilon}$ -norm.*

Unlike the other assumptions, assumption (D) is fundamentally a requirement on a linear basis of  $T$ , not on the concrete regularity structure itself. It may then happen that one basis of  $T$  satisfies it whereas another does not. Satisfying assumption (D) thus means the existence of a linear basis satisfying this assumption. It happens that the class of concrete BHZ regularity structures introduced by Bruned, Hairer and Zambotti in [7] for the study of singular stochastic PDEs all satisfy assumption (D), despite the fact that their canonical bases do not satisfy it. We refer the reader to Section 4.3 for the notations  $\mathfrak{t} \in \mathfrak{L}$  and  $|\mathfrak{t}|$ .



**THEOREM 7.** — Assume that the set  $\{|\mathbf{t}|\}_{\mathbf{t} \in \mathfrak{L}} \cup \{1\}$  is rationally independent. Then the BHZ concrete regularity structures satisfy assumptions (A–D).

*BHZ regularity structures vs general regularity structures.* — Readers familiar with the use of regularity structures for the study of singular stochastic PDEs may feel uncomfortable at the idea of working with regularity structures that do not come from a (system of) singular stochastic PDE(s) and with models where the maps  $\mathbf{g}$  and  $\Pi$  are unrelated, unlike in the former setting. This freedom is useful, and Hoshino showed for instance in [16, 17] how this leads to a clear understanding of a number of fundamental continuity results for iterated correctors introduced in Bailleul & Bernicot’s work [3] on high order paracontrolled calculus, from a regularity structures point of view. As a further illustration of the use of this freedom, let us see how Theorem 1 gives back a proof of the continuity of the product map  $(a, b) \in \mathcal{C}^\alpha(\mathbb{R}^d) \times \mathcal{C}^\beta(\mathbb{R}^d) \mapsto ab \in \mathcal{C}^\beta(\mathbb{R}^d)$ , for  $\alpha \in (0, 1)$ ,  $\beta < 0$ , and  $\alpha + \beta > 0$ ; this is another formulation of [14, Prop. 4.14]. Indeed, consider the concrete regularity structure  $T = ((T^+, \Delta^+), (T, \Delta))$  with

$$T^+ = \text{span}(\mathbf{1}_+, A), \quad T = \text{span}(B, C),$$

with  $|\mathbf{1}_+| := 0$ ,  $|A| := \alpha$ ,  $|B| := \beta$ ,  $|C| := \alpha + \beta$ , and splitting maps

$$\Delta^+ \mathbf{1}_+ = \mathbf{1}_+ \otimes \mathbf{1}_+, \quad \Delta^+ A = A \otimes \mathbf{1}_+ + \mathbf{1}_+ \otimes A,$$

and

$$\Delta B = B \otimes \mathbf{1}_+, \quad \Delta C = C \otimes \mathbf{1}_+ + B \otimes A.$$

Theorem 1 tells us that the model  $(\mathbf{g}, \Pi)$ , with  $\mathbf{g}(\mathbf{1}_+) = 1$ , is uniquely characterized by the two inputs

$$\mathbf{g}(A) = a \in \mathcal{C}^\alpha(\mathbb{R}^d), \quad \Pi B = b \in \mathcal{C}^\beta(\mathbb{R}^d).$$

The distribution  $c := \Pi C$  is in particular determined by  $a$  and  $b$ . We see that  $\Pi$  provides an extension of the product map  $(a, b) \mapsto ab$ , by noting that for smooth inputs  $a, b$ , the identity  $(\Pi_x^{\mathfrak{g}} C)(x) = 0$ , implies in that case  $c(x) = a(x)b(x)$ , for all  $x \in \mathbb{R}^d$ .

As a matter of fact, working with models with unrelated  $\mathbf{g}$  and  $\Pi$  should somehow be easier than working with admissible models, where  $\mathbf{g}$  and  $\Pi$  are entangled with one another so as to satisfy the admissibility condition.

As far as working with general regularity structures rather than just working regularity structures associated with singular stochastic PDEs is concerned, we would like to encourage the reader to think about general regularity structures as mathematical models of rough ‘media’ within which one still has a calculus. Rough medias have no reason to be associated with any PDE on a general basis.

The following additional remarks put further our results in perspective.

– In the theory of regularity structures, the solution map of a singular stochastic PDE has the following structure

$$M_{\text{rap}}(T, \mathbb{R}^d) \longrightarrow \mathcal{D}_{\text{rap}}^\gamma(T, \mathbf{g}) \xrightarrow{\mathbf{R}} \mathcal{C}^{\beta_0}(\mathbb{R}^d).$$

The first arrow associates to a model the solution in  $\mathcal{D}_{\text{rap}}^\gamma(T, \mathfrak{g})$  of the regularity structure counterpart of the equation; the second arrow involves the model-dependent reconstruction map  $\mathbf{R}$ . The composition of these two maps defines a locally Lipschitz map. Theorem 1 implies that the solution map actually has the structure

$$(1.4) \quad \prod_{\sigma \in \mathcal{G}_\bullet^+} \mathcal{C}_{\text{rap}}^{|\sigma|}(\mathbb{R}^d) \times \prod_{\tau \in \mathcal{B}_\bullet, |\tau| < 0} \mathcal{C}_{\text{rap}}^{|\tau|}(\mathbb{R}^d) \longrightarrow \mathcal{C}^{\beta_0}(\mathbb{R}^d).$$

The map (1.4) is a general form of the solution maps constructed in the previous works [13, 3] on paracontrolled calculus. Since the ansatz on solutions were given by hand in those papers, it was very hard to extend the argument to a whole class of equations. Our results reveal the relation between such handmade ansatz and the sophisticated algebraic structure in Hairer’s theory, showing that it is possible to apply paracontrolled calculus to more general equations in an automatic way, like the works [14, 7, 9, 6].

– The map (1.4) provides interesting insights on parts of the theory of regularity structures. For example, one of the difficult part of the theory is the continuity result for the model-dependent multi-level extension

$$\mathcal{K}^M : \mathcal{D}^\gamma(T, \mathfrak{g}) \longrightarrow \mathcal{D}^{\gamma+2}(T, \mathfrak{g}),$$

of the resolution map  $\mathcal{L}^{-1}$ , with the property that  $\mathbf{R}^M(\mathcal{K}^M \mathbf{f}) = \mathcal{L}^{-1}(\mathbf{R}^M \mathbf{f})$ , for any modeled distributions  $\mathbf{f} \in \mathcal{D}^\gamma(T, \mathfrak{g})$  – its very definition is non-obvious, see [14, §5]. From the paracontrolled point of view, we take profit from the fact that the classical resolution map  $\mathcal{L}^{-1}$  preserves the paracontrolled structure

$$\mathcal{L}^{-1} : \sum \mathbf{P}_{f_\tau}[\tau] + \llbracket \mathbf{f} \rrbracket \longmapsto \sum \bar{\mathbf{P}}_{f_\tau}(\mathcal{L}^{-1}[\tau]) + \mathcal{L}^{-1}[\llbracket \mathbf{f} \rrbracket],$$

up to the introduction of the modified paraproduct  $\bar{\mathbf{P}}_f g := \mathcal{L}^{-1} \mathbf{P}_f(\mathcal{L}g)$  – see [3]. The main results in the present paper can be applied to such a modified paraproduct. The map  $\mathcal{K}^M$  can be obtained directly from Theorem 5 by giving first a paracontrolled representation of an element of  $\mathcal{D}^\gamma(T, \mathfrak{g})$ , then applying  $\mathcal{L}^{-1}$ , using the modified paraproduct, and finally using Theorem 5 again to get back an element of  $\mathcal{D}^{\gamma+2}(T, \mathfrak{g})$ . We do not give the details here and leave it to a future work.

– The local Lipschitz parameterizations of the sets of models and modeled distributions from Theorem 1 and Theorem 5 offer the possibility to define dynamics in these spaces by solving ordinary (or controlled/rough) differential equations driven by vector fields on the parameterization spaces. In the setting of pathspace analysis on manifolds, this kind of pathwise dynamics provided a clean understanding of Driver’s flow equation on pathspace, in relation with quasi-invariance questions for Wiener measure on pathspace over a compact Riemannian manifold [12, 19, 2]. One may also make sense of classical stochastic PDEs on the space of models or modeled distributions, as in Liu, Prömel and Teichmann’s work [18].

Notice that we considered function spaces whose elements decrease rapidly at infinity mainly for a technical reason. Our assumption is related to localizing a singular PDE. Indeed, we can consider a class of models on a bounded domain vanishing on the

boundary, via a diffeomorphism between  $\mathbb{R}^d$  and that domain. Instead, the following modifications are also possible.

- Theorems 1 and 5 above hold also with the spaces  $M_{\text{slow}}$  and  $\mathcal{D}_{\text{slow}}^\gamma$  of slowly growing models and modeled distributions, respectively, with the spaces  $\mathcal{C}_{\text{rap}}^\alpha$  replaced by  $\mathcal{C}_{\text{slow}}^\alpha$ . Such a modification is important because temporally or spatially stationary models belong to  $M_{\text{slow}}$  but not to  $M_{\text{rap}}$ . More details can be found in Appendix C.

- If the elements in  $\mathcal{G}_\circ^+$  and  $\mathcal{B}_\bullet$  all have homogeneities smaller than 1, then Theorem 1 and Theorem 5 above hold for the unweighted spaces  $M$  and  $\mathcal{D}^\gamma$ , with the spaces  $\mathcal{C}_{\text{rap}}^\alpha$  replaced by usual Hölder spaces  $\mathcal{C}^\alpha$ . An important example is the space of branched rough paths. As said above, Tapia and Zambotti proved in [24] an analogue of Theorem 1 for the space of branched rough paths by a different approach.

Like in our previous work [5], we work here with the usual isotropic Hölder space rather than with anisotropic spaces. All results given here hold true in that more general setting, with identical proofs. The reader will find relevant technical details in the work [22] of Martin and Perkowski.

Section 2 is dedicated to describing different functional spaces and operators. Section 3 is dedicated to giving paracontrolled representations of models and the reconstruction of modeled distributions in terms of data in paracontrolled systems, proving part of Theorem 1. The latter is proved in Section 4, where the main work consists in providing a parameterization of  $\mathfrak{g}$ -maps by paracontrolled representations, Theorem 17. Theorem 5 and Theorem 7 are proved in Section 4.2 and 4.3, respectively. Appendix A gives back the setting of concrete regularity structures introduced in [5], while Appendix B gives a number of technical details that are variations on corresponding results from [5].

#### NOTATION

- We use exclusively the letters  $\alpha, \beta, \gamma$  to denote real numbers that play the role of regularity exponents, and use the letters  $\sigma, \tau, \mu, \nu$  to denote elements of  $T$  or  $T^+$ .

- We agree to use the shorthand notation  $\mathfrak{s}^{(+)}$  to mean both the statement  $\mathfrak{s}$  and the statement  $\mathfrak{s}^+$ .

- We use the pairing notation  $\langle \cdot, \cdot \rangle$  for duality between a finite dimensional vector space and its dual space.

- We adopt the notations and terminology of the work [5], and write in particular  $\Pi_x^\xi$  and  $\widehat{\mathfrak{g}}_{yx}$ , for what is denoted by  $\Pi_x$  and  $\Gamma_{xy}$  in Hairer's terminology.

## 2. FUNCTIONAL SETTING

We describe in this section different function spaces we shall work with and introduce a modified paraproduct. For  $x \in \mathbb{R}^d$ , set

$$|x|_* := 1 + |x|, \quad x \in \mathbb{R}^d.$$

The weight function  $|x|_*$  satisfies the inequalities

$$|x + y|_* \leq |x|_* |y|_*, \quad |x/\lambda|_* \leq |x|_*,$$

for any  $\lambda > 1$ .

Let  $(\rho_i)_{-1 \leq i < \infty}$  be a dyadic decomposition of unity on  $\mathbb{R}^d$ , i.e.,  $\rho_i : \mathbb{R}^d \rightarrow [0, 1]$  is a compactly supported smooth radial function with the following properties.

- $\text{supp}(\rho_{-1}) \subset \{x \in \mathbb{R}^d; |x| < 4/3\}$  and  $\text{supp}(\rho_0) \subset \{x \in \mathbb{R}^d; 3/4 < |x| < 8/3\}$ .
- $\rho_i(x) = \rho_0(2^{-i}x)$  for any  $x \in \mathbb{R}^d$  and  $i > 0$ .
- $\sum_{i=-1}^{\infty} \rho_i(x) = 1$  for any  $x \in \mathbb{R}^d$ .

We define the *Littlewood-Paley blocks*  $(\Delta_i)_{-1 \leq i < \infty}$  by  $\Delta_i f := \rho_i(\nabla)f := \mathcal{F}^{-1}(\rho_i \mathcal{F}f)$ , where  $\mathcal{F}$  is a Fourier transform on  $\mathbb{R}^d$  and  $\mathcal{F}^{-1}$  is its inverse. For  $j > -1$ , set

$$S_j := \sum_{i < j-1} \Delta_i.$$

Denote by  $Q_i$  and  $P_j$  the integral kernels associated with  $\Delta_i$  and  $S_j$

$$\Delta_i f(x) := \int_{\mathbb{R}^d} Q_i(x-y)f(y)dy, \quad S_j f(x) := \int_{\mathbb{R}^d} P_j(x-y)f(y)dy.$$

- For any measurable function  $f : \mathbb{R}^d \rightarrow \mathbb{R}$ , set

$$\|f\|_{L_a^\infty(\mathbb{R}^d)} := \|\cdot |_*^a f\|_{L^\infty(\mathbb{R}^d)},$$

and define the corresponding space  $L_a^\infty(\mathbb{R}^d)$  of functions with finite  $\|\cdot\|_{L_a^\infty(\mathbb{R}^d)}$ -norm. Set

$$L_{\text{rap}}^\infty(\mathbb{R}^d) := \bigcap_{a=1}^{\infty} L_a^\infty(\mathbb{R}^d), \quad L_{\text{slow}}^\infty(\mathbb{R}^d) := \bigcup_{a=1}^{\infty} L_{-a}^\infty(\mathbb{R}^d).$$

- For any distribution  $\xi \in \mathcal{S}'(\mathbb{R}^d)$ , set

$$\|\xi\|_{\mathcal{C}_a^\alpha(\mathbb{R}^d)} := \sup_{j > -1} 2^{j\alpha} \|\Delta_j \xi\|_{L_a^\infty(\mathbb{R}^d)}.$$

and define the corresponding space  $\mathcal{C}_a^\alpha(\mathbb{R}^d)$  of functions with finite  $\|\cdot\|_{\mathcal{C}_a^\alpha(\mathbb{R}^d)}$ -norm. We have  $\mathcal{C}_0^\alpha(\mathbb{R}^d) = \mathcal{C}^\alpha(\mathbb{R}^d)$ , with the usual definition of the Hölder space  $\mathcal{C}^\alpha(\mathbb{R}^d)$  as the Besov space  $\mathcal{B}_{\infty, \infty}^\alpha(\mathbb{R}^d)$  – see e.g. Bahouri, Chemin and Danchin’s book [1]. Set

$$\mathcal{C}_{\text{rap}}^\alpha(\mathbb{R}^d) := \bigcap_{a=1}^{\infty} \mathcal{C}_a^\alpha(\mathbb{R}^d), \quad \mathcal{C}_{\text{slow}}^\alpha(\mathbb{R}^d) := \bigcup_{a=1}^{\infty} \mathcal{C}_{-a}^\alpha(\mathbb{R}^d).$$

- For any two-parameter function  $F : \mathbb{R}^d \times \mathbb{R}^d \rightarrow \mathbb{R}$  and  $\alpha > 0$ , set

$$\|F\|_{\mathcal{C}_{(2), a}^\alpha(\mathbb{R}^d \times \mathbb{R}^d)} := \sup_{x, y \in \mathbb{R}^d} (|x|_*^\alpha \wedge |y|_*^\alpha) \frac{|F(x, y)|}{|x - y|^\alpha}.$$

Define the corresponding space  $\mathcal{C}_{(2), a}^\alpha(\mathbb{R}^d \times \mathbb{R}^d)$  of functions with finite  $\|\cdot\|_{\mathcal{C}_{(2), a}^\alpha(\mathbb{R}^d \times \mathbb{R}^d)}$ -norm. Set also

$$\mathcal{C}_{(2)}^\alpha(\mathbb{R}^d \times \mathbb{R}^d) := \mathcal{C}_{(2), 0}^\alpha(\mathbb{R}^d \times \mathbb{R}^d), \quad \mathcal{C}_{(2), \text{rap}}^\alpha(\mathbb{R}^d \times \mathbb{R}^d) := \bigcap_{a=1}^{\infty} \mathcal{C}_{(2), a}^\alpha(\mathbb{R}^d \times \mathbb{R}^d).$$

- For any  $\mathbb{R}^d$ -indexed family of distributions  $\Lambda = (\Lambda_x)_{x \in \mathbb{R}^d} \subset \mathcal{S}'(\mathbb{R}^d)$  on  $\mathbb{R}^d$ , and  $\alpha \in \mathbb{R}$ , set

$$\|\Lambda\|_{D_a^\alpha} := \sup_{x \in \mathbb{R}^d} \sup_{j > -1} |x|_*^\alpha 2^{j\alpha} |\langle \Lambda_x, P_j(x - \cdot) \rangle|.$$

Set

$$D^\alpha := D_0^\alpha, \quad D_{\text{rap}}^\alpha := \bigcap_{a=1}^{\infty} D_a^\alpha.$$

(In Hairer's seminal work [14], models are assumed to satisfy a  $(\lambda, \varphi)$ -uniform regularity condition

$$|(\Pi_x^{\mathbf{g}} \tau)(\varphi_x^\lambda)| \leq \lambda^{|\tau|},$$

locally uniformly in  $x$ . Requiring  $(\Pi_x^{\mathbf{g}} \tau)_{x \in \mathbb{R}^d} \in D^{|\tau|}$  is equivalent to the above uniform estimate – see e.g. Lemma 6.6 of Gubinelli, Imkeller and Perkowski's seminal work [13] on paracontrolled distributions.)

For any distributions  $f, g \in \mathcal{S}'(\mathbb{R}^d)$ , we define the paraproduct

$$P_f g := \sum_{j=1}^{\infty} (S_j f)(\Delta_j g),$$

and resonant operator

$$\Pi(f, g) := \sum_{|i-j| \leq 1} (\Delta_i f)(\Delta_j g).$$

For any  $g \in \mathcal{S}'(\mathbb{R}^d)$ , set

$$(2.1) \quad Sg := g - P_1 g = (\Delta_{-1} + \Delta_0)g \in C^\infty(\mathbb{R}^d).$$

(The letter S is chosen for 'smooth'.) The following continuity result is an elementary variation on the classical continuity results for the paraproduct and resonant operators. We refer for instance the reader to Lemma 2.1.34 in J. Martin's thesis [21] for a reference.

**PROPOSITION 8.** — *Let  $\alpha, \beta \in \mathbb{R}$ ,  $a, b \in \mathbb{Z}$ .*

- *If  $\alpha \neq 0$ , then  $\mathcal{C}_a^\alpha(\mathbb{R}^d) \times \mathcal{C}_b^\beta(\mathbb{R}^d) \ni (f, g) \mapsto P_f g \in \mathcal{C}_{a+b}^{\alpha \wedge 0 + \beta}(\mathbb{R}^d)$ , is continuous.*
- *If  $\alpha + \beta > 0$ , then  $\mathcal{C}_a^\alpha(\mathbb{R}^d) \times \mathcal{C}_b^\beta(\mathbb{R}^d) \ni (f, g) \mapsto \Pi(f, g) \in \mathcal{C}_{a+b}^{\alpha + \beta}(\mathbb{R}^d)$ , is continuous.*
- *If  $\alpha, \beta \neq 0$  and  $\alpha + \beta > 0$ , then  $\mathcal{C}_a^\alpha(\mathbb{R}^d) \times \mathcal{C}_b^\beta(\mathbb{R}^d) \ni (f, g) \mapsto f \cdot g \in \mathcal{C}_{a+b}^{\alpha \wedge \beta}(\mathbb{R}^d)$ , is continuous.*

As a consequence of the last item, the product  $fg$ , of  $f \in \mathcal{S}(\mathbb{R}^d)$  and  $g \in \mathcal{C}^\alpha(\mathbb{R}^d)$ , belongs to  $\mathcal{C}_{\text{rap}}^\alpha(\mathbb{R}^d)$ , for any  $\alpha \in \mathbb{R}$  – so the space  $\mathcal{C}_{\text{rap}}^\alpha(\mathbb{R}^d)$  is in particular not empty.

We use a modified paraproduct in Section 3.1.3. Note that

$$|\nabla|^m f := \mathcal{F}^{-1}(|\cdot|^m \mathcal{F} f),$$

for  $m \in \mathbb{Z}$ , is well-defined for functions  $f \in \mathcal{S}(\mathbb{R}^d)$  whose Fourier transform have support in an annulus. For  $m \in \mathbb{N}$  and  $\alpha \in \mathbb{R}$ , the map  $|\nabla|^m$  sends continuously  $\mathcal{C}_{\text{rap}}^\alpha(\mathbb{R}^d)$  into  $\mathcal{C}_{\text{rap}}^{\alpha-m}(\mathbb{R}^d)$ . For  $m \in \mathbb{N}$ , we define the *modified paraproduct*

$$P_f^m g := |\nabla|^m (P_f |\nabla|^{-m} g) = \sum_{j=1}^{\infty} |\nabla|^m (S_j f \cdot |\nabla|^{-m} \Delta_j g).$$

Note that  $P^0 = P$ . The first item of Proposition 8 also holds for the modified paraproduct  $P^m$ . This modified paraproduct will play a pivotal role in the proof of Lemma 18, along the proof of Theorem 17. The latter provides the construction of a  $\mathbf{g}$ -map from bracket data.

### 3. FROM REGULARITY STRUCTURES AND MODELS TO PARACONTROLLED SYSTEMS

This section sets the scene and contains a proof of the first part of Theorem 1. We work in the setting of concrete regularity structures, a special case of regularity structures introduced in [5]. Their definition is recalled in Appendix A. As we work a priori with the most general concrete regularity structures, we need to identify a number of conditions that serve our purpose in Section 3.1. Assumption (A) is a harmless assumption on how polynomials sit inside  $T$  and  $T^+$ . Assumption (B) is a very mild requirement on the splitting map  $\Delta : T \rightarrow T \otimes T^+$ . Both assumptions are met by the regularity structures used in the study of singular PDEs. This is all we need to get a representation of models and reconstructions of modeled distributions by paracontrolled systems. Before embarking on the journey, recall from [5] that we use the notations

$$\Delta\sigma = \sum_{\mu \dot{\subset} \sigma} \mu \otimes \sigma/\mu, \quad \Delta^+\tau = \sum_{\nu \dot{\subset}^+ \tau} \nu \otimes \tau/^+\nu$$

to denote the action of the splitting map  $\Delta$  on  $T$  and the coproduct  $\Delta^+$  on  $T^+$  – see the comments following assumption (A). The notation  $\mu < \sigma$  will mean  $\mu \dot{\subset} \sigma$  and  $\mu \neq \sigma$ ; we shall make a similar use of the expression  $\nu <^+ \tau$ .

We shall introduce along the way three assumptions (A), (B), (C) on general regularity structures. Their meaning is to be understood in the light of what regularity structures are useful for: They encode the algebra at hand in the pointwise description of ‘irregular’ functions. One will for instance read assumption (A) as saying that the classically regular part of functions behave as in the classical Taylor calculus. Interpretations of assumptions (B) and (C) are given after their statement.

**3.1. A BASIC ASSUMPTION.** — Appendix A recalls elementary properties of concrete regularity structures. Let  $\overline{T} = ((T^+, \Delta^+), (T, \Delta))$  be a concrete regularity structure with  $T^+ = \bigoplus_{\alpha \in A^+} T_\alpha^+$  and  $T = \bigoplus_{\beta \in A} T_\beta$ . Write  $\mathbf{1}_+$  for the unit of the algebra  $T^+$ . Recall that we agree to use the shorthand notation  $\mathfrak{s}^{(+)}$  to mean both the statement  $\mathfrak{s}$  and the statement  $\mathfrak{s}^+$ .

**ASSUMPTION A.** — *The spaces  $T^+$  and  $T$  have linear bases  $\mathcal{B}^+$  and  $\mathcal{B}$ , respectively, with the following properties.*

(1)  $\mathcal{B}^+$  is a commutative monoid freely generated by a finite set  $\mathcal{B}_\circ^+$  and Taylor monomials  $X_1, \dots, X_d$ . Each element  $\tau \in \mathcal{B}_\circ^+$  has a positive homogeneity. For general elements in  $\mathcal{B}^+$ , homogeneities are defined by  $|X_i| = 1$ , and multiplicativity

$$|\tau\sigma| = |\tau| + |\sigma|.$$

(2) *The action of  $\Delta^+$  on polynomials is characterized by its action on the monomials*

$$(3.1) \quad \Delta^+ X_i = X_i \otimes \mathbf{1}_+ + \mathbf{1}_+ \otimes X_i,$$

that holds for all  $1 \dot{\subset} i \dot{\subset} d$ . Denote by  $\mathcal{B}_X^+$  the submonoid generated by  $X_1, \dots, X_d$ .

(3) There exists a subset  $\mathcal{B}_\bullet \subset \mathcal{B}$ , such that  $\mathcal{B}$  is in bijection with  $\mathbb{N}^d \times \mathcal{B}_\bullet$ . An element  $(k, \sigma) \in \mathbb{N}^d \times \mathcal{B}_\bullet$ , is denoted by  $\underline{X}^k \sigma$ , and assigned a homogeneity

$$|\underline{X}^k \sigma| := |k| + |\sigma|.$$

(4) If  $\mathcal{B}_\bullet$  contains an element  $\mathbf{1}$  with homogeneity 0, then it is unique and satisfies the identity

$$\Delta \mathbf{1} = \mathbf{1} \otimes \mathbf{1}_+.$$

Write  $\underline{X}^k$  for  $\underline{X}^k \mathbf{1}$ . Set

$$\mathcal{B}_X := \{\underline{X}^k\}_{k \in \mathbb{N}^d} \subset \mathcal{B}.$$

The coproduct  $\Delta$  on  $\underline{X}^k$  is characterized by its action on the monomials

$$\Delta X_i = X_i \otimes \mathbf{1}_+ + \mathbf{1} \otimes X_i,$$

where  $X_i = X^{e_i}$  and  $e_i = (\delta_{ij})_{j=1}^d \in \mathbb{N}^d$ , that holds for all  $1 \leq i \leq d$ , and by requiring multiplicativity on  $\mathcal{B}_X$ . For general elements, one has the multiplicative formula

$$\Delta(\underline{X}^k \sigma) = (\Delta \underline{X}^k)(\Delta \sigma).$$

For later use, denote by  $\{\tau'\}_{\tau \in \mathcal{B}}$  the dual basis of  $\mathcal{B}$ . Following [5], for  $\sigma, \tau \in \mathcal{B}^{(+)}$ , write  $\sigma \circ^{(+)} \tau$ , if  $\sigma$  appears in the left hand side of the tensor products in the optimal expansion of  $\Delta^{(+)} \tau$ , so we have the unique representation

$$\Delta^{(+)} \tau = \sum_{\substack{\sigma \in \mathcal{B}^{(+)} \\ \sigma \circ^{(+)} \tau}} \sigma \otimes (\tau /^{(+)} \sigma),$$

where  $\tau /^{(+)} \sigma \in T^+ \setminus \{0\}$ . The relation  $\circ$  needs not to be transitive. Using the coassociativity

$$(\Delta^{(+)} \otimes \text{Id}) \Delta^{(+)} \tau = (\text{Id} \otimes \Delta^{(+)}) \Delta^{(+)} \tau,$$

we obtain the chain formula

$$(3.2) \quad \Delta^+(\tau /^{(+)} \mu) = \sum_{\mu \circ^{(+)} \nu \circ^{(+)} \tau} (\nu /^{(+)} \mu) \otimes (\tau /^{(+)} \nu).$$

Write  $\sigma <^{(+)} \tau$ , if  $\sigma \circ^{(+)} \tau$  and  $\sigma \neq \tau$ . The notations  $\tau /^{(+)} \sigma$  and  $\sigma <^{(+)} \tau$  are only used for  $\tau$  and  $\sigma$  in  $\mathcal{B}^{(+)}$ . Be careful! The notations  $\circ, <$ , etc. are basis-dependent – like the matrix of a linear map.

The following structural assumption simplifies some arguments in this paper.

#### ASSUMPTION B

- (1) For each  $\tau, \sigma \in \mathcal{B}$  with  $\sigma < \tau$ , either  $\tau / \sigma \in \text{span}(\mathcal{B}_X^+)$  or  $\tau / \sigma \in \text{span}(\mathcal{B}^+ \setminus \mathcal{B}_X^+)$ .
- (2) For any  $\tau \in \mathcal{B}^+ \setminus \mathcal{B}_X^+$  and  $\sigma \in \mathcal{B}_X^+$ ,  $\tau / \sigma \in \text{span}(\mathcal{B}^+ \setminus \mathcal{B}_X^+)$ .
- (3) For any  $\tau \in \mathcal{B} \setminus \mathcal{B}_X$  and  $\sigma \in \mathcal{B}_X$ ,  $\tau / \sigma \in \text{span}(\mathcal{B}^+ \setminus \mathcal{B}_X^+)$ .

Assumption (B) is about the distinction between polynomial and non-polynomial elements. Assumption B(1) means that, in the expansion of  $\Delta \tau$ , there is no term of the form  $\sigma \otimes (\eta + X^k)$ , with  $\eta \in \text{span}(\mathcal{B}^+ \setminus \mathcal{B}_X^+)$  and  $k \in \mathbb{N}^d$ . It is used to justify the quantity  $\llbracket \mu / \tau \rrbracket^{\#}$  in the formula (3.7) below. Assumptions B(2) and B(3) are needed in the proof of Theorem 12. BHZ regularity structure satisfies assumption (B), since

polynomial and non-polynomial elements are obviously distinguished by the number of their edges. See Section 4.3 for details.

A natural way to ensure (B) is to give homogeneities not in  $\mathbb{N}$  for the non-polynomial elements. Hence the following is one of the sufficient conditions.

**ASSUMPTION B'.** — *Homogeneities of elements in  $\mathcal{B}^+ \rceil \mathcal{B}_X^+$  and  $\mathcal{B} \rceil \mathcal{B}_X$  are not nonnegative integers.*

This is a kind of natural assumption on regularity structures associated with PDEs; [14, Ass. 5.3] is a part of assumption (B') for elements in  $\mathcal{B} \rceil \mathcal{B}_X$ . Under assumption (B) write, for  $\tau \in \mathcal{B}^+$ ,

$$\begin{aligned}
 \Delta^+ \tau &= \sum_{\sigma \in \mathcal{B}^+ \rceil \mathcal{B}_X^+} \sigma \otimes (\tau /^+ \sigma) + \sum_k X^k \otimes (\tau /^+ X^k) \\
 (3.3) \quad &=: \sum_{\sigma \in \mathcal{B}^+ \rceil \mathcal{B}_X^+} \sigma \otimes (\tau /^+ \sigma) + \sum_k \frac{X^k}{k!} \otimes D^k \tau.
 \end{aligned}$$

Extend by linearity the map  $D^k$  from  $T_\alpha^+$  to  $T_{\alpha-|k|}^+$ , for all  $\alpha \in A$ .

**LEMMA 9.** — *Under assumptions (A) and (B) one has, for all  $\sigma, \tau \in T^+$  and all  $k, \ell \in \mathbb{N}^d$ ,*

- (a)  $D^0 \tau = \tau$ ,
- (b) if  $\tau \in \mathcal{B}^+ \rceil \mathcal{B}_X^+$ , then  $D^k \tau \in \text{span}(\mathcal{B}^+ \rceil \mathcal{B}_X^+)$ ; moreover,  $D^k \tau \neq 0$  only if  $|k| < |\tau|$ ,
- (c)  $D^k D^\ell \tau = D^{k+\ell} \tau$ ,
- (d)  $D^k X^\ell = \mathbf{1}_{k \triangleleft \ell} \frac{\ell!}{(\ell-k)!} X^{\ell-k}$ ,
- (e)  $D^k(\tau\sigma) = \sum_{k'} \binom{k}{k'} D^{k'} \tau D^{k-k'} \sigma$  — Leibniz rule,

*Proof.* — Item (a) comes from the property (A.1) satisfied by coproducts in Hopf algebras, recalled in Appendix A. The former part of item (b) is a consequence of assumption B(2). The latter part comes from the property (A.1). Since  $|\tau| > 0$  by the definition of concrete regularity structures, the term  $(X^k/k!) \otimes D^k \tau$  appears in the expansion of  $\Delta^+ \tau$  only if  $|k| < |\tau|$ . Item (c) is a consequence of the coassociativity property

$$(\Delta^+ \otimes \text{Id})\Delta^+ = (\text{Id} \otimes \Delta^+)\Delta^+$$

of the coproduct  $\Delta^+$ . Expanding both sides at  $\tau \in \mathcal{B}^+$ , we have

$$\begin{aligned}
 &\sum_{\sigma, \eta \in \mathcal{B}^+ \rceil \mathcal{B}_X^+} \eta \otimes (\sigma /^+ \eta) \otimes (\tau /^+ \sigma) \\
 &+ \sum_{\substack{\sigma \in \mathcal{B}^+ \rceil \mathcal{B}_X^+ \\ k \in \mathbb{N}^d}} \frac{X^k}{k!} \otimes D^k \sigma \otimes (\tau /^+ \sigma) + \sum_{k, \ell \in \mathbb{N}^d} \frac{X^k}{k!} \otimes \frac{X^\ell}{\ell!} \otimes D^{k+\ell} \tau \\
 &= \sum_{\sigma \in \mathcal{B}^+ \rceil \mathcal{B}_X^+} \sigma \otimes \Delta^+(\tau /^+ \sigma) + \sum_{k \in \mathbb{N}^d} \frac{X^k}{k!} \otimes \Delta^+ D^k \tau.
 \end{aligned}$$



It gives indeed the identity

$$(3.4) \quad \Delta^+ D^k \tau = \sum_{\sigma \zeta^+ \tau, \sigma \notin \mathcal{B}_x^+} D^k \sigma \otimes (\tau / \zeta^+ \sigma) + \sum_{\ell \in \mathbb{N}^d} \frac{X^\ell}{\ell!} \otimes D^{k+\ell} \tau,$$

this means (c). Item (d) is a direct consequence of the Leibniz formula for the polynomials, which follows from identity (3.1) giving the action of  $\Delta^+$  on  $X_i$  and the multiplicativity property of  $\Delta^+$ . Item (e) is again a consequence of the multiplicativity property of  $\Delta^+$ .

**3.2. FROM MODELS TO PARACONTROLLED SYSTEMS.** — We recall in this section some of the results proved in [5], stated here in the slightly more general setting of the present work. The proofs of these extensions are given in Appendix B. These results are proved in [5, §§2 & 3] without any extra assumptions about ‘bounded polynomials’ and interaction between  $T^+$  and  $T$ . Hence the proofs are completely parallel to the proofs in [5], except for the use here of the modified paraproduct and the weight.

Given Fréchet spaces  $E$  and  $F$ , denote by  $L(E, F)$  the space of continuous linear maps from  $E$  into  $F$ . Recall  $G^+$  stands for the set of characters of the Hopf algebra  $T^+$ . Given maps

$$\mathbf{g} : \mathbb{R}^d \longrightarrow G^+, \quad \Pi \in L(T, \mathcal{S}'(\mathbb{R}^d)),$$

and  $x, y \in \mathbb{R}^d$ , set

$$\mathbf{g}_{yx} := (\mathbf{g}_y \otimes \mathbf{g}_x^{-1}) \Delta^+ \in G^+,$$

and

$$\Pi_x^{\mathbf{g}} := (\Pi \otimes \mathbf{g}_x^{-1}) \Delta \in L(T, \mathcal{S}'(\mathbb{R}^d)).$$

Set

$$\beta_0 := \min A,$$

where  $A$  is a homogeneity set of  $T = \bigoplus_{\beta \in A} T_\beta$ .

**DEFINITION 10.** — Let a concrete regularity structure  $T$  satisfying assumption (A) be given. We denote by

$$\mathcal{M}_{\text{rap}}(T, \mathbb{R}^d),$$

the set of pairs of maps

$$\mathbf{g} : \mathbb{R}^d \longrightarrow G^+, \quad \Pi \in L(T, \mathcal{S}'(\mathbb{R}^d)),$$

such that

- (a) one has  $\mathbf{g}_x(X^k) = x^k$ , for all  $x \in \mathbb{R}^d, k \in \mathbb{N}^d$ ;
- (b) for any  $\tau \in \mathcal{B}_0^+$ , the function  $x \mapsto \mathbf{g}_x(\tau)$  belongs to  $L_{\text{rap}}^\infty(\mathbb{R}^d)$ , and the function

$$(x, y) \longmapsto \mathbf{g}_{yx}(\tau),$$

belongs to  $\mathcal{C}_{(2), \text{rap}}^{|\tau|}(\mathbb{R}^d \times \mathbb{R}^d)$ ;

- (c) one has  $(\Pi \underline{X}^k \sigma)(x) = x^k (\Pi \sigma)(x)$  and  $(\Pi \mathbf{1})(x) = 1$ ;

(d) for any  $\tau \in \mathcal{B}_\bullet \setminus \{\mathbf{1}\}$ , one has  $\Pi \tau \in \mathcal{C}_{\text{rap}}^{\beta_0}(\mathbb{R}^d)$ , and the  $\mathbb{R}^d$ -indexed family of distributions  $(\Pi_x^{\mathbf{g}} \tau)_{x \in \mathbb{R}^d}$  belongs to  $D_{\text{rap}}^{|\tau|}$ .

The pair  $(\mathbf{g}, \Pi)$  is called a *rapidly decreasing model* on the concrete regularity structure  $T$ .

We define metrics on the space of rapidly decreasing models on  $T$  setting

$$d_a((\mathbf{g}, \Pi), (\mathbf{g}', \Pi')) := \sup_{\tau \in \mathcal{B}_\bullet^+} \left( \|(\mathbf{g} - \mathbf{g}')(\tau)\|_{L_a^\infty(\mathbb{R}^d)} + \|(\mathbf{g} - \mathbf{g}')(\tau)\|_{\mathcal{C}_{(2),a}^{|\tau|}(\mathbb{R}^d \times \mathbb{R}^d)} \right) + \sup_{\sigma \in \mathcal{B}_\bullet} \left( \|(\Pi - \Pi')\sigma\|_{\mathcal{C}_a^{\beta_0}(\mathbb{R}^d)} + \|((\Pi^\mathbf{g} - (\Pi')^\mathbf{g}')\sigma)(\cdot)\|_{D_a^{|\sigma|}} \right).$$

With a slight abuse of notations, we write

$$\mathbf{g}_x(\tau) \in L_{\text{rap}}^\infty(\mathbb{R}^d), \quad \mathbf{g}_{yx}(\tau) \in \mathcal{C}_{(2),\text{rap}}^{|\tau|}(\mathbb{R}^d \times \mathbb{R}^d).$$

The definition of a model depends on the choice of subspaces  $\text{span}(\mathcal{B}^+ \cup \mathcal{B}_X^+)$  and  $\text{span}(\mathcal{B} \cup \mathcal{B}_X)$ , but not on the choice of their bases. Indeed, since

$$\mathbf{g}_x(X^k) \in L_{\text{slow}}^\infty(\mathbb{R}^d), \quad \mathbf{g}_{yx}(X^k) \in \mathcal{C}_{(2)}^{|k|}(\mathbb{R}^d \times \mathbb{R}^d)$$

and since

$$L_{\text{slow}}^\infty(\mathbb{R}^d) \cdot L_{\text{rap}}^\infty(\mathbb{R}^d) \subset L_{\text{rap}}^\infty(\mathbb{R}^d)$$

and 
$$\mathcal{C}_{(2)}^\alpha(\mathbb{R}^d \times \mathbb{R}^d) \cdot \mathcal{C}_{(2),\text{rap}}^\beta(\mathbb{R}^d \times \mathbb{R}^d) \subset \mathcal{C}_{(2),\text{rap}}^{\alpha+\beta}(\mathbb{R}^d \times \mathbb{R}^d),$$

for all non-negative  $\alpha, \beta$ , condition (b) holds for any  $\tau \in \mathcal{B}^+ \cup \mathcal{B}_X^+$ . Recall that the set  $\mathcal{B}_\bullet$  in item (d) of Definition 10 stands for the index parameterizing the non-polynomial part of the basis of  $T$ . It is not so obvious to see whether condition (d) holds or not for any  $\tau \in \mathcal{B} \cup \mathcal{B}_X$ ; however, the following lemma holds.

**LEMMA 11**

Assume (A) and (B). Under the condition (c), the estimate  $(\Pi_x^\mathbf{g}\tau)_{x \in \mathbb{R}^d} \in D_{\text{rap}}^{|\tau|}$  holds for any  $\tau \in \mathcal{B} \cup \mathcal{B}_X$ .

*Proof.* — We prove the estimate for  $\underline{X}^k \tau$ , with  $\tau \in \mathcal{B}_\bullet$ ,  $k \in \mathbb{N}^d \cup \{0\}$ . Because of the multiplicative property in assumption A(4) and item (c) in Definition 10, we have

$$\Pi_x^\mathbf{g} \underline{X}^k \tau = (\cdot - x)^k \Pi_x^\mathbf{g} \tau.$$

Recalling the notations at the beginning of Section 2, we have

$$S_j(\Pi_x^\mathbf{g} \underline{X}^k \tau)(x) = \int P_j(x - y)(y - x)^k (\Pi_x^\mathbf{g} \tau)(y) dy = \int P_j^k(x - y)(\Pi_x^\mathbf{g} \tau)(y) dy,$$

where  $P_j^k(x) := (-x)^k P_j(x)$ . Hence  $P_{j+1} * P_j^k = P_j^k$  by the property of support of Fourier transform, and we have

$$S_j(\Pi_x^\mathbf{g} \underline{X}^k \tau)(x) = P_{j+1} * P_j^k * (\Pi_x^\mathbf{g} \tau)(x) = \int P_j^k(x - y) S_{j+1}(\Pi_x^\mathbf{g} \tau)(y) dy.$$

By induction, we assume the required estimate for any  $\sigma \dot{\subset} \tau$  (hence  $\sigma = \tau$  or  $\sigma = \underline{X}^k \eta$  with some  $k \neq 0$  and  $\eta \in \mathcal{B}_\bullet$  such that  $|\eta| < |\tau|$ ). Since either of  $\sigma$  or  $\tau/\sigma$  is non-polynomial by assumption (B), for any  $a \in \mathbb{N}$ ,

$$\begin{aligned} |S_{j+1}(\Pi_x^{\mathbf{g}} \tau)(y)| &= |S_{j+1}(\Pi_y^{\mathbf{g}} \widehat{\mathbf{g}}_{yx} \tau)(y)| \dot{\subset} \sum_{\sigma \dot{\subset} \tau} |\mathbf{g}_{yx}(\tau/\sigma)| |S_{j+1}(\Pi_y^{\mathbf{g}} \sigma)(y)| \\ &\cdot (|x|_*^{-a} + |y|_*^{-a}) \sum_{\sigma \dot{\subset} \tau} |y-x|^{|\tau|-|\sigma|} 2^{-j|\sigma|}. \end{aligned}$$

By using the scaling property  $P_j^k(x) = 2^{j(d-|k|)} P_j^k(2^j x)$  and by a similar argument to Lemma 29 in Appendix B, we can conclude that

$$|S_j(\Pi_x^{\mathbf{g}} \underline{X}^k \tau)(x)| \cdot |x|_*^{-a} 2^{-j(|\tau|+|k|)},$$

hence  $(\Pi_x^{\mathbf{g}} \underline{X}^k \tau)_{x \in \mathbb{R}^d} \in D_{\text{rap}}^{|\underline{X}^k \tau|}$ .

The next statement is a variation on [5, Prop. 12], where we use now the usual polynomials and polynomial weights, and the modified paraproducts  $\mathbf{P}^m$  instead of the bounded polynomials, no weights and the usual paraproduct  $\mathbf{P}$ . Its proof is given in Appendix B.

**THEOREM 12.** — *Let  $\mathcal{T}$  stand for a regularity structure satisfying assumptions (A) and (B). Pick  $m \in \mathbb{N}$ . For any model  $\mathbf{M} = (\mathbf{g}, \Pi) \in \mathcal{M}_{\text{rap}}(\mathcal{T}, \mathbb{R}^d)$ , there exists a family*

$$\left( ([\tau]^{m, \mathbf{g}} \in \mathcal{C}_{\text{rap}}^{|\tau|}(\mathbb{R}^d))_{\tau \in \mathcal{B}^+ \cap \mathcal{B}_X^+}, ([\sigma]^{m, \mathbf{M}} \in \mathcal{C}_{\text{rap}}^{|\sigma|}(\mathbb{R}^d))_{\sigma \in \mathcal{B} \cap \mathcal{B}_X} \right)$$

such that one has, for any  $\tau \in \mathcal{B}^+ \cap \mathcal{B}_X^+$  and  $\sigma \in \mathcal{B} \cap \mathcal{B}_X$ , the identities

$$(3.5) \quad \mathbf{g}(\tau) = \sum_{\substack{\mathbf{1}_+ < \nu < \tau \\ \nu \in \mathcal{B}^+ \cap \mathcal{B}_X^+}} \mathbf{P}_{\mathbf{g}(\tau/\nu)}^m [[\nu]^{m, \mathbf{g}} + [\tau]^{m, \mathbf{g}}],$$

$$(3.6) \quad \Pi \sigma = \sum_{\substack{\mu < \sigma \\ \mu \in \mathcal{B} \cap \mathcal{B}_X}} \mathbf{P}_{\mathbf{g}(\sigma/\mu)}^m [[\mu]^{m, \mathbf{M}} + [\sigma]^{m, \mathbf{M}}].$$

Moreover, the mapping

$$\mathbf{M} \mapsto \left( ([\tau]^{m, \mathbf{g}} \in \mathcal{C}_{\text{rap}}^{|\tau|}(\mathbb{R}^d))_{\tau \in \mathcal{B}^+ \cap \mathcal{B}_X^+}, ([\sigma]^{m, \mathbf{M}} \in \mathcal{C}_{\text{rap}}^{|\sigma|}(\mathbb{R}^d))_{\sigma \in \mathcal{B} \cap \mathcal{B}_X} \right)$$

is locally Lipschitz continuous.

This version of the statement, with  $m > 1$ , will be used in the proof of Theorem 17 given in Section 4.1. Write  $[[\tau]^{m, \mathbf{g}}$  and  $[[\sigma]^{m, \mathbf{M}}$  instead of  $[\tau]^{m, \mathbf{g}}$  and  $[\sigma]^{m, \mathbf{M}}$ , when  $m = 0$ .

Given a model  $\mathbf{M} \in \mathcal{M}_{\text{rap}}(\mathcal{T}, \mathbb{R}^d)$  on a regularity structure  $\mathcal{T}$ , and  $\gamma \in \mathbb{R}$ , define the space  $\mathcal{D}_{\text{rap}}^\gamma(\mathcal{T}, \mathbf{g})$  of rapidly decreasing modeled distributions as the set of functions

$$\mathbf{f} : \mathbb{R}^d \longrightarrow \bigoplus_{\beta < \gamma} T_\beta,$$

such that, for each  $\tau \in \mathcal{B}$ , the function  $\langle \tau', \mathbf{f}(\cdot) \rangle$  belongs to  $L_{\text{rap}}^\infty(\mathbb{R}^d)$ , and the function

$$(x, y) \mapsto \langle \tau', \mathbf{f}(y) - \widehat{\mathbf{g}}_{yx} \mathbf{f}(x) \rangle$$

belongs to  $\mathcal{C}_{(2),\text{rap}}^{\gamma-|\tau|}(\mathbb{R}^d \times \mathbb{R}^d)$ . The reconstruction  $\mathbf{Rf}$  of  $\mathbf{f} \in \mathcal{D}_{\text{rap}}^\gamma(T, \mathbf{g})$  is an element of  $S'(\mathbb{R}^d)$  satisfying the condition

$$(\mathbf{Rf} - \Pi_x^\mathbf{g} \mathbf{f}(x))_{x \in \mathbb{R}^d} \in D_{\text{rap}}^\gamma.$$

If  $\gamma > 0$ , there exists exactly one reconstruction. If  $\gamma < 0$ , there are infinitely many reconstructions and two reconstructions are equal modulo  $\mathcal{C}_{\text{rap}}^\gamma(\mathbb{R}^d)$ . (This is a key point to prove Proposition 15.) In what follows, we assume  $\gamma \neq 0$  and denote  $\mathbf{Rf}$  by the one defined in Corollary 33 in Appendix B. If  $\gamma = 0$ , existence of the reconstruction is not ensured in general. See [8, Ex. 5.5].

The next statement was proved in [5, Th. 14] in the unweighted setting; its extension to the present setting is given in Appendix B.

**THEOREM 13.** — *Let  $T$  be a regularity structure satisfying assumptions (A) and (B). Let a regularity exponent  $\gamma \in \mathbb{R} \setminus \{0\}$  and a model  $\mathbf{M} = (\mathbf{g}, \Pi) \in M_{\text{rap}}(T, \mathbb{R}^d)$  on  $T$  be given. For any modeled distribution*

$$\mathbf{f} = \sum_{|\sigma| < \gamma} f_\sigma \sigma \in \mathcal{D}_{\text{rap}}^\gamma(T, \mathbf{g}),$$

each coefficient  $f_\sigma$  has a paracontrolled representation

$$(3.7) \quad f_\sigma = \sum_{\substack{\sigma < \mu \\ \mu/\sigma \in \text{span}(\mathbb{B}^{+\Gamma} \mathbb{B}_X^+)}} P_{f_\mu} \llbracket \mu/\sigma \rrbracket^\mathbf{g} + \llbracket f_\sigma \rrbracket^\mathbf{g},$$

where  $\llbracket f_\sigma \rrbracket^\mathbf{g} \in \mathcal{C}_{\text{rap}}^{\gamma-|\sigma|}(\mathbb{R}^d)$ . (The quantity  $\llbracket \mu/\sigma \rrbracket^\mathbf{g}$  is defined as a linear extension of the symbols  $\llbracket \tau \rrbracket^\mathbf{g}$  in Theorem 12.) Moreover, there exists a distribution  $\llbracket \mathbf{f} \rrbracket^\mathbf{M} \in \mathcal{C}_{\text{rap}}^\gamma(\mathbb{R}^d)$  such that

$$(3.8) \quad \mathbf{Rf} = \sum_{\sigma \in \mathbb{B}^+ \mathbb{B}_X} P_{f_\sigma} \llbracket \sigma \rrbracket^\mathbf{M} + \llbracket \mathbf{f} \rrbracket^\mathbf{M}.$$

The mapping

$$(\mathbf{f} \in \mathcal{D}_{\text{rap}}^\gamma(T, \mathbf{g})) \longmapsto \left( \llbracket \mathbf{f} \rrbracket^\mathbf{M}, (\llbracket f_\sigma \rrbracket^\mathbf{g})_{\sigma \in \mathbb{B}} \right) \in \mathcal{C}_{\text{rap}}^\gamma(\mathbb{R}^d) \times \prod_{\sigma \in \mathbb{B}} \mathcal{C}_{\text{rap}}^{\gamma-|\sigma|}(\mathbb{R}^d)$$

is locally Lipschitz continuous.

A similar statement with  $P^m$  used in place of  $P$  holds true. We end this section with three useful formulas involving  $\mathbf{g}$ , that will be used in the proof of Theorem 17. The reader can skip this statement now and come back to it at the moment where it is needed. Recall  $D^k \tau = 0$ , for  $|k| > |\tau|$ . Let  $P_X : T^+ \rightarrow T_X^+$ , stand for the canonical projection map on  $T_X^+$ , and for  $\tau \in \mathbb{B}^+$  set

$$f_x(\tau) := -(\mathbf{g}_x \otimes \mathbf{g}_x^{-1})(P_X \otimes \text{Id})(\Delta^+ \tau) = - \sum_{\ell} \frac{x^\ell}{\ell!} \mathbf{g}_x^{-1}(D^\ell \tau).$$

For  $\tau \neq \mathbf{1}_+$ , we also have

$$f_x(\tau) := (\mathbf{g}_x \otimes \mathbf{g}_x^{-1})((\text{Id} - P_X) \otimes \text{Id})(\Delta^+ \tau) = \sum_{\substack{\sigma \dot{\subset}^+ \tau \\ \sigma \notin \mathcal{B}_X^+}} \mathbf{g}_x(\sigma) \mathbf{g}_x^{-1}(\tau / + \sigma).$$

LEMMA 14. — *Let  $T$  be a regularity structure satisfying assumption (A) and (B). For any  $\tau \in \mathcal{B}^+ \cap \mathcal{B}_X^+$  and any  $k \in \mathbb{N}^d$ , we have*

$$(3.9) \quad \mathbf{g}_x(D^k \tau) = \sum_{\substack{\sigma \dot{\subset}^+ \tau \\ \sigma \notin \mathcal{B}_X^+}} \mathbf{g}_x(\tau / + \sigma) f_x(D^k \sigma).$$

and

$$(3.10) \quad \mathbf{g}_{yx}(D^k \tau) = \sum_{\substack{\sigma \dot{\subset}^+ \tau \\ \sigma \notin \mathcal{B}_X^+}} \mathbf{g}_{yx}(\tau / + \sigma) f_y(D^k \sigma) - \sum_{\ell} \frac{(y-x)^\ell}{\ell!} f_x(D^{k+\ell} \tau),$$

and

$$(3.11) \quad \begin{aligned} f_x(D^k \tau) &= \mathbf{1}_{|k| < |\tau|} \partial_y^k \left\{ (\mathbf{g}_y \otimes \mathbf{g}_x^{-1})((\text{Id} - P_X) \otimes \text{Id}) \Delta^+ \tau \right\} \Big|_{y=x} \\ &= \mathbf{1}_{|k| < |\tau|} \partial_y^k \left\{ \sum_{\substack{\sigma \dot{\subset}^+ \tau \\ \sigma \notin \mathcal{B}_X^+}} \mathbf{g}_y(\sigma) \mathbf{g}_x^{-1}(\tau / + \sigma) \right\} \Big|_{y=x}. \end{aligned}$$

Note that one cannot interchange in (3.11) the derivative operator with the sum, as a given function  $\mathbf{g}_y(\sigma)$  may not be sufficiently regular to be differentiated  $k$  times. Note that formula (3.10) does not have the classical feature of a Taylor-type expansion formula, which would rather involve an  $x$ -dependent term in front of  $\mathbf{g}_{yx}(\tau / + \sigma)$ , in the first term of the right hand side.

*Proof.* — — Note first that formula (3.4) for  $\Delta^+(D^k \tau)$  gives

$$(3.12) \quad f_x(D^k \tau) = (\mathbf{g}_x \otimes \mathbf{g}_x^{-1})((\text{Id} - P_X) \otimes \text{Id}) \Delta^+ D^k \tau = \sum_{\substack{\nu \dot{\subset}^+ \tau \\ \nu \notin \mathcal{B}_X^+}} \mathbf{g}_x(D^k \nu) \mathbf{g}_x^{-1}(\tau / + \nu).$$

Formula (3.9) is an inversion formula for the preceding identity. One obtains the former from the latter by writing

$$\begin{aligned} \sum_{\substack{\sigma \dot{\subset}^+ \tau \\ \sigma \notin \mathcal{B}_X^+}} \mathbf{g}_x(\tau / + \sigma) f_x(D^k \sigma) &= \sum_{\substack{\nu \dot{\subset}^+ \sigma \dot{\subset}^+ \tau \\ \sigma, \nu \notin \mathcal{B}_X^+}} \mathbf{g}_x(\tau / + \sigma) \mathbf{g}_x^{-1}(\sigma / + \nu) \mathbf{g}_x(D^k \nu) \\ &= \sum_{\substack{\nu \dot{\subset}^+ \sigma \dot{\subset}^+ \tau \\ \nu \notin \mathcal{B}_X^+}} \mathbf{g}_x(\tau / + \sigma) \mathbf{g}_x^{-1}(\sigma / + \nu) \mathbf{g}_x(D^k \nu) \\ &= \sum_{\substack{\nu \dot{\subset}^+ \tau \\ \nu \notin \mathcal{B}_X^+}} (\mathbf{g}_x^{-1} \otimes \mathbf{g}_x)(\tau / + \nu) \mathbf{g}_x(D^k \nu) = \mathbf{g}_x(D^k \tau). \end{aligned}$$

(In the second equality, we can remove the condition " $\sigma \notin \mathcal{B}_X^+$ " because  $\nu \lessdot^+ X^k$  implies that  $\nu \in \mathcal{B}_X^+$ . In the last equality, we use the property of the antipode.)

– Applying  $\mathfrak{g}_y \otimes \mathfrak{g}_x^{-1}$  to (3.4), we have

$$\begin{aligned}
 \mathfrak{g}_{yx}(D^k \tau) &= \sum_{\substack{\mu \lessdot^+ \tau \\ \mu \notin \mathcal{B}_X^+}} \mathfrak{g}_y(D^k \mu) \mathfrak{g}_x^{-1}(\tau / + \mu) + \sum_{\ell} \frac{y^\ell}{\ell!} \mathfrak{g}_x^{-1}(D^{k+\ell} \tau) \\
 (3.13) \quad &= \sum_{\substack{\mu \lessdot^+ \nu \lessdot^+ \tau \\ \mu \notin \mathcal{B}_X^+}} \mathfrak{g}_y(D^k \mu) \mathfrak{g}_y^{-1}(\nu / + \mu) \mathfrak{g}_{yx}(\tau / + \nu) - \sum_{\ell'} \frac{(y-x)^{\ell'}}{\ell'!} \mathfrak{f}_x(D^{k+\ell'} \tau),
 \end{aligned}$$

where we use the formula (3.2) in the expansion of  $\mathfrak{g}_x^{-1}(\tau / + \mu)$ . Identity (3.10) follows from (3.13) using (3.12). Note that  $\mu \lessdot^+ \nu$  and  $\mu \notin \mathcal{B}_X^+$  implies that  $\nu \notin \mathcal{B}_X^+$ .

– Formula (3.11) comes from identity (3.12) by rewriting the terms  $\mathfrak{g}_x(D^k \nu)$  for  $\nu \in \mathcal{B}^+ \cap \mathcal{B}_X^+$  in an appropriate form. As a preliminary remark, notice that applying  $\mathfrak{g}_{yx} \otimes \mathfrak{g}_x$  to the defining identity (3.3) for the  $D^k \nu$ , we have

$$\mathfrak{g}_y(\nu) = \sum_{\sigma \lessdot^+ \nu, \sigma \notin \mathcal{B}_X^+} \mathfrak{g}_{yx}(\sigma) \mathfrak{g}_x(\nu / + \sigma) + \sum_k \mathfrak{g}_x(D^k \nu) \frac{(y-x)^k}{k!}.$$

Since one has

$$\partial_y^k \mathfrak{g}_{yx}(\sigma) \Big|_{y=x} = 0,$$

for any  $x \in \mathbb{R}^d$ , whenever  $|k| < |\sigma|$ , one then has

$$(3.14) \quad \mathfrak{g}_x(D^k \nu) = \mathbf{1}_{|k| < |\nu|} \partial_y^k \left\{ \mathfrak{g}_y(\nu) - \sum_{\substack{\sigma \lessdot^+ \nu \\ \sigma \notin \mathcal{B}_X^+ \\ |\sigma| \lessdot |k|}} \mathfrak{g}_{yx}(\sigma) \mathfrak{g}_x(\nu / + \sigma) \right\} \Big|_{y=x}.$$

At the same time, one has

$$\begin{aligned}
 \mathfrak{g}_y(\nu) &= \sum_{\mu \lessdot^+ \nu} (\mathfrak{g}_x^{-1} * \mathfrak{g}_x)(\nu / + \mu) \mathfrak{g}_y(\mu) = \sum_{\substack{\mu \lessdot^+ \nu \\ \mu \notin \mathcal{B}_X^+}} (\mathfrak{g}_x^{-1} * \mathfrak{g}_x)(\nu / + \mu) \mathfrak{g}_y(\mu) \\
 &= \sum_{\substack{\mu \lessdot^+ \sigma \lessdot^+ \nu \\ \mu, \sigma \notin \mathcal{B}_X^+}} \mathfrak{g}_x(\nu / + \sigma) \mathfrak{g}_x^{-1}(\sigma / + \mu) \mathfrak{g}_y(\mu).
 \end{aligned}$$

In the second equality, we use assumption B(2) to derive  $(\mathfrak{g}_x^{-1} * \mathfrak{g}_x)(\nu / + X^k) = 0$  for any  $k \in \mathbb{N}^d$ . In the third equality, we use that  $\mu \lessdot^+ \sigma$  and  $\mu \notin \mathcal{B}_X^+$  implies  $\sigma \notin \mathcal{B}_X^+$ . Furthermore, since  $\mu \lessdot^+ \sigma \notin \mathcal{B}_X^+$ ,  $\mu \in \mathcal{B}_X^+$ , and  $|\sigma| \lessdot |k|$  implies  $\mu \lessdot^+ \sigma$  (hence  $|\mu| < |k|$ ), we have

$$\sum_{\substack{\mu \lessdot^+ \sigma \lessdot^+ \nu \\ \mu, \sigma \notin \mathcal{B}_X^+, |\sigma| \lessdot |k|}} \mathfrak{g}_x(\nu / + \sigma) \mathfrak{g}_x^{-1}(\sigma / + \mu) \mathfrak{g}_y(\mu) = \sum_{\substack{\sigma \lessdot^+ \nu \\ \sigma \notin \mathcal{B}_X^+, |\sigma| \lessdot |k|}} \mathfrak{g}_x(\nu / + \sigma) \mathfrak{g}_{yx}(\sigma) + p_{<|k|}(y),$$

where  $p_{<|k|}$  is a polynomial of degree less than  $|k|$ , hence  $\partial_y^k p_{<|k|} = 0$ . We thus obtain from formula (3.14), that

$$\mathbf{g}_x(D^k \nu) = \mathbf{1}_{|k| < |\nu|} \partial_y^k \left\{ \sum_{\substack{\mu \in \mathcal{B}_X^+ \\ \sigma \in \mathcal{B}_X^+ \\ |\sigma| > |k|}} \mathbf{g}_x(\nu / + \sigma) \mathbf{g}_x^{-1}(\sigma / + \mu) \mathbf{g}_y(\mu) \right\} \Big|_{y=x}.$$

Inserting this expression in formula (3.12) one gets,

$$\begin{aligned} \mathbf{f}_x(D^k \tau) &= \sum_{\substack{\nu \in \mathcal{B}_X^+ \\ \nu \notin \mathcal{B}_X^+}} \mathbf{g}_x^{-1}(\tau / + \nu) \mathbf{g}_x(D^k \nu) \\ &= \partial_y^k \left\{ \sum_{\substack{\mu, \sigma, \nu \in \mathcal{B}_X^+ \\ |\sigma| > |k|}} \mathbf{g}_x^{-1}(\tau / + \nu) \mathbf{g}_x(\nu / + \sigma) \mathbf{g}_x^{-1}(\sigma / + \mu) \mathbf{g}_y(\mu) \right\} \Big|_{y=x} \\ &= \partial_y^k \left\{ \sum_{\substack{\mu \in \mathcal{B}_X^+ \\ |\sigma| > |k|}} \mathbf{g}_x^{-1}(\tau / + \nu) \mathbf{g}_x(\nu / + \sigma) \mathbf{g}_x^{-1}(\sigma / + \mu) \mathbf{g}_y(\mu) \right\} \Big|_{y=x} \\ &= \partial_y^k \left\{ \sum_{\substack{\mu, \sigma \in \mathcal{B}_X^+ \\ |\sigma| > |k|}} \mathbf{g}_x * \mathbf{g}_x^{-1}(\tau / + \sigma) \mathbf{g}_x^{-1}(\sigma / + \mu) \mathbf{g}_y(\mu) \right\} \Big|_{y=x} \\ &= \mathbf{1}_{|k| < |\tau|} \partial_y^k \left\{ \sum_{\mu \in \mathcal{B}_X^+} \mathbf{g}_x^{-1}(\tau / + \mu) \mathbf{g}_y(\mu) \right\} \Big|_{y=x}. \end{aligned}$$

In the third line, we can omit the condition  $\sigma, \nu \notin \mathcal{B}_X^+$  because of  $\mu \notin \mathcal{B}_X^+$ . In the last line, we use that  $(\mathbf{g}_x * \mathbf{g}_x^{-1})(\tau / + \sigma) = 1$  if and only if  $\sigma = \tau$ .

#### 4. FROM PARACONTROLLED SYSTEMS TO MODELS AND MODELED DISTRIBUTIONS

We prove the main results of this work in this section. Theorem 1 gives a parameterization of the space of models by ‘bracket’ data in paracontrolled representations. The main part of the work consists in building a  $\mathbf{g}$ -map from a paracontrolled representation for it on a minimal subset of a linear basis of  $T^+$ . Assumption (C) below gives a structural assumption on  $T^+$  that identifies this minimal set. The proofs of Corollary 2 on the density of smooth models, Corollary 3 and Corollary 4 on extension problems, are proved in Section 4.4.

Theorem 5 provides a parameterization of the space of modeled distributions of regularity  $\gamma$ , for a fixed  $\gamma \in \mathbb{R}$ , by a product of Hölder spaces. It is proved in Section 4.2. On a technical level, one brings back the proof of Theorem 5 to an extension problem for the  $\mathbf{g}$ -map from the Hopf algebra  $T^+$  to a larger Hopf algebra  $T_{\mathbf{F}}^+$ . This allows to see Theorem 5 as a corollary of Theorem 1 under the additional assumption (D).

Unlike the other assumptions, assumption (D) is about a basis  $\mathcal{B}$  of  $T$  rather than about  $T$  itself. It is thus possible that a given basis satisfies assumption (D) whereas another does not. This flexibility is at the heart of the proof of Theorem 7, dealing with the case of BHZ regularity structures, investigated in Section 4.3. Those regularity

structures introduced by Bruned, Hairer and Zambotti in [7] provide the universal model of regularity structures associated with a subcritical singular stochastic PDE.

4.1. FROM PARACONTROLLED SYSTEMS TO MODELS. — The following claim is the same as [5, Cor. 15], with the modified paraproduct  $P^m$  in the role of  $P$ . Recall from Theorem 12 the definition of the reference distributions  $[[\sigma]]^{m,M}$ , in the paracontrolled representation of the  $\Pi$  operator of a model  $M$ , using the modified paraproduct  $P^m$ .

PROPOSITION 15. — *Let  $T$  be a regularity structure satisfying assumptions (A) and (B). Pick  $m \in \mathbb{N}$ , and assume we are given a map  $g : \mathbb{R}^d \rightarrow G^+$ , such that conditions (a) and (b) in Definition 10 are satisfied. Then for any family  $([[\tau]] \in \mathcal{C}_{\text{rap}}^{|\tau|}(\mathbb{R}^d))_{\tau \in \mathcal{B}_\bullet, |\tau| < 0}$ , there exists a unique model  $M = (g, \Pi) \in M_{\text{rap}}(T, \mathbb{R}^d)$  such that*

$$(4.1) \quad \Pi_\tau = \sum_{\sigma < \tau} P_{g(\tau/\sigma)}^m [[\sigma]]^{m,M} + [[\tau]], \quad \forall \tau \in \mathcal{B}_\bullet, |\tau| < 0.$$

The map

$$\left( g, \left( [[\tau]] \in \mathcal{C}_{\text{rap}}^{|\tau|}(\mathbb{R}^d) \right)_{\tau \in \mathcal{B}_\bullet, |\tau| < 0} \right) \mapsto M \in M_{\text{rap}}(T, \mathbb{R}^d)$$

is continuous.

Note that the distributions  $[[\sigma]]^{m,M}$  in (4.1) are recursively defined by application of Theorem 12 to the subspace  $\bigoplus_{\beta < |\tau|} T_\beta$ . If  $\sigma \in \mathcal{B}_\bullet$  with  $|\sigma| < 0$ , then  $[[\sigma]]^{m,M} = [[\sigma]]$ .

Proof

– Recall there is no other element than  $\mathbf{1}$  of zero homogeneity in the present setting, and pick a basis vector  $\tau \in \mathcal{B}_\bullet$  with  $|\tau| < 0$ , and assume that  $(g, \Pi)$  is a model on the sector  $T_{<|\tau|}$ . Set for all  $x \in \mathbb{R}^d$

$$h_\tau(x) := \sum_{\sigma < \tau} g_x(\tau/\sigma)\sigma;$$

this defines a modeled distribution in  $\mathcal{D}_{\text{rap}}^{|\tau|}(T, g)$ . Then the bound  $(\Pi_x^g \tau)_{x \in \mathbb{R}^d} \in D_{\text{rap}}^{|\tau|}$  is equivalent to that  $\Pi_\tau$  is one of the reconstructions of  $h_\tau$ . From the version of Theorem 13 with the modified paraproduct  $P^m$ , the distribution

$$R h_\tau = \sum_{\sigma < \tau} P_{g(\tau/\sigma)}^m [[\sigma]]^{m,M} + [[h_\tau]]^{m,M}$$

is a reconstruction of  $h_\tau$ . Since

$$\Pi_\tau - R h_\tau = [[\tau]] - [[h_\tau]]^{m,M} \in \mathcal{C}_{\text{rap}}^{|\tau|}(\mathbb{R}^d),$$

the distribution  $\Pi_\tau$  appears then as another reconstruction of  $h_\tau$ .

– If one picks now a basis vector  $\mu \in \mathcal{B}$ , with  $|\mu| > 0$ , then  $h_\mu \in \mathcal{D}_{\text{rap}}^{|\mu|}(T, g)$  has a unique reconstruction, equal to  $\Pi_\mu$ , that is characterized by the data

$$(\Pi_x^g \sigma, g_x(\mu/\sigma) ; x \in \mathbb{R}^d, \sigma < \mu),$$



from the defining property of a reconstruction. An elementary induction then shows the existence of a unique extension of  $\Pi$  to  $T$  that satisfies the property  $\Pi\tau = \mathbf{R}h_\tau$ , for every  $\tau \in \mathcal{B}$  with positive homogeneity.

The fact that this statement holds not only for the paraproduct  $P$  but also for the modified paraproduct  $P^m$  will play a pivotal role in the proof of Lemma 18 below. The proof of Proposition 15 makes it clear that the above parameterization of the set of  $\Pi$  maps is related to the non-uniqueness of the reconstruction map on the set of modeled distributions of negative regularity exponent. This statement leaves us with the task of giving a parameterization of the set of characters  $g$  on  $T^+$  by their paracontrolled representation. We need for that purpose to make the following assumptions on the Hopf algebra  $(T^+, \Delta^+)$  and the basis  $\mathcal{B}^+$  of  $T^+$ . Recall that  $D^k : T_\alpha^+ \rightarrow T_{\alpha-|k|}^+$  is a linear map satisfying the recursive rules from Lemma 9. Recall that a pre-order  $\mathbb{E}$  is a reflexive transitive binary relation. Write  $\sigma \triangleleft \tau$  if  $\sigma \mathbb{E} \tau$  and  $\tau \not\mathbb{E} \sigma$ .

**ASSUMPTION C**

(1) *There exists a finite subset  $\mathcal{G}_0^+$  of  $\mathcal{B}_0^+$  such that  $\mathcal{B}_0^+$  is of the form*

$$\mathcal{B}_0^+ = \bigsqcup_{\tau \in \mathcal{G}_0^+} \{D^k \tau ; k \in \mathbb{N}^d, |\tau| - |k| > 0\}.$$

(2) *There exists a preorder  $\mathbb{E}$  on the set  $\mathcal{B}^+$  such that, for each  $\tau \in \mathcal{G}_0^+$ , the coproduct  $\Delta^+ \tau$  is of the form*

$$(4.2) \quad \Delta^+ \tau = \tau \otimes \mathbf{1} + \sum_{\sigma \triangleleft \tau, \sigma \notin \mathcal{B}_X^+} \sigma \otimes (\tau / \sigma) + \sum_k \frac{X^k}{k!} \otimes D^k \tau,$$

with  $\sigma \in \mathcal{B}^+(\tau^-)$  and  $\tau / \sigma \in \text{span}(\mathcal{B}^+(\tau^-))$ , for each  $\sigma$  in the above sum, where for each  $\tau \in \mathcal{B}^+$ , denote by  $\mathcal{B}^+(\tau^-)$  the submonoid of  $\mathcal{B}^+$  generated by

$$\{X_1, \dots, X_d\} \cup \bigsqcup_{\sigma \in \mathcal{G}_0^+, \sigma \triangleleft \tau} \{D^k \sigma ; k \in \mathbb{N}^d, |\sigma| - |k| > 0\}.$$

(3) *For any element  $\sigma \in \mathcal{B}^+ \cap \mathcal{B}_X^+$  such that there exists  $\tau \in \mathcal{G}_0^+$  and  $\sigma \triangleleft \tau$ , the homogeneity of  $\sigma$  is non-integer.*

Note the disjoint union in the description of  $\mathcal{B}_0^+$ . Assumption C(1) identifies a set of generators, modulo the action of the  $D$  operator. Assumption C(2) provides a useful induction structure. Assumption C(3) is a part of assumption B' and it is used at the end of the proof of Theorem 17. If one understands the coproduct  $\Delta^+$  as giving the elementary pieces of any given element, assumption (C) as a whole provides an inductive description of  $\mathcal{B}^+$ .

As discussed in Section 4.3, BHZ regularity structure satisfies assumption (C). Indeed, we can choose  $\mathcal{G}_0^+$  as a set of all conforming trees of the form  $I_0^t(\tau)$ , and the operator  $D^k$  appears as the form  $D^k I_0^t(\tau) = I_k^t(\tau)$ . In the BHZ regularity structure, one of the examples of  $\mathbb{E}$  is the binary relation based on the scale of graphs. Since  $\sigma$  and  $\tau / \sigma$  in (4.2) are subtree and quotient tree of  $\tau$  respectively, it follows from

definition of  $\Delta^+$  that  $\sigma, \tau / + \sigma \triangleleft \tau$ . The last assumption is true, if the types  $\{\mathfrak{t}\}$  are assigned rationally independent homogeneities  $\{|\mathfrak{t}|\}$ . See Theorem 26 for details. We leave now this special setting and come back to our general setting.

LEMMA 16. — Denote  $T^+(\tau^-) := \text{span}(\mathcal{B}^+(\tau^-))$ . For any  $\tau \in \mathcal{G}_\circ^+$  and any  $\sigma \in \mathcal{B}^+(\tau^-)$ , one has

$$(4.3) \quad \Delta^+ \sigma \in T^+(\tau^-) \otimes T^+(\tau^-).$$

Proof. — By multiplicativity it is sufficient to show the case  $\sigma = D^k \eta \in \mathcal{B}^+(\tau^-)$  with  $\eta \in \mathcal{G}_\circ^+$  and  $k \in \mathbb{N}^d$ . If  $k = 0$ , (4.3) follows because of the transitivity of  $\triangleleft$ . By the formula (3.4) for  $\Delta^+ D^k \eta$  and the property  $D^k : T^+(\tau^-) \rightarrow T^+(\tau^-)$  that is proved by Lemma 9,  $D^k \eta$  also satisfies (4.3).

Recall from formula (3.14) that if we are given characters  $(\mathfrak{g}_x)_{x \in \mathbb{R}^d}$  on  $T^+$  as in Definition 10, then

$$(4.4) \quad \mathfrak{g}_x(D^k \tau) = \mathbf{1}_{|k| < |\tau|} \partial_y^k \left\{ \mathfrak{g}_y(\tau) - \sum_{\substack{\sigma \triangleleft^+ \tau \\ \sigma \notin \mathcal{B}_X^+ \\ |\sigma| \triangleleft |k|}} \mathfrak{g}_{yx}(\sigma) \mathfrak{g}_x(\tau / + \sigma) \right\} \Big|_{y=x}.$$

The induction structure from assumption C(2) restricts the above sum and shows that the family of all  $\mathfrak{g}_x(D^k \tau)$  is uniquely determined by the preceding formula. It follows then from assumption C(1) that the character  $\mathfrak{g}$  on  $T^+$  is entirely determined by the datum of the  $\mathfrak{g}(\tau)$ , for  $\tau \in \mathcal{G}_\circ^+$ . We have in particular, if  $\tau \in \mathcal{G}_\circ^+$  is minimal (i.e., there is no  $\sigma \in \mathcal{G}_\circ^+$  such that  $\sigma \triangleleft \tau$ ) then

$$\mathfrak{g}_y(\tau) = \mathfrak{g}_{yx}(\tau) + \sum_{|k| < |\tau|} \frac{(y-x)^k}{k!} \mathfrak{g}_x(D^k \tau),$$

since  $\mathcal{B}^+(\tau^-) = \mathcal{B}_X^+$ , so for  $|k| < |\tau|$ , one has

$$(4.5) \quad \mathfrak{g}_x(D^k \tau) = \partial_y^k \mathfrak{g}_y(\tau) \Big|_{y=x},$$

and

$$\mathfrak{f}_x(D^k \tau) = \mathfrak{g}_x(D^k \tau),$$

and

$$(4.6) \quad \mathfrak{g}_{yx}(D^k \tau) = \mathfrak{g}_y(D^k \tau) - \sum_{\ell} \frac{(y-x)^\ell}{\ell!} \mathfrak{g}_x(D^{k+\ell} \tau).$$

Recall that, given a concrete regularity structure  $\mathcal{T}$ ,

$$\mathcal{T}^+ = ((T^+, \Delta^+), (T^+, \Delta^+))$$

is also a concrete regularity structure, and that for a  $\mathfrak{g}$  map as in Definition 10 one defines a model  $M^\mathfrak{g} = (\mathfrak{g}, \Pi^\mathfrak{g})$  on  $\mathcal{T}^+$  setting

$$(\Pi^\mathfrak{g} \tau)(y) = \mathfrak{g}_y(\tau).$$

**THEOREM 17.** — *Let  $T$  stand for a concrete regularity structure satisfying assumptions (A–C). Then, for any family  $(\llbracket \tau \rrbracket \in C_{\text{rap}}^{|\tau|}(\mathbb{R}^d))_{\tau \in \mathcal{G}_\circ^+}$ , there exists a unique model  $\mathbf{M}^{\mathfrak{g}} = (\mathfrak{g}, \Pi^{\mathfrak{g}})$  on  $T^+$  such that*

$$(4.7) \quad \mathfrak{g}(\tau) = \sum_{\substack{\sigma < +\tau \\ \sigma \in \mathcal{B}^+ \cap \mathcal{B}_X^+}} P_{\mathfrak{g}(\tau/+ \sigma)} \llbracket \sigma \rrbracket^{\mathbf{M}^{\mathfrak{g}}} + \llbracket \tau \rrbracket, \quad \forall \tau \in \mathcal{G}_\circ^+.$$

The map

$$(4.8) \quad (\llbracket \tau \rrbracket \in C_{\text{rap}}^{|\tau|}(\mathbb{R}^d))_{\tau \in \mathcal{G}_\circ^+} \longmapsto \mathbf{M}^{\mathfrak{g}} \in M_{\text{rap}}(T^+, \mathbb{R}^d)$$

is locally Lipschitz continuous.

Note that one uses the paraproduct  $P$  and the brackets  $\llbracket \cdot \rrbracket^{\mathbf{M}^{\mathfrak{g}}}$  in the statement. The modified paraproduct  $P^m$  is only used in the proof of Lemma 18, where we construct a model on an intermediate regularity structure introduced along the proof. The injectivity of the map (4.8) is elementary, so Theorem 17 and Proposition 15, with Theorem 12, prove all together Theorem 1.

The remaining of this section is dedicated to proving Theorem 17. The proof is done by induction on the preorder  $E$ .

*Initialization of the induction.* — If  $\tau \in \mathcal{G}_\circ^+$  is a minimal element, then set

$$\mathfrak{g}(\tau) := \llbracket \tau \rrbracket,$$

and define  $\mathfrak{g}(D^k \tau)$  and  $\mathfrak{g}_{yx}(D^k \tau)$  by (4.5) and (4.6). It is clear on these formulas that they define elements of the spaces  $C_{\text{rap}}^{|\tau|-|k|}(\mathbb{R}^d) \subset L_{\text{rap}}^\infty(\mathbb{R}^d)$  and  $\mathcal{C}_{(2),\text{rap}}^{|\tau|-|k|}(\mathbb{R}^d \times \mathbb{R}^d)$ , respectively.

*Induction step.* — Fix  $\tau \in \mathcal{G}_\circ^+$  and assume that  $\mathfrak{g}$  has been constructed on the submonoid  $\mathcal{B}^+(\tau^-)$  as a continuous function of the bracket data – so all the functions  $\llbracket \sigma \rrbracket^{\mathbf{M}^{\mathfrak{g}}}$  and  $\mathfrak{g}(\tau/+ \sigma)$  make sense as elements of their natural spaces. Define  $\mathfrak{g}(\tau)$  by identity (4.7), and define  $\mathfrak{g}(D^k \tau)$  by (4.4), for all  $k \in \mathbb{N}^d$  with  $|k| < |\tau|$ . The induction step consists in proving that  $\mathfrak{g}_x(D^k \tau) \in L_{\text{rap}}^\infty(\mathbb{R}^d)$  and  $\mathfrak{g}_{yx}(D^k \tau) \in \mathcal{C}_{(2),\text{rap}}^{|\tau|-|k|}(\mathbb{R}^d \times \mathbb{R}^d)$ , as one can use for  $\alpha, \beta$  non-negative the inclusions

$$L_{\text{slow}}^\infty(\mathbb{R}^d) \cdot L_{\text{rap}}^\infty(\mathbb{R}^d) \subset L_{\text{rap}}^\infty(\mathbb{R}^d)$$

and

$$\mathcal{C}_{(2)}^\alpha(\mathbb{R}^d \times \mathbb{R}^d) \cdot \mathcal{C}_{(2),\text{rap}}^\beta(\mathbb{R}^d \times \mathbb{R}^d) \subset \mathcal{C}_{(2),\text{rap}}^{\alpha+\beta}(\mathbb{R}^d \times \mathbb{R}^d),$$

to get the regularity properties of  $\mathfrak{g}_x(\mu D^k \tau)$  and  $\mathfrak{g}_{yx}(\mu D^k \tau)$ , for  $\mu \in \mathcal{B}^+(\tau^-)$ .

Choose  $m \in \mathbb{N}$ , with  $m > |\tau|$ . We introduce a regularity structure  $T^m(\tau)$  with Hopf algebra part  $T^+(\tau^-)$  and  $T$ -space defined as follows. Consider the formal symbols

$$\sigma^{(m)}$$

indexed by  $\sigma \in \mathcal{B}^+ \cap \mathcal{B}_X^+$ , with homogeneity

$$|\sigma^{(m)}| := |\sigma| - m.$$

Set

$$T^m(\tau) := \text{span}(\{\sigma^{(m)} ; \sigma <^+ \tau, \sigma \notin \mathcal{B}_X^+\} \cup \{\tau^{(m)}\}),$$

so all elements of  $T^m(\tau)$  have negative homogeneity. Lemma 16 ensures that we can define a coassociative coproduct

$$\delta : T^m(\tau) \longrightarrow T^m(\tau) \otimes T^+(\tau^-)$$

setting

$$\delta(\sigma^{(m)}) := \sum_{\substack{\mu \dot{<}^+ \sigma, \mu \notin \mathcal{B}_X^+}} \mu^{(m)} \otimes (\sigma /^+ \mu)$$

for each basis element of  $T^m(\tau)$ . Lemma 16 also ensures that

$$\Delta^+(T^+(\tau^-)) \subset T^+(\tau^-) \otimes T^+(\tau^-),$$

so

$$\mathcal{T}^m(\tau) := ((T^+(\tau^-), \Delta^+), (T^m(\tau), \delta))$$

is a concrete regularity structure.

We build a model  $(\mathbf{g}, \Lambda)$  on  $\mathcal{T}^m(\tau)$ , from  $\mathbf{g} : T^+(\tau^-) \rightarrow \mathbb{R}$  given by an induction assumption and an operator  $\Lambda : T^m(\tau) \rightarrow \mathcal{S}'(\mathbb{R}^d)$  defined by

$$\Lambda(\sigma^{(m)}) := |\nabla|^m \mathbf{g}(\sigma),$$

where  $|\nabla|^m$  is the Fourier multiplier operator  $|\nabla|^m \zeta = \mathcal{F}^{-1}(|\cdot|^m \mathcal{F} \zeta)$ . The pair  $(\mathbf{g}, \Lambda)$  turns out to be a model by Lemma 18 below. Then formula (3.11) giving  $\mathbf{f}_x(D^k \sigma)$  can be interpreted in terms of that model, under the form of identities

$$\mathbf{f}_x(D^k \sigma) = \mathbf{J}^{k,m} \left( \Lambda_x^{\mathbf{g}}(\sigma^{(m)}) \right) (x)$$

for operators  $\mathbf{J}^{k,m}$  on distributions defined below. The identity

$$\Lambda_x^{\mathbf{g}} = \Lambda_y^{\mathbf{g}} \circ \widehat{\mathbf{g}}_{yx}^\delta,$$

where  $\widehat{\mathbf{g}}^\delta := (\text{Id} \otimes \mathbf{g})\delta$ , is then used crucially to obtain estimates on  $\mathbf{f}_x(D^k \sigma)$ , that eventually give information on  $\mathbf{g}_x(D^k \tau)$  and  $\mathbf{g}_{yx}(D^k \tau)$  via formulas (3.9) and (3.10).

**LEMMA 18.** — *The pair  $(\mathbf{g}, \Lambda)$  is a rapidly decreasing model on the regularity structure  $\mathcal{T}^m(\tau)$ .*

*Proof.* — Since we have the identity

$$\Lambda(\sigma^{(m)}) = |\nabla|^m \mathbf{g}(\sigma) = \sum_{\substack{\mu < \sigma \\ \mu \notin \mathcal{B}_X^+}} P_{\mathbf{g}(\sigma /^+ \mu)}^m |\nabla|^m \llbracket \mu \rrbracket^{\mathbf{g}} + |\nabla|^m \llbracket \sigma \rrbracket^{\mathbf{g}},$$

for all  $\sigma \in \mathcal{B}^+ \cap \mathcal{B}_X^+$  with  $\sigma \dot{<}^+ \tau$ , from the intertwining relation defining  $P^m$  and the induction assumption, the operator  $\Lambda$  is the unique model on  $\mathcal{T}^m(\tau)$  associated by Proposition 15 to the inputs

$$\llbracket \sigma^{(m)} \rrbracket := |\nabla|^m \llbracket \sigma \rrbracket^{\mathbf{g}} \in \mathcal{C}_{\text{rap}}^{|\sigma| - m}(\mathbb{R}^d),$$

since all elements of  $T^m(\tau)$  have negative homogeneity.

Note that it follows from identity (3.11) in Lemma 14 that the model  $\Pi$  and the function  $f(D^k\sigma)$  are related by the identity

$$\begin{aligned}
 (4.9) \quad f_x(D^k\sigma) &= \partial_y^k \left\{ \sum_{\substack{\mu \in \mathcal{B}^+_{\sigma} \\ \mu \notin \mathcal{B}^+_{\mathcal{X}}}} \mathfrak{g}_x^{-1}(\sigma/+\mu) \mathfrak{g}_y(\mu) \right\} \Big|_{y=x} \\
 &= \partial_y^k \left\{ |\nabla_y|^{-m} \sum_{\substack{\mu \in \mathcal{B}^+_{\sigma} \\ \mu \notin \mathcal{B}^+_{\mathcal{X}}}} \mathfrak{g}_x^{-1}(\sigma/+\mu) \Lambda(\mu^{(m)})(y) \right\} \Big|_{y=x} \\
 &= \partial_y^k \left\{ |\nabla_y|^{-m} \Lambda_x^{\mathfrak{g}}(\sigma^{(m)})(y) \right\} \Big|_{y=x} \\
 &=: \sum_j \mathbf{J}_j^{k,m}(\Lambda_x^{\mathfrak{g}}(\sigma^{(m)}))(x),
 \end{aligned}$$

where the operators  $\mathbf{J}_j^{k,m}$  are defined by

$$\mathbf{J}_j^{k,m}(\zeta) := \partial^k |\nabla|^{-m} \Delta_j \zeta,$$

for an appropriate distribution  $\zeta \in \mathcal{S}'(\mathbb{R}^d)$ . If  $j > 0$ , since the Fourier transform of  $\Delta_j \zeta$  is supported on an annulus, the function  $\mathbf{J}_j^{k,m}(\zeta)$  is always well-defined; this is not the case of  $\mathbf{J}_{-1}^{k,m}(\zeta)$ . However, we only use in this section distributions  $\zeta$  of the form  $\zeta = |\nabla|^m \xi$  (where such  $\xi$  is unique in the class of rapidly decreasing functions), so  $\mathbf{J}_{-1}^{k,m}(\zeta) = \partial^k \Delta_{-1} \xi$ , in our setting.

**LEMMA 19.** — *Under assumptions (A–C), for any  $\sigma \in \mathcal{B}^+ \cap \mathcal{B}_{\mathcal{X}}^+$  with  $\sigma \in \mathcal{B}^+ \cap \mathcal{B}_{\mathcal{X}}^+$ ,  $k \in \mathbb{N}^d$ , and  $a \in \mathbb{N}$ , we have*

$$\begin{aligned}
 & \left| \mathbf{J}_j^{k,m}(\Lambda_x^{\mathfrak{g}}(\sigma^{(m)}))(x) \right| \cdot |x|_*^{-a} 2^{-j(|\sigma|-|k|)}, \\
 & \left| \mathbf{J}_j^{k,m}(\Lambda_x^{\mathfrak{g}}(\sigma^{(m)}))(y) \right| \cdot |y|_*^{-a} \sum_{\substack{\mu \in \mathcal{B}^+_{\sigma} \\ \mu \notin \mathcal{B}^+_{\mathcal{X}}}} |y-x|^{|\sigma|-|\mu|} 2^{-j(|\mu|-|k|)}.
 \end{aligned}$$

Consequently,  $f_x(D^k\sigma) \in L_{\text{rap}}^\infty$ .

*Proof.* — For the first estimate, since

$$\mathbf{J}_{-1}^{k,m}(\Lambda_x^{\mathfrak{g}}(\sigma^{(m)}))(x) = \sum_{\substack{\mu \in \mathcal{B}^+_{\sigma} \\ \mu \notin \mathcal{B}^+_{\mathcal{X}}}} \mathfrak{g}_x^{-1}(\sigma/+\mu) \partial_x^k \Delta_{-1} \mathfrak{g}_x(\mu) \in L_{\text{rap}}^\infty,$$

by assumption, it is sufficient to consider the case  $j > 0$ . By the property of  $\rho_j$ , there exists a smooth function  $\tilde{\rho}$  supported on an annulus, and such that setting  $\tilde{\rho}_j(\cdot) := \tilde{\rho}(2^{-j}\cdot)$ , one has  $\tilde{\rho}_j \rho_j = \rho_j$ . Set

$$\tilde{Q}_j^{k,m} := \partial^k |\nabla|^{-m} (\mathcal{F}^{-1} \tilde{\rho}_j),$$

and note the scaling property

$$\tilde{Q}_j^{k,m}(\cdot) = 2^{j(d+|k|-m)} \tilde{Q}_0^{k,m}(2^j \cdot).$$

We now use the fact that  $(g, \Lambda)$  is a model to write

$$\begin{aligned} \mathbf{J}_j^{k,m}(\Lambda_x^g(\sigma^{(m)}))(x) &= \int \tilde{Q}_j^{k,m}(x-y)\Delta_j(\Lambda_x^g(\sigma^{(m)}))(y)dy \\ &= \int \tilde{Q}_j^{k,m}(x-y)\Delta_j(\Lambda_y^g \circ \widehat{g}_{yx}^\delta(\sigma^{(m)}))(y)dy \\ &= \sum_{\mu \in \sigma^+} \int \tilde{Q}_j^{k,m}(x-y)g_{yx}(\sigma^+/\mu)\Delta_j(\Lambda_y^g(\mu^{(m)}))(y)dy. \end{aligned}$$

Recall that  $|x+y|_* \leq |x|_*|y|_*$ , for all  $x, y \in \mathbb{R}^d$ . By Lemma 18, for any  $a \in \mathbb{N}$  we have

$$\begin{aligned} &|x|_*^a \left| \mathbf{J}_j^{k,m}(\Lambda_x^g(\sigma^{(m)}))(x) \right| \\ &\leq \sum_{\mu \in \sigma^+} \int |x-y|_*^a |\tilde{Q}_j^{k,m}(x-y)| |y-x|^{|\sigma|-|\mu|} |y|_*^a |\Delta_j(\Lambda_y^g(\mu^{(m)}))(y)| dy \\ &\leq \sum_{\mu \in \sigma^+} 2^{-j(|\mu|-m)} \int |z|_*^a |\tilde{Q}_j^{k,m}(z)| |z|^{|\sigma|-|\mu|} dz \\ &\leq \sum_{\mu \in \sigma^+} 2^{-j(|\mu|-m)} 2^{j(|k|-m-|\sigma|+|\mu|)} \int |z|_*^a |\tilde{Q}_0^{k,m}(z)| |z|^{|\sigma|-|\mu|} dz \\ &\leq 2^{-j(|\sigma|-|k|)}. \end{aligned}$$

We get the second estimate from the first using once again the fact that  $(g, \Lambda)$  is a model, writing

$$\begin{aligned} \mathbf{J}_j^{k,m}(\Lambda_x^g(\sigma^{(m)}))(y) &= \mathbf{J}_j^{k,m}(\Lambda_y^g(\widehat{g}_{yx}^\delta(\sigma^{(m)})))(y) \\ &= \sum_{\substack{\mu \in \sigma^+ \\ \mu \notin \mathcal{B}_x^+}} g_{yx}(\sigma^+/\mu) \mathbf{J}_j^{k,m}(\Lambda_y^g(\mu^{(m)}))(y). \end{aligned}$$

We can now prove that  $g_x(D^k\tau) \in L_{\text{rap}}^\infty(\mathbb{R}^d)$  and  $g_{yx}(D^k\tau) \in \mathcal{C}_{(2),\text{rap}}^{|\tau|-|k|}(\mathbb{R}^d \times \mathbb{R}^d)$ , and close the induction step. We use the formulas from Lemma 14 for that purpose. First, since

$$g_x(D^k\tau) = \sum_{\substack{\sigma \in \sigma^+ \\ \sigma \notin \mathcal{B}_x^+}} g_x(\tau^+/\sigma) f_x(D^k\sigma),$$

with  $g_x(\tau^+/\sigma) \in L_{\text{slow}}^\infty(\mathbb{R}^d)$  and  $f_x(D^k\sigma) \in L_{\text{rap}}^\infty(\mathbb{R}^d)$ , from Lemma 19, we have indeed  $g_x(D^k\tau) \in L_{\text{rap}}^\infty(\mathbb{R}^d)$ . Second, one can rewrite the identity

$$g_{yx}(D^k\tau) = \sum_{\substack{\sigma \in \sigma^+ \\ \sigma \notin \mathcal{B}_x^+}} g_{yx}(\tau^+/\sigma) f_y(D^k\sigma) - \sum_{\ell} \frac{(y-x)^\ell}{\ell!} f_x(D^{k+\ell}\tau),$$

from Lemma 14, using identity (4.9) for the f-terms. This gives for  $\mathbf{g}_{yx}(D^k\tau)$  the formula

$$\sum_j \left\{ \sum_{\substack{\sigma \in \mathcal{B}_x^+ \\ |k| < |\sigma|}} \mathbf{g}_{yx}(\tau/+\sigma) \mathbf{J}_j^{k,m}(\Lambda_y^{\mathbf{g}}(\sigma^{(m)}))(y) - \sum_{|k+\ell| < |\tau|} \frac{(y-x)^\ell}{\ell!} \mathbf{J}_j^{k+\ell,m}(\Lambda_x^{\mathbf{g}}(\tau^{(m)}))(x) \right\} =: \sum_j \mathbf{g}_{yx}^j(D^k\tau).$$

Given  $x, y \in \mathbb{R}^d$ , set  $j_0 = -1$ , if  $|y-x| > 2$ , and pick otherwise  $j_0 > -1$  such that  $|y-x| \simeq 2^{-j_0}$ . One uses the first estimate from Lemma 19 to bound above the sum over  $j > j_0$

$$(4.10) \quad \begin{aligned} & |x|_*^a \sum_{j > j_0} |\mathbf{g}_{yx}^j(D^k\tau)| \cdot \sum_{j > j_0} \sum_{\substack{\sigma \in \mathcal{B}_x^+ \\ |k| < |\sigma|}} |y-x|^{|\tau|-|\sigma|} 2^{-j(|\sigma|-|k|)} \\ & \quad + \sum_{j > j_0} \sum_{|k+\ell| < |\tau|} |y-x|^{|\ell|} 2^{-j(|\tau|-|k|-|\ell|)} \\ & \quad \cdot \sum_{\substack{\sigma \in \mathcal{B}_x^+ \\ |k| < |\sigma|}} |y-x|^{|\tau|-|\sigma|} 2^{-j_0(|\sigma|-|k|)} + \sum_{|k+\ell| < |\tau|} |y-x|^{|\ell|} 2^{-j_0(|\tau|-|k|-|\ell|)} \\ & \quad \cdot |y-x|^{|\tau|-|k|}. \end{aligned}$$

To consider the sum over  $j < j_0$ , assume now that  $|y-x| < 2$ . Then, since  $(\mathbf{g}, \Lambda)$  is a model and

$$\Lambda_x^{\mathbf{g}}(\tau^{(m)}) = \Lambda_y^{\mathbf{g}}(\widehat{\mathbf{g}}_{yx}^\delta \tau^{(m)}) = \sum_{\sigma \in \mathcal{B}_x^+} \mathbf{g}_{yx}(\tau/+\sigma) \Lambda_y^{\mathbf{g}}(\sigma^{(m)}),$$

we have for  $\mathbf{g}_{yx}^j(D^k\tau)$  the formula

$$\begin{aligned} \mathbf{g}_{yx}^j(D^k\tau) &= \mathbf{J}_j^{k,m}(\Lambda_x^{\mathbf{g}}(\tau^{(m)}))(y) - \sum_{\substack{\sigma \in \mathcal{B}_x^+ \\ |k| > |\sigma|}} \mathbf{g}_{yx}(\tau/+\sigma) \mathbf{J}_j^{k,m}(\Lambda_y^{\mathbf{g}}(\sigma^{(m)}))(y) \\ & \quad - \sum_{|k+\ell| < |\tau|} \frac{(y-x)^\ell}{\ell!} \mathbf{J}_j^{k+\ell,m}(\Lambda_x^{\mathbf{g}}(\tau^{(m)}))(x) \\ &= [b] \sum_{|k'|=[b]} \frac{(y-x)^{k'}}{k'!} \int_0^1 (1-t)^{[b]} \mathbf{J}_j^{k+k'}(\Lambda_x^{\mathbf{g}}(\tau^{(m)}))(x+t(y-x)) dt \\ & \quad - \sum_{\substack{\sigma \in \mathcal{B}_x^+ \\ |k| > |\sigma|}} \mathbf{g}_{yx}(\tau/\sigma) \mathbf{J}_j^{k,m}(\Lambda_y^{\mathbf{g}}(\sigma^{(m)}))(y), \end{aligned}$$

where  $b := |\tau| - |k|$ , by the multivariable Taylor remainder formula. Note that  $|\tau|, |\sigma| \notin \mathbb{N}$  in the above formula, by assumption C(3). Since  $|y-x| < 2$ ,  $|x+t(y-x)|_* \simeq |x|_*$ .

It follows then from Lemma 19 that  $\sum_{-1 \leq j < j_0} |\mathbf{g}_{yx}^j(D^k \tau)|$  is bounded above by

$$\begin{aligned} & \sum_{j < j_0} \sum_{|k'|=|b|} \sum_{\substack{\sigma \leq \sigma + \tau \\ \sigma \notin \mathcal{B}_X^+}} |y - x|^{|k'|+|\tau|-\sigma} |x|_*^{-a} 2^{-j(|\sigma|-|k|-|k'|)} \\ & \quad + \sum_{j < j_0} \sum_{\substack{\sigma \leq \sigma + \tau \\ \sigma \notin \mathcal{B}_X^+ \\ |k| > |\sigma|}} |y - x|^{|\tau|-\sigma} |y|_*^{-a} 2^{-j(|\sigma|-|k|)} \\ & \quad \cdot |x|_*^{-a} \sum_{\substack{\sigma \leq \sigma + \tau \\ \sigma \notin \mathcal{B}_X^+}} |y - x|^{|k'|+|\tau|-\sigma} 2^{-j_0(|\sigma|-|k|-|k'|)} \\ & \quad + |y|_*^{-a} \sum_{\substack{\sigma \leq \sigma + \tau \\ \sigma \notin \mathcal{B}_X^+ \\ |k| > |\sigma|}} |y - x|^{|\tau|-\sigma} 2^{-j_0(|\sigma|-|k|)} \\ & \quad \cdot (|x|_*^{-a} + |y|_*^{-a}) |y - x|^{|\tau|-|k|}. \end{aligned}$$

Together with inequality (4.10), the preceding upper bound tells us that

$$\mathbf{g}_{yx}(D^k \tau) \in \mathcal{C}_{(2), \text{rap}}^{|\tau|-|k|}(\mathbb{R}^d \times \mathbb{R}^d).$$

This closes the induction step.

**REMARKS**

(1) *On branched rough paths.* The setting of  $\mathbb{R}^\ell$ -valued branched rough paths provides an example of regularity structure where Theorem 5 applies, giving an alternative point of view on the results of Tapia and Zambotti in [24]. The Hopf algebra  $(T^+, \Delta^+)$  is in that case the Butcher-Connes-Kreimer Hopf algebra. We recall the details for the reader as it also sets the scene for part of the results of Section 4.4.

The set  $T^+$  is the free commutative unital algebra generated by the set  $\mathcal{B}_\circ^+$  of non-planar rooted trees with node decorations in a finite set  $\{1, \dots, \ell\}$  and no decoration on the edges. The empty tree plays the role of the unit in  $T^+$ . A product of decorated trees is called a forest, so generic elements of  $T^+$  are linear combinations of forests. The splitting map  $\Delta^+$  is the algebra morphism defined on trees as follows. Given a labeled rooted decorated tree  $\tau$ , denote by  $\text{Sub}(\tau)$  the set of subtrees of  $\tau$  with the same root as  $\tau$ , and induced decoration. Given such a subtree  $s$ , we obtain a collection  $\tau_1, \dots, \tau_n$  of decorated rooted trees by removing  $s$  and all the adjacent edges to  $s$  from  $\tau$ , and keeping the node decoration inherited from  $\tau$ . Write  $\tau/s$  for the monomial  $\tau_1 \dots \tau_n$ . One defines a linear multiplicative map  $\Delta^+ : T^+ \rightarrow T^+ \otimes T^+$ , defining it on decorated trees by the formula

$$\Delta^+ \tau = \sum_{s \in \text{Sub}(\tau)} s \otimes (\tau/s),$$

An explicit formula for the antipode was first given by Connes and Kreimer in their celebrated work [11]; see [10] for a simple and enlightening proof. Each node decoration  $i \in \{1, \dots, \ell\}$  is assigned a homogeneity  $\alpha \in (0, 1)$ , and each decorated tree  $\tau$  is equipped with the homogeneity  $\alpha(\#\tau)$ , where  $\#\tau$  denotes the number of nodes contained in  $\tau$ . The homogeneity of a forest is the sum of the homogeneities



of its decorated trees. It is elementary to check that  $(T^+, \Delta^+)$  is indeed a Hopf algebra. To avoid polynomials and derivatives, we consider the subalgebra of trees with homogeneities smaller than 1. Thus assumptions (A–C) without polynomials  $X^k$  and derivatives  $D^k$  hold, but it does not matter here. Branched rough paths are  $\mathfrak{g}$ -maps on  $(T^+, \Delta^+)$  over a fixed time interval  $[0, T]$  in place of  $\mathbb{R}^d$ . Theorem 17 applies then in this setting and provides a parameterization of the set of branched rough paths by the product space  $\prod_{\tau \in \mathcal{B}_\circ^+} C^{|\tau|}([0, T], \mathbb{R}^\ell)$ , in accordance with Tapia and Zambotti's main result [24, Th. 1.2 & Cor. 1.3]. Our parameterization is different from their identification of the space of branched rough paths as a principal homogeneous space over the preceding product of Hölder spaces.

(Theorem 1 cannot be applied in a finite region  $[0, T]$  directly. To overcome this point, we extend a function  $f \in C([0, T], \mathbb{R}^\ell)$  to  $[-T, T]$  symmetrically, and extend it to  $[(2n - 1)T, (2n + 1)T]$  for any  $n \in \mathbb{Z}$  periodically. Then for any  $\alpha \in (0, 1)$ , the Hölder space  $C^\alpha([0, T], \mathbb{R}^\ell)$  is identified with the space

$$C_{p,s}^\alpha(\mathbb{R}, \mathbb{R}^\ell) := \left\{ f \in C^\alpha(\mathbb{R}, \mathbb{R}^\ell); f(t) = f(-t), f(t + 2T) = f(t) \text{ for any } t \in \mathbb{R} \right\}.$$

Note that Littlewood-Paley blocks  $\Delta_i$  preserve the symmetry and periodicity, so such spaces are closed under paradiﬀerential operators (paraproduct, its two-parameter extension, etc.) used in this paper. Hence we can apply Theorem 1 to such spaces.)

(2) *On the signature of arbitrary models.* We mentioned in the introduction that admissible models on regularity structures built from integration operators have a well-defined signature – that is a unique extension to the full regularity structure with elements of arbitrary large positive homogeneity. This comes from the fact that such models are determined uniquely by the definition of the  $\Pi$  map on elements of the regularity structure of negative homogeneity. Extending a regularity structure with additional elements of positive homogeneity the initial datum of the restriction of  $\Pi$  on the elements of negative homogeneity still defines a unique admissible model on the extended regularity structure. Such an automatic extension result does not hold for general models, with unrelated  $\mathfrak{g}$  and  $\Pi$  maps. Indeed, Theorem 17 tells us that the set of  $\mathfrak{g}$ -maps is parametrized by a set of functions indexed by  $\mathcal{G}_\circ^+$ . Embedding a regularity structure into a larger regularity structure will a priori embed the set  $\mathcal{G}_\circ^+$  into a larger set, implying the non-uniqueness of an extension of the  $\mathfrak{g}$ -map, from Theorem 17 again. The following statement follows nonetheless from Theorem 1 while it is beyond the scope of [5, Th. 21]. See [5, §4] for the definition of admissible model.

**COROLLARY 20.** — *Let  $\mathcal{T}$  be a regularity structure satisfying assumptions (A–C). Let  $\mathcal{T}' \subset \mathcal{T}$  be a sub-regularity structure of  $\mathcal{T}$  satisfying these assumptions as well, and such that  $\mathcal{T}'$  contains all the elements of  $\mathcal{T}$  of negative homogeneity. Then any admissible model on  $\mathcal{T}'$  has a unique extension into an admissible model on  $\mathcal{T}$ .*

Like with Lyons' extension theorem, it is important to notice that the extension map is a continuous map. So even in a stochastic setting where the construction of a

model may require stochastic analysis arguments, once this is done, the extension of this model to a larger structure no longer involves probability arguments.

4.2. FROM PARACONTROLLED SYSTEMS TO MODELED DISTRIBUTIONS. — We prove Theorem 5 in this section. Let  $T$  be a regularity structure satisfying assumptions (A–C). Pick  $\gamma \in \mathbb{R}$ , and  $\mathbf{M} = (\mathbf{g}, \Pi) \in M_{\text{rap}}(T, \mathbb{R}^d)$ .

The key observation is that proving Theorem 5 is equivalent to an extension problem for the map  $\mathbf{g}$ . Consider indeed the commutative algebra  $T_{\mathbf{F}}^{\pm}$  generated by  $\mathcal{B}^+$  and new symbols

$$(\mathbf{F}_{\tau})_{\tau \in \mathcal{B}, |\tau| < \gamma}.$$

Define the homogeneity of the symbol  $\mathbf{F}_{\tau}$  by

$$|\mathbf{F}_{\tau}| := \gamma - |\tau|.$$

The coproduct  $\Delta_{\mathbf{F}}^{\pm}$  on  $T_{\mathbf{F}}^{\pm}$  extending  $\Delta^+$  and such that

$$(4.11) \quad \Delta^+(\mathbf{F}_{\tau}) = (\mathbf{F}_{\tau}) \otimes \mathbf{1} + \sum_{\tau \dot{\subset} \mu} (\mu/\tau) \otimes (\mathbf{F}_{\mu}),$$

is coassociative and turns  $T_{\mathbf{F}}^{\pm}$  into a Hopf algebra. It satisfies assumptions (A) and (B) with

$$\mathcal{B}_{\mathbf{F}, \circ}^{\pm} := \mathcal{B}_{\circ}^{\pm} \cup \{\mathbf{F}_{\tau}; |\tau| < \gamma\}$$

in the role of  $\mathcal{B}_{\circ}^{\pm}$ . Note that  $T_{\mathbf{F}}^{\pm}$  does not satisfy assumption (C) in general, since the  $D^k \mathbf{F}_{\tau}$  have no reason to be independent from the  $\{\mathbf{F}_{\mu}\}_{\mu}$ . The elementary proof of the next statement is left to the reader.

LEMMA 21. — Given a family  $(f_{\tau})_{\tau \in \mathcal{B}}$  of continuous functions on  $\mathbb{R}^d$ , set  $\mathbf{f} := \sum_{\tau \in \mathcal{B}} f_{\tau} \tau$ , and

$$\mathbf{g}_x(\mathbf{F}_{\tau}) := f_{\tau}(x).$$

Then

$$\langle \tau', \mathbf{f}(y) - \widehat{\mathbf{g}}_{yx} \mathbf{f}(x) \rangle = \mathbf{g}_{yx}(\mathbf{F}_{\tau}).$$

Defining a modeled distribution  $\mathbf{f} \in \mathcal{D}_{\text{rap}}^{\gamma}(T, \mathbf{g})$  is thus equivalent to extending the map  $\mathbf{g}$  from  $T^+$  to  $T_{\mathbf{F}}^{\pm}$  in such a way that the extended map on  $(T_{\mathbf{F}}^{\pm}, \Delta_{\mathbf{F}}^{\pm})$  still satisfies the regularity constraints from Definition 10.

Recall from assumption (B) that either  $\mu/\tau \in \text{span}(\mathcal{B}^+ \cap \mathcal{B}_X^{\pm})$  or  $\mu/\tau \in \text{span}(\mathcal{B}_X^{\pm})$ , for  $\tau, \mu \in \mathcal{B}$ . If  $\mu/\tau \in \text{span}(\mathcal{B}_X^{\pm})$ , set

$$\mu/\tau =: \sum_{k \in \mathbb{N}^d} c_{\tau}^{\mu}(k) \frac{X^k}{k!},$$

and define

$$(4.12) \quad D^k \mathbf{F}_{\tau} := \sum_{\substack{\tau \dot{\subset} \mu \\ \mu/\tau \in \text{span}(\mathcal{B}_X^{\pm})}} c_{\tau}^{\mu}(k) \mathbf{F}_{\mu}.$$

Then we have

$$\Delta^+ \mathbf{F}_\tau = \mathbf{F}_\tau \otimes \mathbf{1} + \sum_{\substack{\tau \triangleleft \mu \\ \mu/\tau \in \text{span}(\mathcal{B}^+ \mathcal{B}_X^+)}} (\mu/\tau) \otimes \mathbf{F}_\mu + \sum_{k \in \mathbb{N}^d} \frac{X^k}{k!} \otimes D^k \mathbf{F}_\tau.$$

**THEOREM 22.** — *Let a concrete regularity structure  $T$  satisfying assumptions (A–C) be given, together with a family  $(\llbracket f_\tau \rrbracket \in \mathcal{C}_{\text{rap}}^{\gamma-|\tau|}(\mathbb{R}^d))_{\tau \in \mathcal{B}, |\tau| < \gamma}$ . Assume that  $\gamma - |\tau| \notin \mathbb{N}$  for any  $\tau \in \mathcal{B}$  with  $|\tau| < \gamma$ . Pick a model  $(\mathbf{g}, \Pi) \in M_{\text{rap}}(T, \mathbb{R}^d)$ . Define by decreasing induction on  $|\tau|$*

$$f_\tau := \sum_{\substack{\tau \triangleleft \mu, |\mu| < \gamma \\ \mu/\tau \in \text{span}(\mathcal{B}^+ \mathcal{B}_X^+)}} P_{f_\mu} \llbracket \mu/\tau \rrbracket^{\mathbf{g}} + \llbracket f_\tau \rrbracket,$$

and

$$(4.13) \quad f_\tau^{(k)}(x) := \partial_y^k \left\{ f_\tau(y) - \sum_{\substack{\tau \triangleleft \mu, |\mu| < \gamma, |\mu/\tau| \leq |k| \\ \mu/\tau \in \text{span}(\mathcal{B}^+ \mathcal{B}_X^+)}} \mathbf{g}_{yx}(\mu/\tau) f_\mu(x) \right\} \Big|_{y=x}.$$

If the structure conditions

$$(4.14) \quad f_\tau^{(k)} = \sum_{\substack{\tau \triangleleft \mu, |\mu| < \gamma \\ \mu/\tau \in \text{span}(\mathcal{B}_X^+)}} c_\tau^\mu(k) f_\mu,$$

holds for any  $\tau \in \mathcal{B}$  and  $k \in \mathbb{N}^d$ , then

$$\mathbf{f} = \sum_{\tau \in \mathcal{B}} f_\tau \tau \in \mathcal{D}_{\text{rap}}^\gamma(T, \mathbf{g}).$$

The structure condition is reminiscent of a condition introduced by Martin and Perkowski in [22] to give a characterization of modeled distributions in terms of Besov type spaces. Given that we see  $f_\tau$  as  $\mathbf{g}(\mathbf{F}_\tau)$ , formula (4.13) is nothing but a formula for  $\mathbf{g}(D^k \mathbf{F}_\tau)$  – the analogue of formula (3.14) in the present setting.

*Proof.* — Consider the extended Hopf algebra  $\text{free}T_{\mathbf{F}}^+$  freely generated by the symbols

$$\{X_1, \dots, X_d\} \cup \mathcal{B}_\circ^+ \cup \{D^k(\mathbf{F}_\tau); \tau \in \mathcal{B}, \gamma > |\tau| + |k|\}.$$

It satisfies assumptions (A–C). By Theorem 17 giving a paracontrolled parameterization of the map  $\mathbf{g}$  by its definition on the  $\mathbf{g}(\tau)$ , with  $\tau \in \mathcal{G}_{\mathbf{F}, \circ}^+ := \mathcal{G}_\circ^+ \cup \{\mathbf{F}_\tau; |\tau| < \gamma\}$ , there exists a unique model  $\mathbf{g}$  on  $\text{free}T_{\mathbf{F}}^+$  that coincides with  $\mathbf{g}$  on  $T^+$ , and such that

$$\mathbf{g}(\mathbf{F}_\tau) := \sum_{\substack{\tau \triangleleft \mu \\ |\mu| < \gamma}} P_{\mathbf{g}(\mathbf{F}_\mu)} \llbracket \mu/\tau \rrbracket^{\mathbf{g}} + \llbracket f_\tau \rrbracket,$$

for all  $\tau \in \mathcal{B}$  with  $|\tau| < \gamma$ . Since  $T_{\mathbf{F}}^+$  is the quotient space of  $\text{free}T_{\mathbf{F}}^+$  by the relations (4.12), and

$$\mathbf{g}(D^k \mathbf{F}_\tau) = \sum_{\substack{\tau \triangleleft \mu, |\mu| < \gamma \\ \mu/\tau \in \text{span}(\mathcal{B}_X^+)}} c_\tau^\mu(k) \mathbf{g}(\mathbf{F}_\mu),$$

from the structure condition (4.14), the map  $\mathfrak{g}$  is consistently defined on the quotient space, where it satisfies the estimates from Definition 10.

One can get rid of the structure condition in some cases.

**ASSUMPTION D.** — *For any  $\tau \in \mathcal{B}_\bullet$ , there is no term of the form  $\sigma \otimes X^k$  with  $k \neq 0$ , in the expansion of  $\Delta\tau$ .*

Under assumption (D), we can show that, given  $\tau \in \mathcal{B}$ , the only  $\mu > \tau$  such that  $\mu/\tau$  has a non-null component on  $X^k$  is  $\mu = \underline{X}^k\tau$ . Indeed, writing  $\mu = \underline{X}^\ell\sigma$  ( $\ell \in \mathbb{N}^d$ ,  $\sigma \in \mathcal{B}_\bullet$ ), by the multiplicativity of  $\Delta$  and by (A.2), we have

$$\begin{aligned} (\text{Id} \otimes P_X)\Delta\mu &= (\text{Id} \otimes P_X)(\Delta\underline{X}^\ell)(\Delta\sigma) \\ &= (\text{Id} \otimes P_X)(\Delta\underline{X}^\ell)(\sigma \otimes \mathbf{1}_+) = \sum_k \binom{\ell}{k} \underline{X}^{\ell-k}\sigma \otimes X^k. \end{aligned}$$

Then  $\mu/\tau \in \text{span}(X^k)$  if and only if  $\tau = \underline{X}^{\ell-k}\sigma$ , thus  $\mu = \underline{X}^k\tau$ . Then (4.12) takes the form

$$D^k F_{\underline{X}^{\ell-k}\sigma} = \frac{\ell!}{(\ell-k)!} F_{\underline{X}^\ell\sigma}, \quad \sigma \in \mathcal{B}_\bullet.$$

Moreover, this reduces to the formula

$$D^k F_\sigma = k! F_{\underline{X}^k\sigma}, \quad \sigma \in \mathcal{B}_\bullet,$$

hence the structure condition (4.14) takes the simple form (4.15) below. Note that the data in the next statement is indexed by  $\mathcal{B}_\bullet$ , unlike in the general case of Theorem 22 where it is indexed by  $\mathcal{B}$ .

**COROLLARY 23.** — *Let  $T$  be a regularity structure satisfying assumptions (A–D), and a family  $([f_\tau] \in \mathcal{C}_{\text{rap}}^{\gamma-|\tau|}(\mathbb{R}^d))_{\tau \in \mathcal{B}_\bullet, |\tau| < \gamma}$  be given. Assume that  $\gamma - |\tau| \notin \mathbb{N}$  for any  $\tau \in \mathcal{B}$  with  $|\tau| < \gamma$ . Pick a model  $(\mathfrak{g}, \Pi) \in \mathcal{M}_{\text{rap}}(T, \mathbb{R}^d)$ . Set, for  $\tau \in \mathcal{B}_\bullet$  with  $|\tau| < \gamma$ ,*

$$f_\tau := \sum_{\substack{\tau \triangleleft \mu, |\mu| < \gamma \\ \mu/\tau \in \text{span}(\mathcal{B}^+ \mathcal{B}_X^+)}} P_{f_\mu} [\mu/\tau]^\sharp + [f_\tau],$$

and, for  $\tau \in \mathcal{B}_\bullet, k \in \mathbb{N}^d \setminus \{0\}$  with  $|k| + |\tau| < \gamma$ ,

$$(4.15) \quad f_{\underline{X}^k\tau}(x) := \frac{1}{k!} \partial_y^k \left\{ f_\tau(y) - \sum_{\substack{\tau \triangleleft \mu, |\mu| < \gamma, |\mu/\tau| \triangleleft k \\ \mu/\tau \in \text{span}(\mathcal{B}^+ \mathcal{B}_X^+)}} \mathfrak{g}_{yx}(\mu/\tau) f_\mu(x) \right\} \Big|_{y=x}.$$

Then

$$\mathbf{f} := \sum_{\sigma \in \mathcal{B}, |\sigma| < \gamma} f_\sigma \sigma + \sum_{\substack{\tau \in \mathcal{B}_\bullet, k \in \mathbb{N}^d \\ |\tau| + |k| < \gamma}} f_{\underline{X}^k\tau} \underline{X}^k \tau \in \mathcal{D}_{\text{rap}}^\gamma(T, \mathfrak{g}).$$

Corollary 23 yields the homeomorphism result from Theorem 5. As stated in the introduction, we can see the further homeomorphism result

$$\begin{aligned} M_{\text{rap}} \cap \mathcal{D}_{\text{rap}}^\gamma &:= \{(M, \mathbf{f}); M \in M_{\text{rap}}(T, \mathbb{R}^d), \mathbf{f} \in \mathcal{D}_{\text{rap}}^\gamma(T, \mathfrak{g})\} \\ &\simeq \prod_{\sigma \in \mathcal{G}_\circ^+} \mathcal{C}_{\text{rap}}^{|\sigma|}(\mathbb{R}^d) \times \prod_{\tau \in \mathcal{B}_\bullet, |\tau| < 0} \mathcal{C}_{\text{rap}}^{|\tau|}(\mathbb{R}^d) \times \prod_{\tau \in \mathcal{B}_\bullet, |\tau| < \gamma} \mathcal{C}_{\text{rap}}^{\gamma-|\tau|}(\mathbb{R}^d), \end{aligned}$$

where the left hand side has a topology induced by the metrics

$$\begin{aligned} d_a((M, \mathbf{f}), (M', \mathbf{f}')) \\ := d_a(M, M') + \sup_{\tau \in \mathcal{B}} \left\| \langle \tau', (\mathbf{f}(y) - \widehat{\mathfrak{g}}_{yx} \mathbf{f}(x)) - (\mathbf{f}'(y) - \widehat{\mathfrak{g}}'_{yx} \mathbf{f}'(x)) \rangle \right\|_{\mathcal{C}_{(2),a}^{\gamma-|\tau|}(\mathbb{R}^d \times \mathbb{R}^d)}. \end{aligned}$$

Note that assumption (D) is an assumption about the basis  $\mathcal{B}$  of  $T$  we choose to work with, not about the regularity structure itself. It is thus possible that a given basis satisfies assumption (D) whereas another does not. This flexibility is at the heart of the proof of Theorem 7 in the next section.

**4.3. MODELED DISTRIBUTIONS OVER BHZ REGULARITY STRUCTURES.** — Bruned, Hairer and Zambotti introduced in [7] class of regularity structures convenient for the study of singular stochastic PDEs. We call these structures *BHZ regularity structures*

$$\mathcal{T}_{\text{BHZ}} = ((T_{\text{BHZ}}^+, \Delta_{\text{BHZ}}^+), (T_{\text{BHZ}}, \Delta_{\text{BHZ}})).$$

Although the canonical basis of these concrete regularity structures do not satisfy assumption (D) the following result holds true.

**THEOREM 24.** — *Assume that the set of homogeneities  $\{|\mathfrak{t}|\}_{\mathfrak{t} \in \mathfrak{L}} \cup \{1\}$  is rationally independent, that is, the only tuple of integers  $\{k_{\mathfrak{t}}\}_{\mathfrak{t}} \cup \{k_1\}$  such that  $\sum_{\mathfrak{t}} k_{\mathfrak{t}} |\mathfrak{t}| + k_1 = 0$  is the trivial solution  $k_{\mathfrak{t}} = k_1 = 0$ . Then the canonical bases  $\mathcal{B}_{\text{BHZ}}^+$  and  $\mathcal{B}_{\text{BHZ}}$  satisfy assumptions (A–C). Moreover, one can construct a basis of  $\mathcal{T}_{\text{BHZ}}$  that satisfies assumptions (A–D).*

The remaining of this section is dedicated to proving this statement. We recall first the elements of the construction of BHZ regularity structures that we need here. These concrete regularity structures are indexed by decorated rooted trees.

Any finite connected graph without loops and with a distinguished vertex is called a rooted tree. For any rooted tree  $\tau$ , denote by  $N_\tau$  the node set, by  $E_\tau$  the edge set, by  $\varrho_\tau \in N_\tau$  the distinguished vertex, called root of  $\tau$ . Let also  $\mathfrak{L}$  be a finite set of types. (Edges will be interpreted differently depending on their type, when given any model on  $\mathcal{T}_{\text{BHZ}}$ . Different types may for instance correspond to different convolution operators.) Denote by  $\mathcal{B}$  the set of rooted decorated trees. Each  $\tau \in \mathcal{B}$  is a rooted tree equipped with the type map  $\mathfrak{t} : E_\tau \rightarrow \mathfrak{L}$  and with the decorations

- $\mathfrak{n} : N_\tau \rightarrow \mathbb{N}^d$ .
- $\mathfrak{o} : N_\tau \rightarrow \mathbb{Z}^d \oplus Z(\mathfrak{L})$ .
- $\mathfrak{e} : E_\tau \rightarrow \mathbb{N}^d$ .

Equivalently, the set  $\mathcal{B}$  is generated recursively by the application of the following operations – see [7, §4.3].

- One has  $\bullet^k \in \mathcal{B}$  for any  $k \in \mathbb{N}^d$ , where  $\bullet^k$  is a tree with only one node  $\bullet$ , with  $\mathbf{n}(\bullet) = k$ , and  $\mathfrak{o}(\bullet) = 0 \oplus 0$ .
- If  $\tau, \sigma \in \mathcal{B}$  then  $\tau\sigma \in \mathcal{B}$ , where  $\tau\sigma$  is called a *tree product*;  $\tau\sigma$  is a graph  $\tau \sqcup \sigma$  divided by the equivalence relation  $\sim$  on  $N_\tau \sqcup N_\sigma$ , where  $x \sim y$  means  $x = y$  or  $x, y \in \{\varrho_\tau, \varrho_\sigma\}$ . On the root  $\varrho_{\tau\sigma}$ , the decorations  $\mathbf{n}(\varrho_{\tau\sigma}) = \mathbf{n}(\varrho_\tau) + \mathbf{n}(\varrho_\sigma)$  and  $\mathfrak{o}(\varrho_{\tau\sigma}) = \mathfrak{o}(\varrho_\tau) + \mathfrak{o}(\varrho_\sigma)$  are given.
- For any  $\mathfrak{t} \in \mathfrak{L}$  and  $k \in \mathbb{N}^d$ ,

$$\tau \in \mathcal{B} \implies I_k^{\mathfrak{t}}(\tau) \in \mathcal{B},$$

where the tree  $I_k^{\mathfrak{t}}(\tau)$  is obtained by adding on  $\tau$  one distinguished node  $\varrho'$  and one edge  $e = (\varrho_\tau, \varrho')$  of type  $\mathfrak{t}$ , with decorations  $\mathfrak{e}(e) = k$  and  $\mathfrak{o}(\varrho') = 0 \oplus 0$ .

- For any  $\alpha \in Z^d \oplus Z(\mathfrak{L})$ , denote by  $R_\alpha$  the operator on decorated rooted trees adding a value  $\alpha$  on the decoration  $\mathfrak{o}$  on  $\varrho_\tau$ . Assume

$$\tau \in \mathcal{B} \implies R_\alpha(\tau) \in \mathcal{B}.$$

By applying the operator  $R_\alpha$  with various  $\alpha$  on each step as above, one can see that, if  $\tau \in \mathcal{B}$  then the same decorated tree with any other  $\mathfrak{o}$ -decoration is also an element of  $\mathcal{B}$ .

Each type  $\mathfrak{t} \in \mathfrak{L}$  is assigned a nonzero real number  $|\mathfrak{t}|$ , the collection of which satisfies the assumption of Theorem 24. One assigns a homogeneity  $|\mathbf{n}|, |\mathfrak{o}|, |\mathfrak{e}|, |\mathfrak{t}|$  to the decorations and edge types of any decorated tree  $\tau$ , and set

$$\begin{aligned} |\tau| &:= |\mathbf{n}| + |\mathfrak{o}| - |\mathfrak{e}| + |\mathfrak{t}| \\ &:= \sum_{n \in N_\tau} |\mathbf{n}(n)| + \sum_{n \in N_\tau} |\mathfrak{o}(n)| - \sum_{e \in E_\tau} |\mathfrak{e}(e)| + \sum_{e \in E_\tau} |\mathfrak{t}(e)|, \end{aligned}$$

where  $|a + \sum_{\mathfrak{t}} a_{\mathfrak{t}} \mathfrak{t}| := |a| + \sum_{\mathfrak{t}} a_{\mathfrak{t}} |\mathfrak{t}|$  for  $a + \sum_{\mathfrak{t}} a_{\mathfrak{t}} \mathfrak{t} \in Z^d \oplus Z(\mathfrak{L})$ . A noise-type object  $\Theta$  is represented by  $I_0^{\mathfrak{t}}(\bullet^0)$ , with  $\mathfrak{t}$  of negative homogeneity.

With each subcritical singular stochastic PDE is associated a notion of conforming and strongly conforming decorated tree. The basis  $\mathcal{B}_{\text{BHZ}}$  of  $T_{\text{BHZ}}$  is made up of the set of elements of  $\mathcal{B}$  that strongly conforms the rule (see [7, §5]), and the basis  $\mathcal{B}_{\text{BHZ}}^+$  of  $T_{\text{BHZ}}^+$  is made up of the elements of the form

$$\bullet^k \prod_{i=1}^N I_{k_i}^{\mathfrak{t}_i}(\tau_i),$$

where  $k, k_i \in \mathbb{N}^d$ ,  $\mathfrak{t}_i \in \mathfrak{L}$ ,  $\tau_i \in \mathcal{B}_{\text{BHZ}}$ , and  $|I_{k_i}^{\mathfrak{t}_i}(\tau_i)| > 0$ . Such a tree is said to conform the rule. We do not need more details here and refer the interested reader to [7, §5]. We do not describe in particular the details of the definition of the splitting maps  $\Delta_{\text{BHZ}}$  and  $\Delta_{\text{BHZ}}^+$ ; we only record the following fact, where we write  $\mathbf{1}$  for  $\bullet^0$ , and  $X^k$  for  $\bullet^k$ .

PROPOSITION 25 ([7, Prop. 4.17]). — *The coproduct  $\Delta = \Delta_{\text{BHZ}} : T_{\text{BHZ}} \rightarrow T_{\text{BHZ}} \otimes T_{\text{BHZ}}^+$  satisfies the following identities*

$$\begin{aligned} \Delta \mathbf{1} &= \mathbf{1} \otimes \mathbf{1}, \quad \Delta X_i = X_i \otimes \mathbf{1} + \mathbf{1} \otimes X_i, \quad \Delta(\tau\sigma) = (\Delta\tau)(\Delta\sigma), \\ \Delta I_k^t(\tau) &= (I_k^t \otimes \text{Id})\Delta\tau + \sum_{|\ell|+|k|<|\tau|+|t|} \frac{X^\ell}{\ell!} \otimes I_{k+\ell}^t(\tau), \quad \Delta R_\alpha(\tau) = (R_\alpha \otimes \text{Id})\Delta\tau. \end{aligned}$$

The coproduct  $\Delta^+ = \Delta_{\text{BHZ}}^+ : T_{\text{BHZ}}^+ \rightarrow T_{\text{BHZ}}^+ \otimes T_{\text{BHZ}}^+$ , satisfies the same identities with  $\Delta$  in the right hand sides replaced by  $\Delta^+$ .

THEOREM 26. — *The bases  $\mathcal{B} = \mathcal{B}_{\text{BHZ}}$  and  $\mathcal{B}^+ = \mathcal{B}_{\text{BHZ}}^+$  satisfy assumptions (A–C).*

*Proof.* — Assumption (A) is satisfied by setting

$$\mathcal{B}_\circ^+ := \{I_k^t(\tau) \in \mathcal{B}^+; \mathfrak{t} \in \mathfrak{L}, k \in \mathbb{N}^d, \tau \in \mathcal{B}\}, \quad \mathcal{B}_\bullet := \{\tau \in \mathcal{B}; \mathfrak{n}(\varrho_\tau) = 0\},$$

and  $X^k = \underline{X}^k = \bullet^k$ . Assumption (B) follows because polynomial elements and non-polynomial elements are distinguished by the number of their edges. Indeed,  $\sharp E_\tau = 0$  if and only if  $\tau \in \mathcal{B}_\bullet^+ = \mathcal{B}_\bullet$ . Assumption C(1) is satisfied by setting

$$\mathcal{G}_\circ^+ := \{I_0^t(\tau) \in \mathcal{B}^+; \mathfrak{t} \in \mathfrak{L}, \tau \in \mathcal{B}\}.$$

Then  $D^k I_0^t(\tau) = I_k^t(\tau)$  follows from Proposition 25. To check C(2), we define the binary relation on  $\mathcal{B}^+$  by denoting  $\sigma \vDash \tau$  if

- $\sharp E_\sigma < \sharp E_\tau$ , or
- $\sharp E_\sigma = \sharp E_\tau$  and  $|\mathfrak{n}_\sigma| \subset |\mathfrak{n}_\tau|$ , where  $\mathfrak{n}_\tau$  (resp.  $\mathfrak{n}_\sigma$ ) denotes the  $\mathfrak{n}$ -decoration given for  $\tau$  (resp.  $\sigma$ ).

This relation is transitive and satisfies the first condition of C(2). The second one in C(2) follows from the graphical definition of  $\Delta^+$  – see [7, §2] for details. Essentially, we have the decomposition

$$\Delta^+ \tau = \sum \sigma \otimes (\tau / {}^+ \sigma),$$

where either of the following holds.

- $\sigma$  is the same graph as  $\tau$  but with  $\mathfrak{n}_\sigma \subset \mathfrak{n}_\tau$ .  $\tau / {}^+ \sigma$  consists of only one node.
- $\sigma$  is a strict subtree of  $\tau$  such that  $\varrho_\sigma = \varrho_\tau$ , and  $\tau / {}^+ \sigma$  is a quotient graph of  $\tau$  obtained by contracting the subgraph  $\sigma$  into one node.

For the first case, if  $\mathfrak{n}_\sigma = \mathfrak{n}_\tau$  then  $\sigma = \tau$  as an element of  $\mathcal{B}$ , and if  $\mathfrak{n}_\sigma < \mathfrak{n}_\tau$  then  $\sigma \triangleleft \tau$ . For the second case, if  $\sharp N_\sigma = 0$  then  $\sigma$  is a polynomial and if  $\sharp N_\sigma > 0$  then  $\sigma, \tau / {}^+ \sigma \triangleleft \tau$ . Hence the formula (4.2) holds. For the last assumption, since the set  $\{|\mathfrak{t}|\}_{\mathfrak{t} \in \mathfrak{L}} \cup \{1\}$  is rationally independent, non-polynomial  $\tau$  (hence  $\tau$  has at least one edge) has non-integer homogeneity. Hence C(3) holds.

The canonical bases  $\mathcal{B}_{\text{BHZ}}$  of BHZ concrete regularity structures do not satisfy assumption (D) since one has

$$\Delta I_0^t(X_i \Theta) = I_0^t(X_i \Theta) \otimes \mathbf{1} + I_0^t(\Theta) \otimes X_i + \sum_{|k|<|\Theta|+1+|t|} \frac{X^k}{k!} \otimes I_k^t(X_i \Theta),$$

for any edge type  $\mathfrak{t}$  with positive homogeneity, but the second term in the right hand side contradicts to assumption (D). Now we define another basis of  $T_{\text{BHZ}}$ . Set

$$\mathbf{T} := \text{span}(\mathbf{B}).$$

The tree product  $(\tau, \sigma) \mapsto \tau\sigma$  and the operators  $I_k^{\mathfrak{t}}$  and  $R_\alpha$  are linearly extended to  $\mathbf{T}$ . For any  $\mathfrak{t} \in \mathfrak{L}$  and  $k, \ell \in \mathbb{N}^d$ , we define the new operator  ${}_\ell I_k^{\mathfrak{t}} : \mathbf{T} \rightarrow \mathbf{T}$ , by

$${}_\ell I_k^{\mathfrak{t}}(\tau) := \sum_{m \in \mathbb{N}^d} \binom{\ell}{m} X^m (-1)^{\ell-m} I_k^{\mathfrak{t}}(X^{\ell-m} \tau).$$

(An operator  ${}_\ell I_k$  represents the convolution with a kernel  $x^\ell (\partial^k K)(x)$ . These operators also appeared in the very recent work [15] of Hairer and Pardoux.) If  $\tau$  is homogeneous, then  ${}_\ell I_k^{\mathfrak{t}}(\tau)$  is also homogeneous and

$$|{}_\ell I_k^{\mathfrak{t}}(\tau)| = |\mathfrak{t}| - |k| + |\ell| + |\tau|.$$

LEMMA 27. — Consider the subset  $\tilde{\mathbf{B}}_\bullet \subset \mathbf{T}$  generated by the following rules.

- $\mathbf{1} \in \tilde{\mathbf{B}}_\bullet$ .
- $\tau \in \tilde{\mathbf{B}}_\bullet \Rightarrow {}_\ell I_k^{\mathfrak{t}}(\tau) \in \tilde{\mathbf{B}}_\bullet$ .
- $\tau \in \tilde{\mathbf{B}}_\bullet \Rightarrow R_\alpha(\tau) \in \tilde{\mathbf{B}}_\bullet$ .
- $\tau, \sigma \in \tilde{\mathbf{B}}_\bullet \Rightarrow \tau\sigma \in \tilde{\mathbf{B}}_\bullet$ .

Set

$$\tilde{\mathbf{B}} := \{X^k \tau; k \in \mathbb{N}^d, \tau \in \tilde{\mathbf{B}}_\bullet\}.$$

Then  $\tilde{\mathbf{B}}$  is a linear basis of  $\mathbf{T}$ , and there exists a basis  $\tilde{\mathcal{B}} = \tilde{\mathcal{B}}_{\text{BHZ}}$  of  $T_{\text{BHZ}}$  such that  $\tilde{\mathcal{B}} \subset \tilde{\mathbf{B}}$ .

Proof. — Assume that  $\tau \in \mathbf{B}$  is expanded by the basis  $\tilde{\mathbf{B}}$ , that is,  $\tau$  is of the form

$$\tau = \sum_i a_i X^{k_i} \sigma_i$$

with  $a_i \in \mathbb{R}$ ,  $k_i \in \mathbb{N}^d$ , and  $\sigma_i \in \tilde{\mathbf{B}}_\bullet$ . Since the commutative property  $R_\alpha(X^k \cdot) = X^k R_\alpha(\cdot)$  holds by the definition,  $R_\alpha(\tau)$  is also expanded by  $\tilde{\mathbf{B}}$ . By the inversion formula

$$I_k^{\mathfrak{t}}(X^\ell \sigma) = \sum_{m \in \mathbb{N}^d} \binom{\ell}{m} X^m (-1)^{\ell-m} {}_{\ell-m} I_k^{\mathfrak{t}}(\sigma),$$

$I_k^{\mathfrak{t}}(\tau)$  is also expanded by  $\tilde{\mathbf{B}}$ . Certainly, if  $\tau, \sigma \in \text{span}(\tilde{\mathbf{B}})$ , then  $\tau\sigma \in \text{span}(\tilde{\mathbf{B}})$ . We can conclude that  $\mathbf{T} = \text{span}(\tilde{\mathbf{B}})$  by the induction on the number of edges on  $\tau$ .

As in the definition of  $\mathcal{B}_{\text{BHZ}}$  from  $\mathbf{B}$ , one obtains  $\tilde{\mathcal{B}}$  by keeping only those elements from  $\tilde{\mathbf{B}}$  that strongly conforms.

The set  $\tilde{\mathcal{B}}$  can be encoded as a set of rooted decorated trees using different decorations from the preceding decorations. Each  $\tau \in \tilde{\mathbf{B}}_\bullet$  is represented by a rooted tree with  $\sigma$  and  $\epsilon$  decorations, together with a new decoration

$$\mathfrak{f} : E_\tau \longrightarrow \mathbb{N}^d.$$

The map  ${}_\ell I_k^{\mathfrak{t}} : \tilde{\mathbf{B}}_\bullet \rightarrow \tilde{\mathbf{B}}_\bullet$ , is defined as follows. For any  $\tau \in \tilde{\mathbf{B}}_\bullet$  with root  $\varrho$ , the tree  ${}_\ell I_k^{\mathfrak{t}}(\tau)$  is obtained by adding to  $\tau$  one node  $\varrho'$  and one edge  $e := (\varrho, \varrho')$ , with



decorations  $\mathfrak{e}(e) = k$  and  $\mathfrak{f}(e) = \ell$ . Each  $\tau = X^k \sigma \in \tilde{\mathfrak{B}}$  is represented by a rooted tree with decorations  $\mathfrak{n}, \mathfrak{o}, \mathfrak{e}, \mathfrak{f}$ , where  $\mathfrak{n}$  vanishes at any node except the root, where it is equal to  $k$ . We call this tree representation of elements of  $\tilde{\mathfrak{B}}$  the *non-canonical representation*. Note that, under such exchange of representations, the shape of trees is preserved.

**THEOREM 28.** — *The basis  $\tilde{\mathfrak{B}}$  of  $T_{\text{BHZ}}$  satisfies assumption (D), where  $\tilde{\mathfrak{B}}_{\bullet} = \tilde{\mathfrak{B}}_{\bullet} \cap \tilde{\mathfrak{B}}$ .*

*Proof.* — The proof is done by the induction on the number of edges on  $\tau$  in its non-canonical representation. In fact, one can conclude a stronger claim; for any  $\tau \in \tilde{\mathfrak{B}}_{\bullet}$ , one has

$$(4.16) \quad \Delta\tau = \sum_{\substack{\sigma \in \tilde{\mathfrak{B}}_{\bullet} \\ \eta \in \tilde{\mathfrak{B}} \setminus \{X^k\}_{k \neq 0}}} c_{\sigma\eta}^{\tau} \sigma \otimes \eta.$$

It is sufficient to show that, if the coproduct of  $\tau \in \tilde{\mathfrak{B}}_{\bullet}$  has such a form, then  ${}_{\ell}I_k^t(\tau)$  also satisfies the same condition. To complete the proof, we compute explicitly the coproduct  $\Delta({}_{\ell}I_k^t(\tau))$ . Since

$$\begin{aligned} \Delta I_k^t(X^a \tau) &= (I_k^t \otimes \text{Id})\Delta(X^a \tau) + \sum_{\ell \in \mathbb{N}^d} \frac{X^{\ell}}{\ell!} \otimes I_{k+\ell}^t(X^a \tau) \\ &= \sum_{\substack{\sigma \in \tau \\ b \in \mathbb{N}^d}} \binom{a}{b} I_k^t(X^b \sigma) \otimes X^{a-b}(\tau/\sigma) + \sum_{\ell \in \mathbb{N}^d} \frac{X^{\ell}}{\ell!} \otimes I_{k+\ell}^t(X^a \tau), \end{aligned}$$

we have

$$\begin{aligned} \Delta({}_a I_k^t(\tau)) &= \sum_{b \in \mathbb{N}^d} \binom{a}{b} (\Delta X^b) (-1)^{a-b} \Delta I_k(X^{a-b} \tau) \\ &= \sum_{\substack{\sigma \in \tau \\ b, c, d \in \mathbb{N}^d}} (-1)^{a-b} \binom{a}{b} \binom{b}{c} \binom{a-b}{d} X^c I_k^t(X^d \sigma) \otimes X^{b-c} X^{a-b-d}(\tau/\sigma) \\ &\quad + \sum_{\ell, b, c \in \mathbb{N}^d} (-1)^{a-b} \binom{a}{b} \binom{b}{c} X^c \frac{X^{\ell}}{\ell!} \otimes X^{b-c} I_{k+\ell}^t(X^{a-b} \tau) \\ &=: \text{(i)} + \text{(ii)}. \end{aligned}$$

The term (ii) does not contain any terms of the form  $\sigma \otimes X^k$  with  $k \neq 0$ . The sum (i) is equal to

$$\begin{aligned} &\sum_{\substack{\sigma \in \tau \\ a=c+c'+d+d'}} (-1)^{d+d'} \frac{a!}{c!c'd!d'd!} X^c I_k^t(X^d \sigma) \otimes X^{c'} X^{d'}(\tau/\sigma) \\ &= \sum_{\substack{\sigma \in \tau \\ a=\alpha+\beta}} \frac{a!}{\alpha!\beta!} \left( \sum_{\alpha=c+d} (-1)^d \frac{\alpha!}{c!d!} X^c I_k^t(X^d \sigma) \right) \otimes \left( \sum_{\beta=c'+d'} (-1)^{d'} \frac{\beta!}{c'd!d!} X^{c'} X^{d'}(\tau/\sigma) \right) \\ &= \sum_{\substack{\sigma \in \tau \\ a=\alpha+\beta}} \binom{a}{\alpha} {}_{\alpha} I_k^t(\sigma) \otimes (X - X)^{\beta}(\tau/\sigma) = \sum_{\sigma \in \tau} {}_a I_k^t(\sigma) \otimes (\tau/\sigma) = ({}_a I_k^t \otimes \text{Id})\Delta\tau. \end{aligned}$$

Since  $\tau$  is assumed in the induction step to have a coproduct (4.16), hence  $\Delta(\ell I_k^t(\tau))$ , enjoys the same property.

**4.4. DENSITY AND EXTENSION COROLLARIES.** — Corollaries 2, 3, 4, and 6 are proved as follows. Note that Schwartz space  $\mathcal{S}(\mathbb{R}^d)$  is dense in the space  $C_{\text{rap}}^\beta(\mathbb{R}^d)$  in the topology of  $C_{\text{rap}}^{\beta-\varepsilon}(\mathbb{R}^d)$  for any  $\varepsilon > 0$ ; for any  $f \in C_{\text{rap}}^\beta(\mathbb{R}^d)$ , the function  $e^{t\Delta}f$  belongs to  $\mathcal{S}(\mathbb{R}^d)$  and satisfies

$$\|e^{t\Delta}f - f\|_{C_a^{\beta-\varepsilon}(\mathbb{R}^d)} \cdot t^{2\varepsilon} \|f\|_{C_a^\beta(\mathbb{R}^d)} \xrightarrow{t \rightarrow 0} 0$$

for any  $a > 0$ .

*Proof of Corollary 2.* — By Theorem 1, the space  $M_{\text{rap}}(T, \mathbb{R}^d)$  is homeomorphic to the product space

$$\prod_{\sigma \in \mathcal{S}_\circ^+} \mathcal{C}_{\text{rap}}^{|\sigma|}(\mathbb{R}^d) \times \prod_{\substack{\tau \in \mathcal{B}_\bullet \\ |\tau| < 0}} \mathcal{C}_{\text{rap}}^{|\tau|}(\mathbb{R}^d).$$

For any  $\varepsilon > 0$ , any elements of this space can be approximated by smooth elements in the topology of the same space with each exponent  $|\tau|$  replaced by  $|\tau| - \varepsilon$ . By the formulas (3.5) and (3.6), it turns out that a smooth element of (1.3) is transferred to a smooth model in  $M_{\text{rap}}(T, \mathbb{R}^d)$ .

The proof of Corollary 6 is completely parallel and left to the reader.

*Proof of Corollaries 3 and 4.* — For Corollary 3, consider the algebra  $T_+$  generated by the set  $\mathcal{B}_\circ^+$  of rooted trees as in Remark 1 of the previous section. Given an  $\mathbb{R}^\ell$ -valued  $\alpha$ -Hölder function  $h = (h_i)_{i=1}^d$ , a lift of the control  $h$  is a branched rough path  $(H^\tau)_{\tau \in \mathcal{B}_\circ^+}$  such that  $H^{\bullet i} = h_i$ , where  $\bullet_i$  denotes a graph with only one node and with node decoration  $i$ . By Theorem 1, such a lift is transferred to an elements of the product space  $\prod_{\tau \in \mathcal{B}_\circ^+} \mathcal{C}_{\text{rap}}^{|\tau|}(\mathbb{R})$  such that  $\llbracket \bullet_i \rrbracket = h_i$ . A trivial extension is defined by  $\llbracket \tau \rrbracket = 0$  if  $\sharp\tau > 2$ , and the associated model is nothing but a trivial lift of  $h$ .

Corollary 4 is proved by a similar argument. By admissibility, the set  $M(T, \mathbb{R}^m)$  is homeomorphic to the space  $\prod_{\tau \in \mathcal{B}_\bullet, |\tau| < 0} \mathcal{C}_{\text{rap}}^{|\tau|}(\mathbb{R}^m)$ . Given a multi-dimensional noise  $(\zeta_j)_{j=1}^\ell$ , a trivial extension is defined by

$$\llbracket \tau \rrbracket^{\mathfrak{g}} = \begin{cases} \zeta_j, & \tau = \bullet_j, \\ 0, & \text{otherwise.} \end{cases}$$

### APPENDIX A. CONCRETE REGULARITY STRUCTURES

We recall in this appendix the setting of concrete regularity structures introduced in [5], and refer the reader to [5, §2] for motivations for the introduction of that setting.

**DEFINITION.** — A concrete regularity structure  $T = (T^+, T)$  is the pair of graded vector spaces

$$T^+ = \bigoplus_{\alpha \in A^+} T_\alpha^+, \quad T = \bigoplus_{\beta \in A} T_\beta$$

such that the following holds.

– The index set  $A^+ \subset \mathbb{R}_+$  contains the point 0, and  $A^+ + A^+ \subset A^+$ ; the index set  $A \subset \mathbb{R}$  is bounded below, and both  $A^+$  and  $A$  have no accumulation points in  $\mathbb{R}$ . Set

$$\beta_0 := \min A.$$

- The vector spaces  $T_\alpha^+$  and  $T_\beta$  are finite dimensional.
- The set  $T^+$  is an algebra with unit  $\mathbf{1}_+$ , with an algebra morphism

$$\Delta^+ : T^+ \longrightarrow T^+ \otimes T^+,$$

such that  $\Delta^+ \mathbf{1}_+ = \mathbf{1}_+ \otimes \mathbf{1}_+$ , and, for  $\tau \in T_\alpha^+$ ,

$$(A.1) \quad \Delta^+ \tau \in \left\{ \tau \otimes \mathbf{1}_+ + \mathbf{1}_+ \otimes \tau + \sum_{0 < \beta < \alpha} T_\beta^+ \otimes T_{\alpha-\beta}^+ \right\},$$

and  $\Delta^+$  satisfies the coassociativity property

$$(\Delta^+ \otimes \text{Id})\Delta^+ = (\text{Id} \otimes \Delta^+)\Delta^+.$$

That is,  $T^+$  has a Hopf structure with coproduct  $\Delta^+$  and counit  $\mathbf{1}'_+$ .

- One has  $T_0^+ = \text{span}(\mathbf{1}_+)$ , and for any  $\alpha, \beta \in A^+$ , one has  $T_\alpha^+ T_\beta^+ \subset T_{\alpha+\beta}^+$ .
- One has a linear splitting map

$$\Delta : T \longrightarrow T \otimes T^+,$$

of the form

$$(A.2) \quad \Delta \tau \in \left\{ \tau \otimes \mathbf{1}_+ + \sum_{\beta < \alpha} T_\beta \otimes T_{\alpha-\beta}^+ \right\}$$

for each  $\tau \in T_\alpha$ , with the right comodule property

$$(\Delta \otimes \text{Id})\Delta = (\text{Id} \otimes \Delta^+)\Delta.$$

Let  $\mathcal{B}_\alpha^+$  and  $\mathcal{B}_\beta$  be bases of  $T_\alpha^+$  and  $T_\beta$ , respectively. We assume  $\mathcal{B}_0^+ = \{\mathbf{1}_+\}$ . Set

$$\mathcal{B}^+ := \bigcup_{\alpha \in A^+} \mathcal{B}_\alpha^+, \quad \mathcal{B} := \bigcup_{\beta \in A} \mathcal{B}_\beta.$$

An element  $\tau$  of  $T_\alpha^{(+)}$  is said to be *homogeneous* and is assigned *homogeneity*  $|\tau| := \alpha$ . The homogeneity of a generic element  $\tau \in T^{(+)}$  is defined as  $|\tau| := \max\{\alpha\}$ , such that  $\tau$  has a non-null component in  $T_\alpha^{(+)}$ . We denote by

$$\mathcal{T} := ((T^+, \Delta^+), (T, \Delta))$$

a concrete regularity structure.

One of the elementary and important examples is the Taylor polynomial ring. Consider symbols  $X_1, \dots, X_d$  and set

$$T_X := \mathbb{R}[X_1, \dots, X_d].$$

For a multi index  $k = (k_i)_{i=1}^d \in \mathbb{N}^d$ , we use the notation

$$X^k := X_1^{k_1} \dots X_d^{k_d}.$$

We define the homogeneity  $|X^k| = |k| := \sum_i k_i$ , and the coproduct

$$(A.3) \quad \Delta X_i = X_i \otimes \mathbf{1} + \mathbf{1} \otimes X_i.$$

Then  $((T_X, \Delta), (T_X, \Delta))$  is a concrete regularity structure.

The set  $G^+$  of characters  $g : T^+ \rightarrow \mathbb{R}$ , i.e., nonzero algebra morphisms, forms a group with the convolution product

$$g_1 * g_2 := (g_1 \otimes g_2)\Delta^+.$$

APPENDIX B. TECHNICAL ESTIMATES

We provide in this appendix a number of technical estimates that are variations on the corresponding results from [5]. Proofs are given for completeness.

LEMMA 29. — *If  $\alpha > 0$  and  $a \in \mathbb{Z}$ , then*

$$\int |P_i(x - y)| |x - y|^\alpha |y|_*^{-a} dy \cdot 2^{-i\alpha} |x|_*^{-a},$$

$$\int |Q_i(x - y)| |x - y|^\alpha |y|_*^{-a} dy \cdot 2^{-i\alpha} |x|_*^{-a}.$$

*Proof.* — Recall the inequalities in the beginning of Section 2. If  $a > 0$ ,

$$\begin{aligned} |x|_*^\alpha \int |P_i(x - y)| |x - y|^\alpha |y|_*^{-a} dy &\cdot \int |P_i(x - y)| |x - y|^\alpha |x - y|_*^a dy \\ &= \int |P_i(y)| |y|^\alpha |y|_*^a dy = \int |P_0(y)| \left| \frac{y}{2^i} \right|^\alpha \left| \frac{y}{2^i} \right|_*^a dy \\ &\lesssim 2^{-i\alpha} \int |P_0(y)| |y|^\alpha |y|_*^a dy \cdot 2^{-i\alpha}. \end{aligned}$$

If  $a < 0$ ,

$$\int |P_i(x - y)| |x - y|^\alpha |y|_*^{-a} dy \cdot |x|_*^{-a} \int |P_i(x - y)| |x - y|^\alpha |x - y|_*^{-a} dy \cdot 2^{-i\alpha} |x|_*^{-a}.$$

As a consequence of Lemma 29, we have the inequality

$$\|\Delta_j f\|_{L_a^\infty} \lesssim \sup_x |x|_*^a \int |Q_j(x - y)| |f(y)| dy \lesssim \|f\|_{L_a^\infty} \sup_x |x|_*^a \int |Q_j(x - y)| |y|_*^{-a} dy \cdot \|f\|_{L_a^\infty}.$$

for any  $a \in \mathbb{Z}$ . This ensures that  $S$  maps  $\mathcal{C}_a^\alpha(\mathbb{R}^d)$  to  $\mathcal{C}_a^\infty(\mathbb{R}^d)$  for any  $\alpha \in \mathbb{R}$ .

Recall the two-parameter extension of the paraproduct, used in [5]. For any distribution  $\Lambda$  on  $\mathbb{R}^d \times \mathbb{R}^d$ , we define

$$(\mathbf{Q}_j \Lambda)(x) := \iint_{\mathbb{R}^d \times \mathbb{R}^d} P_j(x - y) Q_j(x - z) \Lambda(y, z) dy dz,$$

$$(\mathbf{P} \Lambda)(x) := \sum_{j > 1} (\mathbf{Q}_j \Lambda)(x).$$

If  $\Lambda(y, z)$  is of the form  $f(y)g(z)$ , then  $\mathbf{P} \Lambda = \mathbf{P}_f g$ .

PROPOSITION 30 ([5, Prop. 8(a)]). — *Fix  $a \in \mathbb{Z}$ .*

(a) *For any  $\Lambda \in \mathcal{S}'(\mathbb{R}^d \times \mathbb{R}^d)$  for which there exists  $\alpha \in \mathbb{R}$  such that  $\|\mathbf{Q}_j \Lambda\|_{L_a^\infty(\mathbb{R}^d)} \cdot 2^{-j\alpha}$ , for all  $j > 1$ , one has  $\mathbf{P} \Lambda \in \mathcal{C}_a^\alpha(\mathbb{R}^d)$  and*

$$\|\mathbf{P} \Lambda\|_{\mathcal{C}_a^\alpha(\mathbb{R}^d)} \cdot \sup_{j > 1} 2^{j\alpha} \|\mathbf{Q}_j \Lambda\|_{L_a^\infty(\mathbb{R}^d)}.$$

(b) *For any  $\alpha > 0$  and  $F \in \mathcal{C}_{(2),a}^\alpha(\mathbb{R}^d \times \mathbb{R}^d)$ , one has  $\mathbf{P} F \in \mathcal{C}_a^\alpha(\mathbb{R}^d)$  and*

$$\|\mathbf{P} F\|_{\mathcal{C}_a^\alpha(\mathbb{R}^d)} \cdot \|F\|_{\mathcal{C}_{(2),a}^\alpha(\mathbb{R}^d \times \mathbb{R}^d)}.$$

*Proof*

(a) Since  $F P_j$  is supported in the annulus  $\{\lambda \in \mathbb{R}^d; |\lambda| < 2^j \times 2/3\}$  and  $F Q_j$  is supported in the annulus  $\{\lambda \in \mathbb{R}^d; 2^j \times 3/4 < |\lambda| < 2^j \times 8/3\}$ , the integral

$$\int Q_i(x-w)P_j(w-y)Q_j(w-z)dw$$

vanishes if  $|i-j| > 5$ . Hence  $\Delta_i(\mathbf{P}\Lambda) = \sum_{|i-j| \leq 4} \Delta_i(\mathbf{Q}_j\Lambda)$  and we have

$$\|\Delta_i(\mathbf{P}\Lambda)\|_{L^\infty_\alpha} \leq \sum_{|i-j| \leq 4} \|\Delta_i(\mathbf{Q}_j\Lambda)\|_{L^\infty_\alpha} \cdot \sum_{|i-j| \leq 4} \|\mathbf{Q}_j\Lambda\|_{L^\infty_\alpha} \cdot \sum_{|i-j| \leq 4} 2^{-\alpha j} \cdot 2^{-\alpha i}.$$

For (b) it is sufficient to show that  $\|\mathbf{Q}_j F\|_{L^\infty(\mathbb{R}^d)} \leq 2^{-j\alpha}$ . By Lemma 29,

$$\begin{aligned} |\mathbf{Q}_j F(x)| &\leq \int |P_j(x-y)Q_j(x-z)| (|y|_*^{-\alpha} + |z|_*^{-\alpha}) |y-z|^\alpha dydz \\ &\quad + \int |P_j(x-y)Q_j(x-z)| (|y|_*^{-\alpha} + |z|_*^{-\alpha}) (|x-y|^\alpha + |x-z|^\alpha) dydz \\ &\leq 2^{-j\alpha} |x|_*^{-\alpha}. \end{aligned}$$

Recall from [3] the definition of the operator

$$\mathbf{R}^\circ(f, g, h) := \mathbf{P}_f \mathbf{P}_g h - \mathbf{P}_{fg} h.$$

This operator is continuous from  $\mathcal{C}^\alpha(\mathbb{R}^d) \times \mathcal{C}^\beta(\mathbb{R}^d) \times \mathcal{C}^\gamma(\mathbb{R}^d)$  into  $\mathcal{C}^{\alpha+\beta+\gamma}(\mathbb{R}^d)$ , for any  $\alpha, \beta \in [0, 1]$  and  $\gamma \in \mathbb{R}$  – see Proposition 14 therein.

**PROPOSITION 31** ([5, Prop. 10]). — *Consider a function  $f \in L^\infty_{\text{slow}}(\mathbb{R}^d)$  and a finite family  $(a_k, b_k)_{1 \leq k \leq N}$  in  $L^\infty_{\text{slow}}(\mathbb{R}^d) \times L^\infty_{\text{slow}}(\mathbb{R}^d)$  such that*

$$f(y) - f(x) = \sum_{k=1}^N a_k(x)(b_k(y) - b_k(x)) + f^\sharp_{yx}, \quad x, y \in \mathbb{R}^d,$$

*with a remainder  $f^\sharp_{yx}$ . Let  $\alpha > 0$  and  $\beta \in \mathbb{R}$  be given. Assume that either of the following assumptions holds.*

- (a)  $f \in L^\infty_{\text{rap}}(\mathbb{R}^d)$ ,  $a_k b_k \in L^\infty_{\text{rap}}(\mathbb{R}^d)$ ,  $f^\sharp \in \mathcal{C}^{\alpha}_{(2), \text{rap}}(\mathbb{R}^d \times \mathbb{R}^d)$ , and  $g \in \mathcal{C}^\beta_{\text{slow}}(\mathbb{R}^d)$ .
- (b)  $f^\sharp \in \mathcal{C}^\alpha_{(2)}(\mathbb{R}^d \times \mathbb{R}^d)$  and  $g \in \mathcal{C}^\beta_{\text{rap}}(\mathbb{R}^d)$ .

*Then one has the estimate*

$$\sum_{k=1}^N \mathbf{R}^\circ(a_k, b_k, g) \in \mathcal{C}^{\alpha+\beta}_{\text{rap}}(\mathbb{R}^d).$$

*Proof.* — Recall from identity (2.1) the definition of the operator  $\mathbf{S}$ . As in the proof of [5, Prop. 10], we see that

$$\sum_k \mathbf{R}^\circ(a_k, b_k, g) = -\mathbf{S}(\mathbf{P}_f g) + \mathbf{P}_f(\mathbf{S}g) - \sum_k \mathbf{P}_{a_k b_k}(\mathbf{S}g) - \mathbf{P}_{x,y}((\mathbf{P}_{f^\sharp_x} g)(y)).$$

The first three terms belong to  $\mathcal{C}_{\text{rap}}^\infty(\mathbb{R}^d)$ , assuming either (a) or (b). Consider the last term. Note that

$$\mathbf{Q}_j\left((P_{f_{\cdot x}^\#}g)(y)\right)(z) = \sum_{|i-j|\leq 4} \int P_j(z-x)Q_j(z-y)(S_i f_{\cdot x}^\#)(y) (\Delta_i gr)(y) dx dy.$$

For case (a), there exists  $b \in \mathbb{N}$  such that  $|\Delta_i gl(y)| \leq 2^{-i\beta} |y|_*^b$ . Since  $f^\# \in \mathcal{C}_{a+b}^\alpha(\mathbb{R}^d \times \mathbb{R}^d)$  for any  $a \in \mathbb{N}$ , one has

$$\begin{aligned} |(S_i f_{\cdot x}^\#)(y)| &\leq \int |P_i(y-u)| |f_{u,x}^\#| du \cdot \int |P_i(y-u)| |u-x|^\alpha (|u|_*^{-a-b} + |x|_*^{-a-b}) du \\ &\cdot \int |P_i(y-u)| (|u-y|^\alpha + |y-x|^\alpha) (|u|_*^{-a-b} + |x|_*^{-a-b}) du \\ &\cdot (|x|_*^{-a-b} + |y|_*^{-a-b}) (2^{-i\alpha} + |y-x|^\alpha) \end{aligned}$$

by Lemma 29. Hence we have

$$\begin{aligned} \left| \mathbf{Q}_j\left((P_{f_{\cdot x}^\#}g)(y)\right)(z) \right| &\cdot \sum_{|i-j|\leq 4} \int |P_j(z-x)| |Q_j(z-y)| |(S_i f_{\cdot x}^\#)(y)| |(\Delta_i gr)(y)| dx dy \\ &\cdot \sum_{|i-j|\leq 4} \int |P_j(z-x)| |Q_j(z-y)| (|x|_*^{-a-b} + |y|_*^{-a-b}) |y|_*^b \\ &\quad \times (2^{-i\alpha} + |y-x|^\alpha) 2^{-i\beta} dx dy \\ &\cdot \sum_{|i-j|\leq 4} \int |P_j(z-x)| |Q_j(z-y)| (|x|_*^{-a-b} |y|_*^b + |y|_*^{-a}) \\ &\quad \times (2^{-i\alpha} + |z-x|^\alpha + |z-y|^\alpha) 2^{-i\beta} dx dy \\ &\cdot \sum_{|i-j|\leq 4} |z|_*^{-a} (2^{-i\alpha} + 2^{-j\alpha}) 2^{-i\beta} \cdot |z|_*^{-a} 2^{-j(\alpha+\beta)}. \end{aligned}$$

For case (b), since  $|\Delta_i gl(y)| \leq 2^{-i\beta} |y|_*^{-a}$  for any  $a \in \mathbb{N}$ , and

$$|(S_i f_{\cdot x}^\#)(y)| \leq \int |P_i(y-u)| |f_{u,x}^\#| du \cdot \int |P_i(y-u)| |u-x|^\alpha du \cdot 2^{-i\alpha} + |y-x|^\alpha,$$

we have

$$\begin{aligned} \left| \mathbf{Q}_j\left((P_{f_{\cdot x}^\#}g)(y)\right)(z) \right| &\cdot \sum_{|i-j|\leq 4} \int |P_j(z-x)| |Q_j(z-y)| |(S_i f_{\cdot x}^\#)(y)| |(\Delta_i gr)(y)| dx dy \\ &\cdot \sum_{|i-j|\leq 4} \int |P_j(z-x)| |Q_j(z-y)| |y|_*^{-a} (2^{-i\alpha} + |y-x|^\alpha) 2^{-i\beta} dx dy \\ &\cdot \sum_{|i-j|\leq 4} \int |P_j(z-x)| |Q_j(z-y)| |y|_*^{-a} (2^{-i\alpha} + |z-x|^\alpha + |z-y|^\alpha) 2^{-i\beta} dx dy \\ &\cdot \sum_{|i-j|\leq 4} |z|_*^{-a} (2^{-i\alpha} + 2^{-j\alpha}) 2^{-i\beta} \cdot |z|_*^{-a} 2^{-j(\alpha+\beta)}. \end{aligned}$$

By Proposition 30, we are done.

**PROPOSITION 32** ([5, Prop. 9]). — *Let  $\gamma \in \mathbb{R}$  and  $\beta_0 \in \mathbb{R}$  be given together with a family  $\Lambda_x$  of distributions on  $\mathbb{R}^d$ , indexed by  $x \in \mathbb{R}^d$ . Assume one has*

$$\sup_{x \in \mathbb{R}^d} |x|_*^a \|\Lambda_x\|_{\mathcal{C}^{\beta_0}} < \infty$$

for any  $a \in \mathbb{Z}$  and one can decompose  $(\Lambda_y - \Lambda_x)$  under the form

$$\Lambda_y - \Lambda_x = \sum_{\ell=1}^L c_{yx}^\ell \Theta_x^\ell$$

for  $L$  finite,  $\mathbb{R}^d$ -indexed distributions  $\Theta_x^\ell$ , and real-valued coefficients  $c_{yx}^\ell$  depending measurably on  $x$  and  $y$ . Assume that for each  $\ell$  there exists  $\beta_\ell < \gamma$  such that either of the following conditions holds.

(a)  $\Theta^\ell \in D_{\text{rap}}^{\beta_\ell}$  and  $c^\ell \in \mathcal{C}_{(2)}^{\gamma-\beta_\ell}(\mathbb{R}^d \times \mathbb{R}^d)$ .

(b)  $\Theta^\ell \in D^{\beta_\ell}$  and  $c^\ell \in \mathcal{C}_{(2),\text{rap}}^{\gamma-\beta_\ell}(\mathbb{R}^d \times \mathbb{R}^d)$ .

Write  $\mathbf{P}(\Lambda)$  for  $\mathbf{P}_{y,z}(\Lambda_y(z))$  below.

(i) If  $\gamma > 0$ , then there exists a unique function  $\lambda \in \mathcal{C}_{\text{rap}}^\gamma(\mathbb{R}^d)$  such that

$$\{(\mathbf{P}(\Lambda) - \lambda) - \Lambda_x\}_{x \in \mathbb{R}^d} \in D_{\text{rap}}^\gamma.$$

(ii) If  $\gamma < 0$ , then

$$\{\mathbf{P}(\Lambda) - \Lambda_x\}_{x \in \mathbb{R}^d} \in D_{\text{rap}}^\gamma.$$

Consequently,  $\mathbf{P}(\Lambda) \in \mathcal{C}_{\text{rap}}^{\beta_0}(\mathbb{R}^d)$ . If furthermore  $\Lambda \in D_{\text{rap}}^\gamma$ , then  $\mathbf{P}(\Lambda) \in \mathcal{C}_{\text{rap}}^\gamma(\mathbb{R}^d)$ .

*Proof.* — In view of [5, Prop. 9], it is sufficient to show that

$$(B.1) \quad \sup_{x \in \mathbb{R}^d} |x|_*^a \left| \Delta_j(\mathbf{P}(\Lambda) - \Lambda_x)(x) \right| \leq 2^{-j\gamma}.$$

We write for that purpose

$$\begin{aligned} \Delta_i(\mathbf{P}(\Lambda) - \Lambda_x)(x) &= \sum_{\substack{j>1 \\ |i-j| \leq 4}} \iiint Q_i(x-y) P_j(y-u) Q_j(y-v) (\Lambda_u - \Lambda_x)(v) dydudv \\ &\quad - \Delta_i \mathbf{S}(\Lambda_x)(x) \\ &=: A + B + C, \end{aligned}$$

where

$$\begin{aligned} A &= \sum_{|i-j| \leq 4} \iiint Q_i(x-y) P_j(y-u) Q_j(y-v) (\Lambda_u - \Lambda_y)(v) dydudv \\ &= \sum_{|i-j| \leq 4} \sum_{\ell=1}^L \iiint Q_i(x-y) P_j(y-u) Q_j(y-v) c_{uy}^\ell \Theta_y(v) dydudv \end{aligned}$$

and

$$\begin{aligned}
 B &= \sum_{|i-j| \leq 4} \iiint Q_i(x-y)P_j(y-u)Q_j(y-v)(\Lambda_y - \Lambda_x)(v) dydudv \\
 &= \sum_{|i-j| \leq 4} \sum_{\ell=1}^L \iiint Q_i(x-y)P_j(y-u)Q_j(y-v)c_{yx}^\ell \Theta_x(v) dydudv.
 \end{aligned}$$

For the C term,

$$\begin{aligned}
 \sup_x |x|_*^a |\Delta_i S(\Lambda_x)(x)| &\cdot 2^{-ir} \sup_x |x|_*^a \|S(\Lambda_x)\|_{\mathcal{C}^r} \\
 &\cdot 2^{-ir} \sup_x |x|_*^a \|\Lambda_x\|_{\mathcal{C}^{\beta_0}} \cdot 2^{-ir}
 \end{aligned}$$

for any  $r > 0$ . For the A term, for any  $a \in \mathbb{Z}$  we have

$$\begin{aligned}
 |A| &\leq \sum_{|i-j| \leq 4} \sum_{\ell=1}^L \iint |Q_i(x-y)||P_j(y-u)|c_{uy}^\ell |\Delta_j \Theta_y(y)| dydu \\
 &\cdot \begin{cases} \sum_{|i-j| \leq 4} \sum_{\ell=1}^L \iint |Q_i(x-y)||P_j(y-u)||u-y|^{\gamma-\beta_\ell} |y|_*^{-a} 2^{-j\beta_\ell} dydu, & \text{if (a)} \\ \sum_{|i-j| \leq 4} \sum_{\ell=1}^L \iint |Q_i(x-y)||P_j(y-u)|(|u|_*^{-a} + |y|_*^{-a})|u-y|^{\gamma-\beta_\ell} 2^{-j\beta_\ell} dydu, & \text{if (b)} \end{cases} \\
 &\cdot \sum_{|i-j| \leq 4} \int |Q_i(x-y)||y|_*^{-a} 2^{-j\gamma} dy \cdot |x|_*^{-a} 2^{-i\gamma}.
 \end{aligned}$$

The B term has the same estimate by a similar argument. So estimate (B.1) follows from Lemma 29.

(i) If  $\gamma > 0$ , the estimate (B.1) implies that the sum

$$\lambda(x) := \sum_{j > -1} \Delta_j(\mathbf{P}(\Lambda) - \Lambda_x)(x)$$

defines an element  $\lambda$  of  $\mathcal{C}_{\text{rap}}^\gamma(\mathbb{R}^d)$ . To show it, we follow the argument in [13, §6]. We decompose  $\lambda = \lambda^{\leq j+1} + \lambda^{> j+1}$ , where

$$\lambda^{\leq j+1}(x) := \sum_{i \leq j+1} \Delta_i(\mathbf{P}(\Lambda) - \Lambda_x)(x) = S_{j+3}(\mathbf{P}(\Lambda) - \Lambda_x)(x).$$

We consider  $\Delta_j \lambda = \Delta_j \lambda^{\leq j+1} + \Delta_j \lambda^{> j+1}$ . For the second term, by the estimate (B.1) one has

$$\|\Delta_j \lambda^{> j+1}\|_{L_a^\infty} \cdot \|\lambda^{> j+1}\|_{L_a^\infty} \cdot \sum_{i > j+1} 2^{-i\gamma} \cdot 2^{-j\gamma}.$$



For the first term, since  $\Delta_j S_{j+3} = \Delta_j$ , one has

$$\begin{aligned} \Delta_j \lambda^{\zeta_{j+1}}(y) &= \int Q_j(y-x) S_{j+3}(\mathbf{P}(\Lambda) - \Lambda_x)(x) dx \\ &= \int Q_j(y-x) S_{j+3} \left( \mathbf{P}(\Lambda) - \Lambda_y + \sum_{\ell} c_{yx}^{\ell} \Theta_x^{\ell} \right) (x) dx \\ &= \Delta_j(\mathbf{P}(\Lambda) - \Lambda_y)(y) + \sum_{\ell} \int Q_j(y-x) c_{yx}^{\ell} (S_{j+3} \Theta_x)(x) dx. \end{aligned}$$

Similarly to above, we can show that  $|\Delta_j \lambda^{\zeta_{j+1}}(y)| \leq |y|_*^{-a} 2^{-j\gamma}$  for any  $a \in \mathbb{Z}$ . In the end we have  $\|\Delta_j \lambda\|_{L_a^{\infty}} \leq 2^{-j\gamma}$ , hence  $\lambda \in \mathcal{C}_{\text{rap}}^{\gamma}(\mathbb{R}^d)$ .

Since  $\sum_{i>-1} \Delta_i(\mathbf{P}(\Lambda) - \Lambda_x - \lambda)(x) = 0$  by definition, we have

$$\begin{aligned} |S_i(\mathbf{P}(\Lambda) - \Lambda_x - \lambda)(x)| &\leq \sum_{j>i-1} |\Delta_j(\mathbf{P}(\Lambda) - \Lambda_x - \lambda)(x)| \\ &\leq |x|_*^{-a} \sum_{j>i-1} 2^{-j\gamma} \leq |x|_*^{-a} 2^{-i\gamma} \end{aligned}$$

for any  $a \in \mathbb{Z}$ .

(ii) If  $\gamma < 0$ , then directly from (B.1),

$$|S_i(\mathbf{P}(\Lambda) - \Lambda_x)(x)| \leq \sum_{j<i-1} |\Delta_j(\mathbf{P}(\Lambda) - \Lambda_x)(x)| \leq |x|_*^{-a} \sum_{j<i-1} 2^{-j\gamma} \leq |x|_*^{-a} 2^{-i\gamma}$$

for any  $a \in \mathbb{Z}$ .

**COROLLARY 33.** — *Given a concrete regularity structure  $\mathcal{T}$  satisfying assumptions (A) and (B) and given a rapidly decreasing model  $\mathbf{M} = (\mathbf{g}, \Pi)$ , we define the map  $\mathbf{R} : \mathcal{D}_{\text{rap}}^{\gamma}(\mathcal{T}, \mathbf{g}) \rightarrow \mathcal{C}_{\text{rap}}^{\beta_0}(\mathbb{R}^d)$ , by*

$$\mathbf{R}\mathbf{f} = \mathbf{P}_{x,y} \left( (\Pi_x^{\mathbf{g}} \mathbf{f}(x))(y) \right).$$

Then one has

$$(\mathbf{R}\mathbf{f} - \Pi_x^{\mathbf{g}} \mathbf{f}(x))_{x \in \mathbb{R}^d} \in D_{\text{rap}}^{\gamma}.$$

*Proof.* — Let  $\Lambda_x = \Pi_x^{\mathbf{g}} \mathbf{f}(x)$ . Since

$$\Lambda_y - \Lambda_x = \sum_{\tau \in \mathcal{B}} \langle \tau', \widehat{\mathbf{g}}_{xy} \mathbf{f}(y) - \mathbf{f}(x) \rangle \Pi_x^{\mathbf{g}} \tau$$

one gets conditions (a) and (b) of Proposition 32 from the definition of a model.

*Proof of Theorem 12.* — We prove the case  $m = 0$  here for simplicity. For general  $m$ , the proof is at the end of this appendix, after we introduce the modified paraproducts.

Consider the first formula (3.5). First we show that, for each  $\tau \in \mathcal{B}^+$  we have

$$(B.2) \quad \mathbf{g}(\tau) = \sum_{\substack{1 < \nu < \tau \\ \nu \in \mathcal{B}^+}} \mathbf{P}_{\mathbf{g}(\tau+\nu)}[\nu]^{\mathbf{g}} + [\tau]^{\mathbf{g}},$$

where  $[\nu]^g \in \mathcal{C}_{\text{rap}}^{|\nu|}(\mathbb{R}^d)$ , if  $\nu \in \mathcal{B}^+ \cap \mathcal{B}_X^+$ , and  $[\nu]^g \in \mathcal{C}_{\text{slow}}^\infty(\mathbb{R}^d)$ , if  $\nu \in \mathcal{B}_X^+$ . If  $\tau = X^k$ , then since  $\Delta^+ X^k = \sum_{\ell} \binom{k}{\ell} X^\ell \otimes X^{k-\ell}$  we have

$$g(X^k) = \sum_{0 < \ell < k} \binom{k}{\ell} P_{g(X^\ell)} [X^{k-\ell}]^g + [X^k]^g.$$

We see for instance that  $[1]^g = 1$ , then  $[X]^g = x$ , since  $g_x(X) = x$ , and since  $g_x(X^2) = x^2$ , one has

$$x^2 = 2P_x x + [X^2]^g.$$

We recognize  $[X^2]^g = \Pi(x, x)$ . More generally, since  $g_x(X^k) = x^k$  is a function belonging to  $\mathcal{C}_{\text{slow}}^\infty(\mathbb{R}^d)$ , by an induction we have  $[X^k]^g \in \mathcal{C}_{\text{slow}}^\infty(\mathbb{R}^d)$ . Now let  $\tau \in \mathcal{B}^+ \cap \mathcal{B}_X^+$ . Recall the formula obtained in [5];

$$\begin{aligned} [\tau]^g &= \text{Sg}(\tau) + P_{x,y}(g_{yx}(\tau)) \\ &+ \sum_{n=1}^\infty (-1)^{n-1} \sum_{\mathbf{1} <^+ \sigma_{n+1} <^+ \dots <^+ \sigma_1 <^+ \tau} R^\circ(g(\tau/^+ \sigma_1) \cdots g(\sigma_{n-1}/^+ \sigma_n), g(\sigma_n/^+ \sigma_{n+1}), [\sigma_{n+1}]^g). \end{aligned}$$

This is obtained from the expansion formula obtained in [5];

$$\begin{aligned} \text{(B.3)} \quad &g_y(\tau/^+ \sigma) - g_x(\tau/^+ \sigma) \\ &= \sum_{n=1}^\infty (-1)^{n-1} \sum_{\sigma <^+ \sigma_n <^+ \dots <^+ \sigma_1 <^+ \tau} g_x(\tau/^+ \sigma_1) \cdots g_x(\sigma_{n-1}/^+ \sigma_n) (g_y(\sigma_n/^+ \sigma) - g_x(\sigma_n/^+ \sigma)) \\ &\quad + g_{yx}(\tau/^+ \sigma) \end{aligned}$$

with  $\sigma = \mathbf{1}_+$  and by definition of the  $R^\circ$  operator. Since  $\tau \in \mathcal{B}^+ \cap \mathcal{B}_X^+$ , we have  $\text{Sg}(\tau) \in \mathcal{C}_{\text{rap}}^\infty(\mathbb{R}^d)$  and  $P_{x,y}(g_{yx}(\tau)) \in \mathcal{C}_{\text{rap}}^{|\tau|}(\mathbb{R}^d)$ . For the  $R^\circ$  terms, we apply Proposition 31 to (B.3). If  $\sigma \in \mathcal{B}_X^+$ , then since  $\tau/^+ \sigma \in \text{span}(\mathcal{B}^+ \cap \mathcal{B}_X^+)$ , by assumption B(2), we have  $g_x(\tau/^+ \sigma) \in L_{\text{rap}}^\infty(\mathbb{R}^d)$  and  $g_{yx}(\tau/^+ \sigma) \in \mathcal{C}_{(2),\text{rap}}^{|\tau|-|\sigma|}(\mathbb{R}^d \times \mathbb{R}^d)$ . For the sum over  $\sigma <^+ \sigma_n <^+ \dots <^+ \sigma_1 <^+ \tau$ , we can see that at least one element among

$$g(\tau/^+ \sigma_1), \quad \dots, \quad g(\sigma_{n-1}/^+ \sigma_n), \quad g(\sigma_n/^+ \sigma)$$

belongs to  $L_{\text{rap}}^\infty(\mathbb{R}^d)$ . Indeed, if  $\sigma_n \notin \mathcal{B}_X^+$  then  $g(\sigma_n/^+ \sigma) \in L_{\text{rap}}^\infty(\mathbb{R}^d)$ . Otherwise, if  $\sigma_{n-1} \notin \mathcal{B}_X^+$  then  $g(\sigma_{n-1}/^+ \sigma_n) \in L_{\text{rap}}^\infty(\mathbb{R}^d)$ . Since  $\tau \notin \mathcal{B}_X^+$ , for at least one  $i$  we have  $g(\sigma_i/^+ \sigma_{i+1}) \in L_{\text{rap}}^\infty(\mathbb{R}^d)$ . Since  $L_{\text{slow}}^\infty(\mathbb{R}^d) \cdot L_{\text{rap}}^\infty(\mathbb{R}^d) \subset L_{\text{rap}}^\infty(\mathbb{R}^d)$ , we can apply Proposition 31(a) to get

$$\sum_{n=1}^\infty (-1)^{n-1} \sum_{\sigma <^+ \sigma_n <^+ \dots <^+ \sigma_1 <^+ \tau} R^\circ(g(\tau/^+ \sigma_1) \cdots g(\sigma_{n-1}/^+ \sigma_n), g(\sigma_n/^+ \sigma), [\sigma]^g) \in \mathcal{C}_{\text{rap}}^{|\tau|}(\mathbb{R}^d).$$

If  $\sigma \notin \mathcal{B}_X^+$ , since  $g_{yx}(\tau/^+ \sigma) \in \mathcal{C}_{(2)}^{|\tau|-|\sigma|}(\mathbb{R}^d \times \mathbb{R}^d)$  and  $[\sigma]^g \in \mathcal{C}_{\text{rap}}^{|\sigma|}(\mathbb{R}^d)$  we can apply Proposition 31(b) to get the same estimate. Hence we obtain the required estimates in the formula (B.2).

To get (3.5) from (B.2), it is sufficient to show

$$\text{(B.4)} \quad \llbracket \tau \rrbracket^g - [\tau]^g \in \mathcal{C}_{\text{rap}}^\infty(\mathbb{R}^d)$$

for any  $\tau \in \mathcal{B}^+ \cap \mathcal{B}_X^+$ . Assume that all  $\nu \in \mathcal{B}^+ \cap \mathcal{B}_X^+$  with  $|\nu| < |\tau|$  satisfy (B.4). Then we have

$$\begin{aligned} \llbracket \tau \rrbracket^{\mathfrak{g}} - [\tau]^{\mathfrak{g}} &= \sum_{\mathbf{1} <^+ \nu <^+ \tau} \mathbb{P}_{\mathfrak{g}(\tau/+ \nu)}[\nu]^{\mathfrak{g}} - \sum_{\substack{\mathbf{1} <^+ \nu <^+ \tau \\ \nu \notin \mathcal{B}_X^+}} \mathbb{P}_{\mathfrak{g}(\tau/+ \nu)}\llbracket \nu \rrbracket^{\mathfrak{g}} \\ &= \sum_{\substack{\mathbf{1} <^+ \nu <^+ \tau \\ \nu \notin \mathcal{B}_X^+}} \mathbb{P}_{\mathfrak{g}(\tau/+ \nu)}([\nu]^{\mathfrak{g}} - \llbracket \nu \rrbracket^{\mathfrak{g}}) + \sum_{k \neq 0} \mathbb{P}_{\mathfrak{g}(\tau/+ X^k)}[X^k]^{\mathfrak{g}}. \end{aligned}$$

The first term belongs to  $\mathcal{C}_{\text{rap}}^{\infty}(\mathbb{R}^d)$  by assumption. For the second term, since  $[X^k]^{\mathfrak{g}} \in \mathcal{C}_{\text{slow}}^{\infty}(\mathbb{R}^d)$  and  $\mathfrak{g}(\tau/+ X^k) \in L_{\text{rap}}^{\infty}(\mathbb{R}^d)$ , we can complete the proof.

One can obtain formula (3.6) in the similar way. As above, we define the quantity  $[\tau]^M$  for each  $\tau \in \mathcal{B}$  by

$$\Pi\tau = \sum_{\nu < \tau, \nu \in \mathcal{B}} \mathbb{P}_{\mathfrak{g}(\tau/\nu)}[\nu]^M + [\tau]^M.$$

Then we can show that  $[\nu]^M \in \mathcal{C}_{\text{rap}}^{|\nu|}(\mathbb{R}^d)$ , if  $\nu \in \mathcal{B} \cap \mathcal{B}_{\underline{X}}$ , and  $[\nu]^M \in \mathcal{C}_{\text{slow}}^{\infty}(\mathbb{R}^d)$ , if  $\nu \in \mathcal{B}_{\underline{X}}$ . The only difference is that, for  $\tau \in \mathcal{B} \cap \mathcal{B}_{\underline{X}}$ , we use the formula obtained in [5]

$$\begin{aligned} [\tau]^{\mathfrak{g}} &= S(\Pi\tau) + \mathbf{P}_{x,y}((\Pi_x^{\mathfrak{g}}\tau)(y)) \\ &\quad + \sum_{n=1}^{\infty} (-1)^{n-1} \sum_{\sigma_{n+1} < \dots < \sigma_1 < \tau} \mathbb{R}^{\circ}(\mathfrak{g}(\tau/\sigma_1) \cdots \mathfrak{g}(\sigma_{n-1}/\sigma_n), \mathfrak{g}(\sigma_n/\sigma_{n+1}), [\sigma_{n+1}]^M), \end{aligned}$$

and use Proposition 32 to get  $\mathbf{P}_{x,y}((\Pi_x^{\mathfrak{g}}\tau)(y)) \in \mathcal{C}_{\text{rap}}^{|\tau|}(\mathbb{R}^d)$ . Since the property (B.4) also holds for the operator  $\llbracket \cdot \rrbracket^M - [\cdot]^M$ , we can conclude (3.6).

*Proof of Theorem 13.* — (3.7) is proved by a similar argument as Theorem 12. See [5, Th. 14] for details. More easily, it is useful to consider the extended algebra  $T_{\mathbf{F}}^+$  defined in Section 4.2. Since a modeled distribution  $\mathbf{f} \in \mathcal{D}^{\gamma}(T, \mathfrak{g})$  defines a  $\mathfrak{g}$ -part of the model on  $T_{\mathbf{F}}^+$  by Lemma 21, we have

$$f_{\sigma} = \mathfrak{g}(\mathbf{F}_{\sigma}) = \sum_{\substack{\sigma < \mu \\ \mu/\sigma \in \text{span}(\mathcal{B}^+ \cap \mathcal{B}_X^+)}} \mathbb{P}_{\mathfrak{g}(\mathbf{F}_{\mu})}\llbracket \mu/\tau \rrbracket^{\mathfrak{g}} + \llbracket \mathbf{F}_{\sigma} \rrbracket^{\mathfrak{g}}.$$

Thus  $\llbracket f_{\sigma} \rrbracket^{\mathfrak{g}} = \llbracket \mathbf{F}_{\sigma} \rrbracket^{\mathfrak{g}} \in \mathcal{C}_{\text{rap}}^{\gamma-|\sigma|}(\mathbb{R}^d)$ .

As for (3.8), a similar interpretation is useful. Consider a symbol  $\mathbf{F}$  and an extended model space  $T_{\mathbf{F}} := T \oplus \text{span}(\mathbf{F})$ . Giving the homogeneity  $|\mathbf{F}| := \gamma$  and the coproduct formula

$$\Delta\mathbf{F} = \mathbf{F} \otimes \mathbf{1}_+ + \sum_{\substack{\tau \in \mathcal{B} \\ |\tau| < \gamma}} \tau \otimes (\mathbf{F}_{\tau}),$$

the pair  $(T_{\mathbf{F}}^+, T_{\mathbf{F}})$  turns out to be a regularity structure. (It is not difficult to check that  $T_{\mathbf{F}}$  is a comodule over  $T_{\mathbf{F}}^+$  by using (4.11).) For given a reconstruction  $\mathbf{R}\mathbf{f}$ , we can

define the model on  $T_{\mathbf{F}}$  by setting  $\Pi_{\mathbf{F}} := \mathbf{R}f$ . Indeed, similarly to Lemma 21, we can show that

$$\Pi_x^{\mathbf{g}} \mathbf{F} = \mathbf{R}f - \Pi_x^{\mathbf{g}} f(x).$$

Then (3.8) follows from (3.6) in Theorem 12.

To complete the proof of Theorem 12, we define here the two-parameter extension  $\mathbf{P}^m$  of the modified paraproduct  $\mathbf{P}^m$ . Note that, there is an annulus  $\mathcal{A} \subset \mathbb{R}^d$  such that the Fourier transform of the function

$$x \mapsto P_j(x - y) Q_j(x - z)$$

is contained in  $2^j \mathcal{A}$  (independently to  $y, z$ ). Let  $\chi$  be a smooth function on  $\mathbb{R}^d$  supported in a larger annulus  $\mathcal{A}'$  and such that  $\chi \equiv 1$  on  $\mathcal{A}$ . Letting  $R_j = \mathcal{F}^{-1}(\chi(2^{-j}\cdot))$ , we have

$$(\mathbf{Q}_j \Lambda)(x) = \iiint_{\mathbb{R}^d \times \mathbb{R}^d \times \mathbb{R}^d} R_j(x - w) P_j(w - y) Q_j(w - z) \Lambda(y, z) dy dz dw.$$

For  $m \in \mathbb{Z}$ , set

$$\begin{aligned} Q_j^{-m} &:= \mathcal{F}^{-1}(|\cdot|^{-m} \rho_j), \\ R_j^m &:= \mathcal{F}^{-1}(|\cdot|^m \chi(2^{-j}\cdot)); \end{aligned}$$

then they are smooth functions such that  $Q_j^{-m} = |\nabla|^{-m} Q_j$  and  $R_j^m = |\nabla|^m R_j$ .

DEFINITION 34. — For any  $m \in \mathbb{N}$  and any two-variable distribution  $\Lambda$  on  $\mathbb{R}^d \times \mathbb{R}^d$ , define

$$\begin{aligned} (\mathbf{Q}_j^m \Lambda)(x) &:= \iiint_{\mathbb{R}^d \times \mathbb{R}^d \times \mathbb{R}^d} R_j^m(x - w) P_j(w - y) Q_j^{-m}(w - z) \Lambda(y, z) dy dz dw, \\ (\mathbf{P}^m \Lambda)(x) &:= \sum_{j > 1} (\mathbf{Q}_j^m \Lambda)(x). \end{aligned}$$

If necessary, we emphasize the integrated variables by writing

$$\mathbf{P}^m \Lambda = \mathbf{P}_{y,z}^m (\Lambda(y, z)).$$

For the special case  $\Lambda(y, z) = f(y)g(z)$ , we have the consistency relation

$$\mathbf{P}^m \Lambda = \mathbf{P}_f^m g.$$

All the above estimates in this appendix still hold for these modified operators. Indeed, because of the scaling properties

$$Q_j^{-m}(x) = 2^{j(d-m)} Q_0^{-m}(2^j x), \quad R_j^m(x) = 2^{j(d+m)} R_0^m(2^j x),$$

we can show the following analogue of Lemma 29; for any  $\alpha > 0$  and  $a \in \mathbb{Z}$ , one has

$$(B.5) \quad \begin{aligned} &\int |R_i^m(x - y)| |x - y|^\alpha |y|_*^{-a} dy \cdot 2^{-i(\alpha+m)} |x|_*^{-a}, \\ &\int |Q_i^{-m}(x - y)| |x - y|^\alpha |y|_*^{-a} dy \cdot 2^{-i(\alpha-m)} |x|_*^{-a}. \end{aligned}$$

Thus we can repeat the argument in this appendix as follows.

– Proposition 30(a) still holds, since  $\Delta_i(\mathbf{P}^m \Lambda) = \sum_{|i-j| \leq N} \Delta_i(\mathbf{Q}_j^m \Lambda)$  for some integer depending only on the support of  $\chi$ .

– Proposition 30(b) still holds, since by the scaling property,

$$\begin{aligned} |\mathbf{Q}_j^m F(x)| &\leq \int |R_j^m(x-w)P_j(w-y)Q_j^{-m}(w-z)| |F(y,z)| dydzdw \\ &\leq \int |R_j^m(x-w)P_j(w-y)Q_j^{-m}(w-z)| (|y|_*^{-a} + |z|_*^{-a}) \\ &\quad \times (|w-y|^\alpha + |w-z|^\alpha) dydzdw \\ &\leq 2^{-j(\alpha-m)} \int |R_j^m(x-w)| |w|_*^{-a} dw \\ &\leq 2^{-j\alpha} |w|_*^{-a}. \end{aligned}$$

– Proposition 31 still holds if  $\mathbf{R}^\circ$  is replaced by

$$\mathbf{R}^m(f, g, h) := \mathbf{P}_f^m \mathbf{P}_g^m h - \mathbf{P}_{fg}^m h,$$

by a parallel argument using (B.5).

Consequently, we can repeat the proof of Theorem 12 for any  $m \in \mathbb{N}$ .

### APPENDIX C. THE SLOWLY GROWING SETTING

In applications of regularity structures to the study of singular stochastic PDEs set in the entire space  $\mathbb{R}^d$  usually involve noises that do not have rapid decrease at infinity, but rather have moderate growth at infinity. Our results can be formulated as follows in this slightly modified setting.

We define the spaces  $\mathcal{M}_{\text{slow}}$  and  $\mathcal{D}_{\text{slow}}^\gamma$  of slowly growing models and modeled distributions, respectively, by replacing ‘rap’ in definitions in Section 3.2 by ‘slow’. We can repeat the same arguments to obtain the variations of Theorems 1 and 5 with the spaces  $\mathcal{M}_{\text{slow}}$  and  $\mathcal{D}_{\text{slow}}^\gamma$ , respectively. All we need is to consider the weight  $|x|_*^{-a}$  for *some* sufficiently large  $a$ , instead of *any*  $a \in \mathbb{Z}$ . Precisely, we need the following minor modifications of the arguments.

– Proposition 31 still holds under the assumption  $f^\sharp \in \mathcal{C}_{(2),\text{slow}}^\alpha(\mathbb{R}^d \times \mathbb{R}^d)$  and  $g \in \mathcal{C}_{\text{slow}}^\beta(\mathbb{R}^d)$ , instead of rapid decrease assumptions.

– Proposition 32 still holds under the assumption

$$\sup_{x \in \mathbb{R}^d} |x|_*^a \|\Lambda_x\|_{\mathcal{C}^{\beta_0}} < \infty$$

for *some*  $a \in \mathbb{Z}$ , and for any  $\ell$ ,  $\Theta^\ell \in D_{\text{slow}}^{\beta_\ell}$  and  $c^\ell \in \mathcal{C}_{(2),\text{slow}}^{\gamma-\beta_\ell}(\mathbb{R}^d \times \mathbb{R}^d)$ .

– Lemma 19 still holds for *some*  $a \in \mathbb{Z}$ , instead of any  $a$ .

Details are left to readers. We end this appendix by writing the precise statements of main theorems.

**THEOREM 35.** — *Let  $T$  be a concrete regularity structure satisfying assumptions (A–C). Then one can construct a locally Lipschitz continuous map*

$$(C.1) \quad \begin{aligned} M_{\text{slow}}(T, \mathbb{R}^d) &\longrightarrow \prod_{\sigma \in \mathcal{B}^+ \uparrow \mathcal{B}_X^+} \mathcal{C}_{\text{slow}}^{|\sigma|}(\mathbb{R}^d) \times \prod_{\tau \in \mathcal{B} \uparrow \mathcal{B}_X} \mathcal{C}_{\text{slow}}^{|\tau|}(\mathbb{R}^d) \\ (\mathbf{g}, \Pi) &\longmapsto ([\sigma]^M, [\tau]^g; \sigma \in \mathcal{B}^+ \uparrow \mathcal{B}_X^+, \tau \in \mathcal{B} \uparrow \mathcal{B}_X) \end{aligned}$$

by giving paracontrolled representations of  $\mathbf{g}$  and  $\Pi$ , for  $(\mathbf{g}, \Pi) \in M_{\text{rap}}(T, \mathbb{R}^d)$ . Furthermore,  $M_{\text{slow}}(T, \mathbb{R}^d)$  is locally bi-Lipschitz homeomorphic to the product space

$$\prod_{\sigma \in \mathcal{G}_\bullet^+} \mathcal{C}_{\text{slow}}^{|\sigma|}(\mathbb{R}^d) \times \prod_{\tau \in \mathcal{B}_\bullet, |\tau| < 0} \mathcal{C}_{\text{slow}}^{|\tau|}(\mathbb{R}^d).$$

**THEOREM 36.** — *Let a concrete regularity structure  $T$  satisfy assumptions (A–D). Pick  $\gamma \in \mathbb{R} \setminus \{0\}$  such that  $\gamma - |\tau| \notin \mathbb{N}$  for any basis element  $\tau$  of  $T$  with  $|\tau| < \gamma$ , and  $M = (\mathbf{g}, \Pi) \in M_{\text{slow}}(T, \mathbb{R}^d)$ . Then one can construct a locally Lipschitz continuous map*

$$\mathcal{D}_{\text{slow}}^\gamma(T, \mathbf{g}) \longrightarrow \prod_{\tau \in \mathcal{B}, |\tau| < \gamma} \mathcal{C}_{\text{slow}}^{\gamma - |\tau|}(\mathbb{R}^d)$$

by giving a paracontrolled representation of elements in  $\mathcal{D}_{\text{slow}}^\gamma(T, \mathbf{g})$ . Furthermore,  $\mathcal{D}_{\text{slow}}^\gamma(T, \mathbf{g})$  is locally bi-Lipschitz homeomorphic to the product space

$$\prod_{\tau \in \mathcal{B}_\bullet, |\tau| < \gamma} \mathcal{C}_{\text{slow}}^{\gamma - |\tau|}(\mathbb{R}^d).$$

### REFERENCES

- [1] H. BAHOURI, J.-Y. CHEMIN & R. DANCHIN – *Fourier analysis and nonlinear partial differential equations*, Grundlehren Math. Wiss., vol. 343, Springer, Heidelberg, 2011.
- [2] I. BAILLEUL – “Regularity of the Itô-Lyons map”, *Confluentes Math.* **7** (2015), no. 1, p. 3–11.
- [3] I. BAILLEUL & F. BERNICOT – “High order paracontrolled calculus”, *Forum Math. Sigma* **7** (2019), article no. e44 (94 pages).
- [4] I. BAILLEUL, F. BERNICOT & D. FREY – “Space-time paraproducts for paracontrolled calculus, 3D-PAM and multiplicative Burgers equations”, *Ann. Sci. École Norm. Sup. (4)* **51** (2018), no. 6, p. 1399–1456.
- [5] I. BAILLEUL & M. HOSHINO – “Paracontrolled calculus and regularity structures I”, *J. Math. Soc. Japan* **73** (2021), no. 2, p. 553–595.
- [6] Y. BRUNED, A. CHANDRA, I. CHEVYREV & M. HAIRER – “Renormalising SPDEs in regularity structures”, *J. Eur. Math. Soc. (JEMS)* **23** (2021), no. 3, p. 869–947.
- [7] Y. BRUNED, M. HAIRER & L. ZAMBOTTI – “Algebraic renormalisation of regularity structures”, *Invent. Math.* **215** (2019), no. 3, p. 1039–1156.
- [8] F. CARAVENNA & L. ZAMBOTTI – “Hairer’s reconstruction theorem without regularity structures”, *EMS Surv. Math. Sci.* **7** (2020), no. 2, p. 207–251.
- [9] A. CHANDRA & M. HAIRER – “An analytic BPHZ theorem for regularity structures”, 2018, arXiv: 1612.08138v5.
- [10] P. CHARTIER, E. HAIRER & G. VILMART – “Algebraic structures of B-series”, *Found. Comput. Math.* **10** (2010), no. 4, p. 407–427.
- [11] A. CONNES & D. KREIMER – “Hopf algebras, renormalization and noncommutative geometry”, *Comm. Math. Phys.* **199** (1998), no. 1, p. 203–242.
- [12] B. K. DRIVER – “A Cameron-Martin type quasi-invariance theorem for Brownian motion on a compact Riemannian manifold”, *J. Funct. Anal.* **110** (1992), no. 2, p. 272–376.

- [13] M. GUBINELLI, P. IMKELLER & N. PERKOWSKI – “Paracontrolled distributions and singular PDEs”, *Forum Math. Pi* **3** (2015), article no. e6 (75 pages).
- [14] M. HAIRER – “A theory of regularity structures”, *Invent. Math.* **198** (2014), no. 2, p. 269–504.
- [15] M. HAIRER & É. PARDOUX – “Fluctuations around a homogenised semilinear random PDE”, *Arch. Rational Mech. Anal.* **239** (2021), no. 1, p. 151–217.
- [16] M. HOSHINO – “Commutator estimates from a viewpoint of regularity structures”, *RIMS Kôkyûroku Bessatsu* **B79** (2020), p. 179–197.
- [17] ———, “Iterated paraproducts and iterated commutator estimates in Besov spaces”, *Adv. Stud. Pure Math.*, vol. 87, Mathematical Society of Japan, Tokyo, 2021, to appear, arXiv: 2001.07414.
- [18] C. LIU, D. J. PRÖMEL & J. TEICHMANN – “Stochastic analysis with modelled distributions”, *Stochastic Partial Differential Equations and Applications* **9** (2021), no. 2, p. 343–379.
- [19] T. LYONS & Z. QIAN – “Flow equations on spaces of rough paths”, *J. Funct. Anal.* **149** (1997), no. 1, p. 135–159.
- [20] T. LYONS & N. VICTOIR – “An extension theorem to rough paths”, *Ann. Inst. H. Poincaré Anal. Non Linéaire* **24** (2007), no. 5, p. 835–847.
- [21] J. MARTIN – “Refinements of the solution theory for singular SPDEs”, PhD Thesis, H.U. Berlin, 2018.
- [22] J. MARTIN & N. PERKOWSKI – “A Littlewood-Paley description of modelled distributions”, *J. Funct. Anal.* **279** (2020), no. 6, p. 108634, 22.
- [23] H. SINGH & J. TEICHMANN – “An elementary proof of the reconstruction theorem”, 2018, arXiv: 1812.03082.
- [24] N. TAPIA & L. ZAMBOTTI – “The geometry of the space of branched rough paths”, *Proc. London Math. Soc. (3)* **121** (2020), no. 2, p. 220–251.
- [25] J. UNTERBERGER – “Hölder-continuous rough paths by Fourier normal ordering”, *Comm. Math. Phys.* **298** (2010), no. 1, p. 1–36.

Manuscript received 24th March 2020

accepted 29th June 2021

ISMAEL BAILLEUL, Univ. Rennes, CNRS, IRMAR - UMR 6625  
 263 avenue du General Leclerc, 35042 Rennes, France  
*E-mail* : ismael.bailleul@univ-rennes1.fr  
*Url* : <https://perso.univ-rennes1.fr/ismael.bailleul>

MASATO HOSHINO, Graduate School of Engineering Science, Osaka University  
 1-3, Machikaneyama, Toyonaka, Osaka, 560-8531, Japan  
*E-mail* : hoshino@sigmath.es.osaka-u.ac.jp  
*Url* : <http://www.sigmath.es.osaka-u.ac.jp/~hoshino>