



Journal de l'École polytechnique

Mathématiques

Mitia DUERINCKX & Antoine GLORIA

Quantitative homogenization theory for random suspensions in steady Stokes flow

Tome 9 (2022), p. 1183-1244.

<https://doi.org/10.5802/jep.204>

© Les auteurs, 2022.



Cet article est mis à disposition selon les termes de la licence
LICENCE INTERNATIONALE D'ATTRIBUTION CREATIVE COMMONS BY 4.0.
<https://creativecommons.org/licenses/by/4.0/>

L'accès aux articles de la revue « Journal de l'École polytechnique — Mathématiques » (<http://jep.centre-mersenne.org/>) implique l'accord avec les conditions générales d'utilisation (<http://jep.centre-mersenne.org/legal/>).

Publié avec le soutien
du Centre National de la Recherche Scientifique



Publication membre du
Centre Mersenne pour l'édition scientifique ouverte
www.centre-mersenne.org

QUANTITATIVE HOMOGENIZATION THEORY FOR RANDOM SUSPENSIONS IN STEADY STOKES FLOW

BY MITIA DUERINCKX & ANTOINE GLORIA

ABSTRACT. — This work develops a quantitative homogenization theory for random suspensions of rigid particles in a steady Stokes flow, and completes recent qualitative results. More precisely, we establish a large-scale regularity theory for this Stokes problem, and we prove moment bounds for the associated correctors and optimal estimates on the homogenization error; the latter further requires a quantitative ergodicity assumption on the random suspension. Compared to the corresponding quantitative homogenization theory for divergence-form linear elliptic equations, substantial difficulties arise from the analysis of the fluid incompressibility and the particle rigidity constraints. Our analysis further applies to the problem of stiff inclusions in (compressible or incompressible) linear elasticity and in electrostatics; it is also new in those cases, even in the periodic setting.

RÉSUMÉ (Homogénéisation quantitative de suspensions aléatoires de particules dans un fluide de Stokes stationnaire)

Ce travail développe une théorie quantitative de l'homogénéisation de suspensions aléatoires de particules rigides dans un fluide de Stokes stationnaire, complétant les résultats qualitatifs récents. Plus précisément, nous établissons une théorie de régularité aux grandes échelles pour ce problème de Stokes et nous montrons des estimations de moments pour les correcteurs associés, ainsi que des estimations optimales de convergence de l'erreur d'homogénéisation (sous des hypothèses quantitatives d'ergodicité de la suspension aléatoire). En comparaison à la théorie pour les équations elliptiques linéaires sous forme divergence, l'incompressibilité du fluide et la rigidité des particules soulèvent des difficultés analytiques additionnelles. Notre analyse couvre également le problème des inclusions rigides en élasticité linéaire (compressible ou incompressible) et en électrostatique ; les résultats sont nouveaux pour ces modèles également, même dans le cas périodique.

CONTENTS

1. Introduction.....	1184
2. Main results.....	1188
3. Perturbative annealed regularity.....	1193
4. Corrector estimates.....	1201

MATHEMATICAL SUBJECT CLASSIFICATION (2020). — 35R60, 76M50, 35Q35, 76D03, 76D07.

KEYWORDS. — Steady Stokes fluid, rigid particles, quantitative stochastic homogenization, large-scale regularity.

M.D. acknowledges financial support from the CNRS-Momentum program, and A.G. from the European Research Council (ERC) under the European Union's Horizon 2020 research and innovation programme (Grant Agreement n° 864066).

5. Large-scale regularity.....	1219
6. Quantitative homogenization.....	1239
References.....	1242

1. INTRODUCTION

We start with the formulation of the steady Stokes model describing a viscous fluid in presence of a random suspension of small rigid particles, see e.g. [18]. Throughout, we denote by $d \geq 2$ the space dimension, we consider a given random set $\mathcal{I} = \bigcup_n I_n \subset \mathbb{R}^d$, where $\{I_n\}_n$ stands for the different particles, and we denote by x_n the barycenter of I_n . Ergodicity, hardcore, and regularity assumptions are listed in Section 2. To model a suspension of small particles, we rescale the random set \mathcal{I} by a small parameter $\varepsilon > 0$ and consider $\varepsilon\mathcal{I} = \bigcup_n \varepsilon I_n$. Next, we view these small particles $\{\varepsilon I_n\}_n$ as suspended in a solvent described by the steady Stokes equation: in a reference domain $U \subset \mathbb{R}^d$, given an internal forcing $f \in L^2(U)^d$, the fluid velocity u_ε satisfies

$$(1.1) \quad -\Delta u_\varepsilon + \nabla P_\varepsilon = f, \quad \operatorname{div}(u_\varepsilon) = 0, \quad \text{in } U \setminus \varepsilon\mathcal{I},$$

with $u_\varepsilon = 0$ on ∂U . (Assume for the moment that no particle intersects the boundary ∂U .) No-slip conditions are imposed at particle boundaries: particles are constrained to have rigid motions, which amounts to extending the velocity field u_ε inside particles in such a way that

$$D(u_\varepsilon) = 0, \quad \text{in } \varepsilon\mathcal{I},$$

where $D(u_\varepsilon)$ denotes the symmetrized gradient of u_ε ; in other words, this condition entails that the velocity field u_ε is a (linearized) rigid motion $V_{\varepsilon,n} + \Theta_{\varepsilon,n}(x - \varepsilon x_n)$ inside each particle εI_n (centered at εx_n), for some $V_{\varepsilon,n} \in \mathbb{R}^d$ and some skew-symmetric matrix $\Theta_{\varepsilon,n} \in \mathbb{R}^{d \times d}$. Finally, assuming that the particles have the same mass density as the fluid, buoyancy forces vanish, and the force and torque balances on each particle take the form

$$(1.2) \quad \int_{\varepsilon\partial I_n} \sigma(u_\varepsilon, P_\varepsilon) \nu = 0,$$

$$(1.3) \quad \int_{\varepsilon\partial I_n} \Theta(x - \varepsilon x_n) \cdot \sigma(u_\varepsilon, P_\varepsilon) \nu = 0, \quad \text{for all skew-symmetric } \Theta \in \mathbb{R}^{d \times d},$$

in terms of the Cauchy stress tensor

$$(1.4) \quad \sigma(u_\varepsilon, P_\varepsilon) = 2D(u_\varepsilon) - P_\varepsilon \operatorname{Id},$$

where ν stands for the outward unit normal vector at the particle boundaries. In the physically relevant 3D case, skew-symmetric matrices $\Theta \in \mathbb{R}^{3 \times 3}$ are equivalent to cross products $\theta \times$ with $\theta \in \mathbb{R}^3$, and equations recover their standard form.

In the companion article [18], we proved that in the macroscopic limit $\varepsilon \downarrow 0$ the velocity and pressure fields $(u_\varepsilon, P_\varepsilon)$ converge weakly to $(\bar{u}, \bar{P} + \bar{\mathbf{b}} : \mathbf{D}(\bar{u}))$, where (\bar{u}, \bar{P}) solves the homogenized equation

$$(1.5) \quad \begin{cases} -\operatorname{div}(2\bar{\mathbf{B}}\mathbf{D}(\bar{u})) + \nabla\bar{P} = (1 - \lambda)f, & \text{in } U, \\ \operatorname{div}(\bar{u}) = 0, & \text{in } U, \\ \bar{u} = 0, & \text{on } \partial U, \end{cases}$$

for some effective viscosity tensor $\bar{\mathbf{B}}$ and some effective matrix $\bar{\mathbf{b}}$, where $\lambda = \mathbb{E}[\mathbb{1}_{\mathcal{S}}]$ denotes the volume fraction of the suspension. The aim of the present contribution is twofold:

(I) Make this qualitative convergence result quantitative by optimally estimating the error between $(u_\varepsilon, P_\varepsilon)$ and a two-scale expansion based on $(\bar{u}, \bar{P} + \bar{\mathbf{b}} : \mathbf{D}(\bar{u}))$ in terms of suitable correctors, cf. Theorem 6 below.

(II) Develop a large-scale regularity theory for the Stokes problem (1.1)–(1.3), which ensures that on large scales the solution u_ε has the same regularity properties as the solution \bar{u} of the limiting equation (1.5) (both in terms of $C^{1,1-}$ Schauder theory and in terms of L^p regularity), cf. Theorems 3, 4, and 5 below.

On the one hand, part (I) provides the optimal quantitative version of [18] by estimating the error in the homogenization process. This is proved under a strong mixing assumption on the random suspension \mathcal{S} , which is conveniently formulated in form of a multiscale variance inequality in the spirit of [15, 16]. On the other hand, part (II) makes precise the intuitive idea that the Stokes problem (1.1)–(1.3) should inherit the regularity properties of the limiting equation (1.5) on sufficiently large scales, which is expressed intrinsically in terms of the growth of correctors. This is qualitatively established under a mere ergodicity assumption, and further quantified assuming the same multiscale variance inequality as above.

Our main motivation to develop a large-scale regularity theory for (1.1)–(1.3) stems from the sedimentation problem for a random suspension in a Stokes flow under a constant gravity field $e \in \mathbb{R}^d$, in which case the force balance (1.2) is replaced by

$$\int_{\varepsilon\partial I_n} \sigma(u_\varepsilon, P_\varepsilon)\nu + e = 0.$$

Since energy is then pumped into the system, naive energy estimates blow up, and the analysis crucially relies on stochastic cancellations. Annealed L^p regularity in form of Theorem 5 below constitutes the main technical ingredient of [19] for our analysis of the sedimentation problem. More precisely, this allows us to prove the celebrated predictions by Batchelor [9], Caffisch and Luke [11], and Koch and Shaqfeh [38] on the effective sedimentation speed and on individual velocity fluctuations, significantly improving on [25].

Although the present contribution primarily focuses on random suspensions of rigid particles in a steady Stokes flow, we point out that our arguments apply more generally to homogenization problems with stiff inclusions. First note that equation (1.1) can

be written in the equivalent form

$$(1.6) \quad -\operatorname{div}(\sigma(u_\varepsilon, P_\varepsilon)) = f, \quad \operatorname{div}(u_\varepsilon) = 0, \quad \text{in } U \setminus \varepsilon\mathcal{I},$$

with $u_\varepsilon = 0$ on ∂U , where we recall that $\sigma(u_\varepsilon, P_\varepsilon)$ denotes the Cauchy stress tensor (1.4), and the equation is completed by the rigidity constraint $\mathbf{D}(u_\varepsilon) = 0$ inside the inclusions $\varepsilon\mathcal{I}$ and by the boundary conditions (1.2)–(1.3). Let us mention a few physical models that can be obtained as a slight modification of the above:

– *Incompressible linear elasticity with stiff inclusions* takes the same form, with the Cauchy stress tensor replaced by $\sigma(u_\varepsilon, P_\varepsilon) = K \mathbf{D}(u_\varepsilon) - P_\varepsilon \operatorname{Id}$, in terms of the constant stiffness tensor K of the background material (satisfying the Legendre–Hadamard condition). Surprisingly, the qualitative homogenization of this problem is quite recent and follows from [18].⁽¹⁾

– *Compressible linear elasticity with stiff inclusions* is obtained by dropping the incompressibility constraint $\operatorname{div}(u_\varepsilon) = 0$ in (1.6), and replacing the Cauchy stress tensor by $\sigma(u_\varepsilon) = K \mathbf{D}(u_\varepsilon)$, in terms of the constant stiffness tensor K of the background material. In this case, qualitative homogenization follows from [36, Chap. 3]; see also [10] for a compactness result in a corresponding nonlinear setting.

– *Linear electrostatics with stiff inclusions* amounts to taking u_ε scalar-valued, dropping the incompressibility constraint in (1.6), and replacing the Cauchy stress tensor by $\sigma(u_\varepsilon) = K \nabla u_\varepsilon$, in terms of the constant conductivity matrix K of the background material. We refer to [36, Chap. 3] for the qualitative homogenization of this problem (under weaker hardcore conditions).

Our present *quantitative* analysis also applies to these models. Our results are all new, even in the periodic setting (that is, when \mathcal{I} is a periodic set), in which case Theorems 2 and 6 below hold with $\mu_d \equiv 1$.

Before turning to precise statements of our results, we discuss the context. The present contribution constitutes a natural extension to the steady Stokes problem (1.1)–(1.3) of the by-now well-developed quantitative homogenization theory for the model case of divergence-form linear elliptic equations with random coefficients. This theory was started in [30, 31, 27, 26, 33, 40], with quantitative statements close to Theorem 6 below under similar mixing conditions. Large-scale regularity was initiated in [7, 8] in the periodic setting, and in [6] in the random setting, which led to a more mature theory of the field. For recent developments, we refer the reader to the recent monograph [3], based on [5, 1, 2], and to the series of works [28, 29, 32, 20, 37]. In the present contribution, we consider for convenience a strong mixing assumption in form of a multiscale variance inequality [15, 16], and we establish large-scale regularity by following the approach of [28, 29, 20] – we believe the approach of [3] could be used as well (see [17, App. B] for some result in this direction). Since we focus on the weakly correlated setting, we may, as in [30] in its efficient reformulation

⁽¹⁾In this problem it might make more sense to include the internal forcing f in the boundary conditions, replacing (1.2) by $\int_{\varepsilon\partial I_n} \sigma(u_\varepsilon, P_\varepsilon)\nu + \int_{\varepsilon I_n} f = 0$. In that case, the forcing term in the homogenized problem (1.5) is f rather than $(1 - \lambda)f$; this is only a minor change in the analysis.

of [37], bypass part of the argument in [28, 29] by appealing to deterministic regularity (in form of Meyers’ perturbative estimates) rather than large-scale regularity, which makes the proof particularly short and elegant. The strongly correlated setting could be treated by following [28], but it would substantially increase both the technicality and the length of the argument.

Compared to the model case of divergence-form linear elliptic equations with random coefficients, we face three additional difficulties in this work:

- the rigidity constraint on the particles makes the canonical structure of fluxes and flux correctors less obvious: as in [14], fluxes are constructed via a nontrivial extension procedure, which is crucial to obtain optimal convergence rates;
- naive two-scale expansions are incompatible with the rigidity constraint on the particles, thus requiring some local surgery;
- the incompressibility of the fluid gives rise to the pressure in the equation and makes many estimates more involved.

NOTATION

– For vector fields u, u' and matrix fields T, T' , we set $(\nabla u)_{ij} = \partial_j u_i$, $\operatorname{div}(T)_i = \partial_j T_{ij}$, $T : T' = T_{ij} T'_{ij}$, $(u \otimes u')_{ij} = u_i u'_j$, where we systematically use Einstein’s summation convention on repeated indices. We also write $\partial_E u = E : \nabla u$ for any matrix E .

– For a vector field u and scalar field P , we denote by $(\mathbf{D}(u))_{ij} = \frac{1}{2}(\partial_j u_i + \partial_i u_j)$ the symmetrized gradient and we recall the notation $\sigma(u, P) = 2\mathbf{D}(u) - P \operatorname{Id}$ for the Cauchy stress tensor. We also recall that ν stands for the outward unit normal vector at particle boundaries.

– We denote by $\mathbb{M}_0 \subset \mathbb{R}^{d \times d}$ the subset of trace-free matrices, by $\mathbb{M}_0^{\operatorname{sym}}$ the subset of symmetric trace-free matrices, and by $\mathbb{M}^{\operatorname{skew}}$ the subset of skew-symmetric matrices.

– We denote by $C \geq 1$ any constant that only depends on dimension d , on the constant $\delta > 0$ in Assumption (\mathbf{H}_δ) below, on the weight π in Assumption (Mix^+) if applicable, and on the reference domain U . We use the notation \lesssim (resp. \gtrsim) for $\leq C \times$ (resp. $\geq (1/C) \times$) up to such a multiplicative constant C . We write \ll (resp. \gg) for $\leq (1/C) \times$ (resp. $\geq C \times$) up to a sufficiently large multiplicative constant C . We add subscripts to $C, \lesssim, \gtrsim, \ll, \gg$ in order to indicate dependence on other parameters.

– The ball centered at x of radius r in \mathbb{R}^d is denoted by $B_r(x)$, and we simply write $B(x) = B_1(x)$, $B_r = B_r(0)$, and $B = B_1(0)$.

– For a function f , we write $[f]_2(x) := (\int_{B(x)} |f|^2)^{1/2}$ for local moving quadratic averages at unit scale.

– We set $\langle r \rangle = (1 + r^2)^{1/2}$ for $r \geq 0$, we set $\langle x \rangle = (1 + |x|^2)^{1/2}$ for $x \in \mathbb{R}^d$, and we similarly write $\langle \nabla \rangle = (1 - \Delta)^{1/2}$.

Acknowledgements. – The authors warmly thank Felix Otto for some enlightening comments on the structure of correctors in connection with Lemma 4.1.

2. MAIN RESULTS

2.1. ASSUMPTIONS. — Given an underlying probability space (Ω, \mathbb{P}) , let $\mathcal{P} = \{x_n\}_n$ be a random point process on \mathbb{R}^d , and consider a collection of random shapes $\{I_n^\circ\}_n$, where each I_n° is a connected random Borel subset of the unit ball B and is centered at 0 in the sense of $\int_{I_n^\circ} x \, dx = 0$. We then define the corresponding inclusions $I_n := x_n + I_n^\circ$ centered at the points of \mathcal{P} , and we consider the random set $\mathcal{I} := \bigcup_n I_n$. We also denote by I_n^+ the convex hull of I_n , hence $I_n \subset I_n^+ \subset B(x_n)$. Throughout, we make the following general assumptions, for some fixed deterministic constant $\delta > 0$.

ASSUMPTION (H_δ) – GENERAL CONDITIONS

- *Stationarity and ergodicity:* The random set \mathcal{I} is stationary and ergodic.
- *Uniform C^2 regularity:* The random shapes $\{I_n^\circ\}_n$ satisfy interior and exterior ball conditions with radius δ almost surely.
- *Uniform hardcore condition:* There holds $(I_n^+ + \delta B) \cap (I_m^+ + \delta B) = \emptyset$ almost surely for all $n \neq m$.

In view of quantitative homogenization results, we need to further consider quantitative ergodicity assumptions, which we make here for convenience in form of the multiscale variance inequality we introduced in [15, 16].

ASSUMPTION (Mix^+) – QUANTITATIVE MIXING CONDITION

There exists a non-increasing weight function $\pi : \mathbb{R}^+ \rightarrow \mathbb{R}^+$ with superalgebraic decay (that is, $\pi(\ell) \leq C_p \ell^{-p}$ for all $p < \infty$) such that the random set \mathcal{I} satisfies, for all $\sigma(\mathcal{I})$ -measurable random variables $Y(\mathcal{I})$,

$$(2.1) \quad \text{Var} [Y(\mathcal{I})] \leq \mathbb{E} \left[\int_0^\infty \int_{\mathbb{R}^d} \left(\partial_{\mathcal{I}, B_\ell(x)}^{\text{osc}} Y(\mathcal{I}) \right)^2 dx \ell^{-d} \pi(\ell) d\ell \right],$$

where the “oscillation” ∂^{osc} of the random variable $Y(\mathcal{I})$ is defined by

$$\begin{aligned} \partial_{\mathcal{I}, B_\ell(x)}^{\text{osc}} Y(\mathcal{I}) := & \sup \text{ess} \left\{ Y(\mathcal{I}') : \mathcal{I}' \cap (\mathbb{R}^d \setminus B_\ell(x)) = \mathcal{I} \cap (\mathbb{R}^d \setminus B_\ell(x)) \right\} \\ & - \inf \text{ess} \left\{ Y(\mathcal{I}') : \mathcal{I}' \cap (\mathbb{R}^d \setminus B_\ell(x)) = \mathcal{I} \cap (\mathbb{R}^d \setminus B_\ell(x)) \right\}. \end{aligned}$$

2.2. CORRECTOR ESTIMATES. — We first recall the suitable definitions of correctors associated with the steady Stokes problem (1.1)–(1.3), as introduced in the companion work [18, Prop. 2.1].

LEMMA 1 (Correctors; [18]). — *Under Assumption (H_δ) , for all $E \in \mathbb{M}_0$, there exists a unique solution (ψ_E, Σ_E) to the following infinite-volume corrector problem:*

– Almost surely, (ψ_E, Σ_E) belongs to $H^1_{\text{loc}}(\mathbb{R}^d)^d \times L^2_{\text{loc}}(\mathbb{R}^d \setminus \mathcal{I})$ and satisfies in the strong sense,

$$(2.2) \quad \begin{cases} -\Delta \psi_E + \nabla \Sigma_E = 0, & \text{in } \mathbb{R}^d \setminus \mathcal{I}, \\ \operatorname{div}(\psi_E) = 0, & \text{in } \mathbb{R}^d \setminus \mathcal{I}, \\ \operatorname{D}(\psi_E + Ex) = 0, & \text{in } \mathcal{I}, \\ \int_{\partial I_n} \sigma(\psi_E + E(x - x_n), \Sigma_E) \nu = 0, & \forall n, \\ \int_{\partial I_n} \Theta(x - x_n) \cdot \sigma(\psi_E + E(x - x_n), \Sigma_E) \nu = 0, & \forall n, \forall \Theta \in \mathbb{M}^{\text{skew}}. \end{cases}$$

– The gradient field $\nabla \psi_E$ and the pressure field $\Sigma_E \mathbb{1}_{\mathbb{R}^d \setminus \mathcal{I}}$ are stationary, they have vanishing expectation, they have finite second moments, and ψ_E satisfies the anchoring condition $\int_B \psi_E = 0$ almost surely.

In addition, the corrector ψ_E is sublinear at infinity, that is, $\varepsilon \psi_E(\frac{\cdot}{\varepsilon}) \rightharpoonup 0$ in $H^1_{\text{loc}}(\mathbb{R}^d)^d$ almost surely as $\varepsilon \downarrow 0$. Note that $(\psi_E, \Sigma_E) = (\psi_{E^{\text{sym}}}, \Sigma_{E^{\text{sym}}})$ where E^{sym} denotes the symmetric part of E .

As a key tool for quantitative homogenization, we establish the following moment bounds on correctors. Inspired by the corresponding strategy for divergence-form linear elliptic equations in [37], the proof is based on the analysis of stochastic cancellations for large-scale averages of the corrector gradient, together with perturbative annealed L^p regularity and a buckling argument. If the weight π in Assumption (Mix^+) has some stretched exponential decay, then the moment bounds below can be upgraded to corresponding stretched exponential moments.

THEOREM 2 (Corrector estimates). — Under Assumptions (H_δ) and (Mix^+) , for all $E \in \mathbb{M}_0$ and $q < \infty$, we have

$$\|[(\nabla \psi_E, \Sigma_E \mathbb{1}_{\mathbb{R}^d \setminus \mathcal{I}})]_2\|_{L^q(\Omega)} \lesssim_q |E|,$$

and

$$(2.3) \quad \|\psi_E\|_{L^q(\Omega)} \lesssim_q |E| \mu_d(|x|), \quad \mu_d(r) := \begin{cases} 1 & : d > 2, \\ \log(2+r)^{1/2} & : d = 2, \\ \langle r \rangle^{1/2} & : d = 1. \end{cases}$$

In particular, in dimension $d > 2$, up to relaxing the anchoring condition, the solution ψ_E of the infinite-volume problem (2.2) can be uniquely constructed itself as a stationary field with vanishing expectation.

REMARK 2.1. — We include the case $d = 1$ in the statements for completeness, in which case the problem is scalar without incompressibility constraint.

2.3. LARGE-SCALE REGULARITY. — Given a random forcing $g \in C^\infty_c(\mathbb{R}^d; L^\infty(\Omega)^{d \times d})$, we consider the unique solution $(\nabla u_g, P_g) \in L^\infty(\Omega; L^2(\mathbb{R}^d)^{d \times d} \times L^2(\mathbb{R}^d \setminus \mathcal{I}))$ of the

following steady Stokes problem,

$$(2.4) \quad \begin{cases} -\Delta u_g + \nabla P_g = \operatorname{div}(g), & \text{in } \mathbb{R}^d \setminus \mathcal{I}, \\ \operatorname{div}(u_g) = 0, & \text{in } \mathbb{R}^d \setminus \mathcal{I}, \\ \mathbf{D}(u_g) = 0, & \text{in } \mathcal{I}, \\ \int_{\partial I_n} (g + \sigma(u_g, P_g)) \nu = 0, & \forall n, \\ \int_{\partial I_n} \Theta(x - x_n) \cdot (g + \sigma(u_g, P_g)) \nu = 0, & \forall n, \forall \Theta \in \mathbb{M}^{\text{skew}}. \end{cases}$$

The energy inequality yields, almost surely,

$$(2.5) \quad \|\nabla u_g\|_{L^2(\mathbb{R}^d)} \leq \|g\|_{L^2(\mathbb{R}^d \setminus \mathcal{I})}.$$

Aside from Meyers' perturbative improvements of this energy inequality, cf. Section 3, and aside from local regularity theory, no other regularity estimates are expected to hold in general in a deterministic form due to the presence of the rigidity constraints on the random set of particles — except in a dilute regime when particles are sufficiently far apart, cf. Remark 2.2. However, in view of homogenization, the heterogeneous Stokes problem (2.4) can be replaced on large scales by a homogenized system as in (1.5). Since standard elliptic regularity theory is available for this large-scale approximation, the solution to (2.4) should enjoy improved regularity properties on large scales. This type of result was pioneered by Avellaneda and Lin [7, 8] in the context of periodic homogenization in the model setting of divergence-form linear elliptic equations. In the stochastic case, while early contributions in form of annealed Green's function estimates appeared in [12, 40], a quenched large-scale regularity theory was first outlined by Armstrong and Smart [6], and later fully developed in [5, 1, 2, 3] and in [28, 29, 32]. We also mention the useful reformulation in form of annealed regularity in [20]. Based on these ideas, we develop corresponding quenched large-scale and annealed regularity theories for the steady Stokes problem (2.4), which constitute the key technical ingredient in our work [19] on sedimentation.

We start with a quenched large-scale Schauder theory. Hölder norms are reformulated à la Campanato in terms of the growth of local integrals, and the latter are restricted to scales $\geq r_*$ for some (well-controlled) random minimal radius r_* . Note that Hölder regularity is naturally measured by replacing Euclidean coordinates $x \mapsto Ex$ by their heterogeneous versions $x \mapsto \psi_E(x) + Ex$ in terms of the corrector ψ_E .

THEOREM 3 (Quenched large-scale Schauder theory). — *Under Assumption (H_δ) , given $\alpha \in (0, 1)$, there exists an almost surely finite stationary random field $r_* \geq 1$ on \mathbb{R}^d , see (5.5), such that the following holds: For all $g \in C_c^\infty(\mathbb{R}^d; L^\infty(\Omega)^{d \times d})$ and $R \geq r_*(0)$, if ∇u_g is a solution of the steady Stokes problem (2.4) in B_R , then the following large-scale Lipschitz estimate holds on scales $\geq r_*(0)$,*

$$(2.6) \quad \sup_{r_*(0) \leq r \leq R} \int_{B_r} |\nabla u_g|^2 \lesssim \int_{B_R} |\nabla u_g|^2 + \sup_{r_*(0) \leq r \leq R} \left(\frac{R}{r}\right)^{2\alpha} \int_{B_r} \left|g - \int_{B_r} g\right|^2,$$

as well as the following large-scale $C^{1,\alpha}$ estimate,

$$\sup_{r_*(0) \leq r \leq R} \frac{1}{r^{2\alpha}} \text{Exc}(\nabla u_g; B_r) \lesssim \frac{1}{R^{2\alpha}} \text{Exc}(\nabla u_g; B_R) + \sup_{r_*(0) \leq r \leq R} \frac{1}{r^{2\alpha}} \int_{B_r} \left| g - \int_{B_r} g \right|^2,$$

where the excess is defined by

$$(2.7) \quad \text{Exc}(h; D) := \inf_{E \in \mathbb{M}_0} \int_D |h - (\nabla \psi_E + E)|^2.$$

Under Assumption (Mix^+) , the so-called minimal radius $r_*(0)$ satisfies $\mathbb{E}[r_*(0)^q] < \infty$ for all $q < \infty$.

As in [4], [3, §7], [28, Cor. 4], or [20, Prop. 6.4], the above large-scale Lipschitz regularity (2.6) can be exploited together with a Calderón–Zygmund argument to deduce the following weighted L^p regularity estimate on scales $\geq r_*$.

THEOREM 4 (Quenched large-scale L^p regularity). — Under Assumption (H_δ) , there exists an almost surely finite stationary random field $r_* \geq 1$ on \mathbb{R}^d as in Theorem 3 such that the following holds: For all $g \in C_c^\infty(\mathbb{R}^d; \mathbb{L}^\infty(\Omega)^{d \times d})$, $1 < p < \infty$, and weight μ in the Muckenhoupt class A_p , the solution $(\nabla u_g, P_g)$ of the steady Stokes problem (2.4) satisfies

$$\left(\int_{\mathbb{R}^d} \left(\int_{B_*(x)} |\nabla u_g|^2 \right)^{p/2} \mu(x) dx \right)^{1/p} \lesssim_p \left(\int_{\mathbb{R}^d} \left(\int_{B_*(x)} |g|^2 \right)^{p/2} \mu(x) dx \right)^{1/p},$$

where we use the short-hand notation $B_*(x) := B_{r_*(x)}(x)$.

As in [19], we establish the following annealed version of the above quenched large-scale L^p regularity statement. The main merit of this estimate is that a stochastic $L^q(\Omega)$ norm appears inside the spatial $L^p(\mathbb{R}^d)$ norm and allows to remove local quadratic averages on the random minimal scale r_* (up to a tiny loss of stochastic integrability), which is particularly convenient for applications.

THEOREM 5 (Annealed L^p regularity). — Under Assumptions (H_δ) and (Mix^+) , for all $g \in C_c^\infty(\mathbb{R}^d; \mathbb{L}^\infty(\Omega)^{d \times d})$, $1 < p, q < \infty$, weight μ in the Muckenhoupt class A_p , and $\eta > 0$, the solution $(\nabla u_g, P_g)$ of the steady Stokes problem (2.4) satisfies

$$(2.8) \quad \|\mu^{1/p}[\nabla u_g]_2\|_{L^p(\mathbb{R}^d; L^q(\Omega))} \lesssim_{p,q,\eta} \|\mu^{1/p}[g]_2\|_{L^p(\mathbb{R}^d; L^{q+\eta}(\Omega))}.$$

In addition, under Assumption (H_δ) (and in particular without Assumption (Mix^+)), a Meyers’ perturbative result holds without loss of stochastic integrability: there exists $C_0 \simeq 1$ such that (2.8) holds with $\eta = 0$ provided $|p - 2|, |q - 2| < 1/C_0$ and $\mu \equiv 1$.

REMARK 2.2 (Deterministic L^p regularity in dilute regime). — In the dilute regime, the recent work of Höfer [34] on the reflection method easily yields the following version of the above; the proof is a direct adaptation of [34] and is omitted. This also constitutes a variant of the dilute Green’s function estimates in [25, Lem. 2.7].

Under assumption (H_δ) , we denote by $\delta(\mathcal{I}) \geq 2\delta$ the minimal interparticle distance in \mathcal{I} . For all p, q with $1 < p, q < \infty$, there exists a constant $\delta_p > 0$ (only depending on d, p) such that, provided \mathcal{I} is dilute enough in the sense of $\delta(\mathcal{I}) \geq \delta_p$, the following

holds: Given a random forcing $g \in L^\infty(\Omega; C_c^\infty(\mathbb{R}^d)^{d \times d})$, the solution $(\nabla u_g, P_g)$ of the steady Stokes problem (2.4) satisfies

$$\|\nabla u_g\|_{L^p(\mathbb{R}^d; L^q(\Omega))} \lesssim_{p,q} \|g\|_{L^p(\mathbb{R}^d; L^q(\Omega))},$$

as well as the following deterministic estimate, almost surely,

$$\|\nabla u_g\|_{L^p(\mathbb{R}^d)} \lesssim_p \|g\|_{L^p(\mathbb{R}^d)}.$$

2.4. QUANTITATIVE HOMOGENIZATION RESULT. — We consider a steady Stokes fluid in a domain $U \subset \mathbb{R}^d$ with some internal forcing and with a dense suspension of small particles, cf. (1.1)–(1.3), and we analyze the fluid velocity in the non-dilute homogenization regime with vanishing particle size but fixed volume fraction. Suspended particles in the fluid act as obstacles and hinder the fluid flow, thus increasing the flow resistance, that is, the viscosity. The system is then expected to behave approximately like an homogeneous Stokes fluid with some effective viscosity, cf. (1.5). This was the basis of Perrin’s celebrated experiment to estimate the Avogadro number as inspired by Einstein’s PhD thesis [21].

Before stating the homogenization result, given a reference domain U , the set of particles must be modified to avoid particles intersecting the boundary: we consider the random set $\mathcal{N}_\varepsilon(U)$ of all indices n such that $\varepsilon(I_n^+ + \delta B) \subset U$, and we define

$$\mathcal{I}_\varepsilon(U) := \bigcup_{n \in \mathcal{N}_\varepsilon(U)} \varepsilon I_n.$$

Particles in this collection are of size $O(\varepsilon)$ and are at distance at least $\varepsilon\delta$ from the boundary ∂U and from one another, cf. (H_δ) . We may now turn to the statement of the optimal quantification of our qualitative homogenization result of [18].

THEOREM 6 (Quantitative homogenization result). — *Under Assumptions (H_δ) and (Mix^+) , given a smooth bounded domain $U \subset \mathbb{R}^d$ and a forcing $f \in W^{1+\alpha, \infty}(U)^d$ for some $\alpha > 0$, consider for all $\varepsilon > 0$ the unique solution*

$$(u_\varepsilon, P_\varepsilon) \in L^\infty(\Omega; H_0^1(U)^d \times L^2(U \setminus \mathcal{I}_\varepsilon(U)))$$

of the steady Stokes problem

$$(2.9) \quad \begin{cases} -\Delta u_\varepsilon + \nabla P_\varepsilon = f, & \text{in } U \setminus \mathcal{I}_\varepsilon(U), \\ \operatorname{div}(u_\varepsilon) = 0, & \text{in } U \setminus \mathcal{I}_\varepsilon(U), \\ u_\varepsilon = 0, & \text{on } \partial U, \\ \mathbf{D}(u_\varepsilon) = 0, & \text{in } \mathcal{I}_\varepsilon(U), \\ \int_{\varepsilon \partial I_n} \sigma(u_\varepsilon, P_\varepsilon) \nu = 0, & \forall n \in \mathcal{N}_\varepsilon(U), \\ \int_{\varepsilon \partial I_n} \Theta(x - x_n) \cdot \sigma(u_\varepsilon, P_\varepsilon) \nu = 0, & \forall n \in \mathcal{N}_\varepsilon(U), \forall \Theta \in \mathbb{M}^{\text{skew}}, \end{cases}$$

with $\int_{U \setminus \mathcal{J}_\varepsilon(U)} P_\varepsilon = 0$. Also consider the unique solution $(\bar{u}, \bar{P}) \in H_0^1(U)^d \times L^2(U)$ of the corresponding homogenized Stokes problem

$$(2.10) \quad \begin{cases} -\operatorname{div}(2\bar{\mathbf{B}}\mathbf{D}(\bar{u})) + \nabla\bar{P} = (1 - \lambda)f, & \text{in } U, \\ \operatorname{div}(\bar{u}) = 0, & \text{in } U, \\ \bar{u} = 0, & \text{on } \partial U, \end{cases}$$

with $\int_U \bar{P} = 0$, where $\lambda := \mathbb{E}[\mathbb{1}_{\mathcal{J}}]$ denotes the volume fraction of the suspension, the effective viscosity tensor $\bar{\mathbf{B}}$ is positive definite on $\mathbb{M}_0^{\operatorname{sym}}$ and is given by

$$(2.11) \quad \bar{\mathbf{B}} := \sum_{E, E' \in \mathcal{E}} (E' \otimes E) \mathbb{E}[(\mathbf{D}(\psi_{E'}) + E') : (\mathbf{D}(\psi_E) + E)],$$

where the sum runs over an orthonormal basis \mathcal{E} of $\mathbb{M}_0^{\operatorname{sym}}$ and the corrector (ψ_E, Σ_E) is defined in Lemma 1. Then, the following quantitative corrector result holds for all $q < \infty$,

$$(2.12) \quad \begin{aligned} & \left\| u_\varepsilon - \bar{u} - \varepsilon \sum_{E \in \mathcal{E}} \psi_E(\cdot/\varepsilon) \partial_E \bar{u} \right\|_{L^q(\Omega; H^1(U))} \\ & + \inf_{\kappa \in \mathbb{R}} \left\| P_\varepsilon - \bar{P} - \bar{\mathbf{b}} : \mathbf{D}(\bar{u}) - \sum_{E \in \mathcal{E}} (\Sigma_E \mathbb{1}_{\mathbb{R}^d \setminus \mathcal{J}})(\cdot/\varepsilon) \partial_E \bar{u} - \kappa \right\|_{L^q(\Omega; L^2(U \setminus \mathcal{J}_\varepsilon(U)))} \\ & \lesssim_{\alpha, q} (\varepsilon \mu_d(1/\varepsilon))^{1/2} \|f\|_{W^{1+\alpha, \infty}(U)}, \end{aligned}$$

where μ_d is defined in (2.3) and the effective matrix $\bar{\mathbf{b}} \in \mathbb{M}_0^{\operatorname{sym}}$ is given by

$$(2.13) \quad \bar{\mathbf{b}} : E := \frac{1}{d} \mathbb{E} \left[\sum_n \frac{\mathbb{1}_{I_n}}{|I_n|} \int_{\partial I_n} (x - x_n) \cdot \sigma(\psi_E + Ex, \Sigma_E) \nu \right].$$

In addition, if f and \bar{u} are compactly supported in U , then boundary layers disappear and the bound (2.12) holds with the optimal convergence rate $\varepsilon \mu_d(1/\varepsilon)$.

3. PERTURBATIVE ANNEALED REGULARITY

This section is devoted to the proof of the Meyers-type perturbative result stated in Theorem 5.

THEOREM 3.1 (Perturbative annealed L^p regularity). — *Under Assumption (\mathbf{H}_δ) , there exists a constant $C_0 \simeq 1$ such that the following holds: For all $g \in C_c^\infty(\mathbb{R}^d; L^\infty(\Omega)^{d \times d})$, the solution $(\nabla u_g, P_g)$ of the Stokes problem (2.4) satisfies for all p, q with $|p - 2|, |q - 2| \leq 1/C_0$,*

$$\|[\nabla u_g]_2\|_{L^p(\mathbb{R}^d; L^q(\Omega))} \lesssim \| [g]_2 \|_{L^p(\mathbb{R}^d; L^q(\Omega))}.$$

3.1. PRELIMINARY. — We start with a number of PDE ingredients that are useful in the proof.

3.1.1. *Whole-space weak formulations.* — The steady Stokes problem (2.4) can be reformulated as an equation on the whole space, where particles generate source terms concentrated at their boundaries. This reformulation is particularly convenient for our computations.

LEMMA 3.2. — *The solution $(\nabla u_g, P_g)$ of the steady Stokes problem (2.4) satisfies in the weak sense in the whole space \mathbb{R}^d ,*

$$(3.1) \quad -\Delta u_g + \nabla(P_g \mathbb{1}_{\mathbb{R}^d \setminus \mathcal{I}}) = \operatorname{div}(g \mathbb{1}_{\mathbb{R}^d \setminus \mathcal{I}}) - \sum_n \delta_{\partial I_n} (g + \sigma(u_g, P_g)) \nu.$$

Likewise, the corrector (ψ_E, Σ_E) in Lemma 1 satisfies in the weak sense in \mathbb{R}^d ,

$$(3.2) \quad -\Delta \psi_E + \nabla(\Sigma_E \mathbb{1}_{\mathbb{R}^d \setminus \mathcal{I}}) = - \sum_n \delta_{\partial I_n} \sigma(\psi_E + Ex, \Sigma_E) \nu.$$

Proof. — We focus on the proof of (3.1), while the argument for (3.2) is similar. Given $\zeta \in C_c^\infty(\mathbb{R}^d)^d$, testing equation (2.4) with ζ and integrating by parts on $\mathbb{R}^d \setminus \mathcal{I}$, we find

$$(3.3) \quad \int_{\mathbb{R}^d \setminus \mathcal{I}} \nabla \zeta : \nabla u - \int_{\mathbb{R}^d \setminus \mathcal{I}} \operatorname{div}(\zeta) P \\ = - \int_{\mathbb{R}^d \setminus \mathcal{I}} \nabla \zeta : g - \sum_n \int_{\partial I_n} (\zeta \otimes \nu) : (g + \nabla u - P \operatorname{Id}).$$

The claim (3.1) follows provided we prove that

$$(3.4) \quad \int_{\mathcal{I}} \nabla \zeta : \nabla u = - \sum_n \int_{\partial I_n} (\nu \otimes \zeta) : \nabla u.$$

Indeed, adding the latter to (3.3) yields the claim (3.1), in view of

$$\int_{\partial I_n} (\nu \otimes \zeta + \zeta \otimes \nu) : \nabla u = \int_{\partial I_n} \zeta \otimes \nu : 2D(u).$$

We turn to the proof of (3.4). Since u is affine in I_n , Stokes' theorem yields

$$\int_{\partial I_n} (\nu \otimes \zeta) : \nabla u = \int_{\partial I_n} \zeta_i \nu \cdot \partial_i u = \int_{I_n} \operatorname{div}(\zeta_i \partial_i u) = \int_{I_n} \nabla \zeta_i \cdot \partial_i u.$$

The relation $D(u) = 0$ on I_n entails that ∇u is skew-symmetric in I_n , so that the above becomes

$$\int_{\partial I_n} (\nu \otimes \zeta) : \nabla u = - \int_{I_n} \nabla \zeta_i \cdot \nabla u_i = - \int_{I_n} \nabla \zeta : \nabla u,$$

and the claim (3.4) follows. □

3.1.2. *Localized pressure estimates.* — We establish the following localized pressure estimate for the steady Stokes problem (2.4). It follows from standard pressure estimates in [22], but as in [18, Proof of Prop. 2.1] some additional care is needed to make it uniform with respect to the size of D although \mathcal{I} consists of an unbounded number of components; a short proof is included for convenience.

LEMMA 3.3 ([22, 18]). — *Given a deterministic point set $\{x_n\}_n$ satisfying the hardcore and regularity conditions in (H_δ) , for all $g \in L^2_{\text{loc}}(\mathbb{R}^d)^{d \times d}$ and all balls $D \subset \mathbb{R}^d$, any solution (u_g, P_g) of the steady Stokes problem (2.4) in D satisfies for all p with $1 < p < \infty$,*

$$\left\| P_g - \int_{D \setminus \mathcal{J}} P_g \right\|_{L^p(D \setminus \mathcal{J})} \lesssim_p \|(\nabla u_g, g)\|_{L^p(D \setminus \mathcal{J})}.$$

Proof. — We split the proof into two steps.

Step 1. Preliminary. — There is a vector field $S \in W^{1,p'}_0(D)^d$ such that $S|_{I_n}$ is constant for all n and such that

$$\begin{aligned} \operatorname{div}(S) &= \left(|Q|Q|^{p-2} - \int_{D \setminus \mathcal{J}} |Q|Q|^{p-2} \right) \mathbb{1}_{D \setminus \mathcal{J}}, & Q &:= P_g - \int_{D \setminus \mathcal{J}} P_g, \\ \|\nabla S\|_{L^{p'}(D)} &\lesssim_p \| |Q|Q|^{p-2} \|_{L^{p'}(D \setminus \mathcal{J})} = \|Q\|_{L^p(D \setminus \mathcal{J})}^{p-1}, \end{aligned}$$

where we emphasize that the prefactor in the last estimate is uniformly bounded independently of D .

By a standard use of the Bogovskii operator in form of [22, Th. III.3.1], there exists a vector field $S^\circ \in W^{1,p'}_0(D)^d$ such that

$$\begin{aligned} \operatorname{div}(S^\circ) &= \left(|Q|Q|^{p-2} - \int_{D \setminus \mathcal{J}} |Q|Q|^{p-2} \right) \mathbb{1}_{D \setminus \mathcal{J}}, \\ \|\nabla S^\circ\|_{L^{p'}(D)} &\lesssim_p \| |Q|Q|^{p-2} \|_{L^{p'}(D \setminus \mathcal{J})}. \end{aligned}$$

We need to modify S° to make it constant in I_n while keeping the divergence-free constraint and controlling the norm. For all n such that $I_n + (\delta/2)B \subset D$, choose an extension $\tilde{S}^\circ_n \in W^{1,p'}_0(I_n + (\delta/2)B)^d$ such that $\tilde{S}^\circ_n = -S^\circ + \int_{I_n} S^\circ$ in I_n and

$$\|\tilde{S}^\circ_n\|_{W^{1,p'}(I_n + (\delta/2)B)} \lesssim \left\| S^\circ - \int_{I_n} S^\circ \right\|_{W^{1,p'}(I_n)}.$$

So defined, $S^\circ + \tilde{S}^\circ_n$ is constant on I_n but not divergence-free. By a standard use of the Bogovskii operator in form of [22, Th. III.3.1], there exists a vector field $\tilde{S}^n \in W^{1,p'}_0((I_n + (\delta/2)B) \setminus I_n)^d$ (extended by 0 in I_n) such that

$$\begin{aligned} \operatorname{div}(\tilde{S}^n) &= -\operatorname{div}(\tilde{S}^\circ_n), \quad \text{in } (I_n + (\delta/2)B) \setminus I_n, \\ \|\nabla \tilde{S}^n\|_{L^{p'}((I_n + (\delta/2)B) \setminus I_n)} &\lesssim_p \|\nabla \tilde{S}^\circ_n\|_{L^{p'}(I_n + (\delta/2)B)}. \end{aligned}$$

We then define $S^n := \tilde{S}^\circ_n + \tilde{S}^n \in W^{1,p'}_0(I_n + (\delta/2)B)^d$, which satisfies $S^n = \tilde{S}^\circ_n = -S^\circ + \int_{I_n} S^\circ$ in I_n and in addition, combining the above with Poincaré’s inequality,

$$(3.5) \quad \begin{aligned} \operatorname{div}(S^n) &= 0, \\ \|\nabla S^n\|_{L^{p'}(I_n + (\delta/2)B)} &\lesssim_p \|\nabla S^\circ\|_{L^{p'}(I_n)}. \end{aligned}$$

For all n such that $(I_n + (\delta/2)B) \cap \partial D \neq \emptyset$, we proceed to a similar construction, replacing $I_n + (\delta/2)B$ by $(I_n + (\delta/2)B) \cap D$, and $\int_{I_n} S^\circ$ by zero. Using Poincaré’s inequality on $(I_n + \delta B) \cap D$, rather than Poincaré’s inequality with vanishing average

on $I_n + (\delta/2)B$, this provides a vector field $S^n \in W_0^{1,p'}((I_n + (\delta/2)B) \cap D)^d$, which satisfies $S^n = -S^\circ$ in $I_n \cap D$ and

$$\begin{aligned} \operatorname{div}(S^n) &= 0, \\ \|\nabla S^n\|_{L^{p'}((I_n + (\delta/2)B) \cap D)} &\lesssim_p \|\nabla S^\circ\|_{L^{p'}((I_n + \delta B) \cap D)}. \end{aligned}$$

Since the fattened inclusions $\{(I_n + \delta B) \cap D\}_n$ are all disjoint, cf. (H_δ) , implicitly extending S^n by 0 outside its domain of definition, the vector field $S := S^\circ + \sum_n S^n$ satisfies all the required properties.

Step 2. Conclusion. — Testing equation (3.1) with S , using that S is constant inside particles, and recalling the boundary conditions for u_g , cf. (2.4), we are led to

$$\int_{D \setminus \mathcal{I}} \operatorname{div}(S) P_g = \int_D \nabla S : \nabla u_g - \int_{D \setminus \mathcal{I}} \nabla S : g.$$

Inserting the definition of $\operatorname{div}(S)$, recalling that ∇S vanishes in \mathcal{I} , and using Hölder’s inequality, we find

$$\|Q\|_{L^p(D \setminus \mathcal{I})}^p \lesssim_p \|\nabla S\|_{L^{p'}(D)} \|(\nabla u_g, g)\|_{L^p(D \setminus \mathcal{I})},$$

and the claim follows from the bound on the norm of ∇S in Step 1. □

3.1.3. Dual Calderón–Zygmund lemma. — As in [20], we shall appeal to the following dual version of the Calderón–Zygmund lemma due to Shen; the present statement is a variant of [41, Th. 3.2] (see also [42, Th. 2.4]). For a ball $D \subset \mathbb{R}^d$, we henceforth set $D = B_{r_D}(x_D)$ and use the abusive short-hand notation $kD := B_{kr_D}(x_D)$ for dilations centered at the same point.

LEMMA 3.4 ([41]). — *Given $1 \leq p_0 < p_1 \leq \infty$, $F, G \in L^{p_0} \cap L^{p_1}(\mathbb{R}^d)$, and $C_0 > 0$, assume that for all balls $D \subset \mathbb{R}^d$ there exist measurable functions $F_{D,0}$ and $F_{D,1}$ such that $|F| \leq |F_{D,0}| + |F_{D,1}|$ and $|F_{D,1}| \leq |F| + |F_{D,0}|$ on D , and such that*

$$\begin{aligned} \left(\int_D |F_{D,0}|^{p_0} \right)^{1/p_0} &\leq C_0 \left(\int_{C_0 D} |G|^{p_0} \right)^{1/p_0}, \\ \left(\int_{(1/C_0)D} |F_{D,1}|^{p_1} \right)^{1/p_1} &\leq C_0 \left(\int_D |F_{D,1}|^{p_0} \right)^{1/p_0}. \end{aligned}$$

Then, for all p with $p_0 < p < p_1$,

$$\left(\int_{\mathbb{R}^d} |F|^p \right)^{1/p} \lesssim_{C_0, p_0, p, p_1} \left(\int_{\mathbb{R}^d} |G|^p \right)^{1/p}.$$

3.1.4. Gehring’s lemma. — We shall appeal to the following version of Gehring’s lemma, which is a mild reformulation of [24, Prop. 5.1].

LEMMA 3.5 ([23, 24]). — *Given $1 < q < s$ and a reference cube $Q_0 \subset \mathbb{R}^d$, let $G \in L^q(Q_0)$ and $F \in L^s(Q_0)$ be nonnegative functions. There exist $\theta_0 > 0$ (only depending*

on d, q, s) with the following property: Given $\theta \leq \theta_0$, if for some $C_0 \geq 1$ the following condition holds for all cubes $Q \subset Q_0$,

$$\left(\int_{(1/C_0)Q} G^q \right)^{1/q} \leq C_0 \int_Q G + C_0 \left(\int_Q F^q \right)^{1/q} + \theta \left(\int_Q G^q \right)^{1/q},$$

then there exists $\eta_0 > 0$ (only depending on C_0, d, q, s) such that for all $q \leq p \leq q + \eta_0$,

$$\left(\int_{(1/C_0)Q_0} G^p \right)^{1/p} \lesssim_{C_0, q, r} \int_{Q_0} G + \left(\int_{Q_0} F^p \right)^{1/p}.$$

3.2. PROOF OF THEOREM 3.1. — The starting point is the following deterministic perturbative result, for which an argument is postponed to Section 3.3.

PROPOSITION 3.6. — Given a deterministic inclusion set \mathcal{I} satisfying the hardcore and regularity conditions in (H_δ) , there exists a constant $C_0 \simeq 1$ such that the following hold.

(i) Meyers-type L^p estimate:

Given $g \in C_c^\infty(\mathbb{R}^d)^{d \times d}$, the solution $(\nabla u_g, P_g)$ of the steady Stokes problem (2.4) satisfies for all p with $2 \leq p \leq 2 + 1/C_0$,

$$\|[\nabla u_g]_2\|_{L^p(\mathbb{R}^d)} \lesssim \|g\|_{L^p(\mathbb{R}^d)}.$$

(ii) Reverse Jensen’s inequality:

For any ball $D \subset \mathbb{R}^d$, if (w, Q) satisfies the following equations in D ,

$$\begin{cases} -\Delta w + \nabla Q = 0, & \text{in } D \setminus \mathcal{I}, \\ \operatorname{div}(w) = 0, & \text{in } D \setminus \mathcal{I}, \\ D(w) = 0, & \text{in } \mathcal{I}, \\ \int_{\partial I_n} \sigma(w, Q)\nu = 0, & \forall n : I_n \subset D, \\ \int_{\partial I_n} \Theta(x - x_n) \cdot \sigma(w, Q)\nu = 0, & \forall n : I_n \subset D, \forall \Theta \in \mathbb{M}^{\text{skew}}, \end{cases}$$

then there holds for all $q \leq p$ with $|p - 2|, |q - 2| \leq 1/C_0$,

$$\left(\int_{(1/C)D} [\nabla w]_2^p \right)^{1/p} \lesssim \left(\int_D [\nabla w]_2^q \right)^{1/q}.$$

We may now proceed with the proof of Theorem 3.1, which follows from the above together with Shen’s dual version of the Calderón–Zygmund lemma, cf. Lemma 3.4.

Proof of Theorem 3.1. — We split the proof into three steps. We start with estimates outside the particles: first for $2 \leq q < p$, and then for $p < q \leq 2$ by a duality argument, so that the full range of exponents is finally reached by interpolation. Next, we extend the estimates inside the particles. Let $C_0 \geq 1$ be fixed as in the statement of Proposition 3.6.

Step 1. Proof that for all p, q with $2 \leq q < p < 2 + 1/C_0$,

$$(3.6) \quad \|\nabla u_g\|_{L^p(\mathbb{R}^d; L^q(\Omega))} \lesssim \|g\|_{L^p(\mathbb{R}^d; L^q(\Omega))}.$$

Let p_0, p_1 with $2 \leq p_0 \leq p_1 \leq 2 + 1/C_0$ be fixed. For balls $D \subset \mathbb{R}^d$, we decompose

$$\nabla u_g = \nabla u_{D,0} + \nabla u_{D,1},$$

where $\nabla u_{D,0} \in L^\infty(\Omega; L^2(\mathbb{R}^d)^{d \times d})$ denotes the unique solution of

$$\begin{cases} -\Delta u_{D,0} + \nabla P_{D,0} = \operatorname{div}(g \mathbb{1}_D), & \text{in } \mathbb{R}^d \setminus \mathcal{I}, \\ \operatorname{div}(u_{D,0}) = 0, & \text{in } \mathbb{R}^d \setminus \mathcal{I}, \\ D(u_{D,0}) = 0, & \text{in } \mathcal{I}, \\ \int_{\partial I_n} (g \mathbb{1}_D + \sigma(u_{D,0}, P_{D,0})) \nu = 0, & \forall n, \\ \int_{\partial I_n} \Theta(x - x_n) \cdot (g \mathbb{1}_D + \sigma(u_{D,0}, P_{D,0})) \nu = 0, & \forall n, \forall \Theta \in \mathbb{M}^{\text{skew}}. \end{cases}$$

On the one hand, for balls D with radius $r_D > 1$, Proposition 3.6(i) applied to the above equation yields

$$\int_D \mathbb{E} \left[\|\nabla u_{D,0}\|_2^{p_0} \right] \leq \mathbb{E} \left[\int_{\mathbb{R}^d} \|\nabla u_{D,0}\|_2^{p_0} \right] \lesssim \mathbb{E} \left[\int_{\mathbb{R}^d} \|g \mathbb{1}_D\|_2^{p_0} \right] \leq \int_{2D} \mathbb{E} \left[\|g\|_2^{p_0} \right],$$

while for balls D with radius $r_D < 1$ we appeal to the plain energy inequality (2.5) in form of

$$\begin{aligned} \int_D \mathbb{E} \left[\|\nabla u_{D,0}\|_2^{p_0} \right] &\leq |D| \mathbb{E} \left[\left(\int_{\mathbb{R}^d} |\nabla u_{D,0}|^2 \right)^{p_0/2} \right] \lesssim |D| \mathbb{E} \left[\left(\int_D |g|^2 \right)^{p_0/2} \right] \\ &\lesssim \int_{2D} \mathbb{E} \left[\|g\|_2^{p_0} \right]. \end{aligned}$$

On the other hand, noting that $\nabla u_{D,1} = \nabla u_g - \nabla u_{D,0}$ satisfies

$$\begin{cases} -\Delta u_{D,1} + \nabla P_{D,1} = 0, & \text{in } D \setminus \mathcal{I}, \\ \operatorname{div}(u_{D,1}) = 0, & \text{in } D \setminus \mathcal{I}, \\ D(u_{D,1}) = 0, & \text{in } \mathcal{I}, \\ \int_{\partial I_n} \sigma(u_{D,1}, P_{D,1}) \nu = 0, & \forall n : I_n \subset D, \\ \int_{\partial I_n} \Theta(x - x_n) \cdot \sigma(u_{D,1}, P_{D,1}) \nu = 0, & \forall n : I_n \subset D, \forall \Theta \in \mathbb{M}^{\text{skew}}, \end{cases}$$

it follows from the Minkowski inequality and from Proposition 3.6(ii) that

$$\begin{aligned} \left(\int_{(1/C)D} \mathbb{E} \left[\|\nabla u_{D,1}\|_2^{p_0} \right]^{p_1/p_0} \right)^{1/p_1} &\leq \mathbb{E} \left[\left(\int_{(1/C)D} \|\nabla u_{D,1}\|_2^{p_1} \right)^{p_0/p_1} \right]^{1/p_0} \\ &\lesssim \mathbb{E} \left[\int_D \|\nabla u_{D,1}\|_2^{p_0} \right]^{1/p_0} = \left(\int_D \mathbb{E} \left[\|\nabla u_{D,1}\|_2^{p_0} \right] \right)^{1/p_0}. \end{aligned}$$

In view of these estimates, appealing to Lemma 3.4 with

$$\begin{aligned} F &:= \mathbb{E} \left[\|\nabla u_g\|_2^{p_0} \right]^{1/p_0}, \quad G := \mathbb{E} \left[\|g\|_2^{p_0} \right]^{1/p_0}, \\ F_{D,0} &:= \mathbb{E} \left[\|\nabla u_{D,0}\|_2^{p_0} \right]^{1/p_0}, \quad F_{D,1} := \mathbb{E} \left[\|\nabla u_{D,1}\|_2^{p_0} \right]^{1/p_0}, \end{aligned}$$

we deduce for all p with $p_0 < p < p_1$,

$$\left(\int_{\mathbb{R}^d} \mathbb{E} [[\nabla u_g]_2^{p_0}]^{p/p_0} \right)^{1/p} \lesssim \left(\int_{\mathbb{R}^d} \mathbb{E} [[g]_2^{p_0}]^p / p_0 \right)^{1/p},$$

and the claim (3.6) follows (with q replaced by p_0).

Step 2. Duality and interpolation: proof that for all p, q with $2 - 1/2C_0 < p < q \leq 2$,

$$(3.7) \quad \| [\mathbb{1}_{\mathbb{R}^d \setminus \mathcal{J}} \nabla u]_2 \|_{L^p(\mathbb{R}^d; L^q(\Omega))} \lesssim \| [g]_2 \|_{L^p(\mathbb{R}^d; L^q(\Omega))}.$$

Combining this with (3.6), we then deduce by interpolation that the same estimate holds for all p, q with $|p - 2|, |q - 2| < 1/8C_0$.

Given a test function $h \in C_c^\infty(\mathbb{R}^d; L^\infty(\Omega)^{d \times d})$, we consider the solution $(\nabla u_h, P_h)$ of the steady Stokes problem (2.4) with g replaced by h . In view of (3.1), there holds in the weak sense in \mathbb{R}^d ,

$$\begin{aligned} -\Delta u_g + \nabla(P_g \mathbb{1}_{\mathbb{R}^d \setminus \mathcal{J}}) &= \nabla \cdot (g \mathbb{1}_{\mathbb{R}^d \setminus \mathcal{J}}) - \sum_n \delta_{\partial I_n} (g + \sigma(u_g, P_g)) \nu, \\ -\Delta u_h + \nabla(P_h \mathbb{1}_{\mathbb{R}^d \setminus \mathcal{J}}) &= \nabla \cdot (h \mathbb{1}_{\mathbb{R}^d \setminus \mathcal{J}}) - \sum_n \delta_{\partial I_n} (h + \sigma(u_h, P_h)) \nu. \end{aligned}$$

Testing the equation for u_h with u_g , and vice versa, and noting that the boundary terms all vanish in view of the respective boundary conditions, we find

$$\int_{\mathbb{R}^d \setminus \mathcal{J}} h : \nabla u_g = - \int_{\mathbb{R}^d} \nabla u_h : \nabla u_g = \int_{\mathbb{R}^d \setminus \mathcal{J}} g : \nabla u_h.$$

Combined with a duality argument, this identity yields

$$\begin{aligned} &\| [\mathbb{1}_{\mathbb{R}^d \setminus \mathcal{J}} \nabla u_g]_2 \|_{L^p(\mathbb{R}^d; L^q(\Omega))} \\ &\lesssim \sup \left\{ \mathbb{E} \left[\int_{\mathbb{R}^d \setminus \mathcal{J}} h : \nabla u_g \right] : \| [h]_2 \|_{L^{p'}(\mathbb{R}^d; L^{q'}(\Omega))} = 1 \right\} \\ &= \sup \left\{ \mathbb{E} \left[\int_{\mathbb{R}^d \setminus \mathcal{J}} g : \nabla u_h \right] : \| [h]_2 \|_{L^{p'}(\mathbb{R}^d; L^{q'}(\Omega))} = 1 \right\} \\ &\leq \| [g]_2 \|_{L^q(\mathbb{R}^d; L^p(\Omega))} \sup \left\{ \| [\nabla u_h]_2 \|_{L^{p'}(\mathbb{R}^d; L^{q'}(\Omega))} : \| [h]_2 \|_{L^{p'}(\mathbb{R}^d; L^{q'}(\Omega))} = 1 \right\}. \end{aligned}$$

Given p, q with $2 - 1/2C_0 < p < q \leq 2$, we may appeal to (3.6) with $2 \leq q' < p' < 2 + 1/C_0$, and the claim (3.7) follows.

Step 3. Conclusion. — In view of Step 2, it remains to show that for all $p, q \geq 1$,

$$(3.8) \quad \| [\mathbb{1}_{\mathcal{J}} \nabla u_g]_2 \|_{L^p(\mathbb{R}^d; L^q(\Omega))} \lesssim \| [\mathbb{1}_{\mathbb{R}^d \setminus \mathcal{J}} \nabla u_g]_2 \|_{L^p(\mathbb{R}^d; L^q(\Omega))}.$$

For all n , since u is affine in I_n , we can write for any constant $c_n \in \mathbb{R}^d$,

$$\| \nabla u_g \|_{L^\infty(I_n)} \lesssim \| u_g - c_n \|_{L^1(\partial I_n)}.$$

By a trace estimate and by Poincaré’s inequality with the choice $c_n := \int_{(I_n + \delta B) \setminus I_n} u_g$, we deduce

$$\| \nabla u_g \|_{L^\infty(I_n)} \lesssim \| u_g - c_n \|_{W^{1,1}((I_n + \delta B) \setminus I_n)} \lesssim \| \nabla u_g \|_{L^1((I_n + \delta B) \setminus I_n)}.$$

We may then estimate pointwise,

$$\mathbb{1}_{\mathcal{I}} |\nabla u_g| \lesssim \sum_n \mathbb{1}_{I_n} \|\nabla u_g\|_{L^1((I_n + \delta B) \setminus I_n)},$$

and the claim (3.8) now follows from the hardcore condition in (H_δ) . □

3.3. PROOF OF PROPOSITION 3.6. — We split the proof into two steps. We start with a Meyers-type perturbative argument based on Caccioppoli’s inequality and Gehring’s lemma, and we conclude in the second step.

Step 1. Meyers-type perturbative argument: there exists $C_0 \geq 1$ (only depending on d, δ) such that for all balls $D \subset \mathbb{R}^d$ and $2 \leq p \leq 2 + 1/C_0$,

$$(3.9) \quad \left(\int_D |\nabla u_g|_2^p \right)^{1/p} \lesssim \left(\int_{C_0 D} |\nabla u_g|_2^{2d/(d+2)} \right)^{(d+2)/2d} + \left(\int_{C_0 D} [g]_2^p \right)^{1/p}.$$

Given a ball $D \subset \mathbb{R}^d$ with radius $r_D \geq 3$, choose a cut-off function χ_D with $\chi_D|_D \equiv 1$, $\chi_D|_{\mathbb{R}^d \setminus 2D} \equiv 0$, and $|\nabla \chi_D| \lesssim 1/r_D$, such that χ_D is constant in I_n for all n . Given arbitrary constants $c_D \in \mathbb{R}^d$ and $c'_D \in \mathbb{R}$, testing the equation (3.1) for u_g with $\chi_D^2(u_g - c_D)$, noting that the boundary terms all vanish, and recalling that $\operatorname{div}(u_g) = 0$, we obtain the following Caccioppoli-type inequality,

$$\begin{aligned} \int_D |\nabla u_g|^2 &\lesssim \frac{1}{r_D^2} \int_{2D} |u_g - c_D|^2 + \int_{2D} |g|^2 \\ &\quad + \left(\frac{1}{r_D^2} \int_{2D} |u_g - c_D|^2 \right)^{1/2} \left(\int_{2D} |P_g - c'_D|^2 \mathbb{1}_{\mathbb{R}^d \setminus \mathcal{I}} \right)^{1/2}. \end{aligned}$$

Hence, for all $K \geq 1$,

$$\int_D |\nabla u_g|^2 \lesssim \frac{K^2}{r_D^2} \int_{2D} |u_g - c_D|^2 + \int_{2D} |g|^2 + \frac{1}{K^2} \int_{2D} |P_g - c'_D|^2 \mathbb{1}_{\mathbb{R}^d \setminus \mathcal{I}}.$$

Using the the Poincaré–Sobolev inequality to estimate the first right-hand side term, with the choice $c_D := \int_{2D} u_g$, and using the localized pressure estimate of Lemma 3.3 to estimate the last right-hand side term, with the choice $c'_D := \int_{2D \setminus \mathcal{I}} P$, we deduce

$$(3.10) \quad \left(\int_D |\nabla u_g|^2 \right)^{1/2} \lesssim K \left(\int_{2D} |\nabla u_g|^{2d/(d+2)} \right)^{(d+2)/2d} + \left(\int_{2D} |g|^2 \right)^{1/2} + \frac{1}{K} \left(\int_{2D} |\nabla u_g|^2 \right)^{1/2}.$$

While this is proved here for all balls D with radius $r_D \geq 3$, taking local quadratic averages allows us to infer for all balls D (with any radius $r_D > 0$) and all $K \geq 1$ that

$$\left(\int_D |\nabla u_g|_2^2 \right)^{1/2} \lesssim K \left(\int_{2D} |\nabla u_g|_2^{2d/(d+2)} \right)^{(d+2)/2d} + \left(\int_{2D} [g]_2^2 \right)^{1/2} + \frac{1}{K} \left(\int_{2D} |\nabla u_g|_2^2 \right)^{1/2}.$$

Choosing K large enough, the claim (3.9) now follows from Gehring’s lemma in form of Lemma 3.5.

Step 2. Conclusion. — We start with the proof of (i). Applying (3.9) together with Jensen’s inequality and with the energy inequality (2.5), we find for all p with $2 \leq p \leq 2 + 1/C_0$,

$$\begin{aligned} \left(\int_D [\nabla u_g]_2^p \right)^{1/p} &\lesssim |D|^{1/p-1/2} \left(\int_{CD} [\nabla u_g]_2^2 \right)^{1/2} + \left(\int_{CD} [g]_2^p \right)^{1/p} \\ &\lesssim |D|^{1/p-1/2} \left(\int_{\mathbb{R}^d} |g|^2 \right)^{1/2} + \left(\int_{CD} [g]_2^p \right)^{1/p}, \end{aligned}$$

hence the conclusion (i) follows for $D \uparrow \mathbb{R}^d$. Next, item (ii) is a consequence of (3.9) with $g = 0$ in CD . □

4. CORRECTOR ESTIMATES

This section is devoted to the proof of Theorem 2. Next to the corrector ψ_E , we further introduce an associated flux corrector ζ_E , which is key to put the equation for two-scale expansion errors into a more favorable form, cf. (6.3). As in [14, Th. 4], motivated by the work of Jikov on homogenization problems with stiff inclusions [43, 35] (see also [36, §3.2]), we start by defining a divergence-free extension J_E of the flux $\sigma(\psi_E + Ex, \Sigma_E) \mathbb{1}_{\mathbb{R}^d \setminus \mathcal{I}}$. Although this extension is not unique, we can choose it as in [14] to coincide with the flux in the corresponding incompressible linear elasticity problem in the limit of inclusions with diverging shear modulus. The flux corrector ζ_E is then defined as a vector potential for this extended flux J_E ; more precisely, equation (4.1) below amounts to choosing the Coulomb gauge. The construction is recalled for convenience in Section 4.1.

LEMMA 4.1 (Extended fluxes and flux correctors; [14]). — *Under Assumption (H_δ) , for all $E \in \mathbb{M}_0$, there is a stationary random 2-tensor field $J_E := \{J_{E;ij}\}_{1 \leq i,j \leq d}$ with finite second moment such that almost surely,*

$$J_E \mathbb{1}_{\mathbb{R}^d \setminus \mathcal{I}} = \sigma(\psi_E + Ex, \Sigma_E) \mathbb{1}_{\mathbb{R}^d \setminus \mathcal{I}}, \quad \operatorname{div}(J_E) = 0.$$

In these terms, there exists a unique random 3-tensor field $\zeta_E = \{\zeta_{E;ijk}\}_{1 \leq i,j,k \leq d}$ that satisfies the following infinite-volume problem:

– *For all i, j, k , almost surely, $\zeta_{E;ijk}$ belongs to $H^1_{\text{loc}}(\mathbb{R}^d)$ and satisfies in the weak sense,*

$$(4.1) \quad -\Delta \zeta_{E,ijk} = \partial_j J_{E,ik} - \partial_k J_{E,ij}.$$

– *The random field $\nabla \zeta_E$ is stationary, has vanishing expectation, has finite second moment, and ζ_E satisfies the anchoring condition $\int_B \zeta_E = 0$ almost surely.*

In addition, the following properties are automatically satisfied:

- (i) ζ_E is skew-symmetric in its last two indices, that is, $\zeta_{E,ijk} = -\zeta_{E,ikj}$ for all i, j, k ;
- (ii) ζ_E is a vector potential for J_E , that is,

$$\operatorname{div}(\zeta_{E,i}) = J_{E,i} - \mathbb{E}[J_{E,i}],$$

in terms of $\zeta_{E,i} = \{\zeta_{E,ijk}\}_{1 \leq j,k \leq d}$ and $J_{E,i} = \{J_{E,ij}\}_{1 \leq j \leq d}$

- (iii) ζ_E is sublinear at infinity, that is, $\varepsilon\zeta_E(\cdot/\varepsilon) \rightarrow 0$ in $H^1_{\text{loc}}(\mathbb{R}^d)$ almost surely as $\varepsilon \downarrow 0$;
- (iv) $\mathbb{E}[J_E] = 2\overline{\mathbf{B}}E + (\overline{\mathbf{b}} : E) \text{Id}$, where we recall that the effective constants $\overline{\mathbf{B}}$ and $\overline{\mathbf{b}}$ are defined in (2.11) and (2.13).

With the above definition, we shall establish the following version of Theorem 2 for the extended corrector (ψ_E, ζ_E) ; the proof is postponed to Section 4.2.

THEOREM 4.2 (Extended corrector estimate). — *Under Assumptions (\mathbf{H}_δ) and (Mix^+) , for all $E \in \mathbb{M}_0$ and $q < \infty$,*

$$(4.2) \quad \|[(\nabla\psi_E, \Sigma_E \mathbb{1}_{\mathbb{R}^d \setminus \mathcal{J}}, \nabla\zeta_E)]_2\|_{L^q(\Omega)} \lesssim_q |E|,$$

and

$$(4.3) \quad \|[(\psi_E, \zeta_E)]_2(x)\|_{L^q(\Omega)} \lesssim_q |E| \mu_d(|x|),$$

where μ_d is defined in (2.3).

4.1. PROOF OF LEMMA 4.1. — Let $E \in \mathbb{M}_0$. We split the proof into two main steps.

Step 1. Construction of the extended flux J_E . — Given a realization of the set of inclusions, we consider for all n the weak solution (ψ_E^n, Σ_E^n) in $H^1(I_n)^d \times L^2(I_n)$ of the following Neumann problem in I_n ,

$$(4.4) \quad \begin{cases} -\Delta\psi_E^n + \nabla\Sigma_E^n = 0, & \text{in } I_n, \\ \text{div}(\psi_E^n) = 0, & \text{in } I_n, \\ \sigma(\psi_E^n, \Sigma_E^n)\nu = \sigma(\psi_E + Ex, \Sigma_E)\nu, & \text{on } \partial I_n. \end{cases}$$

Note that ψ_E^n is defined only up to a rigid motion, which is fixed by choosing $\int_{I_n} \psi_E^n = 0$ and $\int_{I_n} \nabla\psi_E^n \in \mathbb{M}_0^{\text{sym}}$, and we prove that (ψ_E^n, Σ_E^n) satisfies

$$(4.5) \quad \|(\nabla\psi_E^n, \Sigma_E^n)\|_{L^2(I_n)} \lesssim \|\sigma(\psi_E + Ex, \Sigma_E)\|_{L^2((I_n + \delta B) \setminus I_n)}.$$

Substep 1.1. Well-posedness of the Neumann problem (4.4) for ψ_E^n . — The weak formulation of (4.4) takes on the following guise: ψ_E^n is divergence-free and satisfies for all divergence-free test functions $\phi \in H^1(I_n)^d$,

$$(4.6) \quad 2 \int_{I_n} \mathbf{D}(\phi) : \mathbf{D}(\psi_E^n) = \mathcal{L}_E(\phi),$$

in terms of the linear functional

$$\mathcal{L}_E(\phi) := \int_{\partial I_n} \phi \cdot \sigma(\psi_E + Ex, \Sigma_E)\nu.$$

In view of the boundary conditions for ψ_E , we can rewrite for any $V \in \mathbb{R}^d$ and $\Theta \in \mathbb{M}^{\text{skew}}$,

$$\mathcal{L}_E(\phi) = \int_{\partial I_n} (\phi - V - \Theta(\cdot - x_n)) \cdot \sigma(\psi_E + Ex, \Sigma_E)\nu.$$

Choose an extension map

$$T_n : \{\phi \in H^1(I_n)^d : \text{div}(\phi) = 0\} \longrightarrow \{\phi \in H^1_0(I_n + \delta B)^d : \text{div}(\phi) = 0\},$$

such that $T_n[\phi]|_{I_n} = \phi|_{I_n}$ and

$$\|T_n[\phi]\|_{H^1(I_n+\delta B)} \lesssim \|\phi\|_{H^1(I_n)}.$$

In these terms, using Stokes' theorem and recalling that $\sigma(\psi_E + Ex, \Sigma_E)$ is symmetric and divergence-free, we can further rewrite

$$\begin{aligned} \mathcal{L}_E(\phi) &= - \int_{(I_n+\delta B) \setminus I_n} \operatorname{div} \left(\sigma(\psi_E + Ex, \Sigma_E) T_n[\phi - V - \Theta(\cdot - x_n)] \right) \\ &= - \int_{(I_n+\delta B) \setminus I_n} \operatorname{D}(T_n[\phi - V - \Theta(\cdot - x_n)]) : \sigma(\psi_E + Ex, \Sigma_E), \end{aligned}$$

and thus, since $\operatorname{D}(T_n[\phi - V - \Theta(\cdot - x_n)])$ is trace-free,

$$(4.7) \quad \mathcal{L}_E(\phi) = -2 \int_{(I_n+\delta B) \setminus I_n} \operatorname{D}(T_n[\phi - V - \Theta(\cdot - x_n)]) : (\operatorname{D}(\psi_E) + E).$$

We deduce that $\phi \mapsto \mathcal{L}_E(\phi)$ is a continuous linear functional on the subspace $\{\phi \in H^1(I_n)^d : \operatorname{div}(\phi) = 0\}$. In addition, for all divergence-free $\phi \in H^1(I_n)^d$, minimizing over V, Θ and appealing to Korn's inequality, we find

$$\begin{aligned} |\mathcal{L}_E(\phi)| &\lesssim \inf_{\substack{V \in \mathbb{R}^d \\ \Theta \in \mathbb{M}^{\text{skew}}}} \|\phi - V - \Theta(\cdot - x_n)\|_{H^1(I_n)} \|\operatorname{D}(\psi_E) + E\|_{L^2((I_n+\delta B) \setminus I_n)} \\ &\lesssim \|\operatorname{D}(\phi)\|_{L^2(I_n)} \|\operatorname{D}(\psi_E) + E\|_{L^2((I_n+\delta B) \setminus I_n)}. \end{aligned}$$

By the Lax-Milgram theorem, we deduce that there exists a unique trace-free gradient-like solution $\operatorname{D}(\psi_E^n) \in L^2(I_n)_{\text{sym}}^{d \times d}$ of (4.6), and it satisfies

$$\|\operatorname{D}(\psi_E^n)\|_{L^2(I_n)} \lesssim \|\operatorname{D}(\psi_E) + E\|_{L^2((I_n+\delta B) \setminus I_n)}.$$

The vector field ψ_E^n is itself defined only up to a rigid motion and is fixed by choosing $\int_{I_n} \psi_E^n = 0$ and $\int_{I_n} \nabla \psi_E^n \in \mathbb{M}_0^{\text{sym}}$, in which case the above becomes by Korn's inequality,

$$(4.8) \quad \|\nabla \psi_E^n\|_{L^2(I_n)} \lesssim \|\operatorname{D}(\psi_E) + E\|_{L^2((I_n+\delta B) \setminus I_n)}.$$

Substep 1.2. Construction of the pressure. — Consider the extended deformation

$$q_E^n := \operatorname{D}(\psi_E) + E + \operatorname{D}(\psi_E^n) \mathbb{1}_{I_n}, \quad \text{in } I_n + \delta B.$$

In view of (4.7), the weak formulation (4.6) yields for all divergence-free test functions $\phi \in C_c^\infty(I_n + \delta B)^d$,

$$2 \int_{\mathbb{R}^d} \operatorname{D}(\phi) : q_E^n = 0.$$

Appealing e.g. to [36, Prop. 12.10], we deduce that there exists an associated pressure field $\Sigma_E^n \in L^2_{\text{loc}}(I_n + \delta B)$, which is unique up to an additive constant, such that for all test functions $\phi \in C_c^\infty(I_n + \delta B)^d$,

$$(4.9) \quad \int_{\mathbb{R}^d} \operatorname{D}(\phi) : (2q_E^n - \Sigma_E^n \operatorname{Id}) = 0.$$

Since for all $\phi \in C_c^\infty((I_n + \delta B) \setminus I_n)^d$ we have

$$\int_{\mathbb{R}^d} \operatorname{D}(\phi) : (2q_E^n - \Sigma_E^n \operatorname{Id}) = \int_{\mathbb{R}^d} \operatorname{D}(\phi) : (2(\operatorname{D}(\psi_E) + E) - \Sigma_E^n \operatorname{Id}) = 0,$$

we deduce that Σ_E^n can be chosen uniquely to coincide with Σ_E on $(I_n + \delta B) \setminus I_n$. The pair (ψ_E^n, Σ_E^n) is then the unique weak solution of the Neumann problem (4.4) with $f_{I_n} \psi_E^n = 0$ and $f_{I_n} \nabla \psi_E^n \in \mathbb{M}_0^{\text{sym}}$.

It remains to prove (4.5). The estimation of $\nabla \psi_E^n$ follows from (4.8) and it remains to estimate the pressure Σ_E^n . For that purpose, using that Σ_E^n coincides with Σ_E on $(I_n + \delta B) \setminus I_n$, we split

$$\begin{aligned} \|\Sigma_E^n\|_{L^2(I_n)} &\lesssim \left\| \Sigma_E^n - \int_{I_n + \delta B} \Sigma_E^n \right\|_{L^2(I_n + \delta B)} + \left| \int_{I_n + \delta B} \Sigma_E^n \right| \\ &\leq \left\| \Sigma_E^n - \int_{I_n + \delta B} \Sigma_E^n \right\|_{L^2(I_n + \delta B)} + \left| \int_{(I_n + \delta B) \setminus I_n} \Sigma_E^n \right| \\ &\quad + \left| \int_{(I_n + \delta B) \setminus I_n} \left(\Sigma_E^n - \int_{I_n + \delta B} \Sigma_E^n \right) \right| \\ &\lesssim \left\| \Sigma_E^n - \int_{I_n + \delta B} \Sigma_E^n \right\|_{L^2(I_n + \delta B)} + \left| \int_{(I_n + \delta B) \setminus I_n} \Sigma_E^n \right|. \end{aligned}$$

Starting from (4.9), a standard argument based on the Bogovskii operator yields

$$\left\| \Sigma_E^n - \int_{I_n + \delta B} \Sigma_E^n \right\|_{L^2(I_n + \delta B)} \lesssim \|q_E^n\|_{L^2(I_n + \delta B)},$$

so that the above becomes

$$\|\Sigma_E^n\|_{L^2(I_n)} \lesssim \|q_E^n\|_{L^2(I_n + \delta B)} + \|\Sigma_E\|_{L^2((I_n + \delta B) \setminus I_n)},$$

and the claim (4.5) follows from (4.8).

Substep 1.3. Construction of the extended flux. — We define the extended deformation and the extended pressure,

$$\tilde{q}_E := D(\psi_E) + E + \sum_n D(\psi_E^n) \mathbb{1}_{I_n}, \quad \tilde{\Sigma}_E := \Sigma_E \mathbb{1}_{\mathbb{R}^d \setminus \mathcal{J}} + \sum_n \Sigma_E^n \mathbb{1}_{I_n},$$

as well as the corresponding extended flux

$$(4.10) \quad J_E := 2\tilde{q}_E - \tilde{\Sigma}_E = \sigma(\psi_E + Ex, \Sigma_E) \mathbb{1}_{\mathbb{R}^d \setminus \mathcal{J}} + \sum_n \sigma(\psi_E^n, \Sigma_E^n) \mathbb{1}_{I_n}.$$

In view of (4.9), together with (3.2), the pair $(\tilde{q}_E, \tilde{\Sigma}_E)$ satisfies for all test functions $\phi \in C_c^\infty(\mathbb{R}^d)^d$,

$$(4.11) \quad \int_{\mathbb{R}^d} D(\phi) : (2\tilde{q}_E - \tilde{\Sigma}_E) = 0,$$

that is, J_E is divergence-free. The uniqueness of the extensions ensures that \tilde{q}_E and $\tilde{\Sigma}_E$ are both stationary, and we now prove that they have finite second moments. Combining the definition of \tilde{q}_E with the estimate (4.8) on ψ_E^n , we find for all $R > 0$,

$$\|\tilde{q}_E\|_{L^2(B_R)} \lesssim \|D(\psi_E) + E\|_{L^2(B_{R+3})},$$

and thus, by stationarity, letting $R \uparrow \infty$, and using the L^2 estimate on ψ_E , cf. Lemma 1,

$$\|\tilde{q}_E\|_{L^2(\Omega)} \lesssim \|D(\psi_E) + E\|_{L^2(\Omega)} \lesssim |E|.$$

For the pressure, starting from (4.11), a standard argument based on the Bogovskii operator yields for all $R > 0$,

$$\left\| \tilde{\Sigma}_E - \int_{B_R} \tilde{\Sigma}_E \right\|_{L^2(B_R)} \lesssim \|\tilde{q}_E\|_{L^2(B_R)},$$

and thus, by stationarity, letting $R \uparrow \infty$ and using the above L^2 estimate on \tilde{q}_E ,

$$\|\tilde{\Sigma}_E - \mathbb{E}[\tilde{\Sigma}_E]\|_{L^2(\Omega)} \lesssim \|\tilde{q}_E\|_{L^2(\Omega)} \lesssim |E|.$$

We conclude that $\|J_E\|_{L^2(\Omega)} \lesssim |E|$. The identity in item (iv) for the expectation $\mathbb{E}[J_E]$ follows from a direct computation, cf. [14, Lem. 4.2], and is not repeated here.

Step 2. Construction of the flux corrector ζ_E . — In view of standard stationary calculus, e.g. [36, §7] (see also [28, Proof of Lemma 1]), equation (4.1) admits a unique stationary gradient solution $\nabla\zeta_E \in L^2_{\text{loc}}(\mathbb{R}^d; L^2(\Omega)^{d \times d})$ with vanishing expectation and with

$$\|\nabla\zeta_E\|_{L^2(\Omega)} \lesssim \|J_E\|_{L^2(\Omega)} \lesssim |E|.$$

Items (i) and (ii) are easy consequences of the definition of ζ_E . As in Lemma 1, the additional sublinearity statement (iii) is a standard result for random fields having a stationary gradient with vanishing expectation, cf. e.g. [36, §7]. \square

4.2. PROOF OF THEOREM 4.2. — We start with the following estimate on the optimal CLT decay for large-scale averages of the extended corrector gradient $(\nabla\psi_E, \nabla\zeta_E)$ and of the pressure Σ_E . Due to the nonlinearity of the corrector equation with respect to randomness, local norms of $(\nabla\psi_E, \Sigma_E)$ also appear in the right-hand side of (4.12), which is a common difficulty in stochastic homogenization; this will be subsequently absorbed by a buckling argument, taking advantage of the CLT scaling.

PROPOSITION 4.3 (CLT scaling). — *Under Assumptions (H_δ) and (Mix^+) , for all $g \in C_c^\infty(\mathbb{R}^d)$, $E \in \mathbb{M}_0$, $R, s \geq 1$, and $1 \ll q < \infty$, we have*

$$(4.12) \quad \left\| \int_{\mathbb{R}^d} g(\nabla\psi_E, \Sigma_E \mathbb{1}_{\mathbb{R}^d \setminus \mathcal{I}}, \nabla\zeta_E) \right\|_{L^{2q}(\Omega)} \lesssim_q \|g\|_{L^2(\mathbb{R}^d)} \left(|E| + \left\| \left(\int_{B_R} [(\nabla\psi_E, \Sigma_E \mathbb{1}_{\mathbb{R}^d \setminus \mathcal{I}})]_2^{2s} \right)^{1/2s} \right\|_{L^{2q}(\Omega)} \right).$$

(Note that the smaller R and s are, the stronger the estimate.) In order to get such a control on stochastic moments, we appeal to the following consequence of the multiscale variance inequality (2.1) in (Mix^+) , cf. [15, Prop. 1.10(ii)].

LEMMA 4.4 (Control of higher moments; [15]). — *If the inclusion process \mathcal{I} satisfies the multiscale variance inequality (2.1) with some weight π , then we have for all q with $1 \leq q < \infty$ and all $\sigma(\mathcal{I})$ -measurable random variables $Y(\mathcal{I})$ with $\mathbb{E}[Y(\mathcal{I})] = 0$,*

$$(4.13) \quad \|Y(\mathcal{I})\|_{L^{2q}(\Omega)}^2 \lesssim q^2 \mathbb{E} \left[\int_0^\infty \left(\int_{\mathbb{R}^d} \left(\partial_{\mathcal{I}, B_\ell(x)}^{\text{osc}} Y(\mathcal{I}) \right)^2 dx \right)^q \langle \ell \rangle^{-dq} \pi(\ell) d\ell \right]^{1/q}.$$

Next, in preparation for the buckling argument, we show how to bound local norms of $(\nabla\psi_E, \Sigma_E \mathbb{1}_{\mathbb{R}^d \setminus \mathcal{J}})$ as appearing in the right-hand side of (4.12) by corresponding large-scale averages. This statement is inspired by [37] in the context of homogenization for divergence-form linear elliptic equations.

PROPOSITION 4.5. — *Choose $\chi \in C_c^\infty(B)$ with $\int_B \chi = 1$, and set $\chi_r(x) := r^{-d}\chi(x/r)$. Under Assumption (H_δ) , for all $E \in \mathbb{M}_0$, r, R with $1 \ll_\chi r \ll_\chi R$ and $r/R \gtrsim_\chi 1$, and $q, s \geq 1$ with $|s - 1| \ll 1$,*

$$\left\| \left(\int_{B_R} [(\nabla\psi_E, \Sigma_E \mathbb{1}_{\mathbb{R}^d \setminus \mathcal{J}})]_2^{2s} \right) \right\|_{L^{2q}(\Omega)}^{1/2s} \lesssim_\chi |E| + \left\| \int_{\mathbb{R}^d} \chi_r (\nabla\psi_E, \Sigma_E \mathbb{1}_{\mathbb{R}^d \setminus \mathcal{J}}) \right\|_{L^{2q}(\Omega)}.$$

(The smaller (resp. larger) R, r (resp. s), the stronger the estimate.) Based on the above two propositions, we are now in position to proceed with the buckling argument and the proof of Theorem 4.2.

Proof of Theorem 4.2. — Let $E \in \mathbb{M}_0$ be fixed with $|E| = 1$. We split the proof into three steps: after some preliminary estimate, we establish the moment bounds (4.2) on $(\nabla\psi_E, \Sigma_E \mathbb{1}_{\mathbb{R}^d \setminus \mathcal{J}}, \nabla\zeta_E)$ by a buckling argument, before deducing the corresponding moment bounds (4.3) on (ψ_E, ζ_E) by integration.

Step 1. Preliminary: proof that for all $R \geq 1$,

$$(4.14) \quad \left\| [(\nabla\psi_E, \Sigma_E \mathbb{1}_{\mathbb{R}^d \setminus \mathcal{J}})]_2 \right\|_{L^{2q}(\Omega)} \lesssim (R^{d/2})^{1-1/q} \left\| \left(\int_{B_R} |(\nabla\psi_E, \Sigma_E \mathbb{1}_{\mathbb{R}^d \setminus \mathcal{J}})|^2 \right)^{1/2} \right\|_{L^{2q}(\Omega)}.$$

For $R, q \geq 1$, in view of local quadratic averages, the discrete $\ell^2 - \ell^{2q}$ inequality yields

$$\begin{aligned} & \left(\int_{B_R(x)} [(\nabla\psi_E, \Sigma_E \mathbb{1}_{\mathbb{R}^d \setminus \mathcal{J}}, \nabla\zeta_E)]_2^{2q} \right)^{1/2q} \\ & \lesssim \left(R^{-d} \sum_{z \in B_{2R}(x) \cap (1/C)\mathbb{Z}^d} [(\nabla\psi_E, \Sigma_E \mathbb{1}_{\mathbb{R}^d \setminus \mathcal{J}}, \nabla\zeta_E)]_2(z)^{2q} \right)^{1/2q} \\ & \lesssim (R^{d/2})^{1-1/q} \left(\int_{B_{4R}(x)} |(\nabla\psi_E, \Sigma_E \mathbb{1}_{\mathbb{R}^d \setminus \mathcal{J}}, \nabla\zeta_E)|^2 \right)^{1/2}. \end{aligned}$$

Taking the $L^{2q}(\Omega)$ norm and using the stationarity of $(\nabla\psi_E, \Sigma_E \mathbb{1}_{\mathbb{R}^d \setminus \mathcal{J}}, \nabla\zeta_E)$, the claim follows.

Step 2. Moment bounds (4.2). — Combining the results of Propositions 4.5 and 4.3, we find for all r, R with $1 \ll_\chi r \ll_\chi R$ and $r/R \gtrsim_\chi 1$, for all $q, s \geq 1$ with $1 \ll q < \infty$

and $|s - 1| \ll 1$,

$$\begin{aligned} & \left\| \left(\int_{B_R} [(\nabla\psi_E, \Sigma_E \mathbb{1}_{\mathbb{R}^d \setminus \mathcal{J}})]_2^{2s} \right)^{1/2s} \right\|_{L^{2q}(\Omega)} \\ & \lesssim_\chi 1 + \left\| \int_{\mathbb{R}^d} \chi_r (\nabla\psi_E, \Sigma_E \mathbb{1}_{\mathbb{R}^d \setminus \mathcal{J}}) \right\|_{L^{2q}(\Omega)} \\ & \lesssim_{q,\chi} 1 + (R/r)^{d/2} R^{-d/2s'} \left\| \left(1 + \int_{B_R} [(\nabla\psi_E, \Sigma_E \mathbb{1}_{\mathbb{R}^d \setminus \mathcal{J}})]_2^{2s} \right)^{1/2s} \right\|_{L^{2q}(\Omega)}. \end{aligned}$$

Letting r, R be fixed with $1 \ll_\chi r \ll_\chi R$ and $r \simeq_\chi R$, and choosing $R \gg_{q,s,\chi} 1$, we deduce

$$\left\| \left(\int_{B_R} [(\nabla\psi_E, \Sigma_E \mathbb{1}_{\mathbb{R}^d \setminus \mathcal{J}})]_2^{2s} \right)^{1/2s} \right\|_{L^{2q}(\Omega)} \lesssim_{q,s} 1.$$

Inserting this into (4.14) and using Jensen’s inequality, the moment bound (4.2) on $\nabla\psi_E$ and Σ_E follows.

It remains to prove the corresponding moment bound on the flux corrector gradient $\nabla\zeta_E$. Starting from equation (4.1) and appealing to localized maximal regularity theory for the Laplace equation, we find for all q with $1 \leq q < \infty$ and $R \geq 1$,

$$\left(\int_{B_R} [\nabla\zeta_E]_2^{2q} \right)^{1/2q} \lesssim_q \left(\int_{B_{2R}} |\nabla\zeta_E|^2 \right)^{1/2} + \left(\int_{B_{2R}} [J_E]_2^{2q} \right)^{1/2q},$$

hence, by the ergodic theorem, letting $R \uparrow \infty$,

$$\|[\nabla\zeta_E]_2\|_{L^{2q}(\Omega)} \lesssim_q \|\nabla\zeta_E\|_{L^2(\Omega)} + \|[J_E]_2\|_{L^{2q}(\Omega)}.$$

Using an energy estimate for (4.1) to bound the first right-hand side term, we are led to

$$\|[\nabla\zeta_E]_2\|_{L^{2q}(\Omega)} \lesssim_q \|[J_E]_2\|_{L^{2q}(\Omega)}.$$

By definition (4.10) of J_E , combined with (4.5), we deduce

$$\|[\nabla\zeta_E]_2\|_{L^{2q}(\Omega)} \lesssim_q 1 + \|[(\nabla\psi_E, \Sigma_E \mathbb{1}_{\mathbb{R}^d \setminus \mathcal{J}})]_2\|_{L^{2q}(\Omega)},$$

and the moment bound (4.2) on $\nabla\zeta_E$ now follows from the result on $\nabla\psi_E, \Sigma_E$.

Step 3. Moment bounds (4.3). — We focus on the bound on ψ_E , while the argument for ζ_E is similar. Poincaré’s inequality in $B(x)$ gives

$$(4.15) \quad \left\| \left[\psi_E - \int_B \psi_E \right]_2(x) \right\|_{L^{2q}(\Omega)} \lesssim \|[\nabla\psi_E]_2\|_{L^{2q}(\Omega)} + \left\| \int_{B(x)} \psi_E - \int_B \psi_E \right\|_{L^{2q}(\Omega)},$$

and it remains to estimate the second right-hand side term. For that purpose, we write

$$\int_{B(x)} \psi_E - \int_B \psi_E = \int_{\mathbb{R}^d} \nabla\psi_E \cdot \nabla h_x,$$

where h_x denotes the unique decaying solution in \mathbb{R}^d of

$$-\Delta h_x = \frac{1}{|B|} (\mathbb{1}_{B(x)} - \mathbb{1}_B).$$

Appealing to Proposition 4.3 together with the moment bounds (4.2), we find for all $q < \infty$,

$$\left\| \int_{\mathbb{R}^d} \nabla \psi_E \cdot \nabla h_x \right\|_{L^{2q}(\Omega)} \lesssim_q \|\nabla h_x\|_{L^2(\mathbb{R}^d)}.$$

A direct computation with Green’s kernel gives

$$\|\nabla h_x\|_{L^2(\mathbb{R}^d)} \lesssim \mu_d(|x|),$$

and thus

$$\left\| \int_{B(x)} \psi_E - \int_B \psi_E \right\|_{L^{2q}(\Omega)} \lesssim_q \mu_d(|x|).$$

Inserting this into (4.15), together with the moment bounds (4.2), the conclusion (4.3) for ψ_E follows. \square

4.3. PROOF OF PROPOSITION 4.3. — Let $E \in \mathbb{M}_0$ be fixed with $|E| = 1$. Applying the version (4.13) of the multiscale variance inequality (2.1) to control higher moments, we find

$$(4.16) \quad \left\| \int_{\mathbb{R}^d} g(\nabla \psi_E, \Sigma_E \mathbb{1}_{\mathbb{R}^d \setminus \mathcal{S}}, \nabla \zeta_E) \right\|_{L^{2q}(\Omega)}^2 \lesssim_q \mathbb{E} \left[\int_0^\infty \left(\int_{\mathbb{R}^d} \left(\partial_{\mathcal{S}, B_\ell(x)}^{\text{osc}} \int_{\mathbb{R}^d} g(\nabla \psi_E, \Sigma_E \mathbb{1}_{\mathbb{R}^d \setminus \mathcal{S}}, \nabla \zeta_E) \right)^2 dx \right)^q \langle \ell \rangle^{-dq} \pi(\ell) d\ell \right]^{1/q},$$

and it remains to estimate the oscillation of $\int_{\mathbb{R}^d} g(\nabla \psi_E, \Sigma_E \mathbb{1}_{\mathbb{R}^d \setminus \mathcal{S}}, \nabla \zeta_E)$ with respect to the inclusion process \mathcal{S} on any ball $B_\ell(x)$. Given $\ell \geq 0$ and $x \in \mathbb{R}^d$, and given a realization of \mathcal{S} , let \mathcal{S}' be a locally finite point set satisfying the hardcore and regularity conditions in (H_δ) , with $\mathcal{S}' \cap (\mathbb{R}^d \setminus B_\ell(x)) = \mathcal{S} \cap (\mathbb{R}^d \setminus B_\ell(x))$, and denote by $(\nabla \psi'_E, \Sigma'_E \mathbb{1}_{\mathbb{R}^d \setminus \mathcal{S}'}, \nabla \zeta'_E)$ the corresponding extended corrector with \mathcal{S} replaced by \mathcal{S}' (this is obviously well-defined in $L^2_{\text{loc}}(\mathbb{R}^d)$ as the perturbation is compactly supported). We split the proof into nine steps.

Step 1. Preliminary: dual test functions and annealed estimates. — As we shall abundantly appeal to duality arguments in the proof, this first step is devoted to the construction of a number of useful dual test functions and to the proof of corresponding annealed estimates:

– Given a test function $g \in C_c^\infty(\mathbb{R}^d; L^\infty(\Omega)^{d \times d})$, we let $\nabla u_g \in L^\infty(\Omega; L^2(\mathbb{R}^d)^{d \times d})$ denote the unique solution of the steady Stokes problem (2.4), and we recall that Theorem 3.1 yields for all q with $|q - 2| \ll 1$,

$$(4.17) \quad \|\nabla u_g\|_{L^2(\mathbb{R}^d; L^q(\Omega))} \lesssim \|g\|_{L^2(\mathbb{R}^d; L^q(\Omega))}.$$

– Given a test function $g \in C_c^\infty(\mathbb{R}^d; L^\infty(\Omega)^d)$, we let $\nabla v_g \in L^\infty(\Omega; L^2(\mathbb{R}^d)^d)$ denote the unique solution of

$$(4.18) \quad -\Delta v_g = \text{div}(g), \quad \text{in } \mathbb{R}^d,$$

which satisfies for all q with $1 < q < \infty$,

$$(4.19) \quad \|\nabla v_g\|_{L^2(\mathbb{R}^d; L^q(\Omega))} \lesssim_q \|g\|_{L^2(\mathbb{R}^d; L^q(\Omega))}.$$

– Given $g \in C_c^\infty(\mathbb{R}^d; L^\infty(\Omega))$, there exists a vector field $s_g \in L^\infty(\Omega; \dot{H}^1(\mathbb{R}^d)^d)$ such that $s_g|_{I_n}$ is constant for all n , and such that for all q with $1 < q < \infty$,

$$(4.20) \quad \begin{aligned} \operatorname{div}(s_g) &= g \mathbb{1}_{\mathbb{R}^d \setminus \mathcal{I}}, \quad \text{in } \mathbb{R}^d, \\ \|[\nabla s_g]_2\|_{L^2(\mathbb{R}^d; L^q(\Omega))} &\lesssim \|[g]_2\|_{L^2(\mathbb{R}^d; L^q(\Omega))}. \end{aligned}$$

– Given $g \in C_c^\infty(\mathbb{R}^d; L^\infty(\Omega)^{d \times d})$, there exists a 2-tensor field

$$h_g \in L^\infty(\Omega; H^1(\mathbb{R}^d)^{d \times d})$$

such that $h_g|_{I_n} = f_{I_n} g$ for all n , and such that for all q with $1 < q < \infty$,

$$(4.21) \quad \begin{aligned} \operatorname{div}(h_g) &= 0, \quad \text{in } \mathbb{R}^d, \\ \| [h_g]_2 \|_{L^2(\mathbb{R}^d; L^q(\Omega))} &\lesssim \| [g]_2 \|_{L^2(\mathbb{R}^d; L^q(\Omega))}. \end{aligned}$$

The existence and uniqueness of ∇v_g is clear, and the annealed bound (4.19) follows from Banach-valued Fourier multiplier theorems, e.g. in form of the extrapolation result in [39, Th. 3.15].

We turn to the construction of s_g . First denote by $s_g^\circ := \nabla w_g \in L^\infty(\Omega; \dot{H}^1(\mathbb{R}^d)^d)$ the solution of

$$\Delta w_g = g \mathbb{1}_{\mathbb{R}^d \setminus \mathcal{I}}, \quad \text{in } \mathbb{R}^d.$$

In view of (4.19), it satisfies for all q with $1 < q < \infty$,

$$\|[\nabla s_g^\circ]_2\|_{L^2(\mathbb{R}^d; L^q(\Omega))} \lesssim_q \|[g]_2\|_{L^2(\mathbb{R}^d; L^q(\Omega))}.$$

Next, as in (3.5), by a standard use of the Bogovskii operator in form of [22, Th. III.3.1], for all n , we can construct a vector field $s_g^n \in H_0^1(I_n + \delta B)^d$ such that $s_g^n = -s_g^\circ + f_{I_n} s_g^\circ$ in I_n , and

$$\begin{aligned} \operatorname{div}(s_g^n) &= 0, \\ \|\nabla s_g^n\|_{L^2((I_n + \delta B) \setminus I_n)} &\lesssim \|\nabla s_g^\circ\|_{L^2(I_n)}. \end{aligned}$$

Since the fattened inclusions $\{I_n + \delta B\}_n$ are disjoint, cf. (H_δ) , the vector field $s_g := s_g^\circ + \sum_n s_g^n$ (where we implicitly extend s_g^n by 0 outside $I_n + \delta B$) is checked to satisfy the required properties.

It remains to construct h_g . As in (3.5), using the Bogovskii operator in form of [22, Th. III.3.1], for all n , we can construct a 2-tensor field $h_g^n \in H_0^1(I_n + \delta B)^{d \times d}$ such that $h_g^n|_{I_n} = f_{I_n} g$, and

$$\begin{aligned} \operatorname{div}(h_g^n) &= 0, \\ \|\nabla h_g^n\|_{L^2((I_n + \delta B) \setminus I_n)} &\lesssim \|g\|_{L^2(I_n)}, \end{aligned}$$

and the tensor field $h_g = \sum_n h_g^n$ then satisfies the required properties.

Step 2. Preliminary: trace estimate. — For later reference, we prove the following general trace estimate: given a symmetric 2-tensor field $H \in L_{\text{loc}}^2(\mathbb{R}^d)^{d \times d}$ such that

$$\begin{cases} \operatorname{div}(H) = 0, & \text{in } (I_n + \delta B) \setminus I_n, \\ \int_{\partial I_n} H \nu = 0, \\ \int_{\partial I_n} \Theta(x - x_n) \cdot H \nu = 0, \quad \text{for all } \Theta \in \mathbb{M}^{\text{skew}}, \end{cases}$$

we have for all $g \in H^1_{\text{loc}}(\mathbb{R}^d)^d$,

$$(4.22) \quad \left| \int_{\partial I_n} g \cdot H\nu \right| \lesssim \left(\int_{(I_n + \delta B) \setminus I_n} |D(g)|^2 \right)^{1/2} \left(\int_{(I_n + \delta B) \setminus I_n} |H|^2 \right)^{1/2}.$$

We start by considering the following auxiliary Neumann problem,

$$\begin{cases} -\Delta z_n + \nabla R_n = 0, & \text{in } I_n, \\ \operatorname{div}(z_n) = 0, & \text{in } I_n, \\ \sigma(z_n, R_n)\nu = H\nu, & \text{on } \partial I_n. \end{cases}$$

Well-posedness for this problem is obtained as for (4.4) thanks to the assumptions on H , and the solution satisfies

$$\|(\nabla z_n, R_n)\|_{L^2(I_n)} \lesssim \|H\|_{L^2((I_n + \delta B) \setminus I_n)}.$$

Using the equation for z_n , Stokes' theorem yields

$$\int_{\partial I_n} g \cdot H\nu = \int_{\partial I_n} g \cdot \sigma(z_n, R_n)\nu = \int_{I_n} D(g) : \sigma(z_n, R_n),$$

and the claim (4.22) follows.

Step 3. Proof of

$$(4.23) \quad \int_{B_{\ell+3}(x)} |\nabla \psi'_E|^2 \lesssim \int_{B_{\ell+3}(x)} (1 + |\nabla \psi_E|^2),$$

$$(4.24) \quad \int_{B_{\ell+3}(x)} |\Sigma'_E|^2 \mathbb{1}_{\mathbb{R}^d \setminus \mathcal{J}'} \lesssim \int_{B_{\ell+3}(x)} (1 + |\nabla \psi_E|^2 + |\Sigma_E|^2 \mathbb{1}_{\mathbb{R}^d \setminus \mathcal{J}}).$$

Equation (3.2) for $\psi_E - \psi'_E$ takes the form

$$(4.25) \quad \begin{aligned} -\Delta(\psi_E - \psi'_E) + \nabla(\Sigma_E \mathbb{1}_{\mathbb{R}^d \setminus \mathcal{J}} - \Sigma'_E \mathbb{1}_{\mathbb{R}^d \setminus \mathcal{J}'}) \\ = -\sum_n \delta_{\partial I_n} \sigma(\psi_E + Ex, \Sigma_E)\nu + \sum_n \delta_{\partial I'_n} \sigma(\psi'_E + Ex, \Sigma'_E)\nu. \end{aligned}$$

Testing this equation with $\psi_E - \psi'_E$, we find

$$\begin{aligned} \int_{\mathbb{R}^d} |\nabla(\psi_E - \psi'_E)|^2 &= -\sum_n \int_{\partial I_n} (\psi_E - \psi'_E) \cdot \sigma(\psi_E + Ex, \Sigma_E)\nu \\ &\quad + \sum_n \int_{\partial I'_n} (\psi_E - \psi'_E) \cdot \sigma(\psi'_E + Ex, \Sigma'_E)\nu, \end{aligned}$$

which, by the boundary conditions, turns into

$$(4.26) \quad \begin{aligned} \int_{\mathbb{R}^d} |\nabla(\psi_E - \psi'_E)|^2 &= \sum_{n: I_n \cap B_\ell(x) \neq \emptyset} \int_{\partial I_n} \psi'_E \cdot \sigma(\psi_E + Ex, \Sigma_E)\nu \\ &\quad + \sum_{n: I'_n \cap B_\ell(x) \neq \emptyset} \int_{\partial I'_n} \psi_E \cdot \sigma(\psi'_E + Ex, \Sigma'_E)\nu. \end{aligned}$$

Note that, by Stokes' theorem, the constraints $\operatorname{div}(\psi_E) = \operatorname{div}(\psi'_E) = 0$ allow to replace the pressures Σ_E and Σ'_E in this identity by $\Sigma_E - c$ and $\Sigma'_E - c'$, respectively, for any constants $c, c' \in \mathbb{R}$. Appealing to the trace estimate (4.22), we are led to

$$\int_{\mathbb{R}^d} |\nabla(\psi_E - \psi'_E)|^2 \lesssim \left(\int_{B_{\ell+3}(x)} (1 + |\nabla\psi_E|^2 + |\Sigma_E - c|^2 \mathbb{1}_{\mathbb{R}^d \setminus \mathcal{J}}) \right)^{1/2} \times \left(\int_{B_{\ell+3}(x)} (1 + |\nabla\psi'_E|^2 + |\Sigma'_E - c'|^2 \mathbb{1}_{\mathbb{R}^d \setminus \mathcal{J}'}) \right)^{1/2}.$$

Choosing $c := \int_{B_{\ell+3}(x) \setminus \mathcal{J}} \Sigma_E$ and $c' := \int_{B_{\ell+3}(x) \setminus \mathcal{J}'} \Sigma'_E$, and using the pressure estimate of Lemma 3.3, we deduce

$$(4.27) \quad \int_{\mathbb{R}^d} |\nabla(\psi_E - \psi'_E)|^2 \lesssim \left(\int_{B_{\ell+3}(x)} (1 + |\nabla\psi_E|^2) \right)^{1/2} \left(\int_{B_{\ell+3}(x)} (1 + |\nabla\psi'_E|^2) \right)^{1/2},$$

and the claim (4.23) follows from the triangle inequality.

Next, we establish the corresponding bound (4.24) on the perturbed pressure. Using the Bogovskii operator as in the construction of s_g in Step 1, we can construct a vector field $S_E \in \dot{H}^1(\mathbb{R}^d)^d$ such that $S_E|_{I_n}$ is constant for all n and such that

$$\operatorname{div}(S_E) = (\Sigma_E \mathbb{1}_{\mathbb{R}^d \setminus \mathcal{J}} - \Sigma'_E \mathbb{1}_{\mathbb{R}^d \setminus \mathcal{J}'}) \mathbb{1}_{\mathbb{R}^d \setminus \mathcal{J}'},$$

$$\|\nabla S_E\|_{L^2(\mathbb{R}^d)} \lesssim \|\Sigma_E \mathbb{1}_{\mathbb{R}^d \setminus \mathcal{J}} - \Sigma'_E \mathbb{1}_{\mathbb{R}^d \setminus \mathcal{J}'}\|_{L^2(\mathbb{R}^d \setminus \mathcal{J}')}.$$

Testing equation (4.25) with S_E and using the boundary conditions, we find

$$\int_{\mathbb{R}^d} \operatorname{div}(S_E) (\Sigma_E \mathbb{1}_{\mathbb{R}^d \setminus \mathcal{J}} - \Sigma'_E \mathbb{1}_{\mathbb{R}^d \setminus \mathcal{J}'}) = \int_{\mathbb{R}^d} \nabla S_E : \nabla(\psi_E - \psi'_E) + \sum_{n: I_n \cap B_\ell(x) \neq \emptyset} \int_{\partial I_n} S_E \cdot \sigma(\psi_E + Ex, \Sigma_E) \nu,$$

which yields, by inserting the value of $\operatorname{div}(S_E)$ and using again the trace estimate (4.22),

$$\int_{\mathbb{R}^d \setminus \mathcal{J}'} |\Sigma_E \mathbb{1}_{\mathbb{R}^d \setminus \mathcal{J}} - \Sigma'_E \mathbb{1}_{\mathbb{R}^d \setminus \mathcal{J}'}|^2 \lesssim \left(\int_{\mathbb{R}^d} |\nabla S_E|^2 \right)^{1/2} \times \left(\int_{\mathbb{R}^d} |\nabla(\psi_E - \psi'_E)|^2 + \int_{B_{\ell+3}(x)} (1 + |\nabla\psi_E|^2 + |\Sigma_E|^2 \mathbb{1}_{\mathbb{R}^d \setminus \mathcal{J}}) \right)^{1/2}.$$

Appealing to the bound on the norm of ∇S_E , this yields

$$\int_{\mathbb{R}^d \setminus \mathcal{J}'} |\Sigma_E \mathbb{1}_{\mathbb{R}^d \setminus \mathcal{J}} - \Sigma'_E \mathbb{1}_{\mathbb{R}^d \setminus \mathcal{J}'}|^2 \lesssim \int_{\mathbb{R}^d} |\nabla(\psi_E - \psi'_E)|^2 + \int_{B_{\ell+3}(x)} (1 + |\nabla\psi_E|^2 + |\Sigma_E|^2 \mathbb{1}_{\mathbb{R}^d \setminus \mathcal{J}}).$$

Combining this with (4.27) and (4.23), the claim (4.24) follows by the triangle inequality.

Step 4. Sensitivity of the corrector gradient outside the inclusions: for all $g \in C_c^\infty(\mathbb{R}^d)^{d \times d}$,

$$(4.28) \quad \left| \int_{\mathbb{R}^d \setminus \mathcal{I}} g : \nabla \psi_E - \int_{\mathbb{R}^d \setminus \mathcal{I}'} g : \nabla \psi'_E \right| \\ \lesssim \left(\int_{B_{\ell+3}(x)} (|g|^2 + |\nabla u_g|^2) \right)^{1/2} \left(\int_{B_{\ell+3}(x)} (1 + |\nabla \psi_E|^2) \right)^{1/2}.$$

Decomposing $\int_{\mathbb{R}^d \setminus \mathcal{I}} - \int_{\mathbb{R}^d \setminus \mathcal{I}'} = \int_{\mathcal{I}' \setminus \mathcal{I}} - \int_{\mathcal{I} \setminus \mathcal{I}'}$, and noting that $(\mathcal{I}' \setminus \mathcal{I}) \cup (\mathcal{I} \setminus \mathcal{I}') \subset B_\ell(x)$, we find

$$(4.29) \quad \left| \int_{\mathbb{R}^d \setminus \mathcal{I}} g : \nabla \psi_E - \int_{\mathbb{R}^d \setminus \mathcal{I}'} g : \nabla \psi'_E \right| \\ \lesssim \left| \int_{\mathbb{R}^d \setminus \mathcal{I}} g : \nabla (\psi_E - \psi'_E) \right| + \left(\int_{B_\ell(x)} |g|^2 \right)^{1/2} \left(\int_{B_\ell(x)} |\nabla \psi'_E|^2 \right)^{1/2}.$$

It remains to examine the first right-hand side term, for which we appeal to a duality argument, in terms of the solution ∇u_g of (2.4). Testing with $\psi_E - \psi'_E$ the equation (3.1) for ∇u_g , and subtracting an arbitrary constant $c_1 \in \mathbb{R}$ to the pressure P_g , we obtain

$$\int_{\mathbb{R}^d \setminus \mathcal{I}} g : \nabla (\psi_E - \psi'_E) = - \int_{\mathbb{R}^d} \nabla u_g : \nabla (\psi_E - \psi'_E) \\ - \sum_n \int_{\partial I_n} (\psi_E - \psi'_E) \cdot (g + \sigma(u_g, P_g - c_1)) \nu,$$

which, in view of the boundary conditions, turns into

$$(4.30) \quad \int_{\mathbb{R}^d \setminus \mathcal{I}} g : \nabla (\psi_E - \psi'_E) = - \int_{\mathbb{R}^d} \nabla u_g : \nabla (\psi_E - \psi'_E) \\ + \sum_{n: I_n \cap B_\ell(x) \neq \emptyset} \int_{\partial I_n} \psi'_E \cdot (g + \sigma(u_g, P_g - c_1)) \nu.$$

Likewise, testing with u_g the equation (4.25) for $\psi_E - \psi'_E$, we get for any constant $c_2 \in \mathbb{R}$,

$$\int_{\mathbb{R}^d} \nabla u_g : \nabla (\psi_E - \psi'_E) = - \sum_n \int_{\partial I_n} u_g \cdot \sigma(\psi_E + Ex, \Sigma_E) \nu \\ + \sum_n \int_{\partial I'_n} u_g \cdot \sigma(\psi'_E + Ex, \Sigma'_E - c_2) \nu,$$

which, in view of the boundary conditions, takes the form

$$\int_{\mathbb{R}^d} \nabla u_g : \nabla (\psi_E - \psi'_E) = \sum_{n: I'_n \cap B_\ell(x) \neq \emptyset} \int_{\partial I'_n} u_g \cdot \sigma(\psi'_E + Ex, \Sigma'_E - c_2) \nu.$$

Combining this with (4.30), we obtain

$$\int_{\mathbb{R}^d \setminus \mathcal{I}} g : \nabla(\psi_E - \psi'_E) = \sum_{n: I_n \cap B_\ell(x) \neq \emptyset} \int_{\partial I_n} \psi'_E \cdot (g + \sigma(u_g, P_g - c_1)) \nu - \sum_{n: I'_n \cap B_\ell(x) \neq \emptyset} \int_{\partial I'_n} u_g \cdot \sigma(\psi'_E + Ex, \Sigma'_E - c_2) \nu.$$

Appealing to the trace estimate (4.22), we deduce

$$\left| \int_{\mathbb{R}^d \setminus \mathcal{I}} g : \nabla(\psi_E - \psi'_E) \right| \lesssim \left(\int_{B_{\ell+3}(x)} (|g|^2 + |\nabla u_g|^2 + |P_g - c_1|^2 \mathbb{1}_{\mathbb{R}^d \setminus \mathcal{I}}) \right)^{1/2} \times \left(\int_{B_{\ell+3}(x)} (1 + |\nabla \psi'_E|^2 + |\Sigma'_E - c_2|^2 \mathbb{1}_{\mathbb{R}^d \setminus \mathcal{I}'}) \right)^{1/2}.$$

Choosing $c_1 := \int_{B_{\ell+3}(x) \setminus \mathcal{I}} P_g$ and $c_2 := \int_{B_{\ell+3}(x) \setminus \mathcal{I}'} \Sigma'_E$, and appealing to the pressure estimate of Lemma 3.3, we deduce

$$\left| \int_{\mathbb{R}^d \setminus \mathcal{I}} g : \nabla(\psi_E - \psi'_E) \right| \lesssim \left(\int_{B_{\ell+3}(x)} (|g|^2 + |\nabla u_g|^2) \right)^{1/2} \left(\int_{B_{\ell+3}(x)} (1 + |\nabla \psi'_E|^2) \right)^{1/2}.$$

Combined with (4.29) and with the result (4.23) of Step 3, this yields the claim (4.28).

Step 5. Sensitivity of the corrector gradient inside the inclusions: for all $g \in C_c^\infty(\mathbb{R}^d)^{d \times d}$,

$$(4.31) \quad \left| \int_{\mathcal{I}} g : \nabla \psi_E - \int_{\mathcal{I}'} g : \nabla \psi'_E \right| \lesssim \left| \int_{\mathbb{R}^d \setminus \mathcal{I}} h_g : \nabla \psi_E - \int_{\mathbb{R}^d \setminus \mathcal{I}'} h_g : \nabla \psi'_E \right| + \left(\int_{B_{\ell+3}(x)} (|g|^2 + |h_g|^2) \right)^{1/2} \left(\int_{B_{\ell+3}(x)} (1 + |\nabla \psi_E|^2) \right)^{1/2}.$$

First decompose

$$\left| \int_{\mathcal{I}} g : \nabla \psi_E - \int_{\mathcal{I}'} g : \nabla \psi'_E \right| \leq \left| \sum_{n: I_n \cap B_\ell(x) = \emptyset} \int_{I_n} g : \nabla(\psi_E - \psi'_E) \right| + \sum_{n: I_n \cap B_\ell(x) \neq \emptyset} \left| \int_{I_n} g : \nabla \psi_E \right| + \sum_{n: I'_n \cap B_\ell(x) \neq \emptyset} \left| \int_{I'_n} g : \nabla \psi'_E \right|.$$

Since ψ_E and ψ'_E are both affine inside inclusions I_n 's with $I_n \cap B_\ell(x) = \emptyset$, we can rewrite

$$\left| \int_{\mathcal{I}} g : \nabla \psi_E - \int_{\mathcal{I}'} g : \nabla \psi'_E \right| \lesssim \left| \sum_n \left(\int_{I_n} g \right) : \int_{I_n} \nabla(\psi_E - \psi'_E) \right| + \left(\int_{B_{\ell+2}(x)} |g|^2 \right)^{1/2} \left(\int_{B_{\ell+2}(x)} (|\nabla \psi_E|^2 + |\nabla \psi'_E|^2) \right)^{1/2},$$

and it remains to analyze the first right-hand side term. In terms of the 2-tensor field h_g defined in (4.21), we can write by means of Stokes' theorem,

$$\begin{aligned} \sum_n \left(\int_{I_n} g \right) : \int_{I_n} \nabla(\psi_E - \psi'_E) &= \sum_n \left(\int_{I_n} g \right) : \int_{\partial I_n} (\psi_E - \psi'_E) \otimes \nu \\ &= \sum_n \int_{\partial I_n} h_g : (\psi_E - \psi'_E) \otimes \nu \\ &= - \int_{\mathbb{R}^d \setminus \mathcal{J}} \partial_i (h_g : (\psi_E - \psi'_E) \otimes e_i) \\ &= - \int_{\mathbb{R}^d \setminus \mathcal{J}} h_g : \nabla(\psi_E - \psi'_E), \end{aligned}$$

where in the last identity we used that $\operatorname{div}(h_g) = 0$. Combining with the above, and using the result (4.23) of Step 3, the claim (4.31) follows.

Step 6. Sensitivity of the corrector pressure: for all $g \in C_c^\infty(\mathbb{R}^d)$,

$$(4.32) \quad \left| \int_{\mathbb{R}^d \setminus \mathcal{J}} g \Sigma_E - \int_{\mathbb{R}^d \setminus \mathcal{J}'} g \Sigma'_E \right| \lesssim \left| \int_{\mathbb{R}^d \setminus \mathcal{J}} \nabla s_g : \nabla \psi_E - \int_{\mathbb{R}^d \setminus \mathcal{J}'} \nabla s_g : \nabla \psi'_E \right| \\ + \left(\int_{B_{\ell+3}(x)} (|g|^2 + |\nabla s_g|^2) \right)^{1/2} \left(\int_{B_{\ell+3}(x)} (1 + |\nabla \psi_E|^2 + |\Sigma_E|^2 \mathbb{1}_{\mathbb{R}^d \setminus \mathcal{J}}) \right)^{1/2}.$$

In terms of the vector field s_g defined in (4.20), we can write

$$\begin{aligned} \int_{\mathbb{R}^d \setminus \mathcal{J}} g \Sigma_E - \int_{\mathbb{R}^d \setminus \mathcal{J}'} g \Sigma'_E &= \int_{\mathbb{R}^d \setminus \mathcal{J}} g (\Sigma_E \mathbb{1}_{\mathbb{R}^d \setminus \mathcal{J}} - \Sigma'_E \mathbb{1}_{\mathbb{R}^d \setminus \mathcal{J}'}) - \int_{\mathcal{J} \setminus \mathcal{J}'} g \Sigma'_E \\ &= \int_{\mathbb{R}^d} \operatorname{div}(s_g) (\Sigma_E \mathbb{1}_{\mathbb{R}^d \setminus \mathcal{J}} - \Sigma'_E \mathbb{1}_{\mathbb{R}^d \setminus \mathcal{J}'}) - \int_{\mathcal{J} \setminus \mathcal{J}'} g \Sigma'_E, \end{aligned}$$

and thus, using the equation (4.25) for $\psi_E - \psi'_E$, the boundary conditions, and the fact that s_g is constant on the inclusion I_n

$$(4.33) \quad \int_{\mathbb{R}^d \setminus \mathcal{J}} g \Sigma_E - \int_{\mathbb{R}^d \setminus \mathcal{J}'} g \Sigma'_E = \int_{\mathbb{R}^d} \nabla s_g : \nabla(\psi_E - \psi'_E) \\ - \sum_{n: I_n \cap B_\ell(x) \neq \emptyset} \int_{\partial I_n'} s_g \cdot \sigma(\psi'_E + Ex, \Sigma'_E) \nu - \int_{\mathcal{J} \setminus \mathcal{J}'} g \Sigma'_E.$$

As $s_g|_{I_n}$ is constant for all n , $\nabla s_g = 0$ in \mathcal{J} , and since $\mathcal{J} \setminus \mathcal{J}' \subset B_\ell(x)$ the first right-hand side term satisfies

$$\begin{aligned} \left| \int_{\mathbb{R}^d} \nabla s_g : \nabla(\psi_E - \psi'_E) - \left(\int_{\mathbb{R}^d \setminus \mathcal{J}} \nabla s_g : \nabla \psi_E - \int_{\mathbb{R}^d \setminus \mathcal{J}'} \nabla s_g : \nabla \psi'_E \right) \right| \\ \leq \left(\int_{B_\ell(x)} |\nabla s_g|^2 \right)^{1/2} \left(\int_{B_\ell(x)} |\nabla \psi'_E|^2 \right)^{1/2}. \end{aligned}$$

Combining this with (4.33), appealing to the trace estimate (4.22), and using (4.23)–(4.24) in Step 3, the claim (4.32) follows.

Step 7. *Sensitivity of the extended flux: for all $g \in C_c^\infty(\mathbb{R}^d)_{\text{sym}}^{d \times d}$,*

$$(4.34) \quad \left| \int_{\mathbb{R}^d} g : (J_E - J'_E) \right| \lesssim \left| \int_{\mathbb{R}^d \setminus \mathcal{J}} g : \nabla \psi_E - \int_{\mathbb{R}^d \setminus \mathcal{J}'} g : \nabla \psi'_E \right| + \left| \int_{\mathbb{R}^d \setminus \mathcal{J}} \text{tr}(g) \Sigma_E - \int_{\mathbb{R}^d \setminus \mathcal{J}'} \text{tr}(g) \Sigma'_E \right| + \left(\int_{B_{\ell+3}(x)} |g|^2 \right)^{1/2} \left(\int_{B_{\ell+3}(x)} (1 + |\nabla \psi_E|^2 + |\Sigma_E|^2 \mathbb{1}_{\mathbb{R}^d \setminus \mathcal{J}}) \right)^{1/2}.$$

The definition (4.10) of J_E yields

$$\int_{\mathbb{R}^d} g : (J_E - J'_E) = 2 \left(\int_{\mathbb{R}^d \setminus \mathcal{J}} g : (\nabla \psi_E + E) - \int_{\mathbb{R}^d \setminus \mathcal{J}'} g : (\nabla \psi'_E + E) \right) - \left(\int_{\mathbb{R}^d \setminus \mathcal{J}} \text{tr}(g) \Sigma_E - \int_{\mathbb{R}^d \setminus \mathcal{J}'} \text{tr}(g) \Sigma'_E \right) + \sum_{n: I_n \cap B_\ell(x) \neq \emptyset} \int_{I_n} g : \sigma(\psi_E^n, \Sigma_E^n) - \sum_{n: I'_n \cap B_\ell(x) \neq \emptyset} \int_{I'_n} g : \sigma(\psi'_E^n, \Sigma'_E^n),$$

and the claim (4.34) then follows by using (4.5) to estimate the last two right-hand side terms.

Step 8. *Sensitivity of the flux corrector: for all $g \in C_c^\infty(\mathbb{R}^d)^d$,*

$$(4.35) \quad \left| \int_{\mathbb{R}^d} g \cdot \nabla (\zeta_{E;ijk} - \zeta'_{E;ijk}) \right| \lesssim \left| \int_{\mathbb{R}^d} \nabla v_g \otimes (J_E - J'_E) \right|.$$

In terms of the auxiliary field ∇v_g defined in (4.18), we can write

$$\int_{\mathbb{R}^d} g \cdot \nabla \zeta_{E;ijk} - \int_{\mathbb{R}^d} g \cdot \nabla \zeta'_{E;ijk} = - \int_{\mathbb{R}^d} \nabla v_g \cdot \nabla \zeta_{E;ijk} + \int_{\mathbb{R}^d} \nabla v_g \cdot \nabla \zeta'_{E;ijk},$$

which, in view of the equation (4.1) for ζ_E , takes the form

$$\int_{\mathbb{R}^d} g \cdot \nabla \zeta_{E;ijk} - \int_{\mathbb{R}^d} g \cdot \nabla \zeta'_{E;ijk} = \int_{\mathbb{R}^d} \partial_j v_g (J_{E;ik} - J'_{E;ik}) - \int_{\mathbb{R}^d} \partial_k v_g (J_{E;ij} - J'_{E;ij}),$$

and the claim (4.35) follows.

Step 9. *Conclusion.* — Iteratively combining the results (4.28), (4.31), (4.32), (4.34), and (4.35) of Steps 4–8, we obtain for all $g \in C_c^\infty(\mathbb{R}^d)$,

$$(4.36) \quad \left| \partial_{\mathcal{J}, B_\ell(x)}^{\text{osc}} \int_{\mathbb{R}^d} g (\nabla \psi_E, \Sigma_E \mathbb{1}_{\mathbb{R}^d \setminus \mathcal{J}}, \nabla \zeta_E) \right| \lesssim \ell^d M_\ell(x) \left(\int_{B_{\ell+3}(x)} (|A[g]|^2 + |\nabla U[A[g]]|^2) \right)^{1/2},$$

where we have set for abbreviation

$$M_\ell(x) := \left(\int_{B_{\ell+3}(x)} (1 + |\nabla \psi_E|^2 + |\Sigma_E|^2 \mathbb{1}_{\mathbb{R}^d \setminus \mathcal{J}}) \right)^{1/2},$$

$$A[g] := (g, H[g], \nabla S[g], \nabla V[g], \nabla S[\nabla V[g]]),$$

in terms of the following linear operators

$$\nabla U[g] := \nabla u_g, \quad \nabla V[g] := \nabla v_g, \quad \nabla S[g] := \nabla s_g, \quad H[g] := h_g,$$

as defined in Step 1. We commit a slight abuse of notation here as we consider a scalar test function g : the above is understood more precisely as $\nabla U[g] := (\nabla u_{g e_i \otimes e_j})_{1 \leq i, j \leq d}$, and similarly for $\nabla V[g]$, $\nabla S[g]$, and $H[g]$. Inserting (4.36) into (4.16), we find for all $q < \infty$,

$$(4.37) \quad \left\| \int_{\mathbb{R}^d} g (\nabla \psi_E, \Sigma_E \mathbb{1}_{\mathbb{R}^d \setminus \mathcal{J}}, \nabla \zeta_E) \right\|_{L^{2q}(\Omega)}^2 \lesssim_q \mathbb{E} \left[\int_0^\infty \left(\int_{\mathbb{R}^d} M_\ell(x)^2 \left(\int_{B_{\ell+3}(x)} (|A[g]|^2 + |\nabla U[A[g]]|^2) dx \right) \langle \ell \rangle^{dq} \pi(\ell) d\ell \right)^{1/q} \right].$$

Before we estimate the right-hand side of (4.37), we smuggle in a spatial average at some arbitrary scale $R \geq 1$: setting $|f|^2 := |A[g]|^2 + |\nabla U[A[g]]|^2$ for shortness,

$$\int_{\mathbb{R}^d} M_\ell(x)^2 \left(\int_{B_{\ell+3}(x)} |f|^2 \right) dx \lesssim \int_{\mathbb{R}^d} \left(\sup_{B_R(y)} M_\ell^2 \right) \left(\int_{B_{\ell+3}(y)} \left(\int_{B_R(x)} |f|^2 \right) dx \right) dy.$$

We then use a duality argument to compute the $L^q(\Omega)$ norm of this expression,

$$\mathbb{E} \left[\left(\int_{\mathbb{R}^d} M_\ell(x)^2 \left(\int_{B_{\ell+3}(x)} |f|^2 \right) dx \right)^{q\gamma} \right]^{1/q} \lesssim \sup_{\|X\|_{L^{2q'}(\Omega)}=1} \mathbb{E} \left[\int_{\mathbb{R}^d} \left(\sup_{B_R(y)} M_\ell^2 \right) \left(\int_{B_{\ell+3}(y)} \left(\int_{B_R(x)} |Xf|^2 \right) dx \right) dy \right],$$

where the supremum runs over random variables X independent of the space variable. By Hölder’s inequality and by stationarity of M_ℓ , we find

$$\mathbb{E} \left[\left(\int_{\mathbb{R}^d} M_\ell(x)^2 \left(\int_{B_{\ell+3}(x)} |f|^2 \right) dx \right)^{q\gamma} \right]^{1/q} \lesssim \left\| \sup_{B_R} M_\ell \right\|_{L^{2q}(\Omega)}^2 \sup_{\|X\|_{L^{2q'}(\Omega)}=1} \int_{\mathbb{R}^d} \mathbb{E} \left[\left(\int_{B_{\ell+3}(y)} \left(\int_{B_R(x)} |Xf|^2 \right) dx \right)^{q'} \right]^{1/q'} dy,$$

which, by Jensen’s inequality, yields

$$(4.38) \quad \mathbb{E} \left[\left(\int_{\mathbb{R}^d} M_\ell(x)^2 \left(\int_{B_{\ell+3}(x)} |f|^2 \right) dx \right)^{q\gamma} \right]^{1/q} \lesssim \left\| \sup_{B_R} M_\ell \right\|_{L^{2q}(\Omega)}^2 \sup_{\|X\|_{L^{2q'}(\Omega)}=1} \| [Xf]_2 \|_{L^2(\mathbb{R}^d; L^{2q'}(\Omega))}^2.$$

Appealing to the annealed estimate in (4.17), we find for $q \gg 1$ (hence $|2q' - 2| \ll 1$),

$$\begin{aligned} \| [X \nabla U[A[g]]]_2 \|_{L^2(\mathbb{R}^d; L^{2q'}(\Omega))} &= \| [\nabla U[A[Xg]]]_2 \|_{L^2(\mathbb{R}^d; L^{2q'}(\Omega))} \\ &\lesssim \| [A[Xg]]_2 \|_{L^2(\mathbb{R}^d; L^{2q'}(\Omega))}, \end{aligned}$$

while the annealed estimates in (4.19), (4.20), and (4.21) yield for $q > 1$,

$$\| [XA[g]]_2 \|_{L^2(\mathbb{R}^d; L^{2q'}(\Omega))} \lesssim \| [Xg]_2 \|_{L^2(\mathbb{R}^d; L^{2q'}(\Omega))} = \| X \|_{L^{2q'}(\Omega)} \| g \|_{L^2(\mathbb{R}^d)}.$$

Using these bounds in combination with (4.37) and (4.38), together with the supralgebraic decay of the weight π in form of Jensen’s inequality, cf. Assumption (Mix⁺), we obtain for all q with $1 \ll q < \infty$,

$$\left\| \int_{\mathbb{R}^d} g(\nabla\psi_E, \Sigma_E \mathbb{1}_{\mathbb{R}^d \setminus \mathcal{J}}, \nabla\zeta_E) \right\|_{L^{2q}(\Omega)}^2 \lesssim_q \sup_{\ell \geq 0} \left\| \sup_{B_R} M_\ell \right\|_{L^{2q}(\Omega)}^2 \| g \|_{L^2(\mathbb{R}^d)}^2.$$

Finally, by stationarity and by the discrete $\ell^{2s} - \ell^\infty$ inequality, the supremum of M_ℓ can be estimated as follows, for all $s \geq 1$,

$$\sup_{\ell \geq 0} \left\| \sup_{B_R} M_\ell \right\|_{L^{2q}(\Omega)} \lesssim \left\| \left(1 + \int_{B_R} [(\nabla\psi_E, \Sigma_E \mathbb{1}_{\mathbb{R}^d \setminus \mathcal{J}})]_2^{2s} \right)^{1/2s} \right\|_{L^{2q}(\Omega)},$$

and the conclusion (4.12) follows. □

4.4. PROOF OF PROPOSITION 4.5. — Let $E \in \mathbb{M}_0$ be fixed with $|E| = 1$. We split the proof into three steps.

Step 1. Meyers-type perturbative argument: for all $s \geq 1$ with $|s - 1| \ll 1$, for all $R, K \geq 1$ and $c_R \in \mathbb{R}^d$,

$$(4.39) \quad \left(\int_{B_R} [\nabla\psi_E]_2^{2s} \right)^{1/s} \lesssim K^2 \left(1 + \frac{1}{R^2} \int_{B_{cR}} |\psi_E - c_R|^2 \right) + \frac{1}{K^2} \int_{B_{cR}} |\nabla\psi_E|^2.$$

Arguing as in (3.10), with u_g replaced by $\psi_E + Ex$ and with $g = 0$, we obtain the following Caccioppoli-type inequality: for all balls $D \subset \mathbb{R}^d$ with radius $r_D \geq 3$, for all $K \geq 1$ and $c_D \in \mathbb{R}^d$,

$$(4.40) \quad \int_D |\nabla\psi_E|^2 \lesssim K^2 \left(1 + \frac{1}{r_D^2} \int_{2D} |\psi_E - c_D|^2 \right) + \frac{1}{K^2} \int_{2D} |\nabla\psi_E|^2.$$

Using the Poincaré-Sobolev inequality to estimate the first right-hand side term, with the choice $c_D := \int_{2D} \psi_E$, we deduce

$$\left(\int_D |\nabla\psi_E|^2 \right)^{1/2} \lesssim K \left(1 + \int_{2D} |\nabla\psi_E|^{2d/(d+2)} \right)^{(d+2)/2d} + \frac{1}{K} \left(\int_{2D} |\nabla\psi_E|^2 \right)^{1/2}.$$

While this is proved for all balls D with radius $r_D \geq 3$, smuggling in local quadratic averages at scale 1 allows to infer that for all balls D (with any radius $r_D > 0$) and $K \geq 1$,

$$\left(\int_D [\nabla\psi_E]_2^2 \right)^{1/2} \lesssim K \left(1 + \int_{2D} [\nabla\psi_E]_2^{2d/(d+2)} \right)^{(d+2)/2d} + \frac{1}{K} \left(\int_{2D} [\nabla\psi_E]_2^2 \right)^{1/2}.$$

Choosing K large enough and applying Gehring’s lemma in form of Lemma 3.5, we deduce the following Meyers-type estimate: for all $s \geq 1$ with $|s - 1| \ll 1$, and all $R > 0$,

$$\left(\int_{B_R} [\nabla\psi_E]_2^{2s} \right)^{1/s} \lesssim 1 + \int_{B_{cR}} [\nabla\psi_E]_2^2.$$

Combining this with (4.40), the claim (4.39) follows.

Step 2. Conclusion on $\nabla\psi_E$: for all r, R with $1 \leq r \ll_{\chi} R$ and $q, s \geq 1$ with $|s - 1| \ll 1$,

$$\left\| \left(\int_{B_R} [\nabla\psi_E]_2^{2s} \right)^{1/2s} \right\|_{L^{2q}(\Omega)} \lesssim_{\chi} 1 + \left\| \int_{\mathbb{R}^d} \chi_r \nabla\psi_E \right\|_{L^{2q}(\Omega)}.$$

For $1 \leq r \leq R$, choosing $c_R := \int_{B_{CR}} \chi_r * \psi_E$, Poincaré's inequality yields

$$\begin{aligned} \int_{B_{CR}} |\psi_E - c_R|^2 &\lesssim \int_{B_{CR}} |\psi_E - \chi_r * \psi_E|^2 + \int_{B_{CR}} |\chi_r * \psi_E - c_R|^2 \\ &\lesssim_{\chi} r^2 \int_{B_{CR}} |\nabla\psi_E|^2 + R^2 \int_{B_{CR}} |\chi_r * \nabla\psi_E|^2. \end{aligned}$$

Inserting this into (4.39), we find

$$\left(\int_{B_R} [\nabla\psi_E]_2^{2s} \right)^{1/s} \lesssim K^2 + \left(K^2 \frac{r^2}{R^2} + \frac{1}{K^2} \right) \int_{B_{CR}} |\nabla\psi_E|^2 + K^2 \int_{B_{CR}} |\chi_r * \nabla\psi_E|^2.$$

Taking the $L^q(\Omega)$ norm, and using that stationarity and Jensen's inequality yield

$$\left\| \int_{B_{CR}} |\nabla\psi_E|^2 \right\|_{L^q(\Omega)} \lesssim \left\| \int_{B_R} [\nabla\psi_E]_2^2 \right\|_{L^q(\Omega)} \lesssim \left\| \left(\int_{B_R} [\nabla\psi_E]_2^{2s} \right)^{1/2s} \right\|_{L^{2q}(\Omega)}^2,$$

and

$$\left\| \int_{B_{CR}} |\chi_r * \nabla\psi_E|^2 \right\|_{L^q(\Omega)} \leq \|\chi_r * \nabla\psi_E\|_{L^{2q}(\Omega)}^2,$$

we deduce

$$\begin{aligned} &\left\| \left(\int_{B_R} [\nabla\psi_E]_2^{2s} \right)^{1/2s} \right\|_{L^{2q}(\Omega)} \\ &\lesssim_{\chi} K + \left(K \frac{r}{R} + \frac{1}{K} \right) \left\| \left(\int_{B_R} [\nabla\psi_E]_2^{2s} \right)^{1/2s} \right\|_{L^{2q}(\Omega)} + K \left\| \int_{\mathbb{R}^d} \chi_r \nabla\psi_E \right\|_{L^{2q}(\Omega)}. \end{aligned}$$

Choosing $K \gg 1$ and $R \gg_{K,\chi} r$, the second right-hand side term can be absorbed into the left-hand side and the claim follows.

Step 3. Conclusion on the pressure Σ_E . — For all $R, s \geq 1$, we decompose

$$\left(\int_{B_R} [\Sigma_E \mathbb{1}_{\mathbb{R}^d \setminus \mathcal{J}}]_2^{2s} \right)^{1/s} \lesssim \left(\int_{B_R} \left[\left(\Sigma_E - \int_{B_R \setminus \mathcal{J}} \Sigma_E \right) \mathbb{1}_{\mathbb{R}^d \setminus \mathcal{J}} \right]_2^{2s} \right)^{1/s} + \left| \int_{B_R \setminus \mathcal{J}} \Sigma_E \right|^2.$$

Appealing to the pressure estimate of Lemma 3.3 to estimate the first right-hand side term, and further decomposing the second term, we obtain for all r, R with $1 \leq r \leq R$, assuming that $\int_{\mathbb{R}^d \setminus \mathcal{J}} \chi_r \simeq \int_{\mathbb{R}^d} \chi_r = 1$ (which holds automatically provided $r \gg_{\chi} 1$ in view of the hardcore assumption, cf. (H $_{\delta}$)),

$$\begin{aligned} \left(\int_{B_R} [\Sigma_E \mathbb{1}_{\mathbb{R}^d \setminus \mathcal{J}}]_2^{2s} \right)^{1/s} &\lesssim 1 + \left(\int_{B_R} [\nabla\psi_E]_2^{2s} \right)^{1/s} + \left| \int_{\mathbb{R}^d} \chi_r \Sigma_E \mathbb{1}_{\mathbb{R}^d \setminus \mathcal{J}} \right|^2 \\ &\quad + \left| \int_{\mathbb{R}^d} \chi_r \left(\Sigma_E - \int_{B_R \setminus \mathcal{J}} \Sigma_E \right) \mathbb{1}_{\mathbb{R}^d \setminus \mathcal{J}} \right|^2. \end{aligned}$$

It remains to estimate the last right-hand side term. By the Cauchy–Schwarz inequality, for $r \ll_\chi R$ such that χ_r is supported in B_R , using again the pressure estimate of Lemma 3.3, we find

$$\begin{aligned} \left| \int_{\mathbb{R}^d} \chi_r \left(\Sigma_E - \fint_{B_R \setminus \mathcal{J}} \Sigma_E \right) \mathbb{1}_{\mathbb{R}^d \setminus \mathcal{J}} \right|^2 &\lesssim \left(R^d \int_{\mathbb{R}^d} |\chi_r|^2 \right) \fint_{B_R} \left| \Sigma_E - \fint_{B_R \setminus \mathcal{J}} \Sigma_E \right|^2 \mathbb{1}_{\mathbb{R}^d \setminus \mathcal{J}} \\ &\lesssim \left(R^d \int_{\mathbb{R}^d} |\chi_r|^2 \right) \left(1 + \fint_{B_R} |\nabla \psi_E|^2 \right). \end{aligned}$$

Since we have $R^d \int_{\mathbb{R}^d} |\chi_r|^2 \lesssim \|\chi\|_{L^2(\mathbb{R}^d)}^2$ provided $r/R \gtrsim 1$, we conclude

$$\left(\fint_{B_R} [\Sigma_E \mathbb{1}_{\mathbb{R}^d \setminus \mathcal{J}}]_2^{2s} \right)^{1/s} \lesssim_\chi 1 + \left(\fint_{B_R} [\nabla \psi_E]_2^{2s} \right)^{1/s} + \left| \int_{\mathbb{R}^d} \chi_r \Sigma_E \mathbb{1}_{\mathbb{R}^d \setminus \mathcal{J}} \right|^2.$$

Combined with the results on $\nabla \psi_E$ in Step 2, the conclusion follows. □

5. LARGE-SCALE REGULARITY

This section is devoted to the development of a large-scale regularity theory for the steady Stokes problem (2.4), and to the proof of Theorems 3, 4, and 5. We take inspiration from the theory recently developed in the model setting of divergence-form linear elliptic equations with random coefficients [6, 5, 1, 2, 3, 28, 20, 37], and we focus more precisely on the formulation in [28, 20].

5.1. STRUCTURE OF THE ARGUMENT. — Recall that for harmonic functions, regularity of the gradient can be proved by controlling the decay of the excess across scales, where the excess is defined by the local L^2 -distance of the gradient to a constant. In the heterogeneous setting of divergence-form operators $-\nabla \cdot a \nabla$, cf. [28], we rather define the excess by the local L^2 -distance of the gradient to the gradient of a -harmonic coordinates (that is, to a constant plus the associated corrector gradient). The key ingredient to large-scale regularity theory is then encapsulated in a perturbative estimate of excess decay, measured in terms of the growth of an extended corrector, cf. [28, Prop. 1]. The following proposition is the extension of such a result in the context of the steady Stokes problem (2.4); the proof is postponed to Section 5.2. Henceforth, we use the short-hand notation $\psi := (\psi_E)_{E \in \mathcal{E}}$, where \mathcal{E} stands for an orthonormal basis of $\mathbb{M}_0^{\text{sym}}$, and similarly for Σ, ζ .

PROPOSITION 5.1 (Perturbative excess decay). — *There exists an exponent $\varepsilon \simeq 1$ such that the following holds: For all $R \gg 1$, if ∇u is a solution of the following free steady Stokes problem in B_R ,*

$$(5.1) \quad \begin{cases} -\Delta u + \nabla P = 0, & \text{in } B_R \setminus \mathcal{J}, \\ \operatorname{div}(u) = 0, & \text{in } B_R, \\ D(u) = 0, & \text{in } \mathcal{J} \cap B_R, \\ \int_{\partial I_n} \sigma(u, P) \nu = 0, & \forall n : I_n \subset B_R, \\ \int_{\partial I_n} \Theta(x - x_n) \cdot \sigma(u, P) \nu = 0, & \forall n : I_n \subset B_R, \forall \Theta \in \mathbb{M}^{\text{skew}}, \end{cases}$$

then there exists a matrix $E_0 \in \mathbb{M}_0$ such that for all r, R with $4 \leq r \leq R$,

$$(5.2) \quad \int_{B_r} |\nabla u - (\nabla \psi_{E_0} + E_0)|^2 \lesssim \left(\left(\frac{r}{R} \right)^2 + \left(\frac{R}{r} \right)^{d+2} (1 \wedge \gamma_R)^{2\varepsilon} \right) \int_{B_R} |\nabla u|^2,$$

where we have set for abbreviation,

$$(5.3) \quad \gamma_R := \sup_{L \geq R} \frac{1}{L} \left(1 + \int_{B_L} |(\psi, \zeta) - \int_{B_L} (\psi, \zeta)|^2 \right)^{1/2}.$$

Moreover, the following non-degeneracy property holds for all $E \in \mathbb{M}_0$,

$$(5.4) \quad (1 - C\gamma_R)|E| \lesssim \left(\int_{B_{R/2}} |\nabla \psi_E + E|^2 \right)^{1/2} \lesssim (1 + \gamma_R)|E|.$$

Although the proof of Proposition 5.1 follows the main steps as the proof of [28, Prop. 1], it differs in two significant respects. First, the natural two-scale expansion is not rigid inside the inclusions, which makes energy estimates more involved and requires some local surgery. Second, a suitable control is needed on the pressure of the two-scale expansion error, which is made particularly subtle due to the crucial use of weighted norms. Weighted pressure estimates are obtained based on the following weighted version of Bogovskii’s standard construction. This statement is a particular case of [13, Th. 5.2], which holds more generally in any John domain.

LEMMA 5.2 (Weighted Bogovskii construction; [13]). — *Let $D \subset B_R$ be a domain that is star-shaped with respect to every point in B_{R_0} , for some $0 < R_0 \leq R$. Consider a weight $\mu \in C^\infty(\mathbb{R}^d; [0, 1])$ that belongs to the Muckenhoupt class A_2 . Then, for all $F \in L^2(D)$ with $\int_D F = 0$, there exists $S \in H_0^1(D)^d$ such that*

$$\begin{aligned} \operatorname{div}(S) &= F, \quad \text{in } D, \\ \int_D \mu |\nabla S|^2 &\lesssim \int_D \mu |F|^2, \end{aligned}$$

where the multiplicative constant only depends on d , on R/R_0 , on the A_2 -norm of μ .

With Proposition 5.1 at hand, we may now turn to the proof of Theorems 3–5, for which we heavily lean on [28, 20]. First, following [28], we encapsulate a quantitative (averaged) control on the sublinear growth of the extended corrector by considering the minimal radius R such that γ_R in (5.3) is small enough: more precisely, given a constant $C_0 \geq 1$ (to be fixed large enough), we define the *minimal radius* r_* as the following random field,

$$(5.5) \quad r_*(x) := \inf \left\{ R > 0 : \frac{1}{\ell^2} \int_{B_\ell(x)} |(\psi, \zeta) - \int_{B_\ell(x)} (\psi, \zeta)|^2 \leq \frac{1}{C_0}, \forall \ell \geq R \right\}.$$

Stationarity of r_* follows from stationarity of $(\nabla \psi, \nabla \zeta)$. Almost sure finiteness of r_* follows from the sublinearity of (ψ, ζ) at infinity, cf. Lemmas 1 and 4.1(iii). Under Assumption (Mix⁺), moment bounds on r_* are a direct consequence of corrector estimates of Theorem 2 together with a union bound; we omit the details.

Next, still following [28], we consider the excess (2.7) of a trace-free 2-tensor field h on a ball D , that is,

$$\text{Exc}(h; D) := \inf_{E \in \mathbb{M}_0} \int_D |h - (\nabla\psi_E + E)|^2,$$

which measures the deviation of h from gradients of corrected coordinates. In these terms, we establish the following consequence of Proposition 5.1, which quantifies the decay of the excess for solutions of the free steady Stokes problem (5.1) from larger to smaller balls. The proof relies on Proposition 5.1 together with a standard Campanato iteration; in particular, since it is oblivious of the underlying PDE, we refer the reader to the proof of [28, Th. 1] in the context of divergence-form linear elliptic equations, which applies without changing a iota.

THEOREM 5.3 (Excess-decay estimate). — *Under Assumption (\mathbf{H}_δ) , for any Hölder exponent $\alpha \in (0, 1)$, there exists a constant $C_\alpha \simeq_\alpha 1$ such that the following holds: Let r_* be defined in (5.5) with constant C_0 replaced by C_α . For all $R \geq r_*(0)$, if ∇u is a solution of the free steady Stokes problem (5.1) in B_R , then the following large-scale Lipschitz estimate holds for all $r_*(0) \leq r \leq R$,*

$$(5.6) \quad \int_{B_r} |\nabla u|^2 \leq C_\alpha \int_{B_R} |\nabla u|^2,$$

as well as the following large-scale $C^{1,\alpha}$ estimate for all r with $r_*(0) \leq r \leq R$,

$$\text{Exc}(\nabla u; B_r) \leq C_\alpha (r/R)^{2\alpha} \text{Exc}(\nabla u; B_R).$$

In addition, the correctors enjoy the following non-degeneracy property for all $r \geq r_*(0)$ and $E \in \mathbb{M}_0$,

$$\frac{1}{C_\alpha} |E|^2 \leq \int_{B_r} |\nabla\psi_E + E|^2 \leq C_\alpha |E|^2.$$

As a direct consequence, we may deduce a corresponding result for solutions of the steady Stokes problem (5.1) with a nontrivial right-hand side, cf. (2.4), as stated in Theorem 3. The proof, which is identical to that of [28, Cor. 3], is omitted as it only relies on Theorem 5.3 together with an energy estimate.

Next, as a second consequence of the above, we may further deduce quenched large-scale L^p regularity estimates as stated in Theorem 4. This can be obtained by combining the large-scale Lipschitz estimate (5.6) together with Shen’s dual Calderón–Zygmund lemma, cf. [41, Th. 3.2] (see also [42, Th. 2.4]), as done in [20, §6.1] in the context of divergence-form linear elliptic equations: since this argument does not rely on the specific PDE at hand, the same applies without changing a iota and we do not reproduce it here. For estimates with Muckenhoupt weights, it suffices to appeal to [41, Th. 3.4 & Rem. 3.5] instead of [41, Th. 3.2]. Note that this approach requires to replace the minimal radius r_* in the above by the largest $\frac{1}{8}$ -Lipschitz lower bound $r_{*,\psi}$, cf. [28, §3.7]: both satisfy the same boundedness properties and we use the same notation “ r_* ” in the statement.

Finally, making a further use of Shen's dual Calderón–Zygmund lemma, cf. [41, Th. 3.2 or 3.4], together with the quenched large-scale L^p regularity theory of Theorem 4 and with the large-scale Lipschitz estimate (5.6), the annealed regularity estimate of Theorem 5 easily follows as in [20] for $2 \leq q \leq p < \infty$. A duality argument yields the corresponding conclusion for $1 < p \leq q \leq 2$, and an interpolation argument allows to conclude for all p, q with $1 < p, q < \infty$. The additional perturbative statement in Theorem 5 is already established in Theorem 3.1.

5.2. PROOF OF PROPOSITION 5.1. — Let $R \gg 1$ be large enough and fixed. As the statement of Proposition 5.1 does not depend on the choice of anchoring of the correctors, we can assume without loss of generality $f_{B_R}(\psi, \zeta, \Sigma \mathbb{1}_{\mathbb{R}^d \setminus \mathcal{J}}) = 0$. Set $\mathcal{N}_R := \{n : I_n^+ + \delta B \subset B_R\}$ and $\mathcal{N}_R^\circ := \{n : (I_n^+ + \delta B) \cap \partial B_R \neq \emptyset\}$, where we recall that I_n^+ stands for the convex hull of I_n , and define

$$D_R := \left(B_{R-\delta/2} \setminus \bigcup_{n \in \mathcal{N}_R^\circ} (I_n^+ + \delta B) \right) + \frac{\delta}{2} B.$$

In view of Assumption (H_δ) , we note that

- D_R is a C^2 domain (uniformly in R);
- any inclusion that intersects D_R is contained in D_R and is at distance at least δ from ∂D_R ;
- $B_{R-2-\delta} \subset D_R \subset B_R$.

Given $4 \leq \rho \leq R/4$ (the choice of which will be optimized later), we choose a smooth cut-off function $\eta_R \in C_c^\infty(\mathbb{R}^d; [0, 1])$ such that $\eta_R = 1$ in $B_{R-2\rho}$, $\eta_R = 0$ outside $B_{R-\rho}$, and $|\nabla \eta_R| \lesssim \rho^{-1}$, and we further choose η_R to be constant in the fattened inclusions $\{I_n + (\delta/2)B\}_{n \in \mathcal{N}_R}$. Note in particular that η_R is supported inside D_R . We split the proof into five main steps.

Step 1. Two-scale expansion and representation of the error. — We split the proof into two further substeps.

Substep 1.1. Construction of two-scale expansions. — Given a weak solution (u, P) to (5.1), let (\hat{u}, \hat{P}) denote the unique weak solution of the following corresponding homogenized equation with Dirichlet data on D_R ,

$$(5.7) \quad \begin{cases} -\operatorname{div}(2\bar{\mathbf{B}}\mathbf{D}(\hat{u})) + \nabla \hat{P} = 0, & \text{in } D_R, \\ \operatorname{div}(\hat{u}) = 0, & \text{in } D_R, \\ \hat{u} = u, & \text{on } \partial D_R, \end{cases}$$

where we recall that the effective viscosity $\bar{\mathbf{B}}$ is defined in (2.11). For definiteness, the pressures P and \hat{P} are chosen with $\int_{D_R} P \mathbb{1}_{\mathbb{R}^d \setminus \mathcal{J}} = \int_{D_R} \hat{P} = 0$. Reformulating this homogenized equation as

$$-\operatorname{div}(2\bar{\mathbf{B}}\mathbf{D}(\hat{u} - u)) + \nabla \hat{P} = \operatorname{div}(2\bar{\mathbf{B}}\mathbf{D}(u)), \quad \text{in } D_R,$$

testing with $\widehat{u} - u \in H_0^1(D_R)^d$, and combining an energy estimate with the triangle inequality, we obtain

$$\int_{D_R} |\mathbf{D}(\widehat{u})|^2 \lesssim \int_{D_R} |\mathbf{D}(u)|^2,$$

and, further using that $\operatorname{div}(\widehat{u} - u) = 0$ implies $\int_{D_R} |\nabla(\widehat{u} - u)|^2 = 2 \int_{D_R} |\mathbf{D}(\widehat{u} - u)|^2$,

$$(5.8) \quad \int_{D_R} |\nabla \widehat{u}|^2 \lesssim \int_{D_R} |\nabla u|^2.$$

We now compare u and P to their respective two-scale expansions,

$$u \rightsquigarrow \widehat{u} + \eta_R \psi_E \partial_E \widehat{u}, \quad P \rightsquigarrow \widehat{P} + \eta_R \bar{\mathbf{b}} : \mathbf{D}(\widehat{u}) + \eta_R \Sigma_E \mathbb{1}_{\mathbb{R}^d \setminus \mathcal{I}} \partial_E \widehat{u},$$

where we use Einstein’s convention of implicit summation on repeated indices and where the index E runs here over an orthonormal basis \mathcal{E} of $\mathbb{M}_0^{\text{sym}}$. Recall that the pressure P is only defined up to a global arbitrary constant on $\mathbb{R}^d \setminus \mathcal{I}$, so that we may choose an arbitrary constant $P_* \in \mathbb{R}$ and consider the pressure $P' = P + P_*$ on $\mathbb{R}^d \setminus \mathcal{I}$. In addition we choose arbitrary constants $\{P_n\}_n \subset \mathbb{R}$ and extend the pressure inside the inclusions by setting $P'|_{I_n} = P_n$. We thus define in the whole domain D_R ,

$$(5.9) \quad P' := (P + P_*) \mathbb{1}_{\mathbb{R}^d \setminus \mathcal{I}} + \sum_{n \in \mathcal{N}_R} P_n \mathbb{1}_{I_n},$$

where the constants P_* and $\{P_n\}_n$ will be suitably chosen later. We then consider the following two-scale expansion errors in D_R ,

$$(5.10) \quad w := u - \widehat{u} - \eta_R \psi_E \partial_E \widehat{u}, \quad Q := P' - \widehat{P} - \eta_R \bar{\mathbf{b}} : \mathbf{D}(\widehat{u}) - \eta_R \Sigma_E \mathbb{1}_{\mathbb{R}^d \setminus \mathcal{I}} \partial_E \widehat{u}.$$

Substep 1.2. Proof that (w, Q) satisfies in the weak sense in D_R

$$(5.11) \quad -\Delta w + \nabla Q = - \sum_{n \in \mathcal{N}_R} \delta_{\partial I_n} \sigma(u, P + P_* - P_n) \nu - \operatorname{div}((\eta_R \partial_E \widehat{u}) J_E \mathbb{1}_{\mathcal{I}}) \\ + \operatorname{div} \left(2(1 - \eta_R)(\operatorname{Id} - \bar{\mathbf{B}}) \mathbf{D}(\widehat{u}) + (2\psi_E \otimes_s - \zeta_E) \nabla(\eta_R \partial_E \widehat{u}) - \operatorname{Id}(\psi_E \cdot \nabla)(\eta_R \partial_E \widehat{u}) \right).$$

By definition of w, Q , expanding the gradient and reorganizing the terms, we find

$$-\Delta w + \nabla Q = -\Delta u + \nabla P' + \Delta \widehat{u} - \nabla \widehat{P} - \nabla(\eta_R \bar{\mathbf{b}} : \mathbf{D}(\widehat{u})) + \operatorname{div}(\psi_E \otimes \nabla(\eta_R \partial_E \widehat{u})) \\ + (\eta_R \partial_E \widehat{u}) \operatorname{div}(\nabla \psi_E - \Sigma_E \mathbb{1}_{\mathbb{R}^d \setminus \mathcal{I}} \operatorname{Id}) + (\nabla \psi_E - \Sigma_E \mathbb{1}_{\mathbb{R}^d \setminus \mathcal{I}} \operatorname{Id}) \nabla(\eta_R \partial_E \widehat{u}).$$

Further using that $\operatorname{div}(\psi_E) = 0$, and using Leibniz’ rule, this can be rewritten as

$$-\Delta w + \nabla Q = -\Delta u + \nabla P' + \Delta \widehat{u} - \nabla \widehat{P} - \nabla(\eta_R \bar{\mathbf{b}} : \mathbf{D}(\widehat{u})) \\ + \operatorname{div}(2\psi_E \otimes_s \nabla(\eta_R \partial_E \widehat{u})) - \nabla(\psi_E \cdot \nabla(\eta_R \partial_E \widehat{u})) \\ + \operatorname{div} \left((\eta_R \partial_E \widehat{u}) (2\mathbf{D}(\psi_E) - \Sigma_E \mathbb{1}_{\mathbb{R}^d \setminus \mathcal{I}} \operatorname{Id}) \right).$$

Since $\operatorname{div}(\widehat{u}) = 0$, we may decompose

$$\Delta \widehat{u} = \operatorname{div}(2\mathbf{D}(\widehat{u})) = \operatorname{div}(2(1 - \eta_R) \mathbf{D}(\widehat{u})) + \operatorname{div}((\eta_R \partial_E \widehat{u}) 2E).$$

Inserting this into the above, and writing $2(\mathbb{D}(\psi_E) + E) - \Sigma_E \mathbb{1}_{\mathbb{R}^d \setminus \mathcal{J}} = J_E \mathbb{1}_{\mathbb{R}^d \setminus \mathcal{J}}$ in terms of the extended flux J_E , cf. Lemma 4.1, we obtain

$$(5.12) \quad -\Delta w + \nabla Q = -\Delta u + \nabla P' + \operatorname{div}(2(1 - \eta_R) \mathbb{D}(\hat{u})) - \nabla \hat{P} - \nabla(\eta_R \bar{\mathbf{b}} : \mathbb{D}(\hat{u})) \\ + \operatorname{div}(2\psi_E \otimes_s \nabla(\eta_R \partial_E \hat{u})) - \nabla(\psi_E \cdot \nabla(\eta_R \partial_E \hat{u})) + \operatorname{div}((\eta_R \partial_E \hat{u}) J_E \mathbb{1}_{\mathbb{R}^d \setminus \mathcal{J}}).$$

Since $\operatorname{div}(J_E) = 0$, we have

$$\operatorname{div}((\eta_R \partial_E \hat{u}) J_E \mathbb{1}_{\mathbb{R}^d \setminus \mathcal{J}}) = J_E \nabla(\eta_R \partial_E \hat{u}) - \operatorname{div}((\eta_E \partial_E \hat{u}) J_E \mathbb{1}_{\mathcal{J}}),$$

and thus, further recalling $\mathbb{E}[J_E] = 2\bar{\mathbf{B}}E + (\bar{\mathbf{b}} : E) \operatorname{Id}$, writing $J_E - \mathbb{E}[J_E] = \operatorname{div}(\zeta_E)$, and using the skew-symmetry of ζ_E , cf. Lemma 4.1, we find

$$\operatorname{div}((\eta_R \partial_E \hat{u}) J_E \mathbb{1}_{\mathbb{R}^d \setminus \mathcal{J}}) = \operatorname{div}(2\eta_R \bar{\mathbf{B}} \mathbb{D}(\hat{u})) + \nabla(\eta_R \bar{\mathbf{b}} : \mathbb{D}(\hat{u})) \\ - \operatorname{div}(\zeta_E \nabla(\eta_R \partial_E \hat{u})) - \operatorname{div}((\eta_R \partial_E \hat{u}) J_E \mathbb{1}_{\mathcal{J}}).$$

Inserting this into (5.12), and recalling that equation (5.7) yields

$$-\operatorname{div}(2\bar{\mathbf{B}} \mathbb{D}(\hat{u})) + \nabla \hat{P} = 0,$$

we deduce

$$-\Delta w + \nabla Q = -\Delta u + \nabla P' + \operatorname{div}(2(1 - \eta_R)(\operatorname{Id} - \bar{\mathbf{B}}) \mathbb{D}(\hat{u})) \\ + \operatorname{div}((2\psi_E \otimes_s - \zeta_E) \nabla(\eta_R \partial_E \hat{u})) - \nabla(\psi_E \cdot \nabla(\eta_R \partial_E \hat{u})) - \operatorname{div}((\eta_R \partial_E \hat{u}) J_E \mathbb{1}_{\mathcal{J}}).$$

Finally, since equation (3.1) for (u, P) implies of (u, P') on D_R

$$-\Delta u + \nabla P' = - \sum_{n \in \mathcal{N}_R} \delta_{\partial I_n} \sigma(u, P + P_* - P_n) \nu,$$

the claim (5.11) follows.

Step 2. Weighted energy estimate for the two-scale expansion error. — Considering the following weight function as in [28],

$$(5.13) \quad \mu_{R,\varepsilon} : B_R \longrightarrow [0, 1] : x \longmapsto (1 - |x|/R)^{\varepsilon/2},$$

we prove, for all $K \gg 1$ and $\varepsilon \ll K^{-1/2}$,

$$(5.14) \quad \int_{D_R} \mu_{R,\varepsilon}^2 |\nabla w|^2 \lesssim \frac{1}{K} \int_{D_R} \mu_{R,\varepsilon}^2 Q^2 + K \int_{D_R} (1 - \eta_R)^2 \mu_{R,\varepsilon}^2 |\nabla \hat{u}|^2 \\ + K \left(\sup_{D_R} |\nabla(\eta_R \nabla \hat{u})|^2 \right) \int_{D_R} (1 + |(\psi, \zeta, \nabla \psi, \Sigma \mathbb{1}_{\mathbb{R}^d \setminus \mathcal{J}})|^2).$$

The main difficulty is that neither \hat{u} nor $\mu_{R,\varepsilon}$ is constant inside the inclusions, which prohibits us from easily taking advantage of the boundary conditions for u and ψ_E in the estimate. To circumvent this issue, we use the following truncation maps T_0, T_1 :

for all $g \in C_b^\infty(D_R)$,

$$\begin{aligned}
 T_0[g](x) &:= (1 - \chi(x))g(x) + \sum_{n \in \mathcal{N}_R} \chi_n(x) \left(\int_{I_n + (\delta/2)B} g \right), \\
 (5.15) \quad T_1[g](x) &:= (1 - \chi(x))g(x) + \sum_{n \in \mathcal{N}_R} \chi_n(x) \left(\left(\int_{I_n + (\delta/2)B} g \right) \right. \\
 &\quad \left. + \left(\int_{I_n + (\delta/2)B} \nabla g \right) (x - x_n) \right),
 \end{aligned}$$

where for all n we have chosen a cut-off function $\chi_n \in C_c^\infty(\mathbb{R}^d; [0, 1])$ with

$$\chi_n|_{I_n + (\delta/4)B} = 1, \quad \chi_n|_{\mathbb{R}^d \setminus (I_n + (\delta/2)B)} = 0, \quad |\nabla \chi_n| + |\nabla^2 \chi_n| \lesssim 1,$$

and where we have set for abbreviation $\chi := \sum_{n \in \mathcal{N}_R} \chi_n$. In these terms, we consider the following modification of the weight $\mu_{R,\varepsilon}$ and of the two-scale expansion error (w, Q) ,

$$\begin{aligned}
 (5.16) \quad \tilde{\mu}_{R,\varepsilon} &:= T_0[\mu_{R,\varepsilon}], \\
 \tilde{w} &:= u - T_1[\hat{u}] - \eta_R \psi_E T_0[\partial_E \hat{u}], \\
 \tilde{Q} &:= P' - T_0[\hat{P}] - \eta_R \bar{\mathbf{b}} : T_0[\mathbf{D}(\hat{u})] - \eta_R \Sigma_E \mathbb{1}_{\mathbb{R}^d \setminus \mathcal{I}} T_0[\partial_E \hat{u}].
 \end{aligned}$$

Note that $T_1[\hat{u}] = \hat{u} = u$ on ∂D_R , and thus $\tilde{w} \in H_0^1(D_R)^d$. Testing equation (5.11) for w with the test function $\tilde{\mu}_{R,\varepsilon}^2 \tilde{w} \in H_0^1(D_R)^d$, we find

$$(5.17) \quad J_0 = J_1 + J_2 + J_3,$$

in terms of

$$\begin{aligned}
 J_0 &:= \int_{D_R} \nabla(\tilde{\mu}_{R,\varepsilon}^2 \tilde{w}) : (\nabla w - Q \text{Id}), \\
 J_1 &:= - \sum_{n \in \mathcal{N}_R} \int_{\partial I_n} \tilde{\mu}_{R,\varepsilon}^2 \tilde{w} \cdot \sigma(u, P + P_* - P_n) \nu + \sum_{n \in \mathcal{N}_R} \int_{I_n} (\eta_R \partial_E \hat{u}) \nabla(\tilde{\mu}_{R,\varepsilon}^2 \tilde{w}) : J_E, \\
 J_2 &:= -2 \int_{D_R} (1 - \eta_R) \nabla(\tilde{\mu}_{R,\varepsilon}^2 \tilde{w}) : (\text{Id} - \bar{\mathbf{B}}) \mathbf{D}(\hat{u}), \\
 J_3 &:= - \int_{D_R} \nabla(\tilde{\mu}_{R,\varepsilon}^2 \tilde{w}) : \left((2\psi_E \otimes_s - \zeta_E) \nabla(\eta_R \partial_E \hat{u}) - \text{Id}(\psi_E \cdot \nabla)(\eta_R \partial_E \hat{u}) \right).
 \end{aligned}$$

It remains to estimate these terms, and we split the proof of (5.14) into four further substeps.

Substep 2.1. Lower bound on J_0 : for all $K \gg 1$ and $0 < \varepsilon \ll K^{-1/2}$,

$$\begin{aligned}
 (5.18) \quad J_0 &\geq \frac{1}{2} \int_{D_R} \tilde{\mu}_{R,\varepsilon}^2 |\nabla w|^2 \\
 &\quad - \frac{1}{K} \int_{D_R} \tilde{\mu}_{R,\varepsilon}^2 Q^2 - K \int_{D_R} \tilde{\mu}_{R,\varepsilon}^2 |\nabla(w - \tilde{w})|^2 - K \int_{D_R} \tilde{\mu}_{R,\varepsilon}^2 \text{div}(\tilde{w})^2.
 \end{aligned}$$

Expanding the gradient in the definition of J_0 yields

$$J_0 = \int_{D_R} \tilde{\mu}_{R,\varepsilon}^2 \nabla \tilde{w} : \nabla w + \int_{D_R} 2\tilde{\mu}_{R,\varepsilon} (\tilde{w} \otimes \nabla \tilde{\mu}_{R,\varepsilon}) : \nabla w - \int_{D_R} \left(\tilde{\mu}_{R,\varepsilon}^2 \operatorname{div}(\tilde{w}) + 2\tilde{\mu}_{R,\varepsilon} \tilde{w} \cdot \nabla \tilde{\mu}_{R,\varepsilon} \right) Q.$$

Adding and subtracting ∇w to $\nabla \tilde{w}$, we deduce by Young’s inequality, for all $K \geq 1$,

$$(5.19) \quad J_0 \geq \left(1 - \frac{1}{K}\right) \int_{D_R} \tilde{\mu}_{R,\varepsilon}^2 |\nabla w|^2 - \frac{1}{K} \int_{D_R} \tilde{\mu}_{R,\varepsilon}^2 Q^2 - 4K \int_{D_R} |\nabla \tilde{\mu}_{R,\varepsilon}|^2 |\tilde{w}|^2 - \frac{K}{2} \int_{D_R} \tilde{\mu}_{R,\varepsilon}^2 |\nabla(w - \tilde{w})|^2 - \frac{K}{2} \int_{D_R} \tilde{\mu}_{R,\varepsilon}^2 \operatorname{div}(\tilde{w})^2.$$

Since $\tilde{\mu}_{R,\varepsilon}$ satisfies for all $x \in B_R$,

$$\tilde{\mu}_{R,\varepsilon}(x) \simeq \mu_{R,\varepsilon}(x), \quad |\nabla \tilde{\mu}_{R,\varepsilon}(x)| \lesssim |\nabla \mu_{R,\varepsilon}(x)| \simeq \frac{\varepsilon}{R} \left(1 - \frac{|x|}{R}\right)^{(\varepsilon/2)-1},$$

the following estimate follows from Hardy’s inequality in form of e.g. [28, Estimate (88)]: given $0 < \varepsilon \leq 1/2$, there holds for all $g \in H_0^1(B_R)$,

$$(5.20) \quad \int_{B_R} |\nabla \tilde{\mu}_{R,\varepsilon}|^2 |g|^2 \lesssim \varepsilon^2 \int_{B_R} \tilde{\mu}_{R,\varepsilon}^2 |\nabla g|^2.$$

Extending \tilde{w} by 0 outside D_R and applying this inequality, we find

$$\int_{D_R} |\nabla \tilde{\mu}_{R,\varepsilon}|^2 |\tilde{w}|^2 \lesssim \varepsilon^2 \int_{D_R} \tilde{\mu}_{R,\varepsilon}^2 |\nabla \tilde{w}|^2.$$

Inserting this into (5.19), the claim (5.18) follows for $K \geq 3$ and $K\varepsilon^2 \ll 1$.

Substep 2.2. Upper bound on J_1 : for all $K \geq 1$,

$$(5.21) \quad |J_1| \lesssim \frac{1}{K} \int_{D_R} \tilde{\mu}_{R,\varepsilon}^2 |\nabla \tilde{w}|^2 + \frac{1}{K} \int_{D_R} \tilde{\mu}_{R,\varepsilon}^2 |\tilde{Q}|^2 + K \left(\sup_{D_R} |\nabla(\eta_R \nabla \hat{u})|^2 \right) \int_{D_R} (1 + |(\nabla \psi, \Sigma \mathbb{1}_{\mathbb{R}^d \setminus \mathcal{J}})|^2) + K \int_{D_R} (1 - \eta_R)^2 \tilde{\mu}_{R,\varepsilon}^2 |\nabla \hat{u}|^2.$$

We examine separately the two terms in the definition of $J_1 = J_{1,1} + J_{1,2}$,

$$J_{1,1} := - \sum_{n \in \mathcal{N}_R} \int_{\partial I_n} \tilde{\mu}_{R,\varepsilon}^2 \tilde{w} \cdot \sigma(u, P + P_* - P_n) \nu,$$

$$J_{1,2} := \sum_{n \in \mathcal{N}_R} \int_{I_n} (\eta_R \partial_E \hat{u}) \operatorname{D}(\tilde{\mu}_{R,\varepsilon}^2 \tilde{w}) : J_E,$$

and we start with $J_{1,1}$. Since $\tilde{\mu}_{R,\varepsilon}$ and η_R are constant in the inclusions, and since for all $n \in \mathcal{N}_R$ we have

$$(5.22) \quad \operatorname{D}(\tilde{w}) = -(1 - \eta_R) \left(\int_{I_n + (\delta/2)B} \operatorname{D}(\hat{u}) \right), \quad \text{in } I_n,$$

we may use the boundary conditions for u to the effect of

$$J_{1,1} = \sum_{n \in \mathcal{N}_R} ((1 - \eta_R) \tilde{\mu}_{R,\varepsilon}^2)(x_n) \left(\int_{I_n + (\delta/2)B} \mathbf{D}(\hat{u}) \right) : \int_{\partial I_n} \sigma(u, P + P_* - P_n) \nu \otimes (x - x_n).$$

Using Stokes' theorem in the form $\int_{\partial I_n} \nu \otimes (x - x_n) = |I_n| \text{Id}$, together with the constraint $\text{div}(\hat{u}) = 0$ that we use in the form $(\int_{I_n + \frac{\delta}{2}B} \mathbf{D}(\hat{u})) : \text{Id} = 0$, we can subtract any constant to the pressure in the above expression, so that in particular

$$(5.23) \quad J_{1,1} = \sum_{n \in \mathcal{N}_R} ((1 - \eta_R) \tilde{\mu}_{R,\varepsilon}^2)(x_n) \left(\int_{I_n + (\delta/2)B} \mathbf{D}(\hat{u}) \right) : \int_{\partial I_n} \sigma(u, P + P_* - T_0[\hat{P}] - \eta_R \bar{\mathbf{b}} : T_0[\mathbf{D}(\hat{u})]) \nu \otimes (x - x_n).$$

We turn to $J_{1,2}$. Decomposing $\partial_E \hat{u} = (\partial_E \hat{u} - T_0[\partial_E \hat{u}]) + T_0[\partial_E \hat{u}]$, using that $T_0[\partial_E \hat{u}]$, $\tilde{\mu}_{R,\varepsilon}$, and η_R are constant in the inclusions, that \tilde{w} is affine in the inclusions, and using (5.22) again, we find

$$J_{1,2} = \sum_{n \in \mathcal{N}_R} \int_{I_n} \eta_R \tilde{\mu}_{R,\varepsilon}^2 (\partial_E \hat{u} - T_0[\partial_E \hat{u}]) \mathbf{D}(\tilde{w}) : J_E - \sum_{n \in \mathcal{N}_R} ((1 - \eta_R) \eta_R \tilde{\mu}_{R,\varepsilon}^2 T_0[\partial_E \hat{u}]) (x_n) \left(\int_{I_n + (\delta/2)B} \mathbf{D}(\hat{u}) \right) : \int_{I_n} J_E.$$

Writing $J_E|_{I_n} = \sigma(\psi_E^n, \Sigma_E^n)$ with (ψ_E^n, Σ_E^n) defined in (4.4), cf. (4.10), using Stokes' theorem, and recalling that $\sigma(\psi_E^n, \Sigma_E^n) \nu = \sigma(\psi_E + Ex, \Sigma_E) \nu$ on ∂I_n , cf. (4.4), we deduce

$$J_{1,2} = \sum_{n \in \mathcal{N}_R} \int_{I_n} \eta_R \tilde{\mu}_{R,\varepsilon}^2 (\partial_E \hat{u} - T_0[\partial_E \hat{u}]) \mathbf{D}(\tilde{w}) : \sigma(\psi_E^n, \Sigma_E^n) - \sum_{n \in \mathcal{N}_R} ((1 - \eta_R) \eta_R \tilde{\mu}_{R,\varepsilon}^2 T_0[\partial_E \hat{u}]) (x_n) \left(\int_{I_n + (\delta/2)B} \mathbf{D}(\hat{u}) \right) : \int_{\partial I_n} \sigma(\psi_E + Ex, \Sigma_E) \nu \otimes (x - x_n).$$

Combining this with (5.23), and reorganizing the terms, we obtain

$$J_1 = J'_{1,1} + J'_{1,2},$$

in terms of

$$J'_{1,1} = \sum_{n \in \mathcal{N}_R} \int_{I_n} \eta_R \tilde{\mu}_{R,\varepsilon}^2 (\partial_E \hat{u} - T_0[\partial_E \hat{u}]) \mathbf{D}(\tilde{w}) : \sigma(\psi_E^n, \Sigma_E^n),$$

$$J'_{1,2} = \sum_{n \in \mathcal{N}_R} ((1 - \eta_R) \tilde{\mu}_{R,\varepsilon}^2)(x_n) \left(\int_{I_n + (\delta/2)B} \mathbf{D}(\hat{u}) \right) : \int_{\partial I_n} \left(\sigma(u, P + P_* - T_0[\hat{P}] - \eta_R \bar{\mathbf{b}} : T_0[\mathbf{D}(\hat{u})]) - \eta_R T_0[\partial_E \hat{u}] \sigma(\psi_E + Ex, \Sigma_E) \right) \nu \otimes (x - x_n).$$

We separately estimate $J'_{1,1}$ and $J'_{1,2}$, and we start with the former. Using (4.5) and noting that $|\nabla\hat{u} - T_0[\nabla\hat{u}]| = |\nabla\hat{u} - \int_{I_n+(\delta/2)B} \nabla\hat{u}| \lesssim \sup_{I_n+(\delta/2)B} |\nabla^2\hat{u}|$ on I_n and that η_R is constant in I_n , we find

$$(5.24) \quad |J'_{1,1}| \lesssim \left(\sup_{D_R} |\nabla(\eta_R \nabla\hat{u})| \right) \left(\int_{D_R} \tilde{\mu}_{R,\varepsilon}^2 |\nabla\tilde{w}|^2 \right)^{1/2} \left(\int_{D_R} (1 + |(\nabla\psi, \Sigma \mathbb{1}_{\mathbb{R}^d \setminus \mathcal{J}})|^2) \right)^{1/2}.$$

We turn to $J'_{1,2}$. Writing for abbreviation

$$H := \sigma(u, P' - T_0[\hat{P}] - \eta_R \bar{\mathbf{b}} : T_0[\mathbf{D}(\hat{u})]) - \eta_R T_0[\partial_E \hat{u}] \sigma(\psi_E + Ex, \Sigma_E),$$

and noting that $\operatorname{div}(H) = 0$ in $(I_n + (\delta/4)B) \setminus I_n$,

$$\int_{\partial I_n} H \nu = 0, \quad \text{and} \quad \int_{\partial I_n} \Theta(x - x_n) \cdot H \nu = 0$$

for all $n \in \mathcal{N}_R$ and $\Theta \in \mathbb{M}^{\text{skew}}$, the trace estimate (4.22) leads to

$$(5.25) \quad |J'_{1,2}| \lesssim \sum_{n \in \mathcal{N}_R} ((1 - \eta_R) \tilde{\mu}_{R,\varepsilon}^2)(x_n) \left(\int_{I_n+(\delta/2)B} |\mathbf{D}(\hat{u})|^2 \right)^{1/2} \left(\int_{(I_n+(\delta/4)B) \setminus I_n} |H|^2 \right)^{1/2}.$$

For all $n \in \mathcal{N}_R$, we can write in the annulus $(I_n + (\delta/4)B) \setminus I_n$ (where $P' = P + P_*$), recalling the definition (5.16) of the modified two-scale expansion error (\tilde{w}, \tilde{Q}) and the definition of truncations,

$$\begin{aligned} H &= 2 \mathbf{D}(u - \eta_R T_1[\hat{u}] - \eta_R \psi_E T_0[\partial_E \hat{u}]) \\ &\quad - (P' - T_0[\hat{P}] - \eta_R \bar{\mathbf{b}} : T_0[\mathbf{D}(\hat{u})] - \eta_R \Sigma_E T_0[\partial_E \hat{u}]) \operatorname{Id} \\ &= \sigma(\tilde{w}, \tilde{Q}) + 2(1 - \eta_R) T_0[\mathbf{D}(\hat{u})]. \end{aligned}$$

Inserting this into (5.25), using that $\sup_{B(x)} \tilde{\mu}_{R,\varepsilon} \simeq \inf_{B(x)} \tilde{\mu}_{R,\varepsilon}$ holds for all $x \in D_R$, and using that η_R is constant in fattened inclusions, we deduce

$$\begin{aligned} |J'_{1,2}| &\lesssim \left(\int_{D_R} (1 - \eta_R)^2 \tilde{\mu}_{R,\varepsilon}^2 |\mathbf{D}(\hat{u})|^2 \right)^{1/2} \\ &\quad \times \left(\int_{D_R} \tilde{\mu}_{R,\varepsilon}^2 (|\mathbf{D}(\tilde{w})|^2 + |\tilde{Q}|^2) + (1 - \eta_R)^2 \tilde{\mu}_{R,\varepsilon}^2 |\mathbf{D}(\hat{u})|^2 \right)^{1/2}. \end{aligned}$$

Combined with the bound (5.24) on $J'_{1,1}$, the claim (5.21) follows by Young's inequality.

Substep 2.3. Upper bound on J_2, J_3 : for all $K \geq 1$,

$$(5.26) \quad |J_2| + |J_3| \lesssim \frac{1}{K} \int_{D_R} \tilde{\mu}_{R,\varepsilon}^2 |\nabla\tilde{w}|^2 + K \int_{D_R} (1 - \eta_R)^2 \tilde{\mu}_{R,\varepsilon}^2 |\nabla\hat{u}|^2 + K \left(\sup_{D_R} |\nabla(\eta_R \nabla\hat{u})|^2 \right) \int_{D_R} |(\psi, \zeta)|^2.$$

Expanding the gradients and using Young’s inequality, we find for all $K \geq 1$,

$$\begin{aligned}
 |J_2| &\lesssim K \int_{D_R} (1 - \eta_R)^2 \tilde{\mu}_{R,\varepsilon}^2 |\nabla \hat{u}|^2 + \frac{1}{K} \int_{D_R} \tilde{\mu}_{R,\varepsilon}^2 |\nabla \tilde{w}|^2 + \frac{1}{K} \int_{D_R} |\nabla \tilde{\mu}_{R,\varepsilon}|^2 |\tilde{w}|^2, \\
 |J_3| &\lesssim K \int_{D_R} \tilde{\mu}_{R,\varepsilon}^2 |(\psi, \zeta)|^2 |\nabla(\eta_R \nabla \hat{u})|^2 + \frac{1}{K} \int_{D_R} \tilde{\mu}_{R,\varepsilon}^2 |\nabla \tilde{w}|^2 + \frac{1}{K} \int_{D_R} |\nabla \tilde{\mu}_{R,\varepsilon}|^2 |\tilde{w}|^2,
 \end{aligned}$$

and Hardy’s inequality (5.20) yields the claim (5.26).

Substep 2.4. Control of truncation errors

$$\begin{aligned}
 (5.27) \quad \int_{D_R} \tilde{\mu}_{R,\varepsilon}^2 |\nabla(w - \tilde{w})|^2 &\lesssim \int_{D_R} (1 - \eta_R)^2 \tilde{\mu}_{R,\varepsilon}^2 |\nabla \hat{u}|^2 \\
 &\quad + \left(\sup_{D_R} |\nabla(\eta_R \nabla \hat{u})|^2 \right) \int_{D_R} (1 + |(\psi, \nabla \psi)|^2),
 \end{aligned}$$

$$\begin{aligned}
 (5.28) \quad \int_{D_R} \tilde{\mu}_{R,\varepsilon}^2 (Q - \tilde{Q})^2 &\lesssim \int_{D_R} (1 - \eta_R)^2 \tilde{\mu}_{R,\varepsilon}^2 |\nabla \hat{u}|^2 \\
 &\quad + \left(\sup_{D_R} |\nabla(\eta_R \nabla \hat{u})|^2 \right) \int_{D_R} (1 + \Sigma^2 \mathbb{1}_{\mathbb{R}^d \setminus \mathcal{J}}).
 \end{aligned}$$

We start with the proof of (5.27). The definition (5.16) of \tilde{w} yields

$$\nabla(w - \tilde{w}) = -\nabla(\hat{u} - T_1[\hat{u}]) - \eta_R(\partial_E \hat{u} - T_0[\partial_E \hat{u}]) \nabla \psi_E - \psi_E \otimes \nabla(\eta_R(\partial_E \hat{u} - T_0[\partial_E \hat{u}])),$$

and thus

$$\begin{aligned}
 (5.29) \quad \int_{D_R} \tilde{\mu}_{R,\varepsilon}^2 |\nabla(w - \tilde{w})|^2 &\lesssim \int_{D_R} \tilde{\mu}_{R,\varepsilon}^2 |\nabla(\hat{u} - T_1[\hat{u}])|^2 \\
 &\quad + \left(\sup_{D_R} |\eta_R(\nabla \hat{u} - T_0[\nabla \hat{u}])|^2 + \sup_{D_R} |\nabla(\eta_R(\nabla \hat{u} - T_0[\nabla \hat{u}]))|^2 \right) \int_{D_R} |(\psi, \nabla \psi)|^2.
 \end{aligned}$$

The definition (5.15) of the truncation maps T_0, T_1 gives

$$\begin{aligned}
 \nabla \hat{u} - T_0[\nabla \hat{u}] &= \sum_{n \in \mathcal{N}_R} \chi_n \left(\nabla \hat{u} - \int_{I_n + (\delta/2)B} \nabla \hat{u} \right), \\
 \nabla(\hat{u} - T_1[\hat{u}]) &= \sum_{n \in \mathcal{N}_R} \chi_n \left(\nabla \hat{u} - \int_{I_n + (\delta/2)B} \nabla \hat{u} \right) \\
 &\quad + \sum_{n \in \mathcal{N}_R} \nabla \chi_n \left(\hat{u} - \left(\int_{I_n + (\delta/2)B} \hat{u} \right) - \left(\int_{I_n + (\delta/2)B} \nabla \hat{u} \right) (x - x_n) \right).
 \end{aligned}$$

Using the properties of $\tilde{\mu}_{R,\varepsilon}$, η_R , and of the cut-off functions $\{\chi_n\}_n$, and appealing to Poincaré’s inequality on the fattened inclusions (on which we recall that η_R is

constant), we find

$$\begin{aligned}
 & \int_{D_R} \tilde{\mu}_{R,\varepsilon}^2 |\nabla(\hat{u} - T_1[\hat{u}])|^2 \\
 & \lesssim \sum_{n \in \mathcal{N}_R} \left(\sup_{I_n + (\delta/2)B} \tilde{\mu}_{R,\varepsilon}^2 \right) \\
 (5.30) \quad & \quad \times \int_{I_n + (\delta/2)B} \left(\eta_R^2 |\nabla(\hat{u} - T_1[\hat{u}])|^2 + (1 - \eta_R)^2 |\nabla(\hat{u} - T_1[\hat{u}])|^2 \right) \\
 & \lesssim \sum_{n \in \mathcal{N}_R} \left(\sup_{I_n + (\delta/2)B} \tilde{\mu}_{R,\varepsilon}^2 \right) \int_{I_n + (\delta/2)B} \left(\eta_R^2 |\nabla^2 \hat{u}|^2 + (1 - \eta_R)^2 |\nabla \hat{u}|^2 \right) \\
 & \lesssim \int_{D_R} |\nabla(\eta_R \nabla \hat{u})|^2 + \int_{D_R} (1 - \eta_R)^2 \tilde{\mu}_{R,\varepsilon}^2 |\nabla \hat{u}|^2,
 \end{aligned}$$

and similarly,

$$\begin{aligned}
 (5.31) \quad & \sup_{D_R} |\eta_R(\nabla \hat{u} - T_0[\nabla \hat{u}])| + \sup_{D_R} |\nabla(\eta_R(\nabla \hat{u} - T_0[\nabla \hat{u}]))| \\
 & \lesssim \sup_{n \in \mathcal{N}_R} \left(\eta_R(x_n) \sup_{I_n + (\delta/2)B} |\nabla^2 \hat{u}| \right) \lesssim \sup_{D_R} |\nabla(\eta_R \nabla \hat{u})|.
 \end{aligned}$$

Inserting these bounds into (5.29), the claim (5.27) follows.

We turn to the proof of (5.28). The definition (5.16) of \tilde{Q} yields

$$Q - \tilde{Q} = -(\hat{P} - T_0[\hat{P}]) - \eta_R \bar{\mathbf{b}} : (\mathbf{D}(\hat{u}) - T_0[\mathbf{D}(\hat{u})]) - \eta_R \Sigma_E \mathbb{1}_{\mathbb{R}^d \setminus \mathcal{J}} (\partial_E \hat{u} - T_0[\partial_E \hat{u}]),$$

and thus

$$\begin{aligned}
 (5.32) \quad & \int_{D_R} \tilde{\mu}_{R,\varepsilon}^2 (Q - \tilde{Q})^2 \lesssim \int_{D_R} \tilde{\mu}_{R,\varepsilon}^2 (\hat{P} - T_0[\hat{P}])^2 \\
 & \quad + \left(\sup_{D_R} |\eta_R(\nabla \hat{u} - T_0[\nabla \hat{u}])|^2 \right) \int_{D_R} (1 + \Sigma^2 \mathbb{1}_{\mathbb{R}^d \setminus \mathcal{J}}).
 \end{aligned}$$

We start by analyzing the first right-hand side term. By definition of T_0 , using the properties of $\tilde{\mu}_{R,\varepsilon}$ and appealing to Poincaré's inequality on the fattened inclusions (on which we recall that η_R is constant), we find

$$\begin{aligned}
 (5.33) \quad & \int_{D_R} \tilde{\mu}_{R,\varepsilon}^2 (\hat{P} - T_0[\hat{P}])^2 \lesssim \sum_{n \in \mathcal{N}_R} \left(\sup_{I_n + (\delta/2)B} \tilde{\mu}_{R,\varepsilon}^2 \right) \int_{I_n + (\delta/2)B} \left(\hat{P} - \fint_{I_n + (\delta/2)B} \hat{P} \right)^2 \\
 & \lesssim \sum_{n \in \mathcal{N}_R} \left(\sup_{I_n + (\delta/2)B} \tilde{\mu}_{R,\varepsilon}^2 \right) \int_{I_n + (\delta/2)B} \left(\eta_R^2 |\nabla \hat{P}|^2 + (1 - \eta_R)^2 \left(\hat{P} - \fint_{I_n + (\delta/2)B} \hat{P} \right)^2 \right).
 \end{aligned}$$

We now appeal to a classical pressure estimates on \hat{P} . On the one hand, since (\hat{u}, \hat{P}) satisfies a steady Stokes equation (5.7) without forcing in D_R , a direct use of the Bogovskii operator in form of e.g. [22, Th. III.3.1] yields for all $n \in \mathcal{N}_R$,

$$(5.34) \quad \int_{I_n + (\delta/2)B} \left(\hat{P} - \fint_{I_n + (\delta/2)B} \hat{P} \right)^2 \lesssim \int_{I_n + (\delta/2)B} |\nabla \hat{u}|^2.$$

On the other hand, since $(\partial_i \hat{u}, \partial_i \hat{P})$ satisfies the same equation in D_R , the same argument yields

$$\int_{I_n+(\delta/2)B} \left| \nabla \hat{P} - \fint_{I_n+(\delta/2)B} \nabla \hat{P} \right|^2 \lesssim \int_{I_n+(\delta/2)B} |\nabla^2 \hat{u}|^2,$$

Further noting that equation (5.7) yields

$$\begin{aligned} \int_{I_n+(\delta/2)B} \nabla \hat{P} &= \int_{I_n+(\delta/2)B} \operatorname{div}(2\bar{\mathbf{B}}\mathbf{D}(\hat{u})) \\ &= 2 \int_{\partial(I_n+(\delta/2)B)} (\bar{\mathbf{B}}\mathbf{D}(\hat{u}))\nu \\ &= 2 \int_{\partial(I_n+(\delta/2)B)} \left(\bar{\mathbf{B}}\mathbf{D}(\hat{u}) - \fint_{I_n+(\delta/2)B} \bar{\mathbf{B}}\mathbf{D}(\hat{u}) \right)\nu, \end{aligned}$$

and thus

$$\left| \int_{I_n+(\delta/2)B} \nabla \hat{P} \right| \lesssim \sup_{I_n+(\delta/2)B} |\nabla^2 \hat{u}|,$$

we deduce

$$\int_{I_n+(\delta/2)B} |\nabla \hat{P}|^2 \lesssim \sup_{I_n+(\delta/2)B} |\nabla^2 \hat{u}|^2.$$

Inserting this together with (5.34) into (5.33), we obtain

$$\int_{D_R} \tilde{\mu}_{R,\varepsilon}^2 (\hat{P} - T_0[\hat{P}])^2 \lesssim |D_R| \left(\sup_{D_R} |\nabla(\eta_R \nabla \hat{u})|^2 \right) + \int_{D_R} (1 - \eta_R)^2 \tilde{\mu}_{R,\varepsilon}^2 |\nabla \hat{u}|^2.$$

Combining this with (5.32) and (5.31), the claim (5.28) follows.

Substep 2.5. Control of the divergence

$$(5.35) \quad \int_{D_R} \tilde{\mu}_{R,\varepsilon}^2 \operatorname{div}(\tilde{w})^2 \lesssim \int_{D_R} (1 - \eta_R)^2 \tilde{\mu}_{R,\varepsilon}^2 |\nabla \hat{u}|^2 + \left(\sup_{D_R} |\nabla(\eta_R \nabla \hat{u})|^2 \right) \int_{D_R} (1 + |\psi|^2).$$

As $\operatorname{div}(u) = \operatorname{div}(\hat{u}) = \operatorname{div}(\psi_E) = 0$, the definition (5.16) of \tilde{w} yields

$$\operatorname{div}(\tilde{w}) = \operatorname{div}(\hat{u} - T_1[\hat{u}]) - \psi_E \cdot \nabla(\eta_R T_0[\partial_E \hat{u}]),$$

and the claim (5.35) follows from the estimates (5.30) and (5.31).

Substep 2.6. Proof of (5.14). — Combining (5.17), (5.18), (5.21), and (5.26), we obtain for all $K \gg 1$ and $0 < \varepsilon \ll K^{-1/2}$,

$$\begin{aligned} \int_{D_R} \tilde{\mu}_{R,\varepsilon}^2 |\nabla w|^2 &\lesssim \frac{1}{K} \int_{D_R} \tilde{\mu}_{R,\varepsilon}^2 |\nabla \tilde{w}|^2 + \frac{1}{K} \int_{D_R} \tilde{\mu}_{R,\varepsilon}^2 (Q^2 + \tilde{Q}^2) \\ &+ K \left(\sup_{D_R} |\nabla(\eta_R \nabla \hat{u})|^2 \right) \int_{D_R} (1 + |(\psi, \zeta, \nabla \psi, \Sigma \mathbb{1}_{\mathbb{R}^d \setminus \mathcal{J}})|^2) + K \int_{D_R} (1 - \eta_R)^2 \tilde{\mu}_{R,\varepsilon}^2 |\nabla \hat{u}|^2 \\ &+ K \int_{D_R} \tilde{\mu}_{R,\varepsilon}^2 |\nabla(w - \tilde{w})|^2 + K \int_{D_R} \tilde{\mu}_{R,\varepsilon}^2 \operatorname{div}(\tilde{w})^2. \end{aligned}$$

Decomposing $\nabla \tilde{w} = \nabla w + \nabla(\tilde{w} - w)$ and $\tilde{Q} = Q + (\tilde{Q} - Q)$, using the bounds (5.27) and (5.28) on the truncation errors $\nabla(w - \tilde{w})$ and $Q - \tilde{Q}$, and using the bound (5.35) on $\operatorname{div}(\tilde{w})$, we find

$$\begin{aligned} \int_{D_R} \tilde{\mu}_{R,\varepsilon}^2 |\nabla w|^2 &\lesssim \frac{1}{K} \int_{D_R} \tilde{\mu}_{R,\varepsilon}^2 |\nabla w|^2 + \frac{1}{K} \int_{D_R} \tilde{\mu}_{R,\varepsilon}^2 Q^2 \\ &\quad + K \left(\sup_{D_R} |\nabla(\eta_R \nabla \hat{u})|^2 \right) \int_{D_R} (1 + |(\psi, \zeta, \nabla \psi, \Sigma \mathbb{1}_{\mathbb{R}^d \setminus \mathcal{J}})|^2) \\ &\quad + K \int_{D_R} (1 - \eta_R)^2 \tilde{\mu}_{R,\varepsilon}^2 |\nabla \hat{u}|^2. \end{aligned}$$

Choosing $K \gg 1$ large enough to absorb the first right-hand side term, and noting that $\tilde{\mu}_{R,\varepsilon} \simeq \mu_{R,\varepsilon}$ on D_R , the conclusion (5.14) follows.

Step 3. Weighted pressure estimate for the two-scale expansion error: for all ε with $0 < \varepsilon < 1$,

$$(5.36) \quad \int_{D_R} \mu_{R,\varepsilon}^2 Q^2 \lesssim \int_{D_R} \mu_{R,\varepsilon}^2 |\nabla w|^2 + \int_{D_R} (1 - \eta_R)^2 \mu_{R,\varepsilon}^2 |\nabla \hat{u}|^2 \\ + \left(\sup_{D_R} |\nabla(\eta_R \nabla \hat{u})|^2 \right) \int_{D_R} (1 + |(\psi, \zeta, \nabla \psi, \Sigma \mathbb{1}_{\mathbb{R}^d \setminus \mathcal{J}})|^2).$$

Combining this with the bound (5.14) on ∇w , and choosing $K \gg 1$ large enough, we deduce for all ε with $0 < \varepsilon \ll 1$,

$$(5.37) \quad \int_{D_R} \mu_{R,\varepsilon}^2 (|\nabla w|^2 + Q^2) \lesssim \int_{D_R} (1 - \eta_R)^2 \mu_{R,\varepsilon}^2 |\nabla \hat{u}|^2 \\ + \left(\sup_{D_R} |\nabla(\eta_R \nabla \hat{u})|^2 \right) \int_{D_R} (1 + |(\psi, \zeta, \nabla \psi, \Sigma \mathbb{1}_{\mathbb{R}^d \setminus \mathcal{J}})|^2).$$

We turn to the proof of (5.36). For that purpose, we shall again appeal to the truncated version \tilde{Q} of Q as in Step 2, cf. (5.16). We also recall the notation (5.9) for P' , where we choose the constants P_* and $\{P_n\}_n$ such that

$$P_n = \int_{I_n + (\delta/2)B} \hat{P} + \eta_R(x_n) \bar{\mathbf{b}} : \int_{I_n + (\delta/2)B} \mathbf{D}(\hat{u}), \quad \int_{D_R} \tilde{\mu}_{R,\varepsilon}^2 \tilde{Q} = 0.$$

Note that this choice entails in particular $\tilde{Q} = 0$ inside inclusions $\{I_n\}_{n \in \mathcal{N}_R}$. With these definitions, we may turn to the proof of (5.36), which we split into three further substeps.

Substep 3.1. Weighted Bogovskii construction: given $0 < \varepsilon < 1$, there exists a vector field $S \in H_0^1(D_R)^d$ such that $S|_{I_n}$ is constant for all $n \in \mathcal{N}_R$ and such that

$$(5.38) \quad \operatorname{div}(S) = -\tilde{\mu}_{R,\varepsilon}^2 \tilde{Q}, \quad \text{in } D_R, \\ \int_{D_R} \tilde{\mu}_{R,\varepsilon}^{-2} |\nabla S|^2 \lesssim \int_{D_R} \tilde{\mu}_{R,\varepsilon}^2 \tilde{Q}^2.$$

Since $\int_{D_R} \tilde{\mu}_{R,\varepsilon}^2 \tilde{Q} = 0$, and since the weight $\tilde{\mu}_{R,\varepsilon}^{-2} \simeq \mu_{R,\varepsilon}^{-2}$ on D_R can be extended to $x \mapsto |1 - |x|/R|^{-\varepsilon}$ on \mathbb{R}^d , which belongs to the Muckenhoupt class A_2 uniformly in R

provided that $\varepsilon < 1$, we may appeal to the weighted Bogovskii construction in form of Lemma 5.2. Note that by definition the set D_R is star-shaped with respect to every point in $B_{R/2}$ as soon as $R \gg 1$. Hence, there exists a vector field $S^\circ \in H_0^1(D_R)^d$ such that

$$\begin{aligned} \operatorname{div}(S^\circ) &= -\tilde{\mu}_{R,\varepsilon}^2 \tilde{Q}, & \text{in } D_R, \\ \int_{D_R} \tilde{\mu}_{R,\varepsilon}^{-2} |\nabla S^\circ|^2 &\lesssim \int_{D_R} \tilde{\mu}_{R,\varepsilon}^2 \tilde{Q}^2. \end{aligned}$$

It remains to modify S° to make it constant inside the inclusions $\{I_n\}_{n \in \mathcal{N}_R}$ without changing its divergence and the bound on its norm. For that purpose, we essentially follow the argument of [18, Proof of Prop. 2.1]; see also the proof of Lemma 3.3. More precisely, for all $n \in \mathcal{N}_R$, recalling that $\operatorname{dist}(I_n, \partial D_R) \geq \delta$ and that $\tilde{Q} = 0$ in I_n , a standard use of the Bogovskii operator allows to construct as in (3.5) a vector field $S^n \in H_0^1(I_n + (\delta/2)B)^d$ such that $S^n = -S^\circ + \int_{I_n} S^\circ$ in I_n and

$$\begin{aligned} \operatorname{div}(S^n) &= 0, & \text{in } I_n + (\delta/2)B, \\ \|\nabla S^n\|_{L^2((I_n + (\delta/2)B) \setminus I_n)} &\lesssim \|\nabla S^\circ\|_{L^2(I_n)}. \end{aligned}$$

Smuggling in the weight $\tilde{\mu}_{R,\varepsilon}^{-1}$ (which is constant on the fattened inclusions), this yields

$$\|\tilde{\mu}_{R,\varepsilon}^{-1} \nabla S^n\|_{L^2((I_n + (\delta/2)B) \setminus I_n)} \lesssim \|\tilde{\mu}_{R,\varepsilon}^{-1} \nabla S^\circ\|_{L^2(I_n)}.$$

Since the fattened inclusions are all disjoint, cf. (H_δ) , extending S^n by 0 in $D_R \setminus (I_n + (\delta/2)B)$ for all $n \in \mathcal{N}_R$, the vector field $S := S^\circ + \sum_{n \in \mathcal{N}_R} S^n$ satisfies all the required properties.

Substep 3.2. Proof of (5.36). — Testing equation (5.11) with the test function $S \in H_0^1(D_R)^d$ constructed in the previous substep yields

$$L_0 = L_1 + L_2 + L_3,$$

in terms of

$$\begin{aligned} L_0 &:= \int_{D_R} \nabla S : \nabla w - \int_{D_R} \operatorname{div}(S) Q, \\ L_1 &:= - \sum_{n \in \mathcal{N}_R} \int_{\partial I_n} S \cdot \sigma(u, P + P_* - P_n) \nu + \sum_{n \in \mathcal{N}_R} \int_{I_n} (\eta_R \partial_E \hat{u}) \nabla S : J_E, \\ L_2 &:= -2 \int_{D_R} (1 - \eta_R) \nabla S : (\operatorname{Id} - \bar{B}) D(\hat{u}), \\ L_3 &:= - \int_{D_R} \nabla S : \left((2\psi_E \otimes_s - \zeta_E) \nabla (\eta_R \partial_E \hat{u}) - \operatorname{Id}(\psi_E \cdot \nabla) (\eta_R \partial_E \hat{u}) \right). \end{aligned}$$

We start by giving a lower bound on L_0 . Using the defining property (5.38) of the test function S in form of

$$\begin{aligned} - \int_{D_R} \operatorname{div}(S) Q &= \int_{D_R} \tilde{\mu}_{R,\varepsilon}^2 Q \tilde{Q} \\ &\geq \int_{D_R} \tilde{\mu}_{R,\varepsilon}^2 \tilde{Q}^2 - \left(\int_{D_R} \tilde{\mu}_{R,\varepsilon}^2 \tilde{Q}^2 \right)^{1/2} \left(\int_{D_R} \tilde{\mu}_{R,\varepsilon}^2 (Q - \tilde{Q})^2 \right)^{1/2}, \end{aligned}$$

and using the bound (5.38) on the weighted norm of ∇S in form of

$$\begin{aligned} \left| \int_{D_R} \nabla S : \nabla w \right| &\leq \left(\int_{D_R} \tilde{\mu}_{R,\varepsilon}^{-2} |\nabla S|^2 \right)^{1/2} \left(\int_{D_R} \tilde{\mu}_{R,\varepsilon}^2 |\nabla w|^2 \right)^{1/2} \\ &\lesssim \left(\int_{D_R} \tilde{\mu}_{R,\varepsilon}^2 \tilde{Q}^2 \right)^{1/2} \left(\int_{D_R} \tilde{\mu}_{R,\varepsilon}^2 |\nabla w|^2 \right)^{1/2}, \end{aligned}$$

we deduce for all $K \geq 1$,

$$(5.39) \quad L_0 \geq \int_{D_R} \tilde{\mu}_{R,\varepsilon}^2 \tilde{Q}^2 - \frac{1}{K} \int_{D_R} \tilde{\mu}_{R,\varepsilon}^2 \tilde{Q}^2 - K \int_{D_R} \tilde{\mu}_{R,\varepsilon}^2 (Q - \tilde{Q})^2 - CK \int_{D_R} \tilde{\mu}_{R,\varepsilon}^2 |\nabla w|^2.$$

Next, recalling that $S|_{I_n}$ is constant for all $n \in \mathcal{N}_R$, and using the boundary conditions for u , we find $L_1 = 0$. It remains to estimate L_2 and L_3 . Smuggling in the weight $\tilde{\mu}_{R,\varepsilon}$, we find for all $K \geq 1$,

$$\begin{aligned} L_2 + L_3 &\lesssim \frac{1}{K} \int_{D_R} \tilde{\mu}_{R,\varepsilon}^{-2} |\nabla S|^2 + K \int_{D_R} (1 - \eta_R)^2 \tilde{\mu}_{R,\varepsilon}^2 |\nabla \hat{u}|^2 \\ &\quad + K \left(\sup_{D_R} |\nabla(\eta_R \nabla \hat{u})|^2 \right) \int_{D_R} (1 + |(\psi, \zeta)|^2). \end{aligned}$$

Using the weighted estimate (5.38) on ∇S to estimate the first right-hand side term, and combining with the lower bound (5.39) on L_0 , we deduce for all $K \geq 1$,

$$\begin{aligned} \int_{D_R} \tilde{\mu}_{R,\varepsilon}^2 \tilde{Q}^2 &\lesssim \frac{1}{K} \int_{D_R} \tilde{\mu}_{R,\varepsilon}^2 \tilde{Q}^2 + K \int_{D_R} \tilde{\mu}_{R,\varepsilon}^2 (Q - \tilde{Q})^2 + K \int_{D_R} \tilde{\mu}_{R,\varepsilon}^2 |\nabla w|^2 \\ &\quad + K \left(\sup_{D_R} |\nabla(\eta_R \nabla \hat{u})|^2 \right) \int_{D_R} (1 + |(\psi, \zeta)|^2) + K \int_{D_R} (1 - \eta_R)^2 \tilde{\mu}_{R,\varepsilon}^2 |\nabla \hat{u}|^2, \end{aligned}$$

Choosing $K \gg 1$ large enough to absorb the first right-hand side term, and decomposing $\tilde{Q} = Q + (\tilde{Q} - Q)$, we obtain

$$\begin{aligned} \int_{D_R} \tilde{\mu}_{R,\varepsilon}^2 Q^2 &\lesssim \int_{D_R} \tilde{\mu}_{R,\varepsilon}^2 (Q - \tilde{Q})^2 + \int_{D_R} \tilde{\mu}_{R,\varepsilon}^2 |\nabla w|^2 \\ &\quad + \left(\sup_{D_R} |\nabla(\eta_R \nabla \hat{u})|^2 \right) \int_{D_R} (1 + |(\psi, \zeta)|^2) + \int_{D_R} (1 - \eta_R)^2 \tilde{\mu}_{R,\varepsilon}^2 |\nabla \hat{u}|^2. \end{aligned}$$

Using the bound (5.28) on the truncation error $Q - \tilde{Q}$, and recalling that $\tilde{\mu}_{R,\varepsilon} \simeq \mu_{R,\varepsilon}$ on D_R , the conclusion (5.36) follows.

Step 4. Conclusion: proof of (5.2). — We split the proof into five further substeps.

Substep 4.1. Caccioppoli-type inequality for homogeneous steady Stokes equation

Given a solution (\bar{v}, \bar{T}) of

$$(5.40) \quad -\operatorname{div}(2\bar{\mathbf{B}}\mathbf{D}(\bar{v})) + \nabla \bar{T} = 0, \quad \operatorname{div}(\bar{v}) = 0, \quad \text{in } B_R,$$

we have for all r with $0 < r < R$ and $K \geq 1$,

$$(5.41) \quad \int_{B_r} |\nabla \bar{v}|^2 \lesssim \frac{1}{K} \int_{B_R} |\nabla \bar{v}|^2 + \frac{K}{(R-r)^2} \int_{B_R} |\bar{v}|^2.$$

Consider a cut-off function $\chi_{r,R} \in C_c^\infty(\mathbb{R}^d)$ such that $\chi_{r,R}|_{B_r} = 1$, $\chi_{r,R}|_{\mathbb{R}^d \setminus B_R} = 0$, and $|\nabla \chi_{r,R}| \lesssim (R-r)^{-1}$. Testing the equation (5.40) with the test function $\chi_{r,R}^2 \bar{v}$, we find

$$\int_{\mathbb{R}^d} \chi_{r,R}^2 \mathbf{D}(\bar{v}) : 2\bar{\mathbf{B}} \mathbf{D}(\bar{v}) = -2 \int_{\mathbb{R}^d} \chi_{r,R} \bar{v} \otimes \nabla \chi_{r,R} : (2\bar{\mathbf{B}} \mathbf{D}(\bar{v}) - \bar{T}),$$

and thus

$$(5.42) \quad \int_{\mathbb{R}^d} \chi_{r,R}^2 |\mathbf{D}(\bar{v})|^2 \lesssim \frac{1}{R-r} \left(\int_{B_R} (|\mathbf{D}(\bar{v})|^2 + \bar{T}^2) \right)^{1/2} \left(\int_{B_R} |\bar{v}|^2 \right)^{1/2}.$$

Since $\operatorname{div}(\bar{v}) = 0$, integration by parts yields

$$\begin{aligned} \int_{\mathbb{R}^d} \chi_{r,R}^2 |\nabla \bar{v}|^2 &= 2 \int_{\mathbb{R}^d} \chi_{r,R}^2 |\mathbf{D}(\bar{v})|^2 - \int_{\mathbb{R}^d} \chi_{r,R}^2 \partial_i \bar{v}_j \partial_j \bar{v}_i \\ &= 2 \int_{\mathbb{R}^d} \chi_{r,R}^2 |\mathbf{D}(\bar{v})|^2 + 2 \int_{\mathbb{R}^d} \chi_{r,R} \nabla \chi_{r,R} \otimes \bar{v} : \nabla \bar{v}, \end{aligned}$$

and thus

$$\int_{\mathbb{R}^d} \chi_{r,R}^2 |\nabla \bar{v}|^2 \lesssim \int_{\mathbb{R}^d} \chi_{r,R}^2 |\mathbf{D}(\bar{v})|^2 + \frac{1}{(R-r)^2} \int_{B_R} |\bar{v}|^2.$$

Combining this with (5.42), we deduce for all $K \geq 1$,

$$\int_{B_r} |\nabla \bar{v}|^2 \lesssim \frac{1}{K} \int_{B_R} (|\mathbf{D}(\bar{v})|^2 + \bar{T}^2) + \frac{K}{(R-r)^2} \int_{B_R} |\bar{v}|^2.$$

As the pressure \bar{T} in (5.40) is only defined up to an additive constant, we may choose without loss of generality $\int_{B_R} \bar{T} = 0$, and we then appeal to a standard pressure estimate: a standard use of the Bogovskii operator in form of e.g. [22, Th. III.3.1] yields

$$\int_{B_R} \bar{T}^2 \lesssim \int_{B_R} |\nabla \bar{v}|^2$$

and the claim (5.41) follows.

Substep 4.2. Interior regularity estimate for homogeneous steady Stokes equation (5.7): for any boundary layer $4 < \rho < R$,

$$(5.43) \quad \rho^{2(n-1)} \sup_{B_{R-\rho}} |\nabla^n \hat{u}|^2 \lesssim_n \left(\frac{\rho}{R}\right)^{-d} \int_{D_R} |\nabla \hat{u}|^2.$$

First consider a solution (\bar{v}, \bar{T}) of the following homogeneous steady Stokes equation,

$$-\operatorname{div}(2\bar{\mathbf{B}} \mathbf{D}(\bar{v})) + \nabla \bar{T} = 0, \quad \operatorname{div}(\bar{v}) = 0, \quad \text{in } B.$$

In view of the standard interior regularity theory for this equation, see [22, Th. IV.4.1], we find for all $n \geq 0$,

$$\int_{(1/2)B} |\langle \nabla \rangle^n \nabla \bar{v}|^2 \lesssim_n \int_B (|\nabla \bar{v}|^2 + |\bar{T}|^2).$$

We then appeal to a pressure estimate for \bar{T} : assuming without loss of generality $\int_B \bar{T} = 0$, a standard use of the Bogovskii operator in form of e.g. [22, Th. III.3.1] yields

$$\int_{(1/2)B} |\langle \nabla \rangle^n \nabla \bar{v}|^2 \lesssim_n \int_B |\nabla \bar{v}|^2.$$

By Sobolev’s embedding, this entails for all $n \geq 1$,

$$\sup_{(1/2)B} |\nabla^n \bar{v}|^2 \lesssim_n \int_B |\nabla \bar{v}|^2.$$

Upon rescaling and translation, this implies for all $\rho < 1$, $x \in B_{1-\rho}$, and $n \geq 1$,

$$\rho^{2(n-1)} |\nabla^n \bar{v}(x)|^2 \lesssim_n \int_{B_\rho(x)} |\nabla \bar{v}|^2,$$

hence, for all $n \geq 1$,

$$\rho^{2(n-1)} \sup_{B_{1-\rho}} |\nabla^n \bar{v}|^2 \lesssim_n \rho^{-d} \int_B |\nabla \bar{v}|^2.$$

Turning back to equation (5.7) and recalling that $B_{R-3} \subset D_R$, the claim (5.43) follows after rescaling.

Substep 4.3. Reduction to the two-scale expansion error: for all r with $4 \leq r \leq \frac{1}{4}R$,

$$(5.44) \quad \int_{B_r} |\nabla u - \nabla \hat{u}(0) - (\partial_E \hat{u})(0) \nabla \psi_E|^2 \lesssim \left(\frac{r}{R}\right)^{-d-2} \int_{D_R} \mu_{R,\varepsilon}^2 |\nabla w|^2 + \left(\left(\frac{r}{R}\right)^2 \int_{B_R} (1 + |\nabla \psi|^2) + \left(\frac{r}{R}\right)^{-d} \frac{1}{R^2} \int_{B_R} |\psi|^2\right) \int_{B_R} |\nabla u|^2.$$

Consider the following local two-scale expansion error centered at the origin,

$$w_\circ := u - \hat{u}(0) - \nabla \hat{u}(0)x - \psi_E \partial_E \hat{u}(0), \quad Q_\circ := P - \Sigma_E \partial_E \hat{u}(0),$$

and note that equations (3.1) and (3.2) yield the following on D_R ,

$$-\Delta w_\circ + \nabla(Q_\circ \mathbb{1}_{\mathbb{R}^d \setminus \mathcal{J}}) = -\sum_n \delta_{\partial I_n} \sigma(w_\circ, Q_\circ) \nu.$$

We appeal to a Caccioppoli-type argument: as in the proof of (5.41), choosing a cut-off function that is constant in the inclusions, and using the boundary conditions for u and ψ_E , we find for all r with $4 \leq r \leq \frac{1}{4}R$ and $K \geq 1$,

$$(5.45) \quad \int_{B_r} |\nabla w_\circ|^2 \lesssim \frac{1}{K} \int_{B_{2r}} |\nabla w_\circ|^2 + \frac{K}{r^2} \int_{B_{2r}} |w_\circ|^2,$$

and it remains to examine the last right-hand side term. Comparing the local error w_\circ to its global version $w = u - \hat{u} - \eta_R \psi_E \partial_E \hat{u}$, cf. (5.10), and recalling that $\eta_R = 1$ on $B_{R-2\rho}$, we obtain from the triangle inequality, for all $r, \rho \leq R/4$ (which entails $B_{2r} \subset B_{R-2\rho}$),

$$\int_{B_{2r}} |w_\circ|^2 \lesssim \int_{B_{2r}} |w|^2 + \left(\sup_{B_{2r}} |\hat{u} - \hat{u}(0) - \nabla \hat{u}(0)x|^2\right) + \left(\sup_{B_{2r}} |\nabla \hat{u} - \nabla \hat{u}(0)|^2\right) \int_{B_{2r}} |\psi|^2.$$

Using Taylor’s formula, the interior regularity estimate (5.43) with $\rho = R/4$, and the energy estimate (5.8), we find for all $r \leq \frac{1}{4}R$,

$$\begin{aligned} \sup_{B_{2r}} |\widehat{u} - \widehat{u}(0) - \nabla \widehat{u}(0)x|^2 + r^2 \sup_{B_{2r}} |\nabla \widehat{u} - \nabla \widehat{u}(0)|^2 \\ \lesssim r^4 \sup_{B_{2r}} |\nabla^2 \widehat{u}|^2 \lesssim r^2 \left(\frac{r}{R}\right)^2 \int_{D_R} |\nabla \widehat{u}|^2 \lesssim r^2 \left(\frac{r}{R}\right)^2 \int_{D_R} |\nabla u|^2, \end{aligned}$$

so that the above becomes

$$(5.46) \quad \int_{B_{2r}} |w_\circ|^2 \lesssim \int_{B_{2r}} |w|^2 + \left(r^2 \left(\frac{r}{R}\right)^2 + \left(\frac{r}{R}\right)^{2-d} \int_{B_R} |\psi|^2\right) \int_{D_R} |\nabla u|^2.$$

It remains to analyze the first right-hand side term in this estimate. By definition of the weight $\mu_{R,\varepsilon}$ in (5.13), appealing to Hardy’s inequality (5.20), we find for all $r \leq \frac{1}{4}R$,

$$\begin{aligned} \int_{B_{2r}} |w|^2 &\lesssim \int_{B_{2r}} \left(1 - \frac{|x|}{R}\right)^{\varepsilon-2} |w|^2 \lesssim \left(\frac{r}{R}\right)^{-d} \int_{D_R} \left(1 - \frac{|x|}{R}\right)^{\varepsilon-2} |w|^2 \\ &\lesssim \varepsilon^{-2} R^2 \left(\frac{r}{R}\right)^{-d} \int_{D_R} |\nabla \mu_{R,\varepsilon}|^2 |w|^2 \\ &\lesssim R^2 \left(\frac{r}{R}\right)^{-d} \int_{D_R} \mu_{R,\varepsilon}^2 |\nabla w|^2. \end{aligned}$$

Combined with (5.45) and (5.46), this yields the following, for all r with $4 \leq r \leq \frac{1}{4}R$ and $K \geq 1$,

$$(5.47) \quad \begin{aligned} \int_{B_r} |\nabla w_\circ|^2 &\lesssim \frac{1}{K} \int_{B_{2r}} |\nabla w_\circ|^2 \\ &+ K \left(\left(\frac{r}{R}\right)^2 + \left(\frac{r}{R}\right)^{-d} \frac{1}{R^2} \int_{B_R} |\psi|^2\right) \int_{D_R} |\nabla u|^2 + K \left(\frac{r}{R}\right)^{-d-2} \int_{D_R} \mu_{R,\varepsilon}^2 |\nabla w|^2. \end{aligned}$$

In order to absorb the first right-hand side term, we proceed by iteration. Let us first rewrite (5.47) as follows: for any $K \geq 1$,

$$f(r) \leq \frac{1}{K} f(2r) + CKg(r), \quad \text{for all } r \text{ with } 4 \leq r \leq \frac{1}{4}R,$$

where we have set for abbreviation,

$$\begin{aligned} f(r) &:= \int_{B_r} |\nabla w_\circ|^2, \\ g(r) &:= \left(\left(\frac{r}{R}\right)^2 + \left(\frac{r}{R}\right)^{-d} \frac{1}{R^2} \int_{D_R} |\psi|^2\right) \int_{D_R} |\nabla u|^2 + \left(\frac{r}{R}\right)^{-d-2} \int_{D_R} \mu_{R,\varepsilon}^2 |\nabla w|^2. \end{aligned}$$

Iterating this estimate yields for all $r \geq 4$ and $n \geq 1$ with $2^n r \leq \frac{1}{4}R$,

$$f(r) \leq CK \sum_{m=0}^{n-1} K^{-m} g(2^m r) + K^{-n} f(2^n r).$$

Noting that $g(2^m r) \leq 4^m g(r)$ and choosing $K = 8$, this entails

$$f(r) \lesssim g(r) + 8^{-n} f(2^n r).$$

Choosing n large enough such that $2^n r \simeq R$, with $2^n r \leq \frac{1}{4}R$, we deduce

$$(5.48) \quad f(r) \lesssim g(r) + \left(\frac{r}{R}\right)^3 f\left(\frac{1}{4}R\right).$$

It remains to estimate the second right-hand side term. By definition of f and of w_\circ , we find

$$f\left(\frac{1}{4}R\right) \lesssim \int_{D_R} |\nabla w_\circ|^2 \lesssim \int_{D_R} |\nabla u|^2 + |\nabla \hat{u}(0)|^2 \int_{D_R} (1 + |\nabla \psi|^2).$$

Using the interior regularity estimate (5.43) with $\rho \simeq R$ and using the energy estimate (5.8), we note that

$$|\nabla \hat{u}(0)|^2 \lesssim \int_{D_R} |\nabla \hat{u}|^2 \lesssim \int_{D_R} |\nabla u|^2,$$

so that the above becomes

$$f\left(\frac{1}{4}R\right) \lesssim \left(\int_{D_R} (1 + |\nabla \psi|^2) \right) \int_{D_R} |\nabla u|^2.$$

Combining this with (5.48), and inserting the definition of f , g , and w_\circ , the claim (5.44) follows.

Substep 4.4. Estimate on the two-scale expansion error: for all ε with $0 < \varepsilon \ll 1$,

$$(5.49) \quad \int_{D_R} \mu_{R,\varepsilon}^2 |\nabla w|^2 \lesssim \left(\left(\frac{\rho}{R}\right)^\varepsilon + \left(\frac{\rho}{R}\right)^{-d-2} \frac{1}{R^2} \int_{D_R} (1 + |(\psi, \zeta, \nabla \psi, \Sigma \mathbb{1}_{\mathbb{R}^d \setminus \mathcal{J}})|^2) \right) \int_{D_R} |\nabla u|^2.$$

Starting point is (5.37): for all ε with $0 < \varepsilon \ll 1$,

$$\int_{D_R} \mu_{R,\varepsilon}^2 |\nabla w|^2 \lesssim \int_{D_R} (1 - \eta_R)^2 \mu_{R,\varepsilon}^2 |\nabla \hat{u}|^2 + \left(\sup_{D_R} |\nabla(\eta_R \nabla \hat{u})|^2 \right) \int_{D_R} (1 + |(\psi, \zeta, \nabla \psi, \Sigma \mathbb{1}_{\mathbb{R}^d \setminus \mathcal{J}})|^2).$$

Noting that the definition of η_R and $\mu_{R,\varepsilon}$ entails $(1 - \eta_R)^2 \mu_{R,\varepsilon}^2 \lesssim (\rho/R)^\varepsilon$, recalling that η_R is supported in $B_{R-\rho}$ and satisfies $|\nabla \eta_R| \lesssim \rho^{-1}$, using the interior regularity estimate (5.43), and using the energy estimate (5.8), the claim (5.49) follows.

Substep 4.5. Proof of (5.2). — Inserting the error bound (5.49) into (5.44), we find for all r, ρ with $4 \leq r, \rho \leq \frac{1}{4}R$,

$$(5.50) \quad \int_{B_r} |\nabla u - \nabla \hat{u}(0) - (\partial_E \hat{u})(0) \nabla \psi_E|^2 \lesssim \left(\left(\frac{r}{R}\right)^2 \int_{B_R} (1 + |\nabla \psi|^2) + \left(\frac{r}{R}\right)^{-d-2} \left(\left(\frac{\rho}{R}\right)^\varepsilon + \left(\frac{\rho}{R}\right)^{-d-2} \frac{1}{R^2} \int_{B_R} (1 + |(\psi, \zeta, \nabla \psi, \Sigma \mathbb{1}_{\mathbb{R}^d \setminus \mathcal{J}})|^2) \right) \right) \int_{B_R} |\nabla u|^2.$$

Next, we slightly reformulate this estimate by removing the dependence on $\nabla\psi$. For that purpose, we appeal to a Caccioppoli-type argument for ψ : arguing as in (5.45), now starting from equation (3.2), we find for all $K, R \geq 1$,

$$\int_{B_R} |\nabla\psi|^2 \lesssim \frac{1}{K} \int_{B_{2R}} |\nabla\psi|^2 + K \left(1 + \frac{1}{R^2} \int_{B_{2R}} \left| \psi - \int_{B_{2R}} \psi \right|^2 \right).$$

Iterating this estimate for some $K \gg 1$ large enough, and recalling that the ergodic theorem yields $\int_{B_R} |\nabla\psi|^2 \rightarrow \mathbb{E}[|\nabla\psi|^2] \lesssim 1$ almost surely as $R \uparrow \infty$, we deduce for all $R \geq 1$,

$$(5.51) \quad \int_{B_R} |\nabla\psi|^2 \lesssim 1 + \gamma_R^2,$$

where we recall that γ_R is defined in (5.3). Recalling the choice $\int_{B_R} (\psi, \zeta, \Sigma \mathbb{1}_{\mathbb{R}^d \setminus \mathcal{I}}) = 0$ in this proof, and appealing to the pressure estimate of Lemma 3.3 to further remove the dependence on Σ in (5.50), we obtain for all r, ρ with $4 \leq r, \rho \leq \frac{1}{4}R$,

$$\begin{aligned} \int_{B_r} |\nabla u - \nabla \hat{u}(0) - (\partial_E \hat{u})(0) \nabla \psi_E|^2 \\ \lesssim \left(\left(\frac{r}{R} \right)^2 + \left(\frac{r}{R} \right)^{-d-2} \left(\left(\frac{\rho}{R} \right)^\varepsilon + \left(\frac{\rho}{R} \right)^{-d-2} \gamma_R^2 \right) \right) \int_{B_R} |\nabla u|^2. \end{aligned}$$

It remains to optimize in ρ . If $\gamma_R \leq 1$, the choice $(\rho/R)^{d+2+\varepsilon} \simeq \gamma_R^2$ yields the conclusion (5.2) with $E_0 = \nabla \hat{u}(0)$ up to renaming ε . If $\gamma_R \geq 1$ or if $\frac{1}{4}R \leq r \leq R$, then the conclusion (5.2) trivially holds with $E_0 = 0$.

Step 5. Proof of the non-degeneracy property (5.4). — The upper bound in (5.4) follows from the Caccioppoli-type inequality (5.51), and it remains to establish the lower bound. Poincaré’s inequality and the triangle inequality yield

$$\begin{aligned} \left(\int_{B_{R/2}} |\nabla \psi_E + E|^2 \right)^{1/2} &\gtrsim \frac{1}{R} \left(\int_{B_{R/2}} \left| (\psi_E + Ex) - \int_{B_{R/2}} (\psi_E + Ex) \right|^2 \right)^{1/2} \\ &\geq \frac{1}{R} \left(\int_{B_{R/2}} |Ex|^2 \right)^{1/2} - \frac{1}{R} \left(\int_{B_{R/2}} \left| \psi_E - \int_{B_{R/2}} \psi_E \right|^2 \right)^{1/2} \\ &\gtrsim (1 - C\gamma_R) |E|, \end{aligned}$$

and the conclusion (5.4) follows. □

6. QUANTITATIVE HOMOGENIZATION

This section is devoted to the proof of Theorem 6.

Proof of Theorem 6. — First consider a cut-off function $\eta_\varepsilon \in C_c^\infty(\mathbb{R}^d; [0, 1])$ supported in U such that η_ε is constant inside the inclusions $\{\varepsilon I_n\}_n$, and $\eta_\varepsilon|_{\varepsilon I_n} = 0$ for all $n \notin \mathcal{N}_\varepsilon(U)$. In particular, $\mathcal{I}_\varepsilon(U)$ coincides with $\varepsilon \mathcal{I}$ in the support of η_ε . In addition, given R with $5 \leq R \leq 1/\varepsilon$ (to be later optimized depending on ε), we assume that $\eta_\varepsilon = 1$ in $U \setminus \partial_{\varepsilon R} U$ and $|\nabla \eta_\varepsilon| \lesssim (\varepsilon R)^{-1}$, where we use the notation $\partial_{\varepsilon R} U := \{x \in U : \text{dist}(x, \partial U) < \varepsilon R\}$ for the fattened boundary.

Step 1. Two-scale expansion and representation of the error. — Let $(u_\varepsilon, P_\varepsilon)$ denote the solution of the heterogeneous Stokes equation (2.9), and let (\bar{u}, \bar{P}) be the solution of the corresponding homogenized equation (2.10). The pressures P_ε and \bar{P} are chosen such that $\int_U P_\varepsilon \mathbb{1}_{\mathbb{R}^d \setminus \mathcal{J}_\varepsilon(U)} = \int_U \bar{P} = 0$. In terms of the corrector (ψ, Σ) , we consider the two-scale expansions

$$u_\varepsilon \rightsquigarrow \bar{u} + \varepsilon \eta_\varepsilon \psi_E(\cdot/\varepsilon) \partial_E \bar{u}, \quad P_\varepsilon \rightsquigarrow \bar{P} + \eta_\varepsilon \bar{\mathbf{b}} : \mathbf{D}(\bar{u}) + \eta_\varepsilon (\Sigma_E \mathbb{1}_{\mathbb{R}^d \setminus \mathcal{J}})(\cdot/\varepsilon) \partial_E \bar{u}.$$

Given arbitrary constants $P_{\varepsilon,*} \in \mathbb{R}$ and $\{P_{\varepsilon,n}\}_n \subset \mathbb{R}$ (that will be made explicit later in the proof), we modify the pressure P_ε into

$$P'_\varepsilon := (P_\varepsilon + P_{\varepsilon,*}) \mathbb{1}_{U \setminus \mathcal{J}_\varepsilon(U)} + \sum_{n \in \mathcal{N}_\varepsilon(U)} P_{\varepsilon,n} \mathbb{1}_{I_n},$$

and we then consider the following two-scale expansion errors in U ,

$$\begin{aligned} w_\varepsilon &:= u_\varepsilon - \bar{u} - \varepsilon \eta_\varepsilon \psi_E(\cdot/\varepsilon) \partial_E \bar{u}, \\ Q_\varepsilon &:= P'_\varepsilon - \bar{P} - \eta_\varepsilon \bar{\mathbf{b}} : \mathbf{D}(\bar{u}) - \eta_\varepsilon (\Sigma_E \mathbb{1}_{\mathbb{R}^d \setminus \mathcal{J}})(\cdot/\varepsilon) \partial_E \bar{u}. \end{aligned}$$

Arguing as in Substep 1.2 of the proof of Proposition 5.1, cf. (5.11), we find that $(w_\varepsilon, Q_\varepsilon)$ satisfies the following equation in the weak sense in U ,

$$\begin{aligned} (6.1) \quad & -\Delta w_\varepsilon + \nabla Q_\varepsilon = (\lambda - \mathbb{1}_{\mathcal{J}_\varepsilon(U)}) f \\ & - \sum_{n \in \mathcal{N}_\varepsilon(U)} \delta_{\varepsilon \partial I_n} \sigma(u_\varepsilon, P_\varepsilon + P_{\varepsilon,*} - P_{\varepsilon,n}) \nu - \operatorname{div}((\eta_\varepsilon \partial_E \bar{u}) J_E(\cdot/\varepsilon) \mathbb{1}_{\varepsilon \mathcal{J}}) \\ & + \operatorname{div} \left(2(1 - \eta_\varepsilon) (\operatorname{Id} - \bar{\mathbf{B}}) \mathbf{D}(\bar{u}) + 2\varepsilon (\psi_E \otimes_s - \zeta_E)(\cdot/\varepsilon) \nabla (\eta_\varepsilon \partial_E \bar{u}) \right. \\ & \left. - \varepsilon \operatorname{Id} (\psi_E(\cdot/\varepsilon) \cdot \nabla) (\eta_\varepsilon \partial_E \bar{u}) \right). \end{aligned}$$

In order to quantify the almost sure weak convergence $\mathbb{1}_{\mathcal{J}_\varepsilon(U)} \rightharpoonup \lambda$ in $L^2(U)$ in the first right-hand side term, we define a new corrector $\theta := \nabla \gamma$ as the unique solution of the following infinite-volume problem:

- Almost surely, $\theta = \nabla \gamma$ belongs to $L^2_{\text{loc}}(\mathbb{R}^d)^d$ and satisfies

$$\operatorname{div}(\theta) = \Delta \gamma = \mathbb{1}_{\mathcal{J}} - \lambda, \quad \text{in } \mathbb{R}^d.$$

- The field $\nabla \theta = \nabla^2 \gamma$ is stationary, has vanishing expectation, has finite second moment, and θ satisfies the anchoring condition $\int_B \theta = 0$ almost surely.

Under the mixing condition (Mix⁺), along the lines of the proof of Theorem 4.2 (but noting that no buckling is needed here as the corrector problem is linear with respect to randomness), the following moment bounds are easily checked to hold for all $q < \infty$,

$$(6.2) \quad \|\nabla \theta\|_{L^q(\Omega)} \lesssim_q 1, \quad \|\theta(x)\|_{L^q(\Omega)} \lesssim_q \mu_d(|x|).$$

In terms of this corrector, recalling that $\mathcal{J}_\varepsilon(U)$ coincides with $\varepsilon \mathcal{J}$ in the support of η_ε , the first right-hand side term in (6.1) can be decomposed as

$$\begin{aligned} (\lambda - \mathbb{1}_{\mathcal{J}_\varepsilon(U)}) f &= (\lambda - \mathbb{1}_{\mathcal{J}_\varepsilon(U)}) (1 - \eta_\varepsilon) f + (\lambda - \mathbb{1}_{\varepsilon \mathcal{J}}) \eta_\varepsilon f \\ &= (\lambda - \mathbb{1}_{\mathcal{J}_\varepsilon(U)}) (1 - \eta_\varepsilon) f - \operatorname{div}(\eta_\varepsilon f \otimes \varepsilon \theta(\cdot/\varepsilon)) + \nabla(\eta_\varepsilon f) \varepsilon \theta(\cdot/\varepsilon). \end{aligned}$$

Inserting this into (6.1), we are led to the following equation for $(w_\varepsilon, Q_\varepsilon)$ on U ,

$$\begin{aligned}
 (6.3) \quad & -\Delta w_\varepsilon + \nabla Q_\varepsilon = (\lambda - \mathbb{1}_{\mathcal{J}_\varepsilon(U)})(1 - \eta_\varepsilon)f + \nabla(\eta_\varepsilon f) \varepsilon \theta(\cdot/\varepsilon) \\
 & - \sum_{n \in \mathcal{N}_\varepsilon(U)} \delta_{\varepsilon \partial I_n} \sigma(u_\varepsilon, P_\varepsilon + P_{\varepsilon,*} - P_{\varepsilon,n}) \nu - \operatorname{div}((\eta_\varepsilon \partial_E \bar{u}) J_E(\cdot/\varepsilon) \mathbb{1}_{\varepsilon \mathcal{J}}) \\
 & + \operatorname{div} \left(2(1 - \eta_\varepsilon)(\operatorname{Id} - \bar{\mathbf{B}}) \operatorname{D}(\bar{u}) - \eta_\varepsilon f \otimes \varepsilon \theta(\cdot/\varepsilon) + 2\varepsilon(\psi_E \otimes_s - \zeta_E)(\cdot/\varepsilon) \nabla(\eta_\varepsilon \partial_E \bar{u}) \right. \\
 & \quad \left. - \varepsilon \operatorname{Id}(\psi_E(\cdot/\varepsilon) \cdot \nabla)(\eta_\varepsilon \partial_E \bar{u}) \right).
 \end{aligned}$$

Step 2. Conclusion. — We repeat the argument for (5.37) in Step 2 of the proof of Proposition 5.1, now without weight, starting from equation (6.3) instead of (5.11). More precisely, we truncate w_ε to make it affine in the inclusions, we test (6.3) with this truncated version of w_ε , we take advantage of boundary conditions, and we estimate the different terms. Compared to equation (5.11), the only new part here stems from the first two right-hand side terms in (6.3), for which we simply appeal to Poincaré’s inequality: as $w_\varepsilon \in H_0^1(U)^d$, we can estimate for any test function $g \in L^2(U)^d$,

$$\left| \int_U g \cdot w_\varepsilon \right| \leq \left(\int_U |g|^2 \right)^{1/2} \left(\int_U |w_\varepsilon|^2 \right)^{1/2} \lesssim \left(\int_U |g|^2 \right)^{1/2} \left(\int_U |\nabla w_\varepsilon|^2 \right)^{1/2}.$$

In this way, for a suitable choice of the constants $P_{\varepsilon,*}$ and $\{P_{\varepsilon,n}\}_n$, we arrive at the following estimate,

$$\begin{aligned}
 (6.4) \quad & \int_U |(\nabla w_\varepsilon, Q_\varepsilon)|^2 \lesssim \int_U (1 - \eta_\varepsilon)^2 |(f, \nabla \bar{u})|^2 + \varepsilon^2 \int_U |\theta(\cdot/\varepsilon)|^2 (|\nabla f|^2 + |\nabla \eta_\varepsilon|^2 |f|^2) \\
 & + \varepsilon^2 \int_U \left(1 + |(\psi, \zeta, \nabla \psi, \Sigma \mathbb{1}_{\mathbb{R}^d \setminus \mathcal{J}})(x/\varepsilon)|^2 \right) \left(\sup_{B_{4\varepsilon}(x)} (|\nabla^2 \bar{u}|^2 + |\nabla \eta_\varepsilon|^2 |\nabla \bar{u}|^2) \right) dx.
 \end{aligned}$$

Taking the $L^q(\Omega)$ norm, using corrector estimates of Theorem 2, as well as (6.2), recalling that $1 - \eta_\varepsilon$ and $\nabla \eta_\varepsilon$ are supported on the fattened boundary $\partial_{\varepsilon R} U$, noting that the latter has volume $|\partial_{\varepsilon R} U| \lesssim \varepsilon R$, and recalling that $|\nabla \eta_\varepsilon| \lesssim (\varepsilon R)^{-1}$, we deduce for all $q < \infty$,

$$(6.5) \quad \mathbb{E} \left[\left(\int_U |(\nabla w_\varepsilon, Q_\varepsilon)|^2 \right)^q \right]^{1/q} \lesssim_q (\varepsilon R + \varepsilon^2 \mu_d (1/\varepsilon)^2 (1/\varepsilon R)) \| (f, \nabla \bar{u}) \|_{W^{1,\infty}(U)}^2.$$

Next, decomposing

$$\begin{aligned}
 w_\varepsilon & := (u_\varepsilon - \bar{u} - \varepsilon \psi_E(\cdot/\varepsilon) \partial_E \bar{u}) + \varepsilon(1 - \eta_\varepsilon) \psi_E(\cdot/\varepsilon) \partial_E \bar{u}, \\
 Q_\varepsilon \mathbb{1}_{U \setminus \mathcal{J}_\varepsilon(U)} & := (P_\varepsilon + P_{\varepsilon,*} - \bar{P} - \bar{\mathbf{b}} : \operatorname{D}(\bar{u}) - (\Sigma_E \mathbb{1}_{\mathbb{R}^d \setminus \mathcal{J}})(\cdot/\varepsilon) \partial_E \bar{u}) \mathbb{1}_{U \setminus \mathcal{J}_\varepsilon(U)} \\
 & \quad + (1 - \eta_\varepsilon) (\bar{\mathbf{b}} : \operatorname{D}(\bar{u}) + (\Sigma_E \mathbb{1}_{\mathbb{R}^d \setminus \mathcal{J}})(\cdot/\varepsilon) \partial_E \bar{u}) \mathbb{1}_{U \setminus \mathcal{J}_\varepsilon(U)},
 \end{aligned}$$

we deduce for all $q < \infty$,

$$\begin{aligned}
 (6.6) \quad & \| u_\varepsilon - \bar{u} - \varepsilon \psi_E(\cdot/\varepsilon) \partial_E \bar{u} \|_{L^q(\Omega; H^1(U))}^2 \\
 & + \inf_{\kappa \in \mathbb{R}} \| P_\varepsilon - \bar{P} - \bar{\mathbf{b}} : \operatorname{D}(\bar{u}) - (\Sigma_E \mathbb{1}_{\mathbb{R}^d \setminus \mathcal{J}})(\cdot/\varepsilon) \partial_E \bar{u} - \kappa \|_{L^q(\Omega; L^2(U \setminus \mathcal{J}_\varepsilon(U)))}^2 \\
 & \lesssim_q (\varepsilon R + \varepsilon^2 \mu_d (1/\varepsilon)^2 (1/\varepsilon R)) \| (f, \nabla \bar{u}) \|_{W^{1,\infty}(U)}^2.
 \end{aligned}$$

Choosing $\varepsilon R = \varepsilon \mu_d(1/\varepsilon)$, and using the regularity theory for the steady Stokes equation (2.10), cf. [22, §IV], this yields the conclusion (2.12).

Finally, if f and \bar{u} are compactly supported in U , the cut-off function η_ε is equal to 1 identically in the support of $(f, \nabla \bar{u})$ for ε small enough. Hence, the terms involving $1 - \eta_\varepsilon$ and $\nabla \eta_\varepsilon$ drop in (6.4), and the bounds (6.5) and (6.6) are replaced by $\varepsilon^2 \mu_d(1/\varepsilon)^2$. \square

REFERENCES

- [1] S. ARMSTRONG, T. KUUSI & J.-C. MOURRAT – “Mesoscopic higher regularity and subadditivity in elliptic homogenization”, *Comm. Math. Phys.* **347** (2016), no. 2, p. 315–361.
- [2] ———, “The additive structure of elliptic homogenization”, *Invent. Math.* **208** (2017), no. 3, p. 999–1154.
- [3] ———, *Quantitative stochastic homogenization and large-scale regularity*, Grundlehren Math. Wiss., vol. 352, Springer, Cham, 2019.
- [4] S. N. ARMSTRONG & J.-P. DANIEL – “Calderón-Zygmund estimates for stochastic homogenization”, *J. Funct. Anal.* **270** (2016), no. 1, p. 312–329.
- [5] S. N. ARMSTRONG & J.-C. MOURRAT – “Lipschitz regularity for elliptic equations with random coefficients”, *Arch. Rational Mech. Anal.* **219** (2016), no. 1, p. 255–348.
- [6] S. N. ARMSTRONG & C. K. SMART – “Quantitative stochastic homogenization of convex integral functionals”, *Ann. Sci. École Norm. Sup. (4)* **49** (2016), no. 2, p. 423–481.
- [7] M. AVELLANEDA & F.-H. LIN – “Compactness methods in the theory of homogenization”, *Comm. Pure Appl. Math.* **40** (1987), no. 6, p. 803–847.
- [8] ———, “ L^p bounds on singular integrals in homogenization”, *Comm. Pure Appl. Math.* **44** (1991), no. 8-9, p. 897–910.
- [9] G. K. BATCHELOR – “Sedimentation in a dilute dispersion of spheres”, *J. Fluid Mech.* **52** (1972), no. 2, p. 245–268.
- [10] A. BRAIDES & A. GARRONI – “Homogenization of periodic nonlinear media with stiff and soft inclusions”, *Math. Models Methods Appl. Sci.* **5** (1995), no. 4, p. 543–564.
- [11] R. E. CAFLISCH & J. H. C. LUKE – “Variance in the sedimentation speed of a suspension”, *Phys. Fluids* **28** (1985), no. 3, p. 759–760.
- [12] T. DELMOTTE & J.-D. DEUSCHEL – “On estimating the derivatives of symmetric diffusions in stationary random environment, with applications to $\nabla \phi$ interface model”, *Probab. Theory Related Fields* **133** (2005), no. 3, p. 358–390.
- [13] L. DIENING, M. RŮŽIČKA & K. SCHUMACHER – “A decomposition technique for John domains”, *Ann. Acad. Sci. Fenn. Math.* **35** (2010), no. 1, p. 87–114.
- [14] M. DUERINCKX – “Effective viscosity of random suspensions without uniform separation”, 2020, [arXiv:2008.13188](https://arxiv.org/abs/2008.13188).
- [15] M. DUERINCKX & A. GLORIA – “Multiscale functional inequalities in probability: Concentration properties”, *ALEA Lat. Am. J. Probab. Math. Stat.* **17** (2020), no. 1, p. 133–157.
- [16] ———, “Multiscale functional inequalities in probability: Constructive approach”, *Ann. H. Lebesgue* **3** (2020), p. 825–872.
- [17] ———, “On Einstein’s effective viscosity formula”, 2020, [arXiv:2008.03837](https://arxiv.org/abs/2008.03837).
- [18] ———, “Corrector equations in fluid mechanics: Effective viscosity of colloidal suspensions”, *Arch. Rational Mech. Anal.* **239** (2021), p. 1025–1060.
- [19] ———, “Sedimentation of random suspensions and the effect of hyperuniformity”, *Ann. PDE* **8** (2022), no. 1, article no. 2 (66 pages).
- [20] M. DUERINCKX & F. OTTO – “Higher-order pathwise theory of fluctuations in stochastic homogenization”, *Stochastic Partial Differ. Equ. Anal. Comput.* **8** (2020), no. 3, p. 625–692.
- [21] A. EINSTEIN – “Eine neue Bestimmung der Moleküldimensionen”, *Ann. Physics* **19** (1906), no. 2, p. 289–306.
- [22] G. P. GALDI – *An introduction to the mathematical theory of the Navier-Stokes equations. Steady-state problems*, second ed., Springer Monographs in Math., Springer, New York, 2011.

- [23] F. W. GEHRING – “The L^p -integrability of the partial derivatives of a quasiconformal mapping”, *Acta Math.* **130** (1973), p. 265–277.
- [24] M. GIAQUINTA & G. MODICA – “Regularity results for some classes of higher order non linear elliptic systems”, *J. reine angew. Math.* **311/312** (1979), p. 145–169.
- [25] A. GLORIA – “A scalar version of the Caffisch-Luke paradox”, *Comm. Pure Appl. Math.* **74** (2021), no. 7, p. 1403–1452.
- [26] A. GLORIA, S. NEUKAMM & F. OTTO – “An optimal quantitative two-scale expansion in stochastic homogenization of discrete elliptic equations”, *ESAIM Math. Model. Numer. Anal.* **48** (2014), no. 2, p. 325–346.
- [27] ———, “Quantification of ergodicity in stochastic homogenization: optimal bounds via spectral gap on Glauber dynamics”, *Invent. Math.* **199** (2015), no. 2, p. 455–515.
- [28] ———, “A regularity theory for random elliptic operators”, *Milan J. Math.* **88** (2020), no. 1, p. 99–170.
- [29] ———, “Quantitative estimates in stochastic homogenization for correlated coefficient fields”, *Anal. PDE* **14** (2021), no. 8, p. 2497–2537.
- [30] A. GLORIA & F. OTTO – “An optimal variance estimate in stochastic homogenization of discrete elliptic equations”, *Ann. Probab.* **39** (2011), no. 3, p. 779–856.
- [31] ———, “An optimal error estimate in stochastic homogenization of discrete elliptic equations”, *Ann. Appl. Probab.* **22** (2012), no. 1, p. 1–28.
- [32] ———, “The corrector in stochastic homogenization: optimal rates, stochastic integrability, and fluctuations”, 2015, [arXiv:1510.08290](https://arxiv.org/abs/1510.08290).
- [33] ———, “Quantitative results on the corrector equation in stochastic homogenization”, *J. Eur. Math. Soc. (JEMS)* **19** (2017), p. 3489–3548.
- [34] R. M. HÖFER – “Convergence of the method of reflections for particle suspensions in Stokes flows”, *J. Differential Equations* **297** (2021), p. 81–109.
- [35] V. V. JIKOV – “Some problems of extension of functions arising in connection with the homogenization theory”, *Differ. Uravn.* **26** (1990), no. 1, p. 39–51.
- [36] V. V. JIKOV, S. M. KOZLOV & O. A. OLEĬNIK – *Homogenization of differential operators and integral functionals*, Springer-Verlag, Berlin, 1994.
- [37] M. JOSIEN & F. OTTO – “The annealed Calderón-Zygmund estimate as convenient tool in quantitative stochastic homogenization”, 2020, [arXiv:2005.08811](https://arxiv.org/abs/2005.08811).
- [38] D. L. KOCH & E. S. G. SHAQFEH – “Screening in sedimenting suspensions”, *J. Fluid Mech.* **224** (1991), p. 275–303.
- [39] P. C. KUNSTMANN & L. WEIS – “Maximal L_p -regularity for parabolic equations, Fourier multiplier theorems and H^∞ -functional calculus”, in *Functional analytic methods for evolution equations*, Lect. Notes in Math., vol. 1855, Springer, Berlin, 2004, p. 65–311.
- [40] D. MARAHRENS & F. OTTO – “Annealed estimates on the Green function”, *Probab. Theory Related Fields* **163** (2015), no. 3-4, p. 527–573.
- [41] Z. SHEN – “The L^p boundary value problems on Lipschitz domains”, *Adv. Math.* **216** (2007), p. 212–254.
- [42] ———, “The Calderón-Zygmund lemma revisited”, in *Lectures on the analysis of nonlinear partial differential equations. Part 2*, Morningside Lect. Math., vol. 2, Int. Press, Somerville, MA, 2012, p. 203–224.
- [43] V. V. ZHIKOV – “Averaging of functionals in the calculus of variations and elasticity”, *Math. USSR-Izv.* **29** (1987), p. 33–66.

Manuscript received 10th March 2021
accepted 11th July 2022

MITIA DUERINCKX, Université Paris-Saclay, CNRS, Laboratoire de Mathématiques d'Orsay
91400 Orsay, France
& University of California, Los Angeles, Department of Mathematics
Los Angeles CA 90095, USA
& Université Libre de Bruxelles, Département de Mathématique
1050 Brussels, Belgium
E-mail : mitia.duerinckx@u-psud.fr

ANTOINE GLORIA, Sorbonne Université, CNRS, Université de Paris, Laboratoire Jacques-Louis Lions
75005 Paris, France
& Institut Universitaire de France
& Université Libre de Bruxelles, Département de Mathématique
1050 Brussels, Belgium
E-mail : gloria@ljl1.math.upmc.fr
Url : <https://www.ljl1.math.upmc.fr/~gloria/index.html>