



# *J*ournal de l'École polytechnique *Mathématiques*


Thierry DE PAUW

Density estimate from below in relation to a conjecture of A. Zygmund on Lipschitz differentiation

Tome 9 (2022), p. 1473-1512.

<https://doi.org/10.5802/jep.211>

© Les auteurs, 2022.



Cet article est mis à disposition selon les termes de la licence  
LICENCE INTERNATIONALE D'ATTRIBUTION CREATIVE COMMONS BY 4.0.  
<https://creativecommons.org/licenses/by/4.0/>

L'accès aux articles de la revue « Journal de l'École polytechnique — Mathématiques » (<http://jep.centre-mersenne.org/>) implique l'accord avec les conditions générales d'utilisation (<http://jep.centre-mersenne.org/legal/>).

Publié avec le soutien  
du Centre National de la Recherche Scientifique



Publication membre du  
Centre Mersenne pour l'édition scientifique ouverte  
[www.centre-mersenne.org](http://www.centre-mersenne.org)  
e-ISSN : 2270-518X

## DENSITY ESTIMATE FROM BELOW IN RELATION TO A CONJECTURE OF A. ZYGMUND ON LIPSCHITZ DIFFERENTIATION

BY THIERRY DE PAUW

ABSTRACT. — Let  $G(n, m)$  be the Grassmannian consisting of  $m$ -dimensional vector subspaces of  $\mathbb{R}^n$ , let  $L^n$  be the Lebesgue measure in  $\mathbb{R}^n$ , let  $H^m$  be the  $m$ -dimensional Hausdorff measure in  $\mathbb{R}^n$ , and let  $\alpha(m) = L^m(\mathbf{B}(0, 1))$  be the Lebesgue measure of the Euclidean unit ball of  $\mathbb{R}^m$ . We establish that, if  $A \subseteq \mathbb{R}^n$  is Borel measurable and  $\mathbf{W}_0 : A \rightarrow G(n, m)$  is Lipschitzian, then

$$\limsup_{r \rightarrow 0^+} \frac{H^m[A \cap \mathbf{B}(x, r) \cap (x + \mathbf{W}_0(x))]}{\alpha(m)r^m} > \frac{1}{2^n},$$

for  $L^n$ -almost every  $x \in A$ . In particular, it follows that  $A$  is  $L^n$ -negligible if and only if  $H^m(A \cap (x + \mathbf{W}_0(x))) = 0$ , for  $L^n$ -almost every  $x \in A$ .

RÉSUMÉ (Borne inférieure de densité en lien avec une conjecture de A. Zygmund sur des bases de dérivation à variation lipschitzienne)

On désigne par  $G(n, m)$  la grassmannienne constituée des sous-espaces vectoriels de dimension  $m$  dans  $\mathbb{R}^n$ , par  $L^n$  la mesure de Lebesgue dans  $\mathbb{R}^n$ , par  $H^m$  la mesure de Hausdorff  $m$ -dimensionnelle dans  $\mathbb{R}^n$  et par  $\alpha(m) = L^m(\mathbf{B}(0, 1))$  la mesure de Lebesgue de la boule euclidienne unité de  $\mathbb{R}^m$ . Nous montrons que si  $A \subseteq \mathbb{R}^n$  est borélien et  $\mathbf{W}_0 : A \rightarrow G(n, m)$  est lipschitzien, alors

$$\limsup_{r \rightarrow 0^+} \frac{H^m[A \cap \mathbf{B}(x, r) \cap (x + \mathbf{W}_0(x))]}{\alpha(m)r^m} > \frac{1}{2^n},$$

pour  $L^n$ -presque tout  $x \in A$ . Il en résulte en particulier que  $A$  est  $L^n$ -négligeable si et seulement si  $H^m(A \cap (x + \mathbf{W}_0(x))) = 0$ , pour  $L^n$ -presque tout  $x \in A$ .

### CONTENTS

1. Foreword.....	1474
2. Sketch of proof.....	1475
3. Preliminaries.....	1479
4. Common setting.....	1487
5. Two fibrations.....	1487
6. Upper bound for $\Upsilon_E \mathbf{W}$ and $Z_E \mathbf{W}$ .....	1498
7. Lower bound for $\Upsilon_E \mathbf{W}$ and $Z_E \mathbf{W}$ .....	1500
8. Proof of the theorems.....	1507
References.....	1511

MATHEMATICAL SUBJECT CLASSIFICATION (2020). — 28A75, 26B15.

KEYWORDS. — Lebesgue measure, Nikodým set, negligible set, derivation basis, Zygmund conjecture, Lipschitz differentiation.

The author was partially supported by the Science and Technology Commission of Shanghai (No. 18dz2271000).

## 1. FOREWORD

Let  $S_x$  be the set of squares centered at  $x \in \mathbb{R}^2$ . The Lebesgue density theorem states that if  $f : \mathbb{R}^2 \rightarrow \mathbb{R}$  is Lebesgue summable, then

$$(*) \quad f(x) = \lim_{\substack{S \in S_x \\ \text{diam } S \rightarrow 0^+}} \int_S f dL^2,$$

for  $L^2$ -almost every  $x \in \mathbb{R}^2$ . This consequence of the Vitali covering theorem fails if  $S_x$  is replaced with  $R_x$ , the set of rectangles centered at  $x$  of arbitrary direction and eccentricity. It fails even for some indicator function  $f = \mathbb{1}_A$ . Indeed, Nikodým [12] defined a set  $A \subseteq [0, 1] \times [0, 1]$  of full measure with the following property: For every  $x \in A$ , there exists a line  $L(x)$  such that  $A \cap L(x) = \{x\}$ . In fact, Nikodým's example shows that the Lebesgue density theorem fails if we replace  $S_x$  with  $R_x^L$ , the set of rectangles of arbitrary eccentricity, centered at  $x$ , and one side of which is parallel to  $L(x)$ . Furthermore [9, Chap. IV, Th. 3.5], replacing  $A$  by a nonnegligible subset of  $A$ , one may choose  $L$  to be continuous with respect to  $x$ . Yet, if  $L$  is constant, then the Lebesgue density theorem with respect to  $R_x^L$  holds, for  $p$ -summable functions  $f$ ,  $1 < p < \infty$ , by virtue of a theorem of Zygmund, though the corresponding version of the Vitali covering theorem fails, according to an example of H. Bohr [9, Chap. IV, Th. 1.1]. This raises the question: What regularity condition of  $x \mapsto L(x)$  guarantees that the Lebesgue density theorem holds with respect to  $R_x^L$  for, say, functions that are square summable? The following version is a conjecture reportedly [11] attributed to Zygmund. *Assume that  $x \mapsto L(x)$  is a Lipschitzian field of lines, with  $x \in L(x)$ , and  $f : \mathbb{R}^2 \rightarrow \mathbb{R}$  is square summable. Is it true that*

$$f(x) = \lim_{r \rightarrow 0^+} \int_{L(x) \cap B(x,r)} f dH^1,$$

for  $L^2$ -almost every  $x \in \mathbb{R}^2$ ? Here,  $H^1$  is the 1-dimensional Hausdorff measure.

E. M. Stein raised the singular integral variant of this conjecture. Both have received much attention from the harmonic analysis community. To the author's knowledge, most results recorded so far in the literature, via the maximal function approach, assume some extra regularity property of  $L$  – namely, that  $L$  be  $C^1$  together with a hypothesis on the variation of the derivative – see, for instance, [1] and [11].

In this paper we offer a novel approach – a kind of change of variable based on an appropriate fibration and the coarea formula. This allows for treating the case when  $L$  is Lipschitzian, without further restriction. We obtain the following lower density bound for indicator functions. *Let  $x \mapsto L(x)$  be a Lipschitzian field of lines, with  $x \in L(x)$ , and let  $A \subseteq \mathbb{R}^2$  be Lebesgue measurable. Then,*

$$\limsup_{r \rightarrow 0^+} \frac{H^1(A \cap B(x,r) \cap L(x))}{2r} > \frac{1}{4},$$

for  $L^2$ -almost every  $x \in A$ . Incidentally, the following corollary – a nonparallel version of Fubini theorem – seems to be new as well. *The set  $A$  is Lebesgue negligible if and only if  $H^1(A \cap L(x)) = 0$ , for  $L^2$ -almost every  $x \in \mathbb{R}^2$ .*

Our results hold in any dimension and codimension.

## 2. SKETCH OF PROOF

Let  $A$  be a subset of Euclidean space  $\mathbb{R}^n$ ,  $n > 2$ , and let  $L^n$  be the Lebesgue outer measure. We start by considering the following weak question: Can one tell whether  $A$  is Lebesgue negligible from the knowledge only of its trace on each member of some given collection of “lower dimensional” subsets  $\Gamma_i \subseteq \mathbb{R}^n$ ,  $i \in I$ . One expects that if  $A \cap \Gamma_i$  is “negligible in the dimension of  $\Gamma_i$ ”, for each  $i \in I$ , then  $L^n(A) = 0$ . Of course, a necessary condition is that the sets  $\Gamma_i$  cover almost all of  $A$ , i.e.,  $L^n(A \cap \bigcup_{i \in I} \Gamma_i) = 0$ . Consider, for instance,  $n = 2$ ,  $I = \mathbb{R}$ , and  $\Gamma_t = \{t\} \times \mathbb{R}$ , for  $t \in \mathbb{R}$ , the collection of all vertical lines in the plane. It is not true in general that if  $A \subseteq \mathbb{R}^2$  and  $A \cap \Gamma_t$  is a singleton, for each  $t \in \mathbb{R}$ , then  $L^2(A) = 0$ . There exist, indeed, functions  $f : \mathbb{R} \rightarrow \mathbb{R}$  whose graph  $A$  satisfies  $L^2(A) > 0$  – see, e.g., [10, Chap. 2, Th. 4] for an example due to Sierpiński. In order to rule out such examples, we will henceforth assume that  $A \subseteq \mathbb{R}^n$  is Borel measurable. In that case, the theorem of Fubini, together with the invariance of the Lebesgue measure under orthogonal transformations, imply the following. Given an integer  $1 \leq m \leq n - 1$ , if  $(\Gamma_i)_{i \in I}$  is the collection of all  $m$ -dimensional affine subspaces of  $\mathbb{R}^n$  of some fixed direction, and if  $H^m(A \cap \Gamma_i) = 0$  for all  $i \in I$ , then  $L^n(A) = 0$ . Here,  $H^m$  denotes the  $m$ -dimensional Hausdorff measure. A special feature of this collection  $(\Gamma_i)_{i \in I}$  is that it partitions  $\mathbb{R}^n$ , its members being the level sets  $f^{-1}\{y\}$ ,  $y \in \mathbb{R}^{n-m}$ , of a “nice map”  $f : \mathbb{R}^n \rightarrow \mathbb{R}^{n-m}$ , indeed, an orthogonal projection. This is an occurrence of the following more general situation when  $f$  and its leaves  $f^{-1}\{y\}$  are allowed to be nonlinear. The coarea formula due to Federer [7] asserts that if  $f : \mathbb{R}^n \rightarrow \mathbb{R}^{n-m}$  is Lipschitzian and  $A \subseteq \mathbb{R}^n$  is Borel measurable, then

$$\int_A Jf(x) dL^n(x) = \int_{\mathbb{R}^{n-m}} H^m(A \cap f^{-1}\{y\}) dL^{n-m}(y).$$

Thus, if the Jacobian coarea factor  $Jf$  is positive,  $L^n$ -almost everywhere in  $A$ , then the collection  $(f^{-1}\{y\})_{y \in \mathbb{R}^{n-m}}$  is suitable for detecting whether or not  $A$  is Lebesgue null. At  $L^n$ -almost all  $x \in \mathbb{R}^n$ , the map  $f$  is differentiable, by virtue of Rademacher’s theorem, and

$$Jf(x) = \sqrt{|\det(Df(x) \circ Df(x)^*)|} = \|\wedge_{n-m} Df(x)\|,$$

see [5, Chap. 3, §4] and [8, 3.2.1 & 3.2.11].

In this paper, we focus on the case when  $\Gamma_i$ ,  $i \in I$ , are affine subspaces of  $\mathbb{R}^n$ , but not necessarily members of a partition of the ambient space. Specifically, we assume that with each  $x \in \mathbb{R}^n$  is associated an  $m$ -dimensional affine subspace  $\mathbf{W}(x)$  of  $\mathbb{R}^n$  containing  $x$ . Given a Borel set  $A \subseteq \mathbb{R}^n$ , the question whether

$$(1) \quad \text{If } H^m(A \cap \mathbf{W}(x)) = 0, \text{ for all } x \in A, \text{ then } L^n(A) = 0,$$

has a negative answer, in view of Nikodým’s set  $A \subseteq \mathbb{R}^2$  evoked in the previous section. Indeed, corresponding to this set  $A$ , there exists a field of lines  $x \mapsto \mathbf{W}(x)$  such that  $A \cap \mathbf{W}(x) = \{x\}$ , for all  $x \in A$ . In this context, a selection theorem due to von Neumann implies that (possibly considering a smaller, non Lebesgue null, Borel

subset of  $A$ ) the correspondence  $x \mapsto \mathbf{W}(x)$  can be chosen to be Borel measurable (see 3.19) and, in turn, it can be chosen to be continuous, according to Lusin’s theorem. Nonetheless, when  $\mathbf{W}$  is Lipschitzian, the situation improves, as illustrated in our theorem below;  $G(n, m)$  is the Grassmannian manifold consisting of  $m$ -dimensional vector subspaces of  $\mathbb{R}^n$ .

**THEOREM.** — Assume  $\mathbf{W}_0 : \mathbb{R}^n \rightarrow G(n, m)$  is Lipschitzian and  $A \subseteq \mathbb{R}^n$  is Borel measurable. The following are equivalent.

- (A)  $L^n(A) = 0$ .
- (B) For  $L^n$  almost every  $x \in A$ ,  $H^m(A \cap (x + \mathbf{W}_0(x))) = 0$ .
- (C) For  $L^n$  almost every  $x \in \mathbb{R}^n$ ,  $H^m(A \cap (x + \mathbf{W}_0(x))) = 0$ .

As should be apparent from the discussion above, one difficulty stands with the fact that the affine  $m$ -planes  $\mathbf{W}(x) = x + \mathbf{W}_0(x)$  may not be disjointed. Nevertheless, they locally are, in the following sense. Given  $x_0 \in A$  there exist a neighborhood  $U$  of  $x_0$  and Lipschitzian maps  $\mathbf{w}_1, \dots, \mathbf{w}_m : U \rightarrow \mathbb{R}^n$  so that  $\mathbf{w}_1(x), \dots, \mathbf{w}_m(x)$  is an orthonormal frame spanning  $\mathbf{W}_0(x)$ , for  $x \in U$ , and the map

$$\Phi : (V_{x_0} \cap U) \times \mathbb{R}^m \longrightarrow \mathbb{R}^n : (\xi, t) \longmapsto \xi + \sum_{i=1}^m t_i \mathbf{w}_i(\xi)$$

is a lipeomorphism of a neighborhood of  $x_0$  onto its image – here,  $V_{x_0} = x_0 + V$  and  $V \in G(n, n - m)$  is close to  $\mathbf{W}_0(x_0)^\perp$ . This, and applications of Fubini’s theorem, yield (B)  $\Rightarrow$  (A) in the theorem above (see 8.2).

However, we aim at obtaining a quantitative version of this result, that the change of variable just described does not seem to provide. A natural route is to reduce the problem to applying the coarea formula, by spreading out the  $\mathbf{W}(x)$ ’s in a disjointed way, in a higher dimensional space – i.e., adding a variable  $u \in \mathbf{W}(x)$  to the given  $x \in \mathbb{R}^n$  and considering  $\mathbf{W}(x)$  as a fiber above the base space  $\mathbb{R}^n$ . We thus define

$$\Sigma = \mathbb{R}^n \times \mathbb{R}^n \cap \{(x, u) : x \in E \text{ and } u \in \mathbf{W}(x)\},$$

where  $E \subseteq \mathbb{R}^n$  is Borel measurable. This set is  $(n + m)$ -rectifiable, owing to the Lipschitz continuity of  $\mathbf{W}$ . It is convenient to assume that  $L^n(E) < \infty$ , so that

$$(2) \quad \phi_E(B) = \int_E H^m(B \cap \mathbf{W}(x)) dL^n(x),$$

$B \subseteq \mathbb{R}^n$ , is a locally finite Borel measure 3.16. Now,  $\Sigma$  was precisely set up so that, for each  $x \in E$ ,

$$H^m(\Sigma \cap \pi_2^{-1}(B) \cap \pi_1^{-1}\{x\}) = H^m(B \cap \mathbf{W}(x)),$$

where  $\pi_1$  and  $\pi_2$  denote the projections of  $\mathbb{R}^n \times \mathbb{R}^n$  to the  $x$  and  $u$  variable, respectively. Abbreviating  $\Sigma_B = \Sigma \cap \pi_2^{-1}(B)$ , the coarea formula yields

$$(3) \quad \phi_E(B) = \int_{\Sigma_B} J_{\Sigma \pi_1} dH^{n+m}.$$

A simple calculation 5.4 shows that  $J_{\Sigma}\pi_2 > 0$ ,  $H^{n+m}$ -almost everywhere on  $\Sigma_B$ . Since also

$$(4) \quad \int_{\Sigma_B} J_{\Sigma}\pi_2 dH^{n+m} = \int_B H^m(\Sigma_B \cap \pi_2^{-1}\{u\}) dL^n(u),$$

the implication (A)  $\Rightarrow$  (C) above should now be clear: Letting  $B = A$  and  $E = \mathbf{B}(0, R)$ , one infers from hypothesis (A) and (4) that  $H^{n+m}(\Sigma_B) = 0$ , whence,  $\phi_{\mathbf{B}(0,R)}(A) = 0$ , by (3), and, in turn, conclusion (C) ensues from (2).

In order to establish that (B)  $\Rightarrow$  (A) by using this new change of variable, we need to observe 5.2 that  $J_{\Sigma}\pi_1 > 0$ ,  $H^{n+m}$ -almost everywhere, and, ideally, to show that  $H^m(\Sigma_B \cap \pi_2^{-1}\{u\}) > 0$ , for almost every  $u \in B$ . This last part involves some difficulty. To understand this, we let  $m = n - 1$ , in order to keep the notations short. Now  $u \in \mathbf{W}(x)$  iff  $u - x \in \mathbf{W}_0(x)$  iff  $\langle \mathbf{v}_0(x), x - u \rangle = 0$ , where  $\mathbf{v}_0(x) \in \mathbf{W}_0(x)^\perp$  is, say, a unit vector. Abbreviating  $g_u(x) = \langle \mathbf{v}_0(x), x - u \rangle$ , we infer that

$$H^m(\Sigma_B \cap \pi_2^{-1}\{u\}) = H^m(E \cap g_u^{-1}\{0\}).$$

The problem remains that two of the nonlinear  $m$ -sets  $E \cap g_u^{-1}\{0\}$  and  $E \cap g_{u'}^{-1}\{0\}$  may intersect, thereby preventing another application of the coarea formula when looking out for their lower bound. Yet, we already know that

$$\phi_E(B) = \int_B Z_E \mathbf{W} dL^n,$$

where  $Z_E \mathbf{W}$  is a Radon-Nikodým derivative (see 5.6) and also that  $(Z_E \mathbf{W})(u)$  is comparable to  $H^m(E \cap g_u^{-1}\{0\})$  (see 5.8). Adding an extra variable  $y$  to the fibered space  $\Sigma$ , 5.10, we improve on this by showing that

$$(Z_E \mathbf{W})(u) > \boldsymbol{\eta}(n, m) \liminf_j \int_{-j^{-1}}^{j^{-1}} H^m(E \cap g_u^{-1}\{y\}) dL^1(y) = \boldsymbol{\eta}(n, m)(Y_E \mathbf{W})(u),$$

where the last equality defines  $Y_E \mathbf{W}$ , and  $\boldsymbol{\eta}(n, m) > 0$  is a lower bound for the coarea Jacobian factor of the restriction to the fibered space of the projection  $(x, u, y) \mapsto (x, y)$  (see 5.12 and 5.16). We are reduced to showing that  $Y_E \mathbf{W} > 0$ , almost everywhere. The reason why this holds is the following. Fix a Borel set  $Z \subseteq \mathbb{R}^n$ ,  $x_0 \in \mathbb{R}^n$  and  $r > 0$ . Let  $\mathbf{C}_{\mathbf{W}}(x_0, r)$  be the cylindrical box consisting of those  $x \in \mathbb{R}^n$  such that  $|P_{\mathbf{W}_0(x_0)}(x - x_0)| \leq r$  and  $|P_{\mathbf{W}_0(x_0)^\perp}(x - x_0)| \leq r$ . We want to find a lower bound for

$$\int_{Z \cap \mathbf{C}_{\mathbf{W}}(x_0, r)} (Y_{Z \cap \mathbf{C}_{\mathbf{W}}(x_0, r)})(u) dL^n(u).$$

To this end, we fix  $z \in \mathbf{W}_0(x_0) \cap \mathbf{B}(0, r)$  and we let

$$V_z = \mathbb{R}^n \cap \{x_0 + z + s\mathbf{v}_0(x_0) : -r \leq s \leq r\}$$

denote the corresponding vertical line segment. According to Fubini's theorem we are reduced to estimating

$$\int_{V_z} (Y_{Z \cap \mathbf{C}_{\mathbf{W}}(x_0, r)})(u) dH^1(u).$$

By virtue of Vitali’s covering theorem, we can find a disjointed family of line segments  $I_1, I_2, \dots$ , covering almost all  $V_z$ , such that the above integral nearly equals

$$\begin{aligned} \sum_k H^1(I_k) \int_{I_k} H^m(Z \cap \mathbf{C}_W(x_0, r) \cap g_{u_k}^{-1}\{y\}) dL^1(y) \\ \cong \sum_k \int_{Z \cap \mathbf{C}_W(x_0, r) \cap g_{u_k}^{-1}(I_k)} |\nabla g_{u_k}(x)| dL^n(x) \cong L^n(Z \cap \mathbf{C}_W(x_0, r)), \end{aligned}$$

where the first near equality follows from the coarea formula, and the second one because  $|\nabla g_{u_k}| \cong 1$  at small scales (see 3.12) and the “nonlinear horizontal stripes”  $g_{u_k}^{-1}(I_k)$  are nearly pairwise disjoint. Verification of these claims takes up sections 6 and 7. Now, we reach a contradiction if  $Z = \mathbb{R}^n \cap \{Y_E \mathbf{W} = 0\}$  is assumed to have  $L^n(Z) > 0$ , and  $x_0$  is a point of density of  $Z$ .

Along this path, we have, in fact, achieved a quantitative version of these observations. Indeed, assuming that

$$\limsup_{r \rightarrow 0^+} \frac{H^m(A \cap \mathbf{B}(x, r) \cap \mathbf{W}(x))}{\alpha(m)r^m} < \frac{\eta(n, m)}{2^m},$$

for  $L^n$  positively many  $x \in A$ , we start by choosing  $E \subseteq A$  with the same property, whose diameter is small enough for the various estimates above to hold, and we refer to Egoroff’s theorem to select  $\varepsilon > 0$  so that the set

$$Z = E \cap \left\{ \limsup_{r \rightarrow 0^+} \frac{H^m(A \cap \mathbf{B}(x, r) \cap \mathbf{W}(x))}{\alpha(m)r^m} < (1 - \varepsilon) \frac{\eta(n, m)}{(2 + \varepsilon)^m} \right\}$$

has positive  $L^n$  measure. Thus, for  $0 < \hat{\varepsilon} \ll \varepsilon$ , there exists  $x_0 \in Z$  and there exists  $r > 0$  as small as we please so that

$$\begin{aligned} (1 - \hat{\varepsilon})L^n(\mathbf{C}_W(x_0, r)) \\ \leq L^n(Z \cap \mathbf{C}_W(x_0, r)) \cdot \int_{Z \cap \mathbf{C}_W(x_0, r)} \frac{Y_{Z \cap \mathbf{C}_W(x_0, r)} \mathbf{W}(u)}{\alpha(m)r^m} dL^n(u) \\ \cdot \frac{1}{\eta(n, m)} \int_{Z \cap \mathbf{C}_W(x_0, r)} \frac{H^m(Z \cap \mathbf{C}_W(x_0, r) \cap \mathbf{W}(x))}{\alpha(m)r^m} dL^n(x) \\ \leq \frac{1}{\eta(n, m)} \int_{Z \cap \mathbf{C}_W(x_0, r)} \frac{H^m(Z \cap \mathbf{B}(x, 2r + o(r)) \cap \mathbf{W}(x))}{\alpha(m)r^m} dL^n(x) \\ \cdot \frac{1}{\eta(n, m)} \int_{Z \cap \mathbf{C}_W(x_0, r)} (1 - \varepsilon) \frac{\eta(n, m)}{(2 + \varepsilon)^m} dL^n(x) \leq (1 - C\varepsilon)L^n(\mathbf{C}_W(x_0, r)), \end{aligned}$$

where the symbol  $\cdot$  means that we loose a multiplicative factor  $\lambda > 1$  as close as we wish to 1, according to the scale  $r$ . The above contradiction proves our main result 8.4, namely,

**THEOREM.** — Assume  $\mathbf{W}_0 : \mathbb{R}^n \rightarrow \mathbb{G}(n, m)$  is Lipschitzian and  $A \subseteq \mathbb{R}^n$  is Borel measurable. Abbreviating  $\mathbf{W}(x) = x + \mathbf{W}_0(x)$ , it follows that

$$\limsup_{r \rightarrow 0^+} \frac{H^m(A \cap \mathbf{B}(x, r) \cap \mathbf{W}(x))}{\alpha(m)r^m} > \frac{1}{2^n},$$

for  $L^n$ -almost every  $x \in A$ .

*Acknowledgements.* — I extend my warm thanks to Jean-Christophe Léger for carefully reading an early version of the manuscript.

### 3. PRELIMINARIES

3.1. — In this paper,  $1 \leq m \leq n - 1$  are integers. The ambient space is  $\mathbb{R}^n$ . The canonical inner product of  $x, x' \in \mathbb{R}^n$  is denoted  $\langle x, x' \rangle$  and the corresponding Euclidean norm of  $x$  is  $|x|$ . If  $S \subseteq \mathbb{R}^n$ , we let  $\mathcal{B}(S)$  be the  $\sigma$ -algebra of Borel measurable subsets of  $S$ .

3.2. HAUSDORFF MEASURE. — We let  $L^n$  be the Lebesgue outer measure in  $\mathbb{R}^n$  and  $\alpha(n) = L^n(\mathbf{B}(0, 1))$ . For  $S \subseteq \mathbb{R}^n$ , we abbreviate  $\zeta^m(S) = \alpha(m)2^{-m}(\text{diam } S)^m$ . Given  $0 < \delta < \infty$ , we call  $\delta$ -cover of  $A \subseteq \mathbb{R}^n$  a finite or countably infinite family  $(S_j)_{j \in J}$  of subsets of  $\mathbb{R}^n$  such that  $A \subseteq \bigcup_{j \in J} S_j$  and  $\text{diam } S_j < \delta$ , for every  $j \in J$ . We define

$$H_\delta^m(A) = \inf \left\{ \sum_{j \in J} \zeta^m(S_j) : (S_j)_{j \in J} \text{ is a } \delta\text{-cover of } A \right\}$$

and  $H^m(A) = \lim_{\delta \rightarrow 0^+} H_\delta^m(A) = \sup_{\delta > 0} H_\delta^m(A)$ . Thus,  $H^m$  is the  $m$ -dimensional Hausdorff outer measure in  $\mathbb{R}^n$ .

(1) *If  $(K_k)_k$  is a sequence of nonempty compact subsets of  $\mathbb{R}^n$  converging in Hausdorff distance to  $K$ , then  $H_\infty^m(K) > \limsup_k H_\infty^m(K_k)$ .*

Given  $\varepsilon > 0$ , choose an  $\infty$ -cover  $(S_j)_{j \in \mathbb{N}}$  of  $K$  such that  $H_\infty^m(K) + \varepsilon > \sum_j \zeta^m(S_j)$ . Since  $\lim_{r \rightarrow 0^+} \zeta^m(\mathbf{U}(S_j, r)) = \zeta^m(S_j)$ , for each  $j \in \mathbb{N}$ , we can choose an open set  $U_j$  containing  $S_j$  such that  $\zeta^m(U_j) < \varepsilon 2^{-j} + \zeta^m(S_j)$ . Since  $U = \bigcup_j U_j$  is open, there exists a positive integer  $k_0$  such that  $K_k \subseteq U$ , whenever  $k > k_0$ . Thus, in that case  $(U_j)_j$  is an  $\infty$ -cover of  $K_k$  and, therefore,  $H_\infty^m(K_k) < \sum_j \zeta^m(U_j) < 2\varepsilon + H_\infty^m(K)$ . Take the limit superior of the left hand side, as  $k \rightarrow \infty$ , and then let  $\varepsilon \rightarrow 0$ .

(2) *For all  $A \subseteq \mathbb{R}^m$ , one has  $L^m(A) = H^m(A) = H_\infty^m(A)$ .*

It suffices to note that  $H^m(A) > H_\infty^m(A) > L^m(A) > H^m(A)$ . The first inequality is trivial; the second one follows from the isodiametric inequality [8, 2.10.33]; the last one is a consequence of the Vitali covering theorem [5, Chap. 2, §2, Th. 2].

(3) *If  $W \subseteq \mathbb{R}^n$  is an  $m$ -dimensional affine subspace and  $A \subseteq W$ , then  $H^m(A) = H_\infty^m(A)$ .*

Let  $\mathfrak{H}^m$  denote the  $m$ -dimensional Hausdorff outer measure in the metric space  $W$ . In other words,

$$\mathfrak{H}_\delta^m(A) = \inf \left\{ \sum_{j \in J} \zeta^m(S_j) : (S_j)_{j \in J} \text{ is a } \delta\text{-cover of } A \text{ and } S_j \subseteq W, \text{ for all } j \in J \right\},$$

and  $\mathfrak{H}^m(A) = \sup_{\delta > 0} \mathfrak{H}_\delta^m(A)$ . It is elementary to observe that  $\mathfrak{H}^m(A) = H^m(A)$  and that  $\mathfrak{H}_\infty^m(A) = H_\infty^m(A)$ . Now, if  $f : W \rightarrow \mathbb{R}^m$  is an isometry, then  $H^m(A) = \mathfrak{H}^m(A) = H^m(f(A)) = H_\infty^m(f(A)) = \mathfrak{H}_\infty^m(A) = H_\infty^m(A)$ , where the third equality follows from claim (2) above.



**3.3. COAREA FORMULA.** — We recall two versions of the coarea formula. First, if  $A \subseteq \mathbb{R}^n$  is  $\mathcal{L}^n$ -measurable and  $f : A \rightarrow \mathbb{R}^{n-m}$  is Lipschitzian, then

$$\mathbb{R}^{n-m} \rightarrow [0, \infty] : y \mapsto \mathcal{H}^m(A \cap f^{-1}\{y\})$$

is  $\mathcal{L}^{n-m}$ -measurable and

$$\int_A Jf(x) d\mathcal{L}^n(x) = \int_{\mathbb{R}^{n-m}} \mathcal{H}^m(A \cap f^{-1}\{y\}) d\mathcal{L}^{n-m}(y).$$

Here, the coarea Jacobian factor is well-defined  $\mathcal{L}^n$ -almost everywhere, according to Rademacher’s theorem, and equals

$$Jf(x) = \sqrt{|\det(Df(x) \circ Df(x)^*)|} = \|\wedge_{n-m} Df(x)\|;$$

see, for instance, [5, Chap. 3, §4].

Secondly, if  $A \subseteq \mathbb{R}^p$  is  $\mathcal{H}^n$ -measurable and countably  $(\mathcal{H}^n, n)$ -rectifiable and if  $f : A \rightarrow \mathbb{R}^{n-m}$  is Lipschitzian, then  $\mathbb{R}^{n-m} \rightarrow [0, \infty] : y \mapsto \mathcal{H}^m(A \cap f^{-1}\{y\})$  is  $\mathcal{L}^{n-m}$ -measurable and

$$\int_A J_A f(x) d\mathcal{H}^n(x) = \int_{\mathbb{R}^{n-m}} \mathcal{H}^m(A \cap f^{-1}\{y\}) d\mathcal{L}^{n-m}(y).$$

In order to state a formula for the coarea Jacobian factor  $J_A f(x)$  of  $f$  relative to  $A$ , we consider a point  $x \in A$ , where  $A$  admits an approximate  $n$ -dimensional tangent space  $T_x A$  and  $f$  is differentiable at  $x$  along  $A$ . Letting  $L : T_x A \rightarrow \mathbb{R}^{n-m}$  denote the derivative of  $f$  at  $x$ , we have

$$J_A f(x) = \sqrt{|\det(L \circ L^*)|} = \|\wedge_{n-m} L\|;$$

see, for instance, [8, 3.2.22].

In both cases, it is useful to recall the following. If  $L : V \rightarrow V'$  is a linear map between two inner product spaces  $V$  and  $V'$ , then

$$(5) \quad \|\wedge_k L\| = \sup\{\langle \wedge_k L, \xi \rangle : \xi \in \wedge_k V \text{ and } |\xi| = 1\}.$$

On the one hand,  $\|\wedge_k L\| \leq \|L\|^k$  [8, 1.7.6] and  $\|L\| \leq \text{Lip } f$ , with  $L$  as above. On the other hand, if  $v_1, \dots, v_k$  are linearly independent vectors of  $V$ , then

$$(6) \quad \|\wedge_k L\| > \frac{|L(v_1) \wedge \dots \wedge L(v_k)|}{|v_1 \wedge \dots \wedge v_k|}.$$

Finally, we observe that both coarea formulas hold true when  $f$  is merely locally Lipschitzian, according to the monotone convergence theorem.

**3.4. GRASSMANNIAN.** — We let  $G(n, m)$  be the set whose members are the  $m$ -dimensional vector subspaces of  $\mathbb{R}^n$ . With  $W \in G(n, m)$ , we associate  $P_W : \mathbb{R}^n \rightarrow \mathbb{R}^n$ , the orthogonal projection on  $W$ . We give  $G(n, m)$  the structure of a compact metric space by letting  $d(W_1, W_2) = \|P_{W_1} - P_{W_2}\|$ , where  $\|\cdot\|$  is the operator norm. If  $W \in G(n, m)$ , then  $W^\perp \in G(n, n - m)$  satisfies  $P_W + P_{W^\perp} = \text{id}_{\mathbb{R}^n}$ , therefore,  $G(n, m) \rightarrow G(n, n - m) : W \mapsto W^\perp$  is an isometry. The bijective correspondence  $\varphi : G(n, m) \rightarrow \text{Hom}(\mathbb{R}^n, \mathbb{R}^n) : W \mapsto P_W$  identifies  $G(n, m)$  with the submanifold

$$M_{n,m} = \text{Hom}(\mathbb{R}^n, \mathbb{R}^n) \cap \{L : L \circ L = L, L^* = L, \text{ and } \text{trace } L = m\}.$$

There exists an open neighborhood  $V$  of  $M_{n,m}$  in  $\text{Hom}(\mathbb{R}^n, \mathbb{R}^n)$  and a Lipschitzian retraction  $\rho : V \rightarrow M_{n,m}$ , according, for instance, to [8, 3.1.20]. Therefore, if  $S \subseteq \mathbb{R}^n$  and  $\mathbf{W}_0 : S \rightarrow \mathbb{G}(n, m)$  is Lipschitzian, then there exist an open neighborhood  $U$  of  $S$  in  $\mathbb{R}^n$  and a Lipschitzian extension  $\widehat{\mathbf{W}}_0 : U \rightarrow \mathbb{G}(n, m)$  of  $\mathbf{W}_0$ . Indeed,  $\varphi \circ \mathbf{W}_0$  admits a Lipschitzian extension  $Y : \mathbb{R}^n \rightarrow \text{Hom}(\mathbb{R}^n, \mathbb{R}^n)$  – see, e.g., [8, 2.10.43] – and it suffices to let  $U = Y^{-1}(V)$  and  $\widehat{\mathbf{W}}_0 = \rho \circ (Y|_U)$ .

**3.5. ORTHONORMAL FRAMES.** — We let  $\mathbb{V}(n, m)$  be the set of orthonormal  $m$ -frames in  $\mathbb{R}^n$ , i.e.,

$$\mathbb{V}(n, m) = (\mathbb{R}^n)^m \cap \{(w_1, \dots, w_m) : \text{the family } w_1, \dots, w_m \text{ is orthonormal}\}.$$

We will consider it as a metric space, with its structure inherited from  $(\mathbb{R}^n)^m$ .

**3.6.** — Let  $V \subseteq \mathbb{G}(n, m)$  be a nonempty closed set such that  $\text{diam } V < 1$ . There exists a Lipschitzian map  $\Xi : V \rightarrow \mathbb{V}(n, m)$  such that  $W = \text{span}\{\Xi_1(W), \dots, \Xi_m(W)\}$ , for every  $W \in V$ .

*Proof.* — Pick arbitrarily  $W_0 \in V$ . If  $W \in V$ , then the map  $W_0 \rightarrow W : w \mapsto P_W(w)$  is bijective: If  $w \in W_0 \cap \{0\}$ , then  $|P_W(w) - w| = |P_W(w) - P_{W_0}(w)| < |w|$ , thus,  $P_W(w) \neq 0$ . Letting  $w_1, \dots, w_m$  be an arbitrary basis of  $W_0$ , it follows that, for each  $W \in V$ , the vectors  $w_i(W) = P_W(w_i)$ ,  $i = 1, \dots, m$ , constitute a basis of  $W$ . Furthermore, the maps  $w_i : V \rightarrow \mathbb{R}^n$  are Lipschitzian:  $|w_i(W) - w_i(W')| = |P_W(w_i) - P_{W'}(w_i)| \leq d(W, W')|w_i|$ . We apply the Gram-Schmidt process:

$$\bar{w}_1(W) = w_1(W)$$

and 
$$\bar{w}_i(W) = w_i(W) - \sum_{j=1}^{i-1} \langle w_i(W), \bar{w}_j(W) \rangle \bar{w}_j(W), \quad i = 2, \dots, m,$$

so that  $\bar{w}_1(W), \dots, \bar{w}_m(W)$  is readily an orthogonal basis of  $W$  depending upon  $W$  in a Lipschitzian way. Since each  $|\bar{w}_i|$  is bounded away from zero on  $V$ , the formula  $\Xi_i(W) = |\bar{w}_i(W)|^{-1} \bar{w}_i(W)$ ,  $i = 1, \dots, m$ , defines  $\Xi$  with the required property.

**3.7.** — There exists a Borel measurable map  $\Xi : \mathbb{G}(n, m) \rightarrow \mathbb{V}(n, m)$  with the property that  $W = \text{span}\{\Xi_1(W), \dots, \Xi_m(W)\}$ , for every  $W \in \mathbb{G}(n, m)$ .

*Proof.* — Since  $\mathbb{G}(n, m)$  is compact, it can be partitioned into finitely many Borel measurable sets  $V_1, \dots, V_J$ , each having diameter bounded by  $1/2$ . Define  $\Xi$ , piecewise, to coincide on  $V_j$  with  $\Xi_j$  associated to  $\text{Clos } V_j$  in 3.6,  $j = 1, \dots, J$ .

**3.8.** — Assume  $S \subseteq \mathbb{R}^n$ ,  $x_0 \in S$ , and  $\mathbf{W}_0 : S \rightarrow \mathbb{G}(n, m)$  is Lipschitzian. There exist an open neighborhood  $U$  of  $x_0$  in  $\mathbb{R}^n$  and Lipschitzian maps  $\mathbf{w}_1, \dots, \mathbf{w}_m, \mathbf{v}_1, \dots, \mathbf{v}_{n-m} : U \rightarrow \mathbb{R}^n$  with the following properties.

(1) For every  $x \in U$ , the family  $\mathbf{w}_1(x), \dots, \mathbf{w}_m(x), \mathbf{v}_1(x), \dots, \mathbf{v}_{n-m}(x)$  is an orthonormal basis of  $\mathbb{R}^n$ .

(2) For every  $x \in S \cap U$ , one has

$$\mathbf{W}_0(x) = \text{span}\{\mathbf{w}_1(x), \dots, \mathbf{w}_m(x)\} \quad \text{and} \quad \mathbf{W}_0(x)^\perp = \text{span}\{\mathbf{v}_1(x), \dots, \mathbf{v}_{n-m}(x)\}.$$

*Proof.* — We let  $\widehat{\mathbf{W}}_0 : \widehat{U} \rightarrow \mathbb{G}(n, m)$  be a Lipschitzian extension of  $\mathbf{W}_0$ , where  $\widehat{U}$  is an open neighborhood of  $S$  in  $\mathbb{R}^n$  (recall 3.4). Abbreviate  $W_0 := \mathbf{W}_0(x_0)$ . Define  $V = \mathbb{G}(n, m) \cap \{W : d(W, W_0) < 1/4\}$  and  $U = \widehat{U} \cap \widehat{\mathbf{W}}_0^{-1}(V)$ . Apply 3.6 to  $\text{Clos } V$  and denote by  $\Xi$  the resulting Lipschitzian map  $V \rightarrow (\mathbb{R}^n)^m$ . Next, define  $V^\perp = \mathbb{G}(n, n-m) \cap \{W^\perp : W \in V\}$ , apply 3.6 to  $\text{Clos } V^\perp$ , and denote by  $\Xi^\perp$  the resulting Lipschitzian map  $V^\perp \rightarrow (\mathbb{R}^n)^{n-m}$ . Letting  $\mathbf{w}_i(x) = (\Xi_i \circ \widehat{\mathbf{W}}_0)(x)$ ,  $i = 1, \dots, m$ , and  $\mathbf{v}_i(x) = (\Xi_i^\perp \circ \widehat{\mathbf{W}}_0)(x)$ ,  $i = 1, \dots, n-m$ , completes the proof.

3.9. — Assume  $\mathbf{W}_0 : \mathbb{R}^n \rightarrow \mathbb{G}(n, m)$  is Borel measurable. There exist Borel measurable maps  $\mathbf{w}_1, \dots, \mathbf{w}_m, \mathbf{v}_1, \dots, \mathbf{v}_{n-m} : \mathbb{R}^n \rightarrow \mathbb{R}^n$  with the following properties.

(1) For every  $x \in \mathbb{R}^n$ , the family  $\mathbf{w}_1(x), \dots, \mathbf{w}_m(x), \mathbf{v}_1(x), \dots, \mathbf{v}_{n-m}(x)$  is an orthonormal basis of  $\mathbb{R}^n$ .

(2) For every  $x \in \mathbb{R}^n$ , one has

$$\mathbf{W}_0(x) = \text{span}\{\mathbf{w}_1(x), \dots, \mathbf{w}_m(x)\} \quad \text{and} \quad \mathbf{W}_0(x)^\perp = \text{span}\{\mathbf{v}_1(x), \dots, \mathbf{v}_{n-m}(x)\}.$$

*Proof.* — Choose  $\Xi : \mathbb{G}(n, m) \rightarrow \mathbb{V}(n, m)$  and  $\Xi^\perp : \mathbb{G}(n, n-m) \rightarrow \mathbb{V}(n, n-m)$  as in 3.7. Letting

$$(\mathbf{w}_1(x), \dots, \mathbf{w}_m(x)) = (\Xi \circ \mathbf{W}_0)(x) \quad \text{and} \quad (\mathbf{v}_1(x), \dots, \mathbf{v}_{n-m}(x)) = (\Xi^\perp \circ \mathbf{W}_0^\perp)(x),$$

for  $x \in \mathbb{R}^n$ , completes the proof.

3.10. DEFINITION OF  $\mathbf{W}(x)$ . — The typical situation that arises in the remaining part of this paper is that we are given a set  $S \subseteq \mathbb{R}^n$ , a Lipschitzian map  $\mathbf{W}_0 : S \rightarrow \mathbb{G}(n, m)$ , and  $x_0 \in S$ . We will represent  $\mathbf{W}_0(x)$  and  $\mathbf{W}_0^\perp(x)$  in a neighborhood  $U$  of  $x_0$  as in 3.8. We will then further reduce the size of  $U$  several times, in order that various conditions be met. With no exception, we will denote as  $\mathbf{W}(x) = x + \mathbf{W}_0(x)$  the affine subspace containing  $x$ , of direction  $\mathbf{W}_0(x)$ , whenever  $\mathbf{W}_0(x)$  is defined.

3.11. DEFINITION OF  $g_{\mathbf{v}_1, \dots, \mathbf{v}_{n-m}, u}$  AND LOWER BOUND OF ITS COAREA FACTOR

Given an open set  $U \subseteq \mathbb{R}^n$ , a Lipschitzian map  $\mathbf{v} : U \rightarrow \mathbb{R}^n$ , and  $u \in \mathbb{R}^n$ , we define  $g_{\mathbf{v}, u} : U \rightarrow \mathbb{R}$  by the formula

$$g_{\mathbf{v}, u}(x) = \langle \mathbf{v}(x), x - u \rangle.$$

Clearly,  $g_{\mathbf{v}, u}$  is locally Lipschitzian. If  $\mathbf{v}$  is differentiable at  $x \in U$ , then so is  $g_{\mathbf{v}, u}$  and, for every  $h \in \mathbb{R}^n$ , one has

$$(7) \quad Dg_{\mathbf{v}, u}(x)(h) = \langle \nabla g_{\mathbf{v}, u}(x), h \rangle = \langle D\mathbf{v}(x)(h), x - u \rangle + \langle \mathbf{v}(x), h \rangle.$$

Next, we assume that we are given Lipschitzian maps  $\mathbf{v}_1, \dots, \mathbf{v}_{n-m} : U \rightarrow \mathbb{R}^n$ . We define  $g_{\mathbf{v}_1, \dots, \mathbf{v}_{n-m}, u} : U \rightarrow \mathbb{R}^{n-m}$  by the formula

$$g_{\mathbf{v}_1, \dots, \mathbf{v}_{n-m}, u}(x) = (g_{\mathbf{v}_1, u}(x), \dots, g_{\mathbf{v}_{n-m}, u}(x)).$$

It is locally Lipschitzian as well. The relevance of  $g_{\mathbf{v}_1, \dots, \mathbf{v}_{n-m}, u}$  stems from the following observation, assuming that  $\mathbf{v}_1, \dots, \mathbf{v}_{n-m}$  are associated with  $\mathbf{W}_0$  and  $\mathbf{W}$  as in 3.8

and 3.10:

$$\begin{aligned}
 (8) \quad u \in \mathbf{W}(x) &\iff u - x \in \mathbf{W}_0(x) \\
 &\iff \langle \mathbf{v}_i(x), u - x \rangle = 0, \text{ for all } i = 1, \dots, n - m \\
 &\iff g_{\mathbf{v}_1, \dots, \mathbf{v}_{n-m}, u}(x) = 0 \\
 &\iff x \in g_{\mathbf{v}_1, \dots, \mathbf{v}_{n-m}, u}^{-1}\{0\}.
 \end{aligned}$$

In fact,  $|g_{\mathbf{v}_1, \dots, \mathbf{v}_{n-m}, u}(x)| = |P_{\mathbf{W}_0(x)^\perp}(x - u)|$ .

Abbreviate  $g = g_{\mathbf{v}_1, \dots, \mathbf{v}_{n-m}, u}$ . If each  $\mathbf{v}_i$  is differentiable at  $x \in U$ , and  $h \in \mathbb{R}^n$ , then

$$Dg(x)(h) = \sum_{i=1}^{n-m} Dg_{\mathbf{v}_i, u}(x)(h)e_i,$$

where, here and elsewhere,  $e_1, \dots, e_{n-m}$  denotes the canonical basis of  $\mathbb{R}^{n-m}$ . Thus, if  $\mathbf{v}_1(x), \dots, \mathbf{v}_{n-m}(x)$  constitute an orthonormal family in  $\mathbb{R}^n$ , then

$$Dg_{\mathbf{v}_i, u}(x)(\mathbf{v}_j(x)) = \delta_{i,j} + \varepsilon_{i,j}(x, u),$$

where

$$(9) \quad |\varepsilon_{i,j}(x, u)| = |\langle D\mathbf{v}_i(x)(\mathbf{v}_j(x)), x - u \rangle| \leq (\text{Lip } \mathbf{v}_i) |x - u|,$$

according to (7), and, in turn,

$$Dg(x)(\mathbf{v}_j(x)) = \sum_{i=1}^{n-m} (\delta_{i,j} + \varepsilon_{i,j}(x, u)) e_i.$$

This allows for a lower bound of the coarea factor of  $g$  at  $x$  as follows.

$$\begin{aligned}
 \|\wedge_{n-m} Dg(x)\| &> |Dg(x)(\mathbf{v}_1(x)) \wedge \dots \wedge Dg(x)(\mathbf{v}_{n-m}(x))| \\
 &= \left| \left( \sum_{i=1}^{n-m} (\delta_{i,1} + \varepsilon_{i,1}(x, u)) e_i \right) \wedge \dots \wedge \left( \sum_{i=1}^{n-m} (\delta_{i,n-m} + \varepsilon_{i,n-m}(x, u)) e_i \right) \right| \\
 &= \left| \det (\delta_{i,j} + \varepsilon_{i,j}(x, u))_{i,j=1, \dots, n-m} \right|.
 \end{aligned}$$

In view of (9), we obtain the next lemma.

3.12. — Given  $\Lambda > 0$  and  $0 < \varepsilon < 1$ , there exists  $\delta_{3.12}(n, \Lambda, \varepsilon) > 0$  with the following property. Assume that

- (1)  $U \subseteq \mathbb{R}^n$  is open and  $u \in \mathbb{R}^n$ ;
- (2)  $\mathbf{v}_1, \dots, \mathbf{v}_{n-m} : U \rightarrow \mathbb{R}^n$  are Lipschitzian;
- (3)  $\mathbf{v}_1(x), \dots, \mathbf{v}_{n-m}(x)$  is an orthonormal family, for every  $x \in U$ .

If

- (4)  $\text{Lip } \mathbf{v}_i \leq \Lambda$ , for each  $i = 1, \dots, n - m$ ;
- (5)  $\text{diam}(U \cup \{u\}) \leq \delta_{3.12}(n, \Lambda, \varepsilon)$ ;

then

$$Jg_{\mathbf{v}_1, \dots, \mathbf{v}_{n-m}, u}(x) > 1 - \varepsilon,$$

at  $L^n$ -almost every  $x \in U$ .

3.13. DEFINITION OF  $\pi_u$  AND ITS RELATION WITH  $g_{\mathbf{v}_1, \dots, \mathbf{v}_{n-m}, u}$ . — With  $u \in \mathbb{R}^n$ , we associate

$$\begin{aligned} \pi_u : \mathcal{V}(n, n - m) \times \mathbb{R}^n &\longrightarrow \mathbb{R}^{n-m} \\ (\xi_1, \dots, \xi_{n-m}, x) &\longmapsto (\langle \xi_1, x - u \rangle, \dots, \langle \xi_{n-m}, x - u \rangle). \end{aligned}$$

When  $(\xi_1, \dots, \xi_{n-m}) \in \mathcal{V}(n, n - m)$  is fixed, we also abbreviate as  $\pi_{\xi_1, \dots, \xi_{n-m}, u}$  the map  $\mathbb{R}^n \rightarrow \mathbb{R}^{n-m}$  defined by  $\pi_{\xi_1, \dots, \xi_{n-m}, u}(x) = \pi_u(\xi_1, \dots, \xi_{n-m}, x)$ . It is then rather useful to observe that, in the context described in 3.8 and 3.10, the following holds:

$$(10) \quad \pi_{\mathbf{v}_1(x), \dots, \mathbf{v}_{n-m}(x), u}^{-1} \{g_{\mathbf{v}_1, \dots, \mathbf{v}_{n-m}, u}(x)\} = \mathbf{W}(x).$$

Indeed,

$$\begin{aligned} h \in \mathbf{W}(x) &\iff h - x \in \mathbf{W}_0(x) \\ &\iff \langle \mathbf{v}_i(x), h - x \rangle = 0, \text{ for all } i = 1, \dots, n - m \\ &\iff \langle \mathbf{v}_i(x), h - u \rangle = \langle \mathbf{v}_i(x), x - u \rangle, \text{ for all } i = 1, \dots, n - m \\ &\iff \pi_{\mathbf{v}_1(x), \dots, \mathbf{v}_{n-m}(x), u}(h) = g_{\mathbf{v}_1, \dots, \mathbf{v}_{n-m}, u}(x). \end{aligned}$$

In the sequel, we will sometimes abbreviate  $\xi = (\xi_1, \dots, \xi_{n-m}) \in \mathcal{V}(n, n - m)$ . It also helps to notice that, for given  $\xi \in \mathcal{V}(n, n - m)$  and  $y \in \mathbb{R}^{n-m}$ , the set  $\pi_{\xi, u}^{-1}\{y\}$  is an  $m$ -dimensional affine subspace of  $\mathbb{R}^n$ .

3.14. — If  $B \in \mathcal{B}(\mathbb{R}^n)$  and  $u \in \mathbb{R}^n$ , then

$$h_B : \mathcal{V}(n, n - m) \times \mathbb{R}^{n-m} \longrightarrow [0, \infty] : (\xi, y) \longmapsto H^m(B \cap \pi_{\xi, u}^{-1}\{y\})$$

is Borel measurable.

*Proof.* — We start by showing that, when  $B$  is compact,  $h_B$  is upper semicontinuous. Thus, if  $((\xi_k, y_k))_k$  is a sequence in  $\mathcal{V}(n, n - m) \times \mathbb{R}^{n-m}$  that converges to  $(\xi, y)$ , we ought to show that

$$(11) \quad H_\infty^m(K) > \limsup_k H_\infty^m(K_k),$$

where  $K = B \cap \pi_{\xi, u}^{-1}\{y\}$  and  $K_k = B \cap \pi_{\xi_k, u}^{-1}\{y_k\}$ . This is indeed equivalent to the same inequality with  $H_\infty^m$  replaced by  $H^m$ , according to 3.2(3) and the last sentence of 3.13. Considering, if necessary, a subsequence of  $(K_k)_k$  we may assume that none of the compact sets  $K_k$  is empty and that the limit superior in (11) is a limit. Since the set of nonempty compact subsets of the compact set  $B$ , equipped with the Hausdorff metric, is compact, the sequence  $(K_k)_k$  admits a subsequence (denoted the same way) converging to a compact set  $L \subseteq B$ . Given  $z \in L$ , there are  $z_k \in K_k$  converging to  $z$ . Thus,  $\pi_u(\xi, z) = \lim_k \pi_u(\xi_k, z_k) = \lim_k y_k = y$ . In other words,  $z \in K$ . Therefore,  $H_\infty^m(K) > H_\infty^m(L)$ , so that (11) follows from 3.2(1).

Next, we fix  $r > 0$  and we abbreviate

$$A = \mathcal{B}(\mathbb{R}^n) \cap \{B : h_{B \cap B(0, r)} \text{ is Borel measurable}\}.$$

Thus, we have just shown that  $A$  contains the collection  $\mathcal{F}(\mathbb{R}^n)$  of all closed subsets of  $\mathbb{R}^n$ . Observe that if  $(B_j)_j$  is a nondecreasing sequence in  $A$  and  $B = \bigcup_j B_j$ , then

$h_{B \cap \mathbf{B}(0,r)} = \lim_j h_{B_j \cap \mathbf{B}(0,r)}$  pointwise, thus,  $B \in A$ . In particular,  $\mathbb{R}^n \in A$ . Furthermore, if  $B, B' \in A$  and  $B' \subseteq B$ , then  $h_{B \cap B'} = h_B - h_{B'}$ , because all measures involved are finite; indeed,  $h_{B \cap \mathbf{B}(0,r)}(\xi, y) \leq \alpha(m)r^m$ , for all  $(\xi, y)$ . Accordingly,  $B \cap B' \in A$ . This means that  $A$  is a Dynkin class. Since  $F(\mathbb{R}^n)$  is a  $\pi$ -system,  $A$  contains the  $\sigma$ -algebra generated by  $F(\mathbb{R}^n)$ , i.e.,  $B(\mathbb{R}^n)$  [3, Th. 1.6.2]. Finally, if  $B \in B(\mathbb{R}^n)$ , then  $h_B = \lim_j h_{B \cap \mathbf{B}(0,j)}$  pointwise, whence  $h_B$  is Borel measurable.

3.15. — Assume  $B \in B(\mathbb{R}^n)$  and  $\mathbf{W}_0 : \mathbb{R}^n \rightarrow G(n, m)$  is Borel measurable. The following function is Borel measurable.

$$\mathbb{R}^n \rightarrow [0, \infty] : x \mapsto H^m(B \cap \mathbf{W}(x)).$$

*Proof.* — Let  $h_{\mathbf{W}, B}$  denote this function. Let  $\mathbf{v}_1, \dots, \mathbf{v}_{n-m} : \mathbb{R}^n \rightarrow \mathbb{R}^n$  be Borel measurable maps associated with  $\mathbf{W}_0$  in 3.9. Fix  $u \in \mathbb{R}^n$  arbitrarily. Define

$$\Upsilon : \mathbb{R}^n \rightarrow \mathbb{V}(n, n-m) \times \mathbb{R}^{n-m} : x \mapsto (\mathbf{v}_1(x), \dots, \mathbf{v}_{n-m}(x), g_{\mathbf{v}_1(x), \dots, \mathbf{v}_{n-m}(x), u}(x))$$

and notice that

$$h_{\mathbf{W}, B} = h_B \circ \Upsilon$$

(where  $h_B$  is the function associated with  $B$  and  $u$  in 3.14), according to (10). One notes that  $\Upsilon$  is Borel measurable; whence, the conclusion ensues from 3.14.

3.16. DEFINITION OF  $\phi_{E, \mathbf{W}}$ . — Let  $\mathbf{W}_0 : \mathbb{R}^n \rightarrow G(n, m)$  be Borel measurable and  $E \in B(\mathbb{R}^n)$  be such that  $L^n(E) < \infty$ . For each  $B \in B(\mathbb{R}^n)$ , we define

$$\phi_{E, \mathbf{W}}(B) = \int_E H^m(B \cap \mathbf{W}(x)) dL^n(x).$$

This is well-defined, according to 3.15. It is easy to check that  $\phi_{E, \mathbf{W}}$  is a locally finite – hence,  $\sigma$ -finite – Borel measure on  $\mathbb{R}^n$ ; indeed,  $\phi_{E, \mathbf{W}}(B) \leq \alpha(m)(\text{diam } B)^m L^n(E)$ .

To close this section, we discuss the relevance of  $\phi_{E, \mathbf{W}}$  to the problem of existence of “nearly Nikodým sets”.

3.17. DEFINITION OF NEARLY NIKODÝM SET. — Let  $E \in B(\mathbb{R}^n)$ . We say that  $B \in B(E)$  is nearly  $m$ -Nikodým in  $E$  if

- (1)  $L^n(B) > 0$ ;
- (2) For  $L^n$ -almost each  $x \in E$ , there is  $W \in G(n, m)$  such that

$$H^m(B \cap (x + W)) = 0.$$

In case  $n = 2, m = 1$ , and  $E = [0, 1] \times [0, 1]$ , the existence of such a  $B$  (with  $L^2(B) = 1$ ) was established by Nikodým [12], see also [9, Chap. 8]. For arbitrary  $n > 2$  and  $m = n - 1$ , the existence of such a  $B$  was established by Falconer [6]. In fact, in both cases, these authors established the stronger condition that, for every  $x \in B, H^m(B \cap (x + W)) = 0$  can be replaced by  $B \cap (x + W) = \{x\}$ . Thus, in case  $1 \leq m < n - 1$ , letting  $B$  be a set exhibited by Falconer, if  $x \in B$  and  $W \subseteq G(n, n - 1)$  is such that  $B \cap (x + W) = \{x\}$ , then picking arbitrarily  $V \in G(n, m)$  such that  $V \subseteq W$ , we see that  $B \cap (x + V) = \{x\}$ . Whence,  $B$  is also nearly  $m$ -Nikodým in  $B$ .

Assuming that  $\mathbf{W}_0 : E \rightarrow G(n, m)$  is Borel measurable, we say that  $B \in \mathcal{B}(E)$  is *nearly  $m$ -Nikodým in  $E$  relative to  $\mathbf{W}$*  if

- (1)  $L^n(B) > 0$ ;
- (2) For  $L^n$ -almost each  $x \in E$ , one has  $H^m(B \cap \mathbf{W}(x)) = 0$ .

**3.18.** — Let  $E \in \mathcal{B}(\mathbb{R}^n)$  have finite  $L^n$  measure and  $\mathbf{W}_0 : \mathbb{R}^n \rightarrow G(n, m)$  be Borel measurable. The following are equivalent.

- (1)  $L^n|_{\mathcal{B}(E)}$  is absolutely continuous with respect to  $\phi_{E, \mathbf{W}}|_{\mathcal{B}(E)}$ .
- (2) There does not exist a nearly  $m$ -Nikodým set in  $E$  relative to  $\mathbf{W}$ .

*Proof.* — A set  $B \in \mathcal{B}(E)$  such that  $\phi_{E, \mathbf{W}}(B) = 0$  and  $L^n(B) > 0$  is, by definition, a nearly  $m$ -Nikodým set relative to  $\mathbf{W}$ . Condition (1) is equivalent to their nonexistence.

**3.19.** — Assume that  $E \in \mathcal{B}(\mathbb{R}^n)$  and that  $B \in \mathcal{B}(E)$  is nearly  $m$ -Nikodým. The following hold.

- (1) There exists  $\mathbf{W}_0 : \mathbb{R}^n \rightarrow G(n, m)$  Borel measurable such that  $B$  is nearly  $m$ -Nikodým in  $E$  relative to  $\mathbf{W}$ .
- (2) There exists  $C \subseteq B$  compact and  $\overline{\mathbf{W}}_0 : \mathbb{R}^n \rightarrow G(n, m)$  continuous such that  $C$  is nearly  $m$ -Nikodým in  $C$  relative to  $\overline{\mathbf{W}}$ .

Recall our convention that  $\mathbf{W}(x) = x + \mathbf{W}_0(x)$ .

*Proof.* — Define a Borel measurable map  $\xi : G(n, m) \rightarrow V(n, n - m)$  by  $\xi(W) = \Xi(W^\perp)$ , where  $\Xi : G(n, n - m) \rightarrow V(n, n - m)$  is as in 3.7. Choose arbitrarily  $u \in \mathbb{R}^n$  and define a Borel measurable map

$$\begin{aligned} \Upsilon : E \times G(n, m) &\longrightarrow V(n, n - m) \times \mathbb{R}^{n-m} \\ (x, W) &\longmapsto (\xi(W), \langle \xi_1(W), x - u \rangle, \dots, \langle \xi_{n-m}(W), x - u \rangle). \end{aligned}$$

Similarly to (10), observe that

$$x + W = \pi_{\xi(W), u}^{-1} \{ (\langle \xi_1(W), x - u \rangle, \dots, \langle \xi_{n-m}(W), x - u \rangle) \},$$

for every  $(x, W) \in E \times G(n, m)$ . We infer from 3.14 that

$$h_B \circ \Upsilon : E \times G(n, m) \longrightarrow [0, \infty] : (x, W) \longmapsto H^m(B \cap (x + W))$$

is Borel measurable. Thus, the set

$$E = E \times G(n, m) \cap \{(x, W) : H^m(B \cap (x + W)) = 0\}$$

is Borel measurable as well. The set  $N = E \cap \{x : E_x = ?\}$  is coanalytic and  $L^n(N) = 0$ , by assumption. By virtue of von Neumann's selection theorem [13, 5.5.3], there exists a universally measurable map  $\widetilde{\mathbf{W}}_0 : E \cap N \rightarrow G(n, m)$  such that  $\widetilde{\mathbf{W}}_0(x) \in E_x$ , for every  $x \in E \cap N$ , i.e.,  $H^m(B \cap (x + \widetilde{\mathbf{W}}_0(x))) = 0$ . We extend  $\widetilde{\mathbf{W}}_0$  to be an arbitrary constant on  $N \cup (\mathbb{R}^n \setminus E)$ . This makes  $\widetilde{\mathbf{W}}_0$  an  $L^n$ -measurable map defined on  $\mathbb{R}^n$ . Therefore, it is equal  $L^n$ -almost everywhere to a Borel measurable map  $\mathbf{W}_0 : \mathbb{R}^n \rightarrow G(n, m)$ . This proves (1).

In order to prove (2), we recall 3.4, specifically, the retraction  $\rho : V \rightarrow M_{n,m}$  and the homeomorphic identification  $\varphi : G(n, m) \rightarrow M_{n,m}$ . Owing to the compactness of  $M_{n,m}$ , there are finitely many open balls  $U_j, j = 1, \dots, J$ , whose closures are contained in  $V$  and whose union contains  $M_{n,m}$ . Since  $L^n(B) > 0$ , there exists  $j \in \{1, \dots, J\}$  such that  $L^n(B \cap E_j) > 0$ , where  $E_j = (\varphi \circ \mathbf{W}_0)^{-1}(U_j)$ . It follows from Lusin's theorem [8, 2.5.3] that there exists a compact set  $C \subseteq B \cap E_j$  such that  $L^n(C) > 0$  and the restriction  $\mathbf{W}_0|_C$  is continuous. The map  $\varphi \circ \mathbf{W}_0|_C$  takes its values in the closed ball  $\text{Clos } U_j$ , therefore, it admits a continuous extension  $Y_0 : \mathbb{R}^n \rightarrow \text{Clos } U_j \subseteq V$ . Letting  $\overline{\mathbf{W}}_0 = \varphi^{-1} \circ \rho \circ Y_0$  completes the proof.

#### 4. COMMON SETTING

4.1. SETTING FOR THE NEXT THREE SECTIONS. — In the next three sections, we shall assume the following.

- (1)  $E \subseteq \mathbb{R}^n$  is Borel measurable and  $L^n(E) < \infty$ .
- (2)  $U \subseteq \mathbb{R}^n$  is open and  $E \subseteq U$ .
- (3)  $B \subseteq \mathbb{R}^n$  is Borel measurable.
- (4)  $\mathbf{W}_0 : U \rightarrow G(n, m)$  is Lipschitzian.
- (5)  $\mathbf{W}(x) = x + \mathbf{W}_0(x)$ , for each  $x \in U$ .
- (6)  $\Lambda > 0$ .
- (7)  $\mathbf{w}_1, \dots, \mathbf{w}_m : U \rightarrow \mathbb{R}^n$  and  $\text{Lip } \mathbf{w}_i \leq \Lambda, i = 1, \dots, m$ .
- (8)  $\mathbf{v}_1, \dots, \mathbf{v}_{n-m} : U \rightarrow \mathbb{R}^n$  and  $\text{Lip } \mathbf{v}_i \leq \Lambda, i = 1, \dots, n - m$ .
- (9)  $\mathbf{W}_0(x) = \text{span}\{\mathbf{w}_1(x), \dots, \mathbf{w}_m(x)\}$ , for every  $x \in U$ .
- (10)  $\mathbf{W}_0(x)^\perp = \text{span}\{\mathbf{v}_1(x), \dots, \mathbf{v}_{n-m}(x)\}$ , for every  $x \in U$ .
- (11)  $\mathbf{w}_1(x), \dots, \mathbf{w}_m(x), \mathbf{v}_1(x), \dots, \mathbf{v}_{n-m}(x)$  constitute an orthonormal basis of  $\mathbb{R}^n$ , for every  $x \in U$ .

#### 5. TWO FIBRATIONS

5.1. A FIBERED SPACE ASSOCIATED WITH  $E, B, \mathbf{w}_1, \dots, \mathbf{w}_m$ . — We define

$$F : E \times \mathbb{R}^m \longrightarrow \mathbb{R}^n \times \mathbb{R}^n : (x, t_1, \dots, t_m) \longmapsto \left(x, x + \sum_{i=1}^m t_i \mathbf{w}_i(x)\right)$$

as well as

$$\Sigma = F(E \times \mathbb{R}^m) = \mathbb{R}^n \times \mathbb{R}^n \cap \{(x, u) : x \in E \text{ and } u \in \mathbf{W}(x)\}.$$

We will oftentimes abbreviate  $(x, t) = (x, t_1, \dots, t_m)$ . It is obvious that  $F$  is locally Lipschitzian and, therefore,  $\Sigma$  is countably  $(n+m)$ -rectifiable and  $H^{n+m}$ -measurable. We also consider the two canonical projections

$$\pi_1 : \mathbb{R}^n \times \mathbb{R}^n \longrightarrow \mathbb{R}^n : (x, u) \longmapsto x \quad \text{and} \quad \pi_2 : \mathbb{R}^n \times \mathbb{R}^n \longrightarrow \mathbb{R}^n : (x, u) \longmapsto u,$$

as well as the set

$$\Sigma_B = \Sigma \cap \pi_2^{-1}(B) = \mathbb{R}^n \times \mathbb{R}^n \cap \{(x, u) : x \in E \text{ and } u \in B \cap \mathbf{W}(x)\},$$



which also is, clearly, countably  $(n + m)$ -rectifiable and  $H^{n+m}$ -measurable. With the prospect of applying the coarea formula to  $\Sigma_B$  and  $\pi_1$ , and to  $\Sigma_B$  and  $\pi_2$ , respectively, we observe that, for each fixed  $x \in E$ ,

$$\Sigma_B \cap \pi_1^{-1}\{x\} = \{x\} \times (\mathbb{R}^n \cap \{u : u \in B \cap \mathbf{W}(x)\}),$$

so that

$$(12) \quad H^m(\Sigma_B \cap \pi_1^{-1}\{x\}) = H^m(B \cap \mathbf{W}(x)),$$

and that, for each fixed  $u \in B$ ,

$$\begin{aligned} \Sigma_B \cap \pi_2^{-1}\{u\} &= (\mathbb{R}^n \cap \{x : x \in E \text{ and } u \in \mathbf{W}(x)\}) \times \{u\} \\ &= (\mathbb{R}^n \cap \{x : x \in E \cap g_{\mathbf{v}_1, \dots, \mathbf{v}_{n-m}, u}^{-1}\{0\}\}) \times \{u\}, \end{aligned}$$

according to (8), so that

$$(13) \quad H^m(\Sigma_B \cap \pi_2^{-1}\{u\}) = H^m(E \cap g_{\mathbf{v}_1, \dots, \mathbf{v}_{n-m}, u}^{-1}\{0\}),$$

whenever  $u \in B$ . It now follows from the coarea formula that

$$(14) \quad \int_{\Sigma_B} J_{\Sigma} \pi_1 dH^{n+m} = \int_E H^m(B \cap \mathbf{W}(x)) dL^n(x) = \phi_{E, \mathbf{W}}(B)$$

and

$$(15) \quad \int_{\Sigma_B} J_{\Sigma} \pi_2 dH^{n+m} = \int_B H^m(E \cap g_{\mathbf{v}_1, \dots, \mathbf{v}_{n-m}, u}^{-1}\{0\}) dL^n(u).$$

For these formulas to be useful, we need to establish bounds for the coarea Jacobian factors  $J_{\Sigma} \pi_1$  and  $J_{\Sigma} \pi_2$ . In order to do so, we notice that if  $\Sigma \ni (x, u) = F(x, t)$ ,  $F$  is differentiable at  $(x, t)$ , i.e., each  $\mathbf{w}_i$  is differentiable at  $x$ ,  $i = 1, \dots, m$ , then the approximate tangent space  $T_{(x,u)}\Sigma$  exists and is generated by the following  $n + m$  vectors of  $\mathbb{R}^n \times \mathbb{R}^n$ :

$$\begin{aligned} \sum_{p=1}^n \langle w_j, e_p \rangle \frac{\partial F}{\partial x_p}(x, t) &= \left( w_j, w_j + \sum_{i=1}^m t_i D\mathbf{w}_i(x)(w_j) \right), \quad j = 1, \dots, n, \\ \frac{\partial F}{\partial t_k}(x, t) &= (0, \mathbf{w}_k(x)), \quad k = 1, \dots, m, \end{aligned}$$

where  $w_1, \dots, w_n$  is an arbitrary basis of  $\mathbb{R}^n$ . As usual,  $e_1, \dots, e_n$  denotes the canonical basis of  $\mathbb{R}^n$ .

5.2. COAREA JACOBIAN FACTOR OF  $\pi_1$ . — For  $H^{n+m}$ -almost every  $(x, u) \in \Sigma$ , one has  $(1 + m^2 \Lambda^2 |x - u|^2)^{-m/2} (2 + 2m\Lambda |x - u| + m^2 \Lambda^2 |x - u|^2)^{-(n-m)/2} \triangleleft J_{\Sigma} \pi_1(x, u) \triangleleft 1$ .

*Proof.* — We recall 3.3. That the right hand inequality be valid follows from  $\text{Lip } \pi_1 = 1$ . Regarding the left hand inequality, fix  $(x, u) = F(x, t)$  such that  $F$  is differentiable at  $(x, t)$  and let  $L : T_{(x,u)}\Sigma \rightarrow \mathbb{R}^n$  denote the restriction of  $\pi_1$  to  $T_{(x,u)}\Sigma$ . Define  $w_j = \mathbf{w}_j(x)$ ,  $j = 1, \dots, m$ , and  $w_j = \mathbf{v}_{j-m}(x)$ ,  $j = m + 1, \dots, n$ . Put

$$v_j = \sum_{p=1}^n \langle w_j, e_p \rangle \frac{\partial F}{\partial x_p}(x, t) - \frac{\partial F}{\partial t_j}(x, t) = \left( w_j, \sum_{i=1}^m t_i D\mathbf{w}_i(x)(w_j) \right),$$

$j = 1, \dots, m$ , and

$$v_j = \sum_{p=1}^n \langle w_j, e_p \rangle \frac{\partial F}{\partial x_p}(x, t) = \left( w_j, w_j + \sum_{i=1}^m t_i D\mathbf{w}_i(x)(w_j) \right),$$

$j = m + 1, \dots, n$ . Recall (6) that

$$J_{\Sigma} \pi_1(x, u) = \|\wedge_n L\| > \frac{1}{|v_1 \wedge \dots \wedge v_n|},$$

since  $L(v_j) = w_j$ ,  $j = 1 \dots, n$ . Now, notice that, for  $j = 1, \dots, m$ ,

$$|v_j|^2 = |w_j|^2 + \left| \sum_{i=1}^m t_i D\mathbf{w}_i(x)(w_j) \right|^2 \leq 1 + m^2 \Lambda^2 |t|^2,$$

whereas, for  $j = m + 1, \dots, n$ ,

$$\begin{aligned} |v_j|^2 &= |w_j|^2 + \left| w_j + \sum_{i=1}^m t_i D\mathbf{w}_i(x)(w_j) \right|^2 \\ &\leq 2 + 2 \left| \sum_{i=1}^m t_i D\mathbf{w}_i(x)(w_j) \right| + \left| \sum_{i=1}^m t_i D\mathbf{w}_i(x)(w_j) \right|^2 \leq 2 + 2m\Lambda|t| + m^2 \Lambda^2 |t|^2. \end{aligned}$$

Since  $u = x + \sum_{i=1}^m t_i \mathbf{w}_i(x)$ , one also has

$$|u - x|^2 = \left| \sum_{i=1}^m t_i \mathbf{w}_i(x) \right|^2 = |t|^2.$$

Finally,

$$|v_1 \wedge \dots \wedge v_n| \leq (1 + m^2 \Lambda^2 |x - u|^2)^{m/2} (2 + 2m\Lambda|x - u| + m^2 \Lambda^2 |x - u|^2)^{(n-m)/2},$$

and the conclusion follows.

5.3. — Let  $1 \leq q \leq n - 1$  be an integer and let  $v_1, \dots, v_q$  be an orthonormal family in  $\mathbb{R}^n$ . There exists  $\lambda \in \Lambda(n, q)$  such that

$$|\det (\langle v_k, e_{\lambda(j)} \rangle)_{j,k=1,\dots,q}| > \binom{n}{q}^{-1/2}.$$

Here,  $\Lambda(n, q)$  denotes the set of increasing maps  $\{1, \dots, q\} \rightarrow \{1, \dots, n\}$ .

*Proof.* — We define a linear map  $L : \mathbb{R}^q \rightarrow \mathbb{R}^n : (s_1, \dots, s_q) \mapsto \sum_{k=1}^q s_k v_k$  and we observe that  $L$  is an isometry. Therefore, its area Jacobian factor  $JL = 1$ , by definition. Now, also

$$(JL)^2 = \sum_{\lambda \in \Lambda(n, q)} |\det (\langle v_k, e_{\lambda(j)} \rangle)_{j,k=1,\dots,q}|^2,$$

according to the Binet-Cauchy formula [5, Chap. 3, §2, Th. 4]. The conclusion easily follows.

5.4. COAREA JACOBIAN FACTOR OF  $\pi_2$ . — *The following hold.*

(1) For  $H^{n+m}$ -almost every  $(x, u) \in \Sigma$ , one has

$$\frac{\binom{n}{n-m}^{-1/2} - m(n-m)\Lambda|x-u|(1+m\Lambda|x-u|)^{n-m-1}}{(2+2m\Lambda|x-u|+m^2\Lambda^2|x-u|^2)^{(n-m)/2}} \circlearrowleft J_\Sigma\pi_2(x, u) \circlearrowright 1.$$

(2) For  $H^{n+m}$ -almost every  $(x, u) \in \Sigma$ , one has  $J_\Sigma\pi_2(x, u) > 0$ .

*Proof.* — Clearly,  $J_\Sigma\pi_2(x, u) \circlearrowleft (\text{Lip } \pi_2)^n \circlearrowright 1$ . Regarding the left hand inequality, fix  $(x, u) = F(x, t)$  such that  $F$  is differentiable at  $(x, t)$  and, this time, let  $L : T_{(x,u)}\Sigma \rightarrow \mathbb{R}^n$  denote the restriction of  $\pi_2$  to  $T_{(x,u)}\Sigma$ . We will now define a family of  $n$  vectors  $v_1, \dots, v_n$  belonging to  $T_{(x,u)}\Sigma$ . We choose  $v_k = \partial F / \partial t_k(x, t) = (0, \mathbf{w}_k(x))$ , for  $k = 1, \dots, m$ . For choosing the  $n - m$  remaining vectors, we proceed as follows. We select  $\lambda \in \Lambda(n, n - m)$  as in 5.3 applied with  $q = n - m$  to  $\mathbf{v}_1(x), \dots, \mathbf{v}_{n-m}(x)$  and we let  $v_{m+j} = \partial F / \partial x_{\lambda(j)}(x, t)$ ,  $j = 1, \dots, n - m$ . Recalling (6), we have

$$J_\Sigma\pi_1(x, u) = \|\wedge_n L\| > \frac{|L(v_1) \wedge \dots \wedge L(v_n)|}{|v_1 \wedge \dots \wedge v_n|}.$$

As in the proof of 5.2, we find that

$$|v_1 \wedge \dots \wedge v_n| \circlearrowleft (2 + 2m\Lambda|x-u| + m^2\Lambda^2|x-u|^2)^{(n-m)/2}$$

and it remains only to find a lower bound for  $|L(v_1) \wedge \dots \wedge L(v_n)|$ . The latter equals the absolute value of the determinant of the matrix of coefficients of  $L(v_i)$ ,  $i = 1, \dots, n$ , with respect to any orthonormal basis of  $\mathbb{R}^n$ . We choose the basis  $\mathbf{w}_1(x), \dots, \mathbf{w}_m(x), \mathbf{v}_1(x), \dots, \mathbf{v}_{n-m}(x)$ . Thus,

$$\begin{aligned} (16) \quad |L(v_1) \wedge \dots \wedge L(v_n)| &= \left| \det \begin{pmatrix} 1 \cdots 0 & & * \\ \vdots & \ddots & \vdots \\ 0 \cdots 1 & & * \\ 0 \cdots 0 & \langle e_{\lambda(j)} + \sum_{i=1}^m t_i \frac{\partial \mathbf{w}_i}{\partial x_{\lambda(j)}}(x), \mathbf{v}_k(x) \rangle & \end{pmatrix} \right| \\ &= \left| \det \left( \left\langle e_{\lambda(j)} + \sum_{i=1}^m t_i \frac{\partial \mathbf{w}_i}{\partial x_{\lambda(j)}}(x), \mathbf{v}_k(x) \right\rangle \right)_{j,k=1, \dots, n-m} \right|. \end{aligned}$$

Abbreviate

$$h_{\lambda(j)} = \sum_{i=1}^m t_i \frac{\partial \mathbf{w}_i}{\partial x_{\lambda(j)}}(x)$$

and observe that  $|h_{\lambda(j)}| \circlearrowleft m\Lambda|t| = m\Lambda|x-u|$ ,  $j = 1, \dots, n - m$  (recall the proof of 5.2). It remains only to remember that  $\lambda$  has been selected in order that

$$\left| \det \left( \langle e_{\lambda(j)}, \mathbf{v}_k(x) \rangle \right)_{j,k=1, \dots, n-m} \right| > \binom{n}{n-m}^{-1/2}$$

and to infer from the multilinearity of the determinant that

$$\begin{aligned} & \left| \det(\langle e_{\lambda(j)}, \mathbf{v}_k(x) \rangle + \langle h_{\lambda(j)}, \mathbf{v}_k(x) \rangle)_{j,k} - \det(\langle e_{\lambda(j)}, \mathbf{v}_k(x) \rangle)_{j,k} \right| \\ & \leq (n - m) \left( \max_{j=1, \dots, n-m} |h_{\lambda(j)}| \right) \left( \max_{j=1, \dots, n-m} |h_{\lambda(j)}| + |e_{\lambda(j)}| \right)^{n-m-1} \\ & \leq (n - m)m\Lambda|x - u| (1 + m\Lambda|x - u|)^{n-m-1}. \end{aligned}$$

This completes the proof of conclusion (1).

Let  $E_0$  denote the subset of  $E$  consisting of those  $x$  such that each  $\mathbf{w}_i, i = 1, \dots, m$ , is differentiable at  $x$ . Thus,  $E_0$  is Borel measurable and so is

$$A = E_0 \times \mathbb{R}^m$$

$$\cap \left\{ (x, t) : \text{rank}(\mathbf{w}_1(x) | \dots | \mathbf{w}_m(x) | e_1 + \sum_{i=1}^m t_i \frac{\partial \mathbf{w}_i}{\partial x_1}(x) | \dots | e_n + \sum_{i=1}^m t_i \frac{\partial \mathbf{w}_i}{\partial x_n}(x)) < n \right\}.$$

If  $(x, u) \in \Sigma \cap F(A)$ , then the restriction of  $\pi_2$  to  $T_{(x,u)}\Sigma$  is surjective and, therefore,  $J_\Sigma \pi_2(x, u) > 0$ . Thus, we ought to show that  $H^{n+m}(F(A)) = 0$ . Since  $F$  is Lipschitzian, it suffices to establish that  $L^{n+m}(A) = 0$ . As  $A$  is Borel measurable, it is enough to prove that  $L^m(A_x) = 0$ , for every  $x \in E_0$ , according to Fubini's theorem. Fix  $x \in E_0$ . As in the proof of conclusion (1), choose  $\lambda \in \Lambda(n, n - m)$  associated with  $\mathbf{v}_1(x), \dots, \mathbf{v}_{n-m}(x)$ , according to 5.3. Based on (16), we see that

$$A_x \subseteq \mathbb{R}^m \cap \left\{ t : \det \left( \left\langle e_{\lambda(j)} + \sum_{i=1}^m t_i \frac{\partial \mathbf{w}_i}{\partial x_{\lambda(j)}}(x), \mathbf{v}_k(x) \right\rangle \right)_{j,k=1, \dots, n-m} = 0 \right\}.$$

The set on the right is of the form  $S_x = \mathbb{R}^m \cap \{(t_1, \dots, t_m) : P_x(t_1, \dots, t_m) = 0\}$ , for some polynomial  $P_x \in \mathbb{R}[T_1, \dots, T_m]$ , and

$$P_x(0, \dots, 0) = \det(\langle e_{\lambda(j)}, \mathbf{v}_k(x) \rangle)_{j,k=1, \dots, n-m} \neq 0.$$

It follows that  $L^m(S_x) = 0$  – see, e.g., [8, 2.6.5] – and the proof of (2) is complete.

**5.5. PROPOSITION.** — *The measure  $\phi_{E, \mathbf{W}}$  is absolutely continuous with respect to  $L^n$ .*

*Proof.* — Let  $B \in \mathcal{B}(\mathbb{R}^n)$  be such that  $L^n(B) = 0$ . It follows from (15) that

$$\int_{\Sigma_B} J_\Sigma \pi_2 dH^{n+m} = 0.$$

It next follows from 5.4(2) that  $H^{n+m}(\Sigma_B) = 0$ . In turn, (14) implies that

$$\phi_{E, \mathbf{W}}(B) = \int_{\Sigma_B} J_\Sigma \pi_1 dH^{n+m} = 0.$$

**5.6. DEFINITION OF  $Z_E \mathbf{W}$ .** — Note that  $\phi_{E, \mathbf{W}}$  is a  $\sigma$ -finite Borel measure on  $\mathbb{R}^n$  (see 3.16) and it is absolutely continuous with respect to  $L^n$  (see 5.5). It then ensues from the Radon-Nikodým theorem that there exists a Borel measurable function

$$Z_E \mathbf{W} : \mathbb{R}^n \rightarrow \mathbb{R}$$

such that, for every  $B \in \mathcal{B}(\mathbb{R}^n)$ , one has

$$\int_E H^m(B \cap \mathbf{W}(x)) dL^n(x) = \phi_{E, \mathbf{W}}(B) = \int_B Z_E \mathbf{W}(u) dL^n(u).$$

Furthermore,  $Z_E \mathbf{W}$  is uniquely defined (only) up to an  $L^n$  null set. This will not affect the reasoning in this paper. Each time we write  $Z_E \mathbf{W}$ , we mean *one* particular Borel measurable function satisfying the above equality, for every  $B \in \mathcal{B}(\mathbb{R}^n)$ .

5.7. DEFINITION OF  $\Upsilon_E^0 \mathbf{W}$ . — We define  $\Upsilon_E^0 \mathbf{W} : \mathbb{R}^n \rightarrow [0, \infty]$  by the formula

$$(17) \quad \Upsilon_E^0 \mathbf{W}(u) = H^m(E \cap g_{v_1, \dots, v_{n-m}, u}^{-1}\{0\}),$$

$u \in \mathbb{R}^n$ . Letting  $B = \mathbb{R}^n$  in (13), one infers from 3.3 that  $\Upsilon_E^0 \mathbf{W}$  is  $L^n$ -measurable. Using the estimates we have established so far regarding coarea Jacobian factors, we now show that  $Z_E \mathbf{W}$  and  $\Upsilon_E^0 \mathbf{W}$  are comparable, when the diameter of  $E$  is not too large.

5.8. PROPOSITION. — Given  $0 < \varepsilon < 1$ , there exists  $\delta_{5.8}(n, \Lambda, \varepsilon) > 0$  with the following property. If  $\text{diam } E \leq \delta_{5.8}(n, \Lambda, \varepsilon)$ , then

$$(1 - \varepsilon)2^{-(n-m)/2} \Upsilon_E^0 \mathbf{W}(u) \leq Z_E \mathbf{W}(u) \leq (1 + \varepsilon)2^{(n-m)/2} \binom{n}{n-m}^{1/2} \Upsilon_E^0 \mathbf{W}(u),$$

for  $L^n$ -almost every  $u \in E$ .

*Proof.* — We readily infer from 5.2 and 5.4(1) that there exists  $\delta(n, \Lambda, \varepsilon) > 0$  such that, for  $H^{n+m}$ -almost all  $(x, u) \in \Sigma$ , if  $|x - u| \leq \delta(n, \Lambda, \varepsilon)$ , then

$$(18) \quad \alpha := (1 - \varepsilon)2^{-(n-m)/2} \leq J_{\Sigma} \pi_1(x, u)$$

and

$$(19) \quad \beta := (1 + \varepsilon)^{-1}2^{-(n-m)/2} \binom{n}{n-m}^{-1/2} \leq J_{\Sigma} \pi_2(x, u),$$

where the above define  $\alpha$  and  $\beta$ .

Assume now that  $\text{diam } E \leq \delta(n, \Lambda, \varepsilon)$ . Given  $B \in \mathcal{B}(E)$ , we infer from (14), 5.2, 5.4(1), (15), and the above lower bounds, that

$$\begin{aligned} \phi_{E, \mathbf{W}}(B) &= \int_{\Sigma_B} J_{\Sigma} \pi_1 dH^{n+m} > \alpha H^{n+m}(\Sigma_B) > \alpha \int_{\Sigma_B} J_{\Sigma} \pi_2 dH^{n+m} \\ &= \alpha \int_B \Upsilon_E^0 \mathbf{W} dL^n \end{aligned}$$

and

$$\begin{aligned} \phi_{E, \mathbf{W}}(B) &= \int_{\Sigma_B} J_{\Sigma} \pi_1 dH^{n+m} \leq H^{n+m}(\Sigma_B) \leq \beta^{-1} \int_{\Sigma_B} J_{\Sigma} \pi_2 dH^{n+m} \\ &= \beta^{-1} \int_B \Upsilon_E^0 \mathbf{W} dL^n. \end{aligned}$$

Thus,

$$\int_B \alpha \Upsilon_E^0 \mathbf{W} dL^n \leq \int_B Z_E \mathbf{W} dL^n \leq \int_B \beta^{-1} \Upsilon_E^0 \mathbf{W} dL^n,$$

for every  $B \in \mathcal{B}(E)$ . The conclusion follows from the  $L^n$ -measurability of both  $Z_E \mathbf{W}$  and  $\Upsilon_E^0 \mathbf{W}$ .

5.9. REST STOP. — The above upper bound for  $Z_E \mathbf{W}$  is already enough to bound it from above, in turn, by a constant times  $(\text{diam } E)^m$  – see 6.4. We next want to establish that  $Z_E \mathbf{W} > 0$ , almost everywhere in  $E$ . Yet, in the definition (17) of  $\Upsilon_E^0 \mathbf{W}(u)$ ,  $u$  does not appear as the covariable of a function whose level sets we are

measuring, thereby preventing the use of the coarea formula in an attempt to estimate  $\Upsilon_E^0 \mathbf{W}(u)$ . This naturally leads to adding a variable  $y \in \mathbb{R}^{n-m}$  to the fibered space  $\Sigma$ , a covariable for  $g_{\mathbf{v}_1, \dots, \mathbf{v}_{n-m}, u}$ .

5.10. A FIBERED SPACE ASSOCIATED WITH  $E, B, \mathbf{w}_1, \dots, \mathbf{w}_m, \mathbf{v}_1, \dots, \mathbf{v}_{n-m}$

Let  $r > 0$ . Abbreviate  $C_r = \mathbb{R}^{n-m} \cap \{y : |y| \leq r\}$ , the Euclidean ball centered at the origin, of radius  $r$ , in  $\mathbb{R}^{n-m}$ . We define

$$\begin{aligned} \widehat{F}_r : E \times \mathbb{R}^m \times C_r &\longrightarrow \mathbb{R}^n \times \mathbb{R}^n \times \mathbb{R}^{n-m} \\ (x, t, y) &\longmapsto \left( x, x + \sum_{i=1}^m t_i \mathbf{w}_i(x) + \sum_{i=1}^{n-m} y_i \mathbf{v}_i(x), y \right) \end{aligned}$$

and

$$\begin{aligned} \widehat{\Sigma}_r &= \widehat{F}_r(E \times \mathbb{R}^m \times C_r) \\ &= \mathbb{R}^n \times \mathbb{R}^n \times C_r \cap \{(x, u, y) : x \in E \text{ and } u \in \mathbf{W}(x) + \sum_{i=1}^{n-m} y_i \mathbf{v}_i(x)\}. \end{aligned}$$

We note that  $\widehat{F}_r$  is locally Lipschitzian and  $\widehat{\Sigma}_r$  is countably  $2n$ -rectifiable and  $H^{2n}$ -measurable. Similarly to 5.1, we define

$$\widehat{\Sigma}_{r,B} = \widehat{\Sigma}_r \cap \pi_2^{-1}(B)$$

which, clearly, is also countably  $2n$ -rectifiable and  $H^{2n}$ -measurable. We aim to apply the coarea formula to  $\widehat{\Sigma}_{r,B}$  and to the two projections

$$\pi_1 \times \pi_3 : \mathbb{R}^n \times \mathbb{R}^n \times \mathbb{R}^{n-m} \longrightarrow \mathbb{R}^n \times \mathbb{R}^{n-m} : (x, u, y) \longmapsto (x, y)$$

and

$$\pi_2 \times \pi_3 : \mathbb{R}^n \times \mathbb{R}^n \times \mathbb{R}^{n-m} \longrightarrow \mathbb{R}^n \times \mathbb{R}^{n-m} : (x, u, y) \longmapsto (u, y).$$

To this end, we notice that

$$\begin{aligned} \widehat{\Sigma}_{r,B} \cap (\pi_1 \times \pi_3)^{-1}\{(x, y)\} \\ = \mathbb{R}^n \times \mathbb{R}^n \times \mathbb{R}^{n-m} \cap \{(x, u, y) : u \in B \cap (\mathbf{W}(x) + \sum_{i=1}^{n-m} y_i \mathbf{v}_i(x))\} \end{aligned}$$

and, thus,

$$H^m(\widehat{\Sigma}_{r,B} \cap (\pi_1 \times \pi_3)^{-1}\{(x, y)\}) = H^m\left(B \cap (\mathbf{W}(x) + \sum_{i=1}^{n-m} y_i \mathbf{v}_i(x))\right),$$

for every  $(x, y) \in E \times C_r$ . We further notice that

$$\begin{aligned} \widehat{\Sigma}_{r,B} \cap (\pi_2 \times \pi_3)^{-1}\{(u, y)\} \\ = \mathbb{R}^n \times \mathbb{R}^n \times \mathbb{R}^{n-m} \cap \{(x, u, y) : x \in E \text{ and } u \in \mathbf{W}(x) + \sum_{i=1}^{n-m} y_i \mathbf{v}_i(x)\} \\ = \mathbb{R}^n \times \mathbb{R}^n \times \mathbb{R}^{n-m} \cap \{(x, u, y) : x \in E \cap g_{\mathbf{v}_1, \dots, \mathbf{v}_{n-m}, u}^{-1}\{y\}\}, \end{aligned}$$

because

$$\begin{aligned}
 u \in \mathbf{W}(x) + \sum_{i=1}^{n-m} y_i \mathbf{v}_i(x) &\iff u - x - \sum_{i=1}^{n-m} y_i \mathbf{v}_i(x) \in \mathbf{W}_0(x) \\
 &\iff \langle \mathbf{v}_j(x), u - x - \sum_{i=1}^{n-m} y_i \mathbf{v}_i(x) \rangle = 0, \text{ for all } j = 1, \dots, n - m \\
 &\iff g_{\mathbf{v}_1, \dots, \mathbf{v}_{n-m}, u}(x) = y
 \end{aligned}$$

and, therefore,

$$H^m(\widehat{\Sigma}_{r,B} \cap (\pi_2 \times \pi_3)^{-1}\{(u, y)\}) = H^m(E \cap g_{\mathbf{v}_1, \dots, \mathbf{v}_{n-m}, u}^{-1}\{y\}),$$

whenever  $u \in B$  and  $y \in C_r$ .

It now follows from the coarea formula and Fubini's theorem that

$$\begin{aligned}
 (20) \quad \int_{\widehat{\Sigma}_{r,B}} J_{\widehat{\Sigma}_r}(\pi_1 \times \pi_3) dH^{2n} &= \int_E dL^n(x) \int_{C_r} H^m(B \cap (\mathbf{W}(x) + \sum_{i=1}^{n-m} y_i \mathbf{v}_i(x))) dL^{n-m}(y)
 \end{aligned}$$

and

$$\begin{aligned}
 (21) \quad \int_{\widehat{\Sigma}_{r,B}} J_{\widehat{\Sigma}_r}(\pi_2 \times \pi_3) dH^{2n} &= \int_B dL^n(u) \int_{C_r} H^m(E \cap g_{\mathbf{v}_1, \dots, \mathbf{v}_{n-m}, u}^{-1}\{y\}) dL^{n-m}(y).
 \end{aligned}$$

5.11. COAREA JACOBIAN FACTORS OF  $\pi_1 \times \pi_3$  AND  $\pi_2 \times \pi_3$ . — *The following inequalities hold, for  $H^{2n}$ -almost every  $(x, u, y) \in \widehat{\Sigma}_r$ .*

$$2^{-(n-m)} (1 + 2n\Lambda|u - x| + 2n^2\Lambda^2|u - x|^2)^{-n/2} \triangleleft J_{\widehat{\Sigma}_r}(\pi_1 \times \pi_3)(x, u, y)$$

and

$$J_{\widehat{\Sigma}_r}(\pi_2 \times \pi_3)(x, u, y) \triangleleft 1.$$

*Proof.* — The second conclusion is obvious, since  $\text{Lip } \pi_2 \times \pi_3 = 1$ . Regarding the first conclusion, we reason similarly to the proof of 5.2. Fix  $(x, u, y) = \widehat{F}_r(x, t, y)$  such that  $\widehat{F}_r$  is differentiable at  $(x, t, y)$  and denote by  $L$  the restriction of  $\pi_1 \times \pi_3$  to  $T_{(x,u,y)}\widehat{\Sigma}_r$ . This tangent space is generated by the following  $2n$  vectors of  $\mathbb{R}^n \times \mathbb{R}^n \times \mathbb{R}^{n-m}$ :

$$\begin{aligned}
 \frac{\partial \widehat{F}_r}{\partial x_j}(x, t, y) &= \left( e_j, e_j + \sum_{i=1}^m t_i \frac{\partial \mathbf{w}_i(x)}{\partial x_j}(x) + \sum_{i=1}^{n-m} y_i \frac{\partial \mathbf{v}_i(x)}{\partial x_j}(x), 0 \right), \quad j = 1, \dots, n, \\
 \frac{\partial \widehat{F}_r}{\partial t_k}(x, t, y) &= (0, \mathbf{w}_k(x), 0), \quad k = 1, \dots, m, \\
 \frac{\partial \widehat{F}_r}{\partial y_\ell}(x, t, y) &= (0, \mathbf{v}_\ell(x), e_\ell), \quad \ell = 1, \dots, n - m.
 \end{aligned}$$

The range of  $\pi_1 \times \pi_3$  being  $(2n - m)$ -dimensional, we need to select  $2n - m$  vectors  $v_1, \dots, v_{2n-m}$  in  $T_{(x,u,y)}\widehat{\Sigma}_r$  in view of obtaining a lower bound

$$(22) \quad J_{\widehat{\Sigma}_r}(\pi_1 \times \pi_3)(x, u, y) = \|\wedge_{2n-m} L\| > \frac{|L(v_1) \wedge \dots \wedge L(v_{2n-m})|}{|v_1 \wedge \dots \wedge v_{2n-m}|}.$$

Our choice of  $v_1, \dots, v_{2n-m}$  is as follows. As in the proof of 5.2, we let  $w_j = \mathbf{w}_j(x)$ , for  $j = 1, \dots, m$ , and  $w_j = \mathbf{v}_{j-m}(x)$ , for  $j = m + 1, \dots, n$ . For  $j = 1, \dots, m$ , we define

$$v_j = \left( \sum_{p=1}^n \langle w_j, e_p \rangle \frac{\partial \widehat{F}_r}{\partial x_p}(x, t, y) \right) - \frac{\partial \widehat{F}_r}{\partial t_j}(x, t, y) \\ = \left( w_j, \sum_{i=1}^m t_i D\mathbf{w}_i(x)(w_j) + \sum_{i=1}^{n-m} y_i D\mathbf{v}_i(x)(w_j), 0 \right),$$

for  $j = m + 1, \dots, n$ , we define

$$v_j = \sum_{p=1}^m \langle w_j, e_p \rangle \frac{\partial \widehat{F}_r}{\partial x_p}(x, t, y) = \left( w_j, w_j + \sum_{i=1}^m t_i D\mathbf{w}_i(x)(w_j) + \sum_{i=1}^{n-m} y_i D\mathbf{v}_i(x)(w_j), 0 \right),$$

and, for  $j = n + 1, \dots, 2n - m$ , we define

$$v_j = \frac{\partial \widehat{F}_r}{\partial y_{j-n}}(x, t, y) = (0, \mathbf{v}_{j-n}(x), e_{j-n}),$$

so that  $L(v_1), \dots, L(v_{2n-m})$  is an orthonormal basis of  $\mathbb{R}^n \times \mathbb{R}^{n-m}$  and, therefore, the numerator in (22) equals 1. In order to determine an upper bound for its denominator, we start by fixing  $j = 1, \dots, n$ , we abbreviate  $a_j(x, t, y) = \sum_{i=1}^m t_i D\mathbf{w}_i(x)(w_j)$  and  $b_j(x, t, y) = \sum_{i=1}^{n-m} y_i D\mathbf{v}_i(x)(w_j)$ , and we notice that  $|a_j(x, t, y)| \leq m\Lambda|t| \leq n\Lambda|t|$ ,  $|b_j(x, t, y)| \leq (n - m)\Lambda|y| \leq n\Lambda|y|$ . Furthermore, since  $u - x = \sum_{i=1}^m t_i \mathbf{w}_i(x) + \sum_{i=1}^{n-m} y_i \mathbf{v}_i(x)$ , one has  $|u - x|^2 = |t|^2 + |y|^2 > \max\{|t|^2, |y|^2\}$ . Therefore, if  $j = 1, \dots, m$ , then

$$|v_j|^2 = |w_j|^2 + |a_j(x, t, y) + b_j(x, t, y)|^2 \\ \leq 1 + |a_j(x, t, y)|^2 + |b_j(x, t, y)|^2 + 2|a_j(x, t, y)||b_j(x, t, y)| \\ \leq 1 + n^2\Lambda^2(|t|^2 + |y|^2 + 2|t||y|) \\ \leq 1 + 2n^2\Lambda^2|u - x|^2,$$

whereas, if  $j = m + 1, \dots, n$ , then

$$|v_j|^2 = |w_j|^2 + |w_j + a_j(x, t, y) + b_j(x, t, y)|^2 \\ \leq 1 + 1 + |a_j(x, t, y)|^2 + |b_j(x, t, y)|^2 \\ \quad + 2|a_j(x, t, y)| + 2|b_j(x, t, y)| + 2|a_j(x, t, y)||b_j(x, t, y)| \\ \leq 2(1 + n^2\Lambda^2|u - x|^2 + 2n\Lambda|u - x|),$$

and, if  $j = n + 1, \dots, 2n - m$ , then

$$|v_j| = \sqrt{2}.$$

We conclude that

$$|v_1 \wedge \dots \wedge v_{2n-m}| \\ \leq 2^{n-m} (1 + 2n^2\Lambda^2|u - x|^2)^{m/2} (1 + n^2\Lambda^2|u - x|^2 + 2n\Lambda|u - x|)^{(n-m)/2} \\ \leq 2^{n-m} (1 + 2n^2\Lambda^2|u - x|^2 + 2n\Lambda|u - x|)^{n/2}$$

and the proof is complete.



5.12. DEFINITION OF  $\Upsilon_E \mathbf{W}$ . — It follows from the coarea theorem that the function

$$\mathbb{R}^n \times \mathbb{R}^{n-m} \longrightarrow [0, \infty] : (u, y) \longrightarrow H^m(E \cap g_{\mathbf{v}_1, \dots, \mathbf{v}_{n-m}, u}^{-1}\{y\})$$

is  $L^n \otimes L^{n-m}$ -measurable (recall 5.10 applied with  $B = \mathbb{R}^n$ ). It now follows from Fubini's theorem that, for each  $r > 0$ , the function

$$\mathbb{R}^n \longrightarrow [0, \infty] : u \longmapsto \int_{C_r} H^m(E \cap g_{\mathbf{v}_1, \dots, \mathbf{v}_{n-m}, u}^{-1}\{y\}) dL^{n-m}(y)$$

is  $L^n$ -measurable. In turn, the function

$$\Upsilon_E \mathbf{W} : \mathbb{R}^n \longrightarrow [0, \infty] : u \longmapsto \liminf_j \int_{C_{j-1}} H^m(E \cap g_{\mathbf{v}_1, \dots, \mathbf{v}_{n-m}, u}^{-1}\{y\}) dL^{n-m}(y)$$

is  $L^n$ -measurable. It is a replacement for  $\Upsilon_E^0 \mathbf{W}$  defined in 5.7. We shall establish, for  $Z_E \mathbf{W}$ , a similar lower bound to that in 5.8, this time involving  $\Upsilon_E \mathbf{W}$ . Before doing so, we notice the rather trivial fact that if  $F \subseteq E$ , then

$$\Upsilon_F \mathbf{W}(u) \leq \Upsilon_E \mathbf{W}(u),$$

for all  $u \in \mathbb{R}^n$ .

5.13. PREPARATORY REMARK FOR THE PROOF OF 5.15. — It follows from the coarea theorem that the function

$$\mathbb{R}^n \times \mathbb{R}^{n-m} \longrightarrow [0, \infty] : (x, y) \longmapsto H^m\left(B \cap (\mathbf{W}(x) + \sum_{i=1}^{n-m} y_i \mathbf{v}_i(x))\right)$$

is  $L^n \otimes L^{n-m}$ -measurable (recall 5.10 applied with  $B = \mathbb{R}^n$ ). It therefore follows from Fubini's theorem as in 5.12 that

$$f_j : \mathbb{R}^n \longrightarrow [0, \infty] : x \longmapsto \int_{C_{j-1}} H^m\left(B \cap (\mathbf{W}(x) + \sum_{i=1}^{n-m} y_i \mathbf{v}_i(x))\right) dL^{n-m}(y)$$

is  $L^n$ -measurable. Furthermore, if  $B$  is bounded, then  $|f_j(x)| \leq \alpha(m)(\text{diam } B)^m$ , for every  $x \in \mathbb{R}^n$ .

5.14. — If  $B$  is compact, then, for every  $x \in E$ , the function

$$\mathbb{R}^{n-m} \longrightarrow \mathbb{R}_+ : y \longmapsto H^m\left(B \cap (\mathbf{W}(x) + \sum_{i=1}^{n-m} y_i \mathbf{v}_i(x))\right)$$

is upper semicontinuous.

*Proof.* — The proof is analogous to that of 3.14. For each  $y \in \mathbb{R}^{n-m}$ , define the compact set  $K_y = B \cap (\mathbf{W}(x) + \sum_{i=1}^{n-m} y_i \mathbf{v}_i(x))$ . If  $(y_k)_k$  is a sequence converging to  $y$ , we ought to show that

$$H_\infty^m(K_y) \geq \limsup_k H_\infty^m(K_{y_k}).$$

Since each  $K_{y_k}$  is a subset of an  $m$ -dimensional affine subspace of  $\mathbb{R}^n$ , this is indeed equivalent to the same inequality with  $H_\infty^m$  replaced by  $H^m$ , according to 3.2(3). Considering, if necessary, a subsequence of  $(y_k)_k$ , we may assume that none of the compact sets  $K_{y_k}$  is empty and that the above limit superior is a limit. Considering yet a further subsequence, we may now assume that  $(K_{y_k})_k$  converges in Hausdorff

distance to some compact set  $L \subseteq B$ . One checks that  $L \subseteq K_y$ . It then follows from 3.2(1) that  $H^m_\infty(K_y) > H^m_\infty(L) > \limsup_k H^m_\infty(K_{y_k})$ .

5.15. PROPOSITION. — Given  $0 < \varepsilon < 1$ , there exists  $\delta_{5.15}(n, \Lambda, \varepsilon) > 0$  with the following property. If  $\text{diam}(E \cup B) \leq \delta_{5.15}(n, \Lambda, \varepsilon)$  and  $B$  is compact, then

$$\int_E H^m(B \cap \mathbf{W}(x)) dL^n(x) > (1 - \varepsilon)2^{-(n-m)} \int_B \chi_E \mathbf{W}(u) dL^n(u).$$

Proof. — We first observe that we can choose  $\delta_{5.15}(n, \Lambda, \varepsilon) > 0$  small enough so that

$$(23) \quad J_{\widehat{\Sigma}_r}(\pi_1 \times \pi_3)(x, u, y) > (1 - \varepsilon)2^{-(n-m)},$$

for  $H^{2n}$ -almost every  $(x, u, y) \in \widehat{\Sigma}_r$ , provided  $|u - x| \leq \delta_{5.15}(n, \Lambda, \varepsilon)$ , according to 5.11. Thus, (23) holds, for  $H^{2n}$ -almost every  $(x, u, y) \in \widehat{\Sigma}_{r,B}$ , under the assumption that  $\text{diam}(E \cup B) \leq \delta_{5.15}(n, \Lambda, \varepsilon)$ . In that case, (20), (21), and 5.11 imply that

$$\begin{aligned} & \int_E dL^n(x) \int_{C_r} H^m\left(B \cap (\mathbf{W}(x) + \sum_{i=1}^{n-m} y_i \mathbf{v}_i(x))\right) dL^{n-m}(y) \\ &= \int_{\widehat{\Sigma}_{r,B}} J_{\widehat{\Sigma}_r}(\pi_1 \times \pi_3) dH^{2n} \\ (24) \quad & > (1 - \varepsilon)2^{-(n-m)} H^{2n}(\widehat{\Sigma}_{r,B}) \\ & > (1 - \varepsilon)2^{-(n-m)} \int_{\widehat{\Sigma}_{r,B}} J_{\widehat{\Sigma}_r}(\pi_2 \times \pi_3) dH^{2n} \\ &= (1 - \varepsilon)2^{-(n-m)} \int_B dL^n(u) \int_{C_r} H^m(E \cap g_{\mathbf{v}_1, \dots, \mathbf{v}_{n-m}, u}^{-1}\{y\}) dL^{n-m}(y). \end{aligned}$$

Fix  $x \in E$  and  $\beta > 0$ . According to 5.14, there exists a positive integer  $j(x, \beta)$  such that if  $j > j(x, \beta)$ , then

$$\begin{aligned} H^m(B \cap \mathbf{W}(x)) + \beta &> \sup_{y \in C_{j-1}} H^m\left(B \cap (\mathbf{W}(x) + \sum_{i=1}^{n-m} y_i \mathbf{v}_i(x))\right) \\ &> \int_{C_{j-1}} H^m\left(B \cap (\mathbf{W}(x) + \sum_{i=1}^{n-m} y_i \mathbf{v}_i(x))\right) dL^{n-m}(y). \end{aligned}$$

Taking the limit superior as  $j \rightarrow \infty$ , on the right hand side, and letting  $\beta \rightarrow 0$ , we obtain

$$(25) \quad H^m(B \cap \mathbf{W}(x)) > \limsup_j \int_{C_{j-1}} H^m\left(B \cap (\mathbf{W}(x) + \sum_{i=1}^{n-m} y_i \mathbf{v}_i(x))\right) dL^{n-m}(y).$$

As this holds for all  $x \in E$ , we may integrate over  $E$  with respect to  $L^n$ . We notice that for every  $j = 1, 2, \dots$ ,  $|f_j| \leq \alpha(m)(\text{diam } B)^m \mathbb{1}_{\mathbb{R}^n}$  (recall the notation of 5.13); the latter being  $L^n \llcorner E$ -summable, this justifies the application of the reverse Fatou lemma below. Thus, the following ensues from (25), the reverse Fatou lemma, (24),

and the Fatou lemma:

$$\begin{aligned}
 & \int_E H^m(B \cap \mathbf{W}(x)) dL^n(x) \\
 & > \int_E dL^n(x) \limsup_j \int_{C_{j-1}} H^m\left(B \cap (\mathbf{W}(x) + \sum_{i=1}^{n-m} y_i \mathbf{v}_i(x))\right) dL^{n-m}(y) \\
 & > \limsup_j \int_E dL^n(x) \int_{C_{j-1}} H^m\left(B \cap (\mathbf{W}(x) + \sum_{i=1}^{n-m} y_i \mathbf{v}_i(x))\right) dL^{n-m}(y) \\
 & > (1 - \varepsilon) 2^{-(n-m)} \limsup_j \int_B dL^n(u) \int_{C_{j-1}} H^m(E \cap g_{\mathbf{v}_1, \dots, \mathbf{v}_{n-m}, u}^{-1}\{y\}) dL^{n-m}(y) \\
 & > (1 - \varepsilon) 2^{-(n-m)} \liminf_j \int_B dL^n(u) \int_{C_{j-1}} H^m(E \cap g_{\mathbf{v}_1, \dots, \mathbf{v}_{n-m}, u}^{-1}\{y\}) dL^{n-m}(y) \\
 & > (1 - \varepsilon) 2^{-(n-m)} \int_B dL^n(u) \liminf_j \int_{C_{j-1}} H^m(E \cap g_{\mathbf{v}_1, \dots, \mathbf{v}_{n-m}, u}^{-1}\{y\}) dL^{n-m}(y) \\
 & = (1 - \varepsilon) 2^{-(n-m)} \int_B \Upsilon_E \mathbf{W}(u) dL^n(u).
 \end{aligned}$$

5.16. COROLLARY. — If  $0 < \varepsilon < 1$  and  $\text{diam } E \leq \delta_{5.15}(n, \Lambda, \varepsilon)$ , then

$$Z_E \mathbf{W}(u) > (1 - \varepsilon) 2^{-(n-m)} \Upsilon_E \mathbf{W}(u),$$

for  $L^n$ -almost every  $u \in E$ .

### 6. UPPER BOUND FOR $\Upsilon_E \mathbf{W}$ AND $Z_E \mathbf{W}$

6.1. BOW TIE LEMMA. — Let  $S \subseteq \mathbb{R}^n$ ,  $W \in G(n, m)$ , and  $0 < \tau < 1$ . Assume that

$$(\forall x \in S)(\forall 0 < \rho \leq \text{diam } S) : S \cap \mathbf{B}(x, \rho) \subseteq \mathbf{B}(x + W, \tau \rho).$$

Then there exists  $F : P_W(S) \rightarrow \mathbb{R}^n$  such that  $S = \text{im } F$  and  $\text{Lip } F \leq 1/\sqrt{1 - \tau^2}$ . In particular,

$$H^m(S) \leq \left(\frac{1}{\sqrt{1 - \tau^2}}\right)^m \alpha(m) (\text{diam } S)^m.$$

*Proof.* — Let  $x, x' \in S$  and define  $\rho = |x - x'| \leq \text{diam } S$ . Thus,  $x' \in S \cap \mathbf{B}(x, \rho)$  and, therefore,  $|P_{W^\perp}(x - x')| \leq \tau \rho = \tau |x - x'|$ . Since  $|x - x'|^2 = |P_W(x - x')|^2 + |P_{W^\perp}(x - x')|^2$ , we infer that

$$(1 - \tau^2) |x - x'|^2 \leq |P_W(x - x')|^2.$$

Therefore,  $P_W|_S$  is injective; and the Lipschitzian bound on  $F = (P_W|_S)^{-1}$  clearly follows from the above inequality. Regarding the second conclusion, we note that

$$H^m(S) = H^m(F(P_W(S))) \leq (\text{Lip } F)^m H^m(P_W(S))$$

and  $P_W(S)$  is contained in a ball of radius  $\text{diam } P_W(S) \leq \text{diam } S$ .

6.2. — Given  $0 < \tau < 1$ , there exists  $\delta_{6.2}(n, \Lambda, \tau) > 0$  with the following property. If

- (1)  $x_0 \in U$  and  $u \in \mathbb{R}^n$ ;
- (2)  $\text{diam}(E \cup \{x_0\} \cup \{u\}) \leq \delta_{6.2}(n, \Lambda, \tau)$ ;

then: For each  $y \in \mathbb{R}^{n-m}$ , each  $x \in E \cap g_{\mathbf{v}_1, \dots, \mathbf{v}_{n-m}, u}^{-1}\{y\}$ , and each  $0 < \rho < \infty$ , one has

$$E \cap g_{\mathbf{v}_1, \dots, \mathbf{v}_{n-m}, u}^{-1}\{y\} \cap \mathbf{B}(x, \rho) \subseteq \mathbf{B}(x + \mathbf{W}_0(x_0), \tau\rho).$$

*Proof.* — We show that  $\delta_{6.2}(n, \Lambda, \tau) = \frac{\tau}{2\Lambda\sqrt{n}}$  will do. Let  $x, x' \in E \cap g_{\mathbf{v}_1, \dots, \mathbf{v}_{n-m}, u}^{-1}\{y\}$ , for some  $y \in \mathbb{R}^{n-m}$ . Thus,  $g_{\mathbf{v}_1, \dots, \mathbf{v}_{n-m}, u}(x) = g_{\mathbf{v}_1, \dots, \mathbf{v}_{n-m}, u}(x')$  and, hence,

$$\begin{aligned} 0 &= |g_{\mathbf{v}_1, \dots, \mathbf{v}_{n-m}, u}(x) - g_{\mathbf{v}_1, \dots, \mathbf{v}_{n-m}, u}(x')| \\ &= \sqrt{\sum_{i=1}^{n-m} |\langle \mathbf{v}_i(x), x - u \rangle - \langle \mathbf{v}_i(x'), x' - u \rangle|^2} \\ &= \sqrt{\sum_{i=1}^{n-m} |\langle \mathbf{v}_i(x), x - x' \rangle - \langle \mathbf{v}_i(x') - \mathbf{v}_i(x), x' - u \rangle|^2} \\ &> \sqrt{\sum_{i=1}^{n-m} |\langle \mathbf{v}_i(x), x - x' \rangle|^2} - \sqrt{\sum_{i=1}^{n-m} |\langle \mathbf{v}_i(x') - \mathbf{v}_i(x), x' - u \rangle|^2}, \end{aligned}$$

thus,

$$\begin{aligned} \sqrt{\sum_{i=1}^{n-m} |\langle \mathbf{v}_i(x), x - x' \rangle|^2} &\leq \sqrt{\sum_{i=1}^{n-m} |\langle \mathbf{v}_i(x') - \mathbf{v}_i(x), x' - u \rangle|^2} \\ &\leq \sqrt{n-m} \Lambda |x - x'| |x' - u| \leq \frac{\tau}{2} |x - x'|. \end{aligned}$$

In turn,

$$\begin{aligned} |P_{\mathbf{W}_0(x_0)^\perp}(x - x')| &= \sqrt{\sum_{i=1}^{n-m} |\langle \mathbf{v}_i(x_0), x - x' \rangle|^2} \\ &\leq \sqrt{\sum_{i=1}^{n-m} |\langle \mathbf{v}_i(x'), x - x' \rangle|^2} + \sqrt{\sum_{i=1}^{n-m} |\langle \mathbf{v}_i(x') - \mathbf{v}_i(x_0), x - x' \rangle|^2} \\ &\leq \frac{\tau}{2} |x - x'| + \sqrt{n-m} \Lambda |x' - x_0| |x - x'| \leq \tau |x - x'|. \end{aligned}$$

**6.3. PROPOSITION.** — *There are  $\delta_{6.3}(n, \Lambda) > 0$  and  $c_{6.3}(m) > 1$  with the following property. If  $u \in \mathbb{R}^n$  and  $\text{diam}(E \cup \{u\}) \leq \delta_{6.3}(n, \Lambda)$ , then*

$$\max \{ \Upsilon_E^0 \mathbf{W}(u), \Upsilon_E \mathbf{W}(u) \} \leq c_{6.3}(m) (\text{diam } E)^m.$$

*Proof.* — Let  $\delta_{6.3}(n, \Lambda) = \delta_{6.2}(n, \Lambda, 1/2)$ . Recall the definitions of  $\Upsilon_E^0 \mathbf{W}$  and  $\Upsilon_E \mathbf{W}$ , 5.7 and 5.12, respectively. If  $E = \{u\}$ , the conclusion is obvious. If not, pick  $x_0 \in E$  arbitrarily. Given any  $y \in \mathbb{R}^{n-m}$ , we see that 6.2 applies with  $\tau = 1/2$  and, in turn, the bow tie lemma 6.1 applies to  $S = E \cap g_{\mathbf{v}_1, \dots, \mathbf{v}_{n-m}, u}^{-1}\{y\}$  and  $W = \mathbf{W}_0(x_0)$ . Thus,

$$H^m(E \cap g_{\mathbf{v}_1, \dots, \mathbf{v}_{n-m}, u}^{-1}\{y\}) \leq \left(\frac{2}{\sqrt{3}}\right)^m \alpha(m) (\text{diam } E)^m.$$

The proposition is proved.

**6.4. COROLLARY.** — *There are  $\delta_{6.4}(n, \Lambda) > 0$  and  $c_{6.4}(n) > 1$  with the following property. If  $\text{diam } E \leq \delta_{6.4}(n, \Lambda)$ , then*

$$Z_E \mathbf{W}(u) \leq c_{6.4}(n) (\text{diam } E)^m,$$

for  $L^n$ -almost every  $u \in E$ .

*Proof.* — Let  $\delta_{6.4}(n, \Lambda) = \min\{\delta_{6.3}(n, \Lambda), \delta_{5.8}(n, \Lambda, 1/2)\}$ .

7. LOWER BOUND FOR  $Y_E \mathbf{W}$  AND  $Z_E \mathbf{W}$

7.1. SETTING FOR THIS SECTION. — We enforce again the exact same assumptions as in 4.1; and as in 5.10, we let  $C_r = \mathbb{R}^{n-m} \cap \{y : |y| \leq r\}$ .

7.2. POLYBALLS. — Given  $x_0 \in \mathbb{R}^n$  and  $r > 0$ , we define

$$\mathbf{C}_{\mathbf{W}}(x_0, r) = \mathbb{R}^n \cap \{x : |P_{\mathbf{W}_0(x_0)}(x - x_0)| \leq r \text{ and } |P_{\mathbf{W}_0(x_0)^\perp}(x - x_0)| \leq r\}.$$

We notice that, if  $x \in \mathbf{C}_{\mathbf{W}}(x_0, r)$ , then  $|x - x_0| \leq r\sqrt{2}$ ; in particular,  $\text{diam } \mathbf{C}_{\mathbf{W}}(x_0, r) \leq 2\sqrt{2}r$ . We also notice that  $L^n(\mathbf{C}_{\mathbf{W}}(x_0, r)) = \alpha(m)\alpha(n - m)r^n$ .

7.3. — Given  $0 < \varepsilon < 1/3$ , there exists  $\delta_{7.3}(n, \Lambda, \varepsilon) > 0$  with the following property. If

- (1)  $0 < r < \delta_{7.3}(n, \Lambda, \varepsilon)$ ;
- (2)  $u \in \mathbf{C}_{\mathbf{W}}(x_0, r) \subseteq U$ ;
- (3)  $|g_{\mathbf{v}_1, \dots, \mathbf{v}_{n-m}, u}(x_0)| \leq (1 - 3\varepsilon)r$ ;
- (4)  $C \subseteq C_{\varepsilon r}$  is closed;

then

$$L^n(\mathbf{C}_{\mathbf{W}}(x_0, r) \cap g_{\mathbf{v}_1, \dots, \mathbf{v}_{n-m}, u}^{-1}(C)) > \frac{1}{1 + \varepsilon} \alpha(m)r^m L^{n-m}(C).$$

7.4. REMARK. — With hopes that the following will help the reader form a geometrical imagery: Under the circumstances 7.3,  $\mathbf{C}_{\mathbf{W}}(x_0, r) \cap g_{\mathbf{v}_1, \dots, \mathbf{v}_{n-m}, u}^{-1}(C)$  may be seen as a “nonlinear stripe”, “horizontal” with respect to  $\mathbf{W}_0(x_0)$ , “at height”  $g_{\mathbf{v}_1, \dots, \mathbf{v}_{n-m}, u}(x_0)$  with respect to  $x_0$ , and of “width”  $C$ .

Proof of 7.3. — Given  $z \in \mathbf{W}_0(x_0) \cap B(0, r)$ , we define

$$V_z = \mathbb{R}^n \cap \{x_0 + z + \sum_{i=1}^{n-m} y_i \mathbf{v}_i(x_0) : y \in C_r\} \subseteq \mathbf{C}_{\mathbf{W}}(x_0, r)$$

and we consider the isometric parametrization  $\gamma_z : C_r \rightarrow V_z$  defined by the formula

$$\gamma_z(y) = x_0 + z + \sum_{i=1}^{n-m} y_i \mathbf{v}_i(x_0).$$

We also abbreviate  $f_{z,u} = g_{\mathbf{v}_1, \dots, \mathbf{v}_{n-m}, u} \circ \gamma_z$ .

CLAIM #1.  $\text{Lip } f_{z,u} \leq (1 + \varepsilon)^1 / (n - m)$ . — Since  $\gamma_z$  is an isometry, it suffices to obtain an upper bound for  $\text{Lip } g_{\mathbf{v}_1, \dots, \mathbf{v}_{n-m}, u}|_{\mathbf{C}_{\mathbf{W}}(x_0, r)}$ . Let  $x, x' \in \mathbf{C}_{\mathbf{W}}(x_0, r)$  and note that

$$\begin{aligned} |g_{\mathbf{v}_1, \dots, \mathbf{v}_{n-m}, u}(x) - g_{\mathbf{v}_1, \dots, \mathbf{v}_{n-m}, u}(x')| &= \sqrt{\sum_{i=1}^{n-m} |\langle \mathbf{v}_i(x), x - u \rangle - \langle \mathbf{v}_i(x'), x' - u \rangle|^2} \\ &\leq \sqrt{\sum_{i=1}^{n-m} (|\langle \mathbf{v}_i(x) - \mathbf{v}_i(x'), x - u \rangle| + |\langle \mathbf{v}_i(x'), x - x' \rangle|)^2} \\ &\leq \sqrt{\sum_{i=1}^{n-m} |\langle \mathbf{v}_i(x) - \mathbf{v}_i(x'), x - u \rangle|^2} + \sqrt{\sum_{i=1}^{n-m} |\langle \mathbf{v}_i(x'), x - x' \rangle|^2} \\ &\leq \sqrt{n - m} \Lambda |x - x'| |x - u| + |P_{\mathbf{W}_0(x')^\perp}(x - x')| \\ &\leq (1 + \sqrt{n - m} \Lambda 2\sqrt{2}r) |x - x'|. \end{aligned}$$

Recalling hypothesis (1), it is now apparent that  $\delta_{7.3}$  can be chosen small enough, according to  $n, \Lambda$ , and  $\varepsilon$ , so that Claim #1 holds.

CLAIM #2. For  $L^{n-m}$ -almost every  $y \in C_r$ , one has  $\|Df_{z,u}(y) - \text{id}_{\mathbb{R}^{n-m}}\| \leq \varepsilon$ .

Let  $y \in C_r$  be such that  $f_{z,u}$  is differentiable at  $y$ . We shall estimate the coefficients of the matrix representing  $Df_{z,u}(y)$  with respect to the canonical basis. Fix  $i, j = 1, \dots, n - m$  and recall (7):

$$\begin{aligned} \frac{\partial}{\partial y_i} \langle f_{u,z}(y), e_j \rangle &= \frac{\partial}{\partial y_i} \langle g_{\mathbf{v}_1, \dots, \mathbf{v}_{n-m}, u}(\gamma_z(y)), e_j \rangle \\ &= \left( \frac{\partial}{\partial y_i} \right) g_{\mathbf{v}_j, u}(\gamma_z(y)) \\ &= \left\langle \nabla g_{\mathbf{v}_j, u}(\gamma_z(y)), \frac{\partial \gamma_z(y)}{\partial y_i} \right\rangle \\ &= \langle D\mathbf{v}_j(\gamma_z(y))(\mathbf{v}_i(x_0)), \gamma_z(y) - u \rangle + \langle \mathbf{v}_j(\gamma_z(y)), \mathbf{v}_i(x_0) \rangle \\ &= \text{I} + \text{II}. \end{aligned}$$

Next, note that

$$\begin{aligned} |\text{II} - \delta_{ij}| &= |\text{II} - \langle \mathbf{v}_j(x_0), \mathbf{v}_i(x_0) \rangle| = |\langle \mathbf{v}_j(\gamma_z(y)) - \mathbf{v}_j(x_0), \mathbf{v}_i(x_0) \rangle| \\ &\leq \Lambda |\gamma_z(y) - x_0| \leq \Lambda 2\sqrt{2}r \leq \frac{\varepsilon}{2(n - m)}, \end{aligned}$$

where the last inequality follows from hypothesis (1), upon choosing  $\delta_{7.3}$  small enough, according to  $n, \Lambda$ , and  $\varepsilon$ . Moreover,

$$|\text{I}| \leq \Lambda |\gamma_z(y) - u| \leq \Lambda 2\sqrt{2}r \leq \frac{\varepsilon}{2(n - m)}.$$

Therefore, if  $(a_{ij})_{i,j=1, \dots, n-m}$  is the matrix representing  $Df_{z,u}(y)$  with respect to the canonical basis, we have shown that  $|a_{ij} - \delta_{ij}| \leq \varepsilon/(n - m)$ , for all  $i, j = 1, \dots, n - m$ . This completes the proof of Claim #2.

CLAIM #3.  $C_{\varepsilon r} \subseteq f_{z,u}(C_r)$ . — We shall show that  $|y - f_{z,u}(y)| \leq (1 - \varepsilon)r$ , for every  $y \in \text{Bdry } C_r$ , and the conclusion will become a consequence of the intermediate value theorem, in case  $m = n - 1$ , or a standard application of homology theory, see, e.g., [4, 4.6.1], in case  $m < n - 1$ . If  $m < n - 1$ , it is clearly enough to establish this inequality only for  $H^{n-m-1}$ -almost every  $y \in \text{Bdry } C_r$ . In that case, owing to the coarea theorem [8, 3.2.22], we may choose such a  $y$  so that that  $f_{z,u}$  is differentiable  $H^1$ -almost everywhere on the line segment  $\mathbb{R}^{n-m} \cap \{sy : 0 \leq s \leq 1\}$ . Whether  $m < n - 1$ , or  $m = n - 1$ , it then follows from Claim #2 that

$$\begin{aligned} |f_{z,u}(y) - f_{z,u}(0) - y| &= \left| \int_0^1 Df_{z,u}(sy)(y) dL^1(s) - y \right| \\ &\leq \int_0^1 |Df_{z,u}(sy)(y) - y| dL^1(s) \leq \varepsilon|y| = \varepsilon r. \end{aligned}$$

Accordingly,

$$|f_{z,u}(y) - y| \leq |f_{z,u}(y) - f_{z,u}(0) - y| + |f_{z,u}(0)| \leq \varepsilon r + |f_{z,u}(0)|$$

and the claim will be established upon showing that  $|f_{z,u}(0)| \leq (1 - 2\varepsilon)r$ . Note that  $f_{z,u}(0) = g_{\mathbf{v}_1, \dots, \mathbf{v}_{n-m}, u}(x_0 + z)$ ; we shall use hypothesis (3) to bound its norm from

above. Given  $j = 1, \dots, n - m$ , recall that  $\langle \mathbf{v}_j(x_0), z \rangle = 0$ , thus,

$$\begin{aligned} |g_{\mathbf{v}_j, u}(x_0 + z) - g_{\mathbf{v}_j, u}(x_0)| &= |\langle \mathbf{v}_j(x_0 + z), x_0 + z - u \rangle - \langle \mathbf{v}_j(x_0), x_0 - u \rangle| \\ &= |\langle \mathbf{v}_j(x_0 + z), x_0 + z - u \rangle - \langle \mathbf{v}_j(x_0), x_0 + z - u \rangle| \leq \Lambda |z| |x_0 + z - u| \\ &\leq \Lambda r 2\sqrt{2}r \leq \frac{\varepsilon r}{\sqrt{n - m}}, \end{aligned}$$

where the last inequality holds, according to hypothesis (1), provided  $\delta_{7.3}$  is chosen sufficiently small. In turn,

$$\begin{aligned} |f_{z, u}(0)| &\leq |g_{\mathbf{v}_1, \dots, \mathbf{v}_{n-m}, u}(x_0 + z) - g_{\mathbf{v}_1, \dots, \mathbf{v}_{n-m}, u}(x_0)| + |g_{\mathbf{v}_1, \dots, \mathbf{v}_{n-m}, u}(x_0)| \\ &\leq \varepsilon r + (1 - 3\varepsilon)r = (1 - 2\varepsilon)r, \end{aligned}$$

by virtue of hypothesis (3).

**CLAIM #4.** For every  $z \in \mathbf{W}_0(x_0) \cap \mathbf{B}(0, r)$  and every closed  $C \subseteq C_{\varepsilon r}$ , one has  $H^{n-m}(C) \leq (1 + \varepsilon)H^{n-m}(g_{\mathbf{v}_1, \dots, \mathbf{v}_{n-m}, u}^{-1}(C) \cap V_z)$ .

First, notice that

$$g_{\mathbf{v}_1, \dots, \mathbf{v}_{n-m}, u}^{-1}(C) \cap V_z = \gamma_z(\gamma_z^{-1}(g_{\mathbf{v}_1, \dots, \mathbf{v}_{n-m}, u}^{-1}(C) \cap V_z)) = \gamma_z(f_{z, u}^{-1}(C))$$

and, therefore,

$$H^{n-m}(g_{\mathbf{v}_1, \dots, \mathbf{v}_{n-m}, u}^{-1}(C) \cap V_z) = H^{n-m}(f_{z, u}^{-1}(C)),$$

since  $\gamma_z$  is an isometry. Now, since  $C \subseteq C_{\varepsilon r} \subseteq f_{z, u}(C_r)$ , according to Claim #3, we have

$$C = f_{z, u}(f_{z, u}^{-1}(C)).$$

It therefore follows from Claim #1 that

$$\begin{aligned} H^{n-m}(C) &\leq (\text{Lip } f_{z, u})^{n-m} H^{n-m}(f_{z, u}^{-1}(C)) \\ &\leq (1 + \varepsilon)H^{n-m}(g_{\mathbf{v}_1, \dots, \mathbf{v}_{n-m}, u}^{-1}(C) \cap V_z). \end{aligned}$$

We are now ready to finish the proof, by an application of Fubini's theorem:

$$\begin{aligned} L^n(\mathbf{C}_{\mathbf{W}}(x_0, r) \cap g_{\mathbf{v}_1, \dots, \mathbf{v}_{n-m}, u}^{-1}(C)) &= \int_{\mathbf{W}_0(x_0) \cap \mathbf{B}(0, r)} dL^m(z) H^{n-m}(g_{\mathbf{v}_1, \dots, \mathbf{v}_{n-m}, u}^{-1}(C) \cap V_z) \\ &> \frac{1}{1 + \varepsilon} \alpha(m) r^m H^{n-m}(C). \end{aligned}$$

**7.5. LOWER BOUND FOR  $Y_E \mathbf{W}$ .** — Given  $0 < \varepsilon < 1/3$ , there exists  $\delta_{7.5}(n, \Lambda, \varepsilon) > 0$  with the following property. If

- (1)  $0 < r < \delta_{7.5}(n, \Lambda, \varepsilon)$ ;
- (2)  $\mathbf{C}_{\mathbf{W}}(x_0, r) \subseteq U$ ;
- (3)  $A \subseteq U$  is closed;
- (4)  $L^n(A \cap \mathbf{C}_{\mathbf{W}}(x_0, r)) > (1 - \varepsilon)L^n(\mathbf{C}_{\mathbf{W}}(x_0, r))$ ;

then

$$\int_{\mathbf{C}_{\mathbf{W}}(x_0,r)} \check{Y}_{A \cap \mathbf{C}_{\mathbf{W}}(x_0,r)} \mathbf{W}(u) dL^{n-m}(u) > (1 - c_{7.5}(n)\varepsilon)\alpha(m)r^m L^{n-m}(\mathbf{C}_{\mathbf{W}}(x_0,r)),$$

where  $c_{7.5}(n)$  is a positive integer depending only upon  $n$ .

*Proof.* — Similarly to the proof of 7.3, we will first establish a lower bound for  $\check{Y}_{A \cap \mathbf{C}_{\mathbf{W}}(x_0,r)} \mathbf{W}$  on “vertical slices”  $V_z$  of the given polyball, followed, next, by an application of Fubini. Given  $z \in \mathbf{W}_0(x_0) \cap \mathbf{B}(0,r)$ , we let  $V_z$  and  $\gamma_z$  be as in 7.3, and we consider the set

$$\check{V}_z = \mathbb{R}^n \cap \{x_0 + z + \sum_{i=1}^{n-m} y_i \mathbf{v}_i(x_0) : y \in C_{(1-3\varepsilon)r}\}$$

(notice  $\check{V}_z$  is slightly smaller than  $V_z$  used in the proof of 7.3) and the isometric parametrization  $\check{\gamma}_z : C_{(1-3\varepsilon)r} \rightarrow \check{V}_z$  defined by

$$\check{\gamma}_z(y) = x_0 + z + \sum_{i=1}^{n-m} y_i \mathbf{v}_i(x_0).$$

For part of the proof, we find it convenient to abbreviate  $E = A \cap \mathbf{C}_{\mathbf{W}}(x_0,r)$ . We also put  $\check{Y}_E \mathbf{W} = (\check{Y}_E \mathbf{W}) \circ \check{\gamma}_z$ .

By definition of  $\check{Y}_E \mathbf{W}$ , for each  $\check{\gamma}_z(y) \in \check{V}_z$ , there exists a collection  $C_y$  of closed balls in  $\mathbb{R}^{n-m}$  with the following properties: For every  $C \in C_y$ ,  $C$  is a ball centered at 0,  $C \subseteq C_{\varepsilon r}$ ,

$$\check{Y}_E \mathbf{W}(\check{\gamma}_z(y)) + \varepsilon > \int_C H^m(E \cap g_{\mathbf{v}_1, \dots, \mathbf{v}_{n-m}, \check{\gamma}_z(y)}^{-1}\{h\}) dL^{n-m}(h),$$

and  $\inf\{\text{diam } C : C \in C_y\} = 0$ . Furthermore,  $\check{Y}_E \mathbf{W}$  being  $L^{n-m}$ -summable, according to 6.3, there exists  $N \subseteq C_{(1-3\varepsilon)r}$  such that  $L^{n-m}(N) = 0$  and every  $y \notin N$  is a Lebesgue point of  $\check{Y}_E \mathbf{W}$ . For such a  $y$ , we may reduce  $C_y$ , if necessary, keeping all the previously stated properties valid, while enforcing also that

$$\int_{y+C} \check{Y}_E \mathbf{W} dL^{n-m} + \varepsilon > (\check{Y}_E \mathbf{W})(y),$$

whenever  $C \in C_y$ . We infer that, for each  $y \in C_{(1-3\varepsilon)r} \cap N$  and each  $C \in C_y$ ,

$$(26) \quad \int_{y+C} \check{Y}_E \mathbf{W} dL^{n-m} + 2\varepsilon L^{n-m}(y+C) > \int_C H^m(E \cap g_{\mathbf{v}_1, \dots, \mathbf{v}_{n-m}, \check{\gamma}_z(y)}^{-1}\{h\}) dL^{n-m}(h).$$

It ensues from the Vitali covering theorem that there exists a sequence  $(y_k)_k$  in  $C_{(1-3\varepsilon)r} \cap N$ , and  $C_k \in C_{y_k}$ , such that the balls  $y_k + C_k$ ,  $k = 1, 2, \dots$ , are pairwise disjoint and  $L^{n-m}(C_{(1-3\varepsilon)r} \cap \bigcup_{k=1}^{\infty} (y_k + C_k)) = 0$ . It therefore follows, from (26) and the fact that  $\gamma_z$  is an isometry, that

$$(27) \quad \int_{V_z} \check{Y}_E \mathbf{W} dH^{n-m} + 2\varepsilon H^{n-m}(V_z) > \sum_{k=1}^{\infty} \int_{C_k} H^m(E \cap g_{\mathbf{v}_1, \dots, \mathbf{v}_{n-m}, u_k}^{-1}\{y\}) dL^{n-m}(y),$$



where we have abbreviated  $u_k = \check{\gamma}_z(y_k)$ . We also abbreviate  $S_k = g_{\mathbf{v}_1, \dots, \mathbf{v}_{n-m}, u_k}^{-1}(C_k)$  and we infer from the coarea formula that, for each  $k = 1, 2, \dots$ ,

$$(28) \quad \int_{C_k} H^m(E \cap g_{\mathbf{v}_1, \dots, \mathbf{v}_{n-m}, u_k}^{-1}\{y\}) dL^{n-m}(y) = \int_{E \cap S_k} Jg_{\mathbf{v}_1, \dots, \mathbf{v}_{n-m}, u_k} dL^n > (1 - \varepsilon)L^n(E \cap S_k),$$

where the last inequality follows from 3.12, applied with  $U = \text{Int } \mathbf{C}_{\mathbf{W}}(x_0, r)$ , provided that  $\delta_{7.5}(n, \Lambda, \varepsilon)$  is chosen smaller than  $(2\sqrt{2})^{-1}\delta_{3.12}(n, \Lambda, \varepsilon)$ . Letting  $S = \bigcup_{k=1}^\infty S_k$  and recalling that  $E = A \cap \mathbf{C}_{\mathbf{W}}(x_0, r)$ , we infer, from (27) and (28), that

$$(29) \quad \int_{V_z} \Upsilon_E \mathbf{W} dH^{n-m} + 2\varepsilon H^{n-m}(V_z) > (1 - \varepsilon)L^n(E \cap S) > (1 - \varepsilon)(L^n(\mathbf{C}_{\mathbf{W}}(x_0, r) \cap S) - L^n(\mathbf{C}_{\mathbf{W}}(x_0, r) \cap A)).$$

Applying 7.3 to each  $S_k$  does not immediately yield a lower bound for

$$L^n(\mathbf{C}_{\mathbf{W}}(x_0, r) \cap S),$$

because the  $S_k$  are not necessarily pairwise disjoint. This is why we now introduce slightly smaller versions of these:

$$\check{C}_k = (1 - \varepsilon)C_k \quad \text{and} \quad \check{S}_k = g_{\mathbf{v}_1, \dots, \mathbf{v}_{n-m}, u_k}^{-1}(\check{C}_k).$$

CLAIM. *The sets  $\check{S}_k \cap \mathbf{C}_{\mathbf{W}}(x_0, r)$ ,  $k = 1, 2, \dots$ , are pairwise disjoint.*

Assume, if possible, that there are  $j \neq k$  and  $x \in \check{S}_j \cap \check{S}_k \cap \mathbf{C}_{\mathbf{W}}(x_0, r)$ . Letting  $\rho_j$  and  $\rho_k$  denote, respectively, the radius of  $C_j$ , and of  $C_k$ , we notice that  $\rho_j + \rho_k < |y_j - y_k|$ , because  $(y_j + C_j) \cap (y_k + C_k) = \emptyset$ . Since  $\check{\gamma}_z$  is an isometry, we have  $|u_j - u_k| = |\check{\gamma}_z(y_j) - \check{\gamma}_z(y_k)| = |y_j - y_k|$  and, therefore, also

$$(30) \quad \begin{aligned} & |g_{\mathbf{v}_1, \dots, \mathbf{v}_{n-m}, u_j}(x) - g_{\mathbf{v}_1, \dots, \mathbf{v}_{n-m}, u_k}(x)| \\ & \leq |g_{\mathbf{v}_1, \dots, \mathbf{v}_{n-m}, u_j}(x)| + |g_{\mathbf{v}_1, \dots, \mathbf{v}_{n-m}, u_k}(x)| \\ & \leq (1 - \varepsilon)\rho_j + (1 - \varepsilon)\rho_k < (1 - \varepsilon)|u_j - u_k|. \end{aligned}$$

We now introduce the following vectors of  $\mathbb{R}^{n-m}$ :

$$h_j = \sum_{i=1}^{n-m} \langle \mathbf{v}_i(x_0), u_j \rangle e_i \quad \text{and} \quad h_k = \sum_{i=1}^{n-m} \langle \mathbf{v}_i(x_0), u_k \rangle e_i$$

and we notice that

$$|h_j - h_k| = |P_{\mathbf{W}_0(x_0)^\perp}(u_j - u_k)| = |u_j - u_k|,$$

where the second equality holds because  $u_j - u_k \in \mathbf{W}_0(x_0)^\perp$ , as clearly follows from the definition of  $\check{\gamma}_z$ . Furthermore,

$$\begin{aligned} & |(g_{\mathbf{v}_1, \dots, \mathbf{v}_{n-m}, u_j}(x) - g_{\mathbf{v}_1, \dots, \mathbf{v}_{n-m}, u_k}(x)) - (h_k - h_j)| \\ & = \sqrt{\sum_{i=1}^{n-m} |\langle \mathbf{v}_i(x) - \mathbf{v}_i(x_0), u_k - u_j \rangle|^2} \\ & \leq \sqrt{n-m} \Lambda \sqrt{2}r |u_j - u_k| \leq \varepsilon |u_j - u_k|, \end{aligned}$$

since we may choose  $\delta_{7.5}(n, \Lambda, \varepsilon)$  to be so small that the last inequality holds, in view of hypothesis (1). Whence,

$$|g_{\mathbf{v}_1, \dots, \mathbf{v}_{n-m}, u_j}(x) - g_{\mathbf{v}_1, \dots, \mathbf{v}_{n-m}, u_k}(x)| > |h_j - h_k| - \varepsilon |u_j - u_k| = (1 - \varepsilon) |u_j - u_k|,$$

contradicting (30). The Claim is established.

Thus,

$$\begin{aligned} L^n(\mathbf{C}_{\mathbf{W}}(x_0, r) \cap S) &= L^n(\mathbf{C}_{\mathbf{W}}(x_0, r) \cap \bigcup_{k=1}^{\infty} S_k) \\ &> L^n(\mathbf{C}_{\mathbf{W}}(x_0, r) \cap \bigcup_{k=1}^{\infty} \check{S}_k) \\ &= \sum_{k=1}^{\infty} L^n(\mathbf{C}_{\mathbf{W}}(x_0, r) \cap \check{S}_k) \\ (31) \qquad &= \sum_{k=1}^{\infty} L^n(\mathbf{C}_{\mathbf{W}}(x_0, r) \cap g_{\mathbf{v}_1, \dots, \mathbf{v}_{n-m}, u_k}^{-1}(\check{C}_k)) \\ &> \frac{1}{1 + \varepsilon} \alpha(m) r^m \sum_{k=1}^{\infty} L^{n-m}(\check{C}_k), \end{aligned}$$

where the last inequality follows from 7.3. We notice that, indeed, 7.3 applies, since  $\check{C}_k \subseteq C_k \subseteq C_{\varepsilon r}$  and  $|g_{\mathbf{v}_1, \dots, \mathbf{v}_{n-m}, u_k}(x_0)| = |P_{\mathbf{W}_0(x_0)^\perp}(u_k - x_0)| = |y_k| \leq (1 - 3\varepsilon)r$ .

Now,

$$\begin{aligned} \sum_{k=1}^{\infty} L^{n-m}(\check{C}_k) &= (1 - \varepsilon)^{n-m} \sum_{k=1}^{\infty} L^{n-m}(C_k) \\ (32) \qquad &= (1 - \varepsilon)^{n-m} \sum_{k=1}^{\infty} L^{n-m}(y_k + C_k) \\ &> (1 - \varepsilon)^{n-m} L^{n-m}(C_{(1-3\varepsilon)r}) > (1 - 3\varepsilon)^{2(n-m)} \alpha(n - m) r^{n-m}. \end{aligned}$$

We infer, from (31) and (32), that

$$L^n(\mathbf{C}_{\mathbf{W}}(x_0, r) \cap S) > \frac{(1 - 3\varepsilon)^{2(n-m)}}{1 + \varepsilon} L^n(\mathbf{C}_{\mathbf{W}}(x_0, r)).$$

It therefore ensues, from (29) and hypothesis (4), that

$$\int_{V_z} \gamma_E \mathbf{W} dH^{n-m} + 2\varepsilon H^{n-m}(V_z) > (1 - \varepsilon) \left( \frac{(1 - 3\varepsilon)^{2(n-m)}}{1 + \varepsilon} - \varepsilon \right) L^n(\mathbf{C}_{\mathbf{W}}(x_0, r)).$$

Integrating over  $z$ , we infer from Fubini's theorem that

$$\begin{aligned} \int_{\mathbf{C}_{\mathbf{W}}(x_0, r)} \gamma_{A \cap \mathbf{C}_{\mathbf{W}}(x_0, r)} \mathbf{W} dL^n &= \int_{\mathbf{W}_0(x_0) \cap \mathbf{B}(0, r)} dL^n(z) \int_{V_z} \gamma_E \mathbf{W} dH^{n-m} \\ &> \left[ (1 - \varepsilon) \left( \frac{(1 - 3\varepsilon)^{2(n-m)}}{1 + \varepsilon} - \varepsilon \right) - 2\varepsilon \right] \alpha(m) r^m L^n(\mathbf{C}_{\mathbf{W}}(x_0, r)). \end{aligned}$$

**7.6. PROPOSITION.** — *Given  $0 < \varepsilon < 1/3$ , there exist  $\delta_{7.6}(n, \Lambda, \varepsilon) > 0$  and  $c_{7.6}(n) > 1$  with the following property. If*

$$(1) \quad 0 < r < \delta_{7.6}(n, \Lambda, \varepsilon);$$

- (2)  $\mathbf{C}_{\mathbf{W}}(x_0, r) \subseteq U$ ;  
 (3)  $A \subseteq U$  is closed;  
 (4)  $L^n(A \cap \mathbf{C}_{\mathbf{W}}(x_0, r)) > (1 - \varepsilon)L^n(\mathbf{C}_{\mathbf{W}}(x_0, r))$ ;

then

$$\int_{A \cap \mathbf{C}_{\mathbf{W}}(x_0, r)} \Upsilon_{A \cap \mathbf{C}_{\mathbf{W}}(x_0, r)} \mathbf{W}(u) dL^n(u) > (1 - c_{7.6}(n)\varepsilon)\alpha(m)r^m L^n(\mathbf{C}_{\mathbf{W}}(x_0, r)).$$

**7.7. REMARK.** — The difference with 7.5 is the domain of integration (being smaller) in the integral, on the left hand side in the conclusion.

*Proof of 7.6.* — The reader will happily check that

$$\delta_{7.6}(n, \Lambda, \varepsilon) = \min\{\delta_{7.5}(n, \Lambda, \varepsilon), (2\sqrt{2})^{-1}\delta_{6.3}(n, \Lambda)\}$$

suits their needs with  $c_{7.6}(n) = \max\{c_{7.6}(n, m) : m = 1, \dots, n\}$  where

$$c_{7.6}(n, m) = c_{7.5}(n) + \frac{(2\sqrt{2})^m c_{6.3}(m)}{\alpha(m)}.$$

**7.8. PROPOSITION.** — *There exists  $\delta_{7.8}(n, \Lambda) > 0$  with the following property.*

*If  $\text{diam } E \subset \delta_{7.8}(n, \Lambda)$ , then*

$$Z_E \mathbf{W}(u) > 0,$$

*for  $L^n$ -almost every  $u \in E$ .*

*Proof.* — We let

$$\delta_{7.8}(n, \Lambda) = \min\{\delta_{7.6}(n, \Lambda, 1/(4c_{7.6}(n))), \delta_{5.16}(n, \Lambda, 1/2)\}.$$

According to 5.16, it suffices to show that  $\Upsilon_E \mathbf{W}(u) > 0$ , for  $L^n$ -almost every  $u \in E$ . Define  $Z = E \cap \{u : \Upsilon_E \mathbf{W}(u) = 0\}$  and assume, if possible, that  $L^n(Z) > 0$ . Since  $Z$  is  $L^n$ -measurable (recall 5.12), there exists a compact set  $A \subseteq Z$  such that  $L^n(A) > 0$ . Observe that the sets  $\mathbf{C}_{\mathbf{W}}(x, r)$ , for  $x \in U$  and  $r > 0$ , form a density basis for  $L^n$ -measurable subsets of  $U$  – because their eccentricity is bounded away from zero – thus, there exists  $x_0 \in A$  and  $r_0 > 0$  such that

$$L^n(A \cap \mathbf{C}_{\mathbf{W}}(x_0, r)) > \left(1 - \frac{1}{4c_{7.6}(n)}\right)L^n(\mathbf{C}_{\mathbf{W}}(x_0, r)),$$

whenever  $0 < r < r_0$ . There is no restriction to assume that  $r_0$  is small enough for  $\mathbf{C}_{\mathbf{W}}(x_0, r_0) \subseteq U$ . Thus, if we let  $r = \min\{r_0, \delta_{7.6}(n, \Lambda, 1/(4c_{7.6}(n)))\}$ , it follows from 7.6 that

$$(33) \quad \int_{A \cap \mathbf{C}_{\mathbf{W}}(x_0, r)} \Upsilon_{A \cap \mathbf{C}_{\mathbf{W}}(x_0, r)} \mathbf{W}(u) dL^n(u) > \left(1 - \frac{1}{4}\right)\alpha(m)r^m L^n(\mathbf{C}_{\mathbf{W}}(x_0, r)) > 0.$$

On the other hand, recalling 5.12 and the fact that  $A \cap \mathbf{C}_{\mathbf{W}}(x_0, r) \subseteq E$ , we infer that  $\Upsilon_{A \cap \mathbf{C}_{\mathbf{W}}(x_0, r)} \mathbf{W}(u) \subset \Upsilon_E \mathbf{W}(u)$ , for all  $u \in \mathbb{R}^n$ . In particular,  $\Upsilon_{A \cap \mathbf{C}_{\mathbf{W}}(x_0, r)} \mathbf{W}(u) = 0$ , for all  $u \in A \cap \mathbf{C}_{\mathbf{W}}(x_0, r) \subseteq Z$ , contradicting (33).

8. PROOF OF THE THEOREMS

8.1. THEOREM. — Assume that  $S \subseteq \mathbb{R}^n$ ,  $\mathbf{W}_0 : S \rightarrow \mathbb{G}(n, m)$  is Lipschitzian, and  $A \subseteq S$  is Borel measurable. The following are equivalent.

- (1)  $L^n(A) = 0$ .
- (2) For  $L^n$  almost every  $x \in A$ ,  $H^m(A \cap \mathbf{W}(x)) = 0$ .
- (3) For  $L^n$  almost every  $x \in S$ ,  $H^m(A \cap \mathbf{W}(x)) = 0$ .

Recall our convention that  $\mathbf{W}(x) = x + \mathbf{W}_0(x)$ .

*Proof.* — Since  $\mathbb{G}(n, m)$  is complete, we can extend  $\mathbf{W}_0$  to the closure of  $S$ . Furthermore, if the theorem holds for  $\text{Clos } S$ , then it also holds for  $S$ . Thus, there is no restriction to assume that  $S$  is closed.

(1)  $\Rightarrow$  (3). It follows from 3.8 that each  $x \in S$  admits an open neighborhood  $U_x$  in  $\mathbb{R}^n$  such that  $\mathbf{W}(x)$  can be associated with a Lipschitzian orthonormal frame satisfying all the conditions of 4.1, for some  $\Lambda_x > 0$ . Since  $S$  is Lindelöf, there are countably many  $x_1, x_2, \dots$  such that  $S \subseteq \bigcup_j U_{x_j}$ . Letting  $E_j = S \cap U_{x_j}$ , we infer from 5.5 that  $\phi_{E_j, \mathbf{W}}$  is absolutely continuous with respect to  $L^n$ . Thus, if  $L^n(A) = 0$ , then  $H^m(A \cap \mathbf{W}(x)) = 0$ , for  $L^n$ -almost every  $x \in E_j$ , by definition of  $\phi_{E_j, \mathbf{W}}$ . Since  $j$  is arbitrary, the proof is complete.

(3)  $\Rightarrow$  (2) is trivial.

(2)  $\Rightarrow$  (1). Let  $A$  satisfy condition (2). It is enough to show that  $L^n(A \cap \mathbf{B}(0, r)) = 0$ , for each  $r > 0$ . Fix  $r > 0$  and define  $S_r = S \cap \mathbf{B}(0, r)$ . Consider the  $U_{x_j}$  defined in the second paragraph of the present proof; since  $S_r$  is compact, finitely many of those, say  $U_{x_1}, \dots, U_{x_N}$ , cover  $S_r$ . Let  $\Lambda = \max_{j=1, \dots, N} \Lambda_{x_j}$ . Partition each  $U_{x_j}$ ,  $j = 1, \dots, N$ , into Borel measurable sets  $E_{j,k}$ ,  $k = 1, \dots, K_j$ , such that  $\text{diam } E_{j,k} \leq \delta_{7.8}(n, \Lambda)$ . It then follows from 7.8 that

$$(34) \quad (Z_{A \cap E_{j,k}} \mathbf{W})(u) > 0,$$

for  $L^n$ -almost every  $u \in A \cap E_{j,k}$ . Now, fix  $j$  and  $k$ . Observe that

$$H^m(A \cap E_{j,k} \cap \mathbf{W}(x)) = 0,$$

for  $L^n$ -almost every  $x \in A \cap E_{j,k}$ . Thus,  $\phi_{A \cap E_{j,k}, \mathbf{W}}(A \cap E_{j,k}) = 0$ . Moreover,

$$0 = \phi_{A \cap E_{j,k}, \mathbf{W}}(A \cap E_{j,k}) = \int_{A \cap E_{j,k}} (Z_{A \cap E_{j,k}} \mathbf{W})(u) dL^n(u).$$

It follows from (34) that  $L^n(A \cap E_{j,k}) = 0$ . Since  $j$  and  $k$  are arbitrary,  $L^n(A) = 0$ .

8.2. REMARK. — Alternatively, one can prove the principal implication (2)  $\Rightarrow$  (1) in two other ways. One way – more involved – consists in applying our main result 8.4 below. A second – simpler – way, along the following lines, avoids reference to the estimates we obtained for the functions  $\gamma_E \mathbf{W}$  and  $Z_E \mathbf{W}$ . Consider  $x \in A$ ,  $\eta > 0$ ,

and  $V \in G(n, n - m)$  such that  $d(V^\perp, \mathbf{W}_0(x)) < \eta$ , and put  $V_x = x + V$ . Define

$$\Phi : (U_x \cap V_x) \times \mathbb{R}^m \longrightarrow \mathbb{R}^n : (\xi, t) \longmapsto \xi + \sum_{i=1}^m t_i \mathbf{w}_i(\xi).$$

$\Phi$  is locally Lipschitzian and one checks that, in fact,  $\Phi$  is a lipeomorphism between  $B$  and  $\Phi(B)$ , where  $B = \mathbf{B}(x, \rho)$ , for some  $\rho > 0$  depending upon  $\eta$  and  $\Lambda_x$ , because its differential is close to the identity. Referring to Fubini’s theorem, one then further checks that

$$L^n(A') = 0 \text{ if and only if } H^m(A' \cap \mathbf{W}(\xi)) = 0, \text{ for } H^{n-m}\text{-almost every } \xi \in V_x.$$

Finally, using Fubini again, with respect to the decomposition  $\mathbb{R}^n = \mathbf{W}_0(x) \oplus \mathbf{W}_0(x)^\perp$ , one shows that  $H^m(A \cap \mathbf{W}(\zeta)) = 0$ , for  $H^{n-m}$ -almost every  $\zeta \in x' + \mathbf{W}_0(x)^\perp$ , where  $x'$  is as close as we wish to  $x$ . Applying the previous construction with  $x'$  replacing  $x$ , using  $V = \mathbf{W}_0(x)^\perp$ , we find that  $L^n(A \cap \mathbf{B}(x, r)) = 0$ , for some  $r > 0$  depending on  $\Lambda_x$ .

Notwithstanding, it seems that the (simpler) change of variable described here is not enough to yield the (stronger) theorem below.

8.3. POLYBALLS. — Recalling 7.2, we notice that

$$\mathbf{C}_{\mathbf{W}}(x_0, r) = \mathbb{R}^n \cap \{\nu_{x_0}(x - x_0) \leq r\},$$

where  $\nu_{x_0}$  is a norm on  $\mathbb{R}^n$  defined by the formula

$$\nu_{x_0}(x) = \max\{|P_{\mathbf{W}_0(x_0)}(x)|, |P_{\mathbf{W}_0(x_0)^\perp}(x)|\},$$

for  $x \in \mathbb{R}^n$ . It is readily observed that  $\text{Lip } \nu_{x_0} \leq 1$ .

(1) *One has  $|\nabla \nu_{x_0}(x)| = 1$ , for  $L^n$ -almost every  $x \in \mathbb{R}^n$ .*

Abbreviate  $P = P_{\mathbf{W}_0(x_0)}$  and  $Q = P_{\mathbf{W}_0(x_0)^\perp}$  and define

$$S = \mathbb{R}^n \cap \{x : |P(x)| = |Q(x)|\},$$

so that  $L^n(S) = 0$ . Let  $x \in \mathbb{R}^n \setminus S$  and notice  $\nu_{x_0}$  is differentiable at  $x$ . We henceforth assume that  $|P(x)| < |Q(x)|$ , whence,  $\nu_{x_0}(x) = |Q(x)|$  – the proof in the other case is similar. Define  $\varepsilon = |Q(x)| - |P(x)| > 0$ ,  $e = Q(x)/|Q(x)|$ , and let  $0 < t < \varepsilon$ . Note that

$$|Q(x + te)| = |Q(x)| \left(1 + \frac{t}{|Q(x)|}\right) > |Q(x)| \quad \text{and} \quad |P(x + te)| \leq |P(x)| + t < |Q(x).$$

Thus,  $\nu_{x_0}(x + te) = |Q(x + te)|$  and, in turn,

$$\nu_{x_0}(x + te) - \nu_{x_0}(x) = |Q(x + te)| - |Q(x)| = t.$$

It follows that  $\langle \nabla \nu_{x_0}(x), e \rangle = 1$ . Since  $e$  is a unit vector and  $\text{Lip } \nu_{x_0} \leq 1$ , we conclude that  $|\nabla \nu_{x_0}(x)| = 1$ .

(2) *Let  $U$ ,  $\mathbf{W}$  and  $\Lambda > 0$  be as in 4.1. Assume  $\mathbf{C}_{\mathbf{W}}(x_0, r) \subseteq U$ ,  $x \in \mathbf{C}_{\mathbf{W}}(x_0, r)$ , and  $0 \leq t \leq 1$  is so that  $\nu_{x_0}(x - x_0) = tr$ . It follows that*

$$\mathbf{C}_{\mathbf{W}}(x_0, r) \cap \mathbf{W}(x) \subseteq \mathbf{B}(x, r(1 + t) + 8m\Lambda r^2) \cap \mathbf{W}(x).$$

First notice that  $|z| \leq \sqrt{2}\nu_{x_0}(z)$ , for every  $z \in \mathbb{R}^n$ , so that

$$|x - x_0| \leq \sqrt{2}\nu_{x_0}(x - x_0) = \sqrt{2}tr$$

and, for every  $x' \in \mathbf{C}_{\mathbf{W}}(x_0, r)$ ,

$$|x - x'| \leq \sqrt{2}(\nu_{x_0}(x - x_0) + \nu_{x_0}(x_0 - x')) \leq \sqrt{2}r(1 + t).$$

Next notice that, for each  $h \in \mathbb{R}^n$ ,

$$\begin{aligned} |(P_{\mathbf{W}_0(x_0)} - P_{\mathbf{W}_0(x)})(h)| &= \left| \sum_{i=1}^m \langle \mathbf{w}_i(x_0), h \rangle \mathbf{w}_i(x_0) - \sum_{i=1}^m \langle \mathbf{w}_i(x), h \rangle \mathbf{w}_i(x) \right| \\ &\leq \sum_{i=1}^m |\langle \mathbf{w}_i(x_0), h \rangle| |\mathbf{w}_i(x_0) - \mathbf{w}_i(x)| + \sum_{i=1}^m |\langle \mathbf{w}_i(x_0) - \mathbf{w}_i(x), h \rangle| |\mathbf{w}_i(x)| \\ &\leq 2m\Lambda |x - x_0| |h|. \end{aligned}$$

We now assume that  $x' \in \mathbf{C}_{\mathbf{W}}(x, r) \cap \mathbf{W}(x)$ , in particular,  $x - x' \in \mathbf{W}_0(x)$ , whence,

$$\begin{aligned} |x - x'| &= |P_{\mathbf{W}_0(x)}(x - x')| \leq |P_{\mathbf{W}_0(x_0)}(x - x')| + |(P_{\mathbf{W}_0(x_0)} - P_{\mathbf{W}_0(x)})(x - x')| \\ &\leq \nu_{x_0}(x - x_0) + \nu_{x_0}(x' - x_0) + 2m\Lambda |x - x_0| |x - x'| \leq r(1 + t) + 4m\Lambda t(1 + t)r^2. \end{aligned}$$

**8.4. THEOREM.** — Assume that  $A \subseteq \mathbb{R}^n$  is Borel measurable and that  $\mathbf{W}_0 : A \rightarrow \mathbb{G}(n, m)$  is Lipschitzian. It follows that

$$\limsup_{r \rightarrow 0^+} \frac{H^m(A \cap \mathbf{B}(x, r) \cap \mathbf{W}(x))}{\alpha(m)r^m} > \frac{1}{2^n},$$

for  $L^n$ -almost every  $x \in A$ .

Recall our convention that  $\mathbf{W}(x) = x + \mathbf{W}_0(x)$ .

*Proof.* — Extend  $\mathbf{W}_0$  to  $\mathbb{R}^n$  in a Borel measurable way, for instance, to be an arbitrary constant outside of  $A$ . It follows from 3.15 that, for each  $r > 0$ , the function

$$\mathbb{R}^n \rightarrow [0, \infty] : x \mapsto \frac{H^m(A \cap \mathbf{B}(x, r) \cap \mathbf{W}(x))}{\alpha(m)r^m}$$

is Borel measurable. Thus, for each  $j = 1, 2, \dots$ , the function  $g_j : \mathbb{R}^n \rightarrow [0, \infty]$  defined by

$$g_j(x) = \sup_{0 < r < 1/j} \frac{H^m(A \cap \mathbf{B}(x, r) \cap \mathbf{W}(x))}{\alpha(m)r^m} = \sup_{\substack{0 < r < 1/j \\ r \text{ rational}}} \frac{H^m(A \cap \mathbf{B}(x, r) \cap \mathbf{W}(x))}{\alpha(m)r^m}$$

is Borel measurable as well, and so is  $g = \lim_j g_j = \inf_j g_j$ .

Abbreviate  $\eta(n, m) = 2^{-(n-m)}$ . Arguing *reductio ad absurdum*, we henceforth assume that  $A$  and  $\mathbf{W}_0$  fail the conclusion of the theorem. Thus, the set

$$Z_0 = A \cap \{x : g(x) < \eta(n, m)/2^m\}$$

is Borel measurable and non Lebesgue null. Accordingly, there exists  $\varepsilon_1 > 0$  such that the set

$$Z'_0 = A \cap \{x : g(x) < (1 - \varepsilon_1)\eta(n, m)/(2 + \varepsilon_1)^m\}$$

is also Borel measurable and of positive Lebesgue measure. It therefore ensues from Egoroff's theorem [8, 2.3.7] that there exists a closed set  $Z \subseteq Z'_0 \subseteq A$  such that  $L^n(Z) > 0$  and that there exists a positive integer  $j_0$  such that

$$(35) \quad \frac{H^m(Z \cap \mathbf{B}(x, r) \cap \mathbf{W}(x))}{\alpha(m)r^m} \leq g_{j_0}(x) < (1 - \varepsilon_1) \frac{\eta(n, m)}{(2 + \varepsilon_1)^m},$$

for each  $x \in Z$  and each  $0 < r < 1/j_0$ . Choose  $0 < \varepsilon_2 < 10$  such that

$$(36) \quad \frac{1 - \varepsilon_1}{1 - \varepsilon_2} \leq 1 - \frac{\varepsilon_1}{2}$$

and choose  $0 < \varepsilon_3 < 1/3$  such that

$$(37) \quad 1 - \frac{\varepsilon_1}{2} < 1 - c_{7.6}(n)\varepsilon_3.$$

As in the proof of 7.8, we recall that the family  $\mathbf{C}_{\mathbf{W}}(x, r)$ , for  $x \in \mathbb{R}^n$  and  $r > 0$ , is a density basis of  $L^n$ -measurable sets. Since  $L^n(Z) > 0$ , there exists  $x_0 \in Z$  such that

$$\lim_{r \rightarrow 0^+} \frac{L^n(Z \cap \mathbf{C}_{\mathbf{W}}(x_0, r))}{L^n(\mathbf{C}_{\mathbf{W}}(x_0, r))} = 1.$$

In particular, there exists  $R > 0$  such that

$$(38) \quad (1 - \varepsilon_3)L^n(\mathbf{C}_{\mathbf{W}}(x_0, r)) \leq L^n(Z \cap \mathbf{C}_{\mathbf{W}}(x_0, r)),$$

whenever  $0 < r < R$ .

We let  $U$  be an open neighborhood of  $x_0$  in  $\mathbb{R}^n$  associated with  $A$  and  $\mathbf{W}_0$  in 3.8, so that  $\mathbf{W}_0$  and  $\mathbf{W}_0^\perp$  are associated with orthonormal frames as in 4.1, for some  $\Lambda > 0$ . Define

$$r_0 = \min \left\{ 1, \frac{1}{j_0(2 + 8m\Lambda)}, \frac{\varepsilon_1}{8m\Lambda}, \frac{\delta_{5.15}(n, \Lambda, \varepsilon_2)}{2\sqrt{2}}, \frac{\text{dist}(x_0, \mathbb{R}^n \setminus U)}{2\sqrt{2}}, \delta_{7.6}(n, \Lambda, \varepsilon_3), R \right\}.$$

Let  $0 < r < r_0$  and observe that

$$(39) \quad (1 - c_{7.6}(n)\varepsilon_3)L^n(\mathbf{C}_{\mathbf{W}}(x_0, r)) \leq \int_{Z \cap \mathbf{C}_{\mathbf{W}}(x_0, r)} \frac{\chi_{Z \cap \mathbf{C}_{\mathbf{W}}(x_0, r)} \mathbf{W}(u)}{\alpha(m)r^m} dL^n(u)$$

(by 7.6 applied with  $\varepsilon = \varepsilon_3$  and  $A = Z \cap \mathbf{C}_{\mathbf{W}}(x_0, r)$ )

$$\leq \frac{1}{(1 - \varepsilon_2)\eta(n, m)} \int_{Z \cap \mathbf{C}_{\mathbf{W}}(x_0, r)} \frac{H^m(Z \cap \mathbf{C}_{\mathbf{W}}(x_0, r) \cap \mathbf{W}(x))}{\alpha(m)r^m} dL^n(x)$$

(by 5.15, applied with  $\varepsilon = \varepsilon_2$  and  $E = B = Z \cap \mathbf{C}_{\mathbf{W}}(x_0, r)$ ).

We also note that

$$(40) \quad \int_{Z \cap \mathbf{C}_{\mathbf{W}}(x_0, r)} \frac{H^m(Z \cap \mathbf{C}_{\mathbf{W}}(x_0, r) \cap \mathbf{W}(x))}{\alpha(m)r^m} dL^n(x) \\ = \int_{Z \cap \{\nu_{x_0} \leq r\}} \frac{H^m(Z \cap \mathbf{C}_{\mathbf{W}}(x_0, r) \cap \mathbf{W}(x))}{\alpha(m)r^m} |\nabla \nu_{x_0}(x)| dL^n(x)$$

(by 8.3(1))

$$\begin{aligned}
&= \int_0^r dL^{-1}(\rho) \int_{Z \cap \{\nu_{x_0}=\rho\}} \frac{H^m(Z \cap \mathbf{C}_W(x_0, r) \cap \mathbf{W}(x))}{\alpha(m)r^m} dH^{n-1}(x) \\
&\text{(by [5, 3.4.3])} \\
&= r \int_0^1 dL^{-1}(t) \int_{Z \cap \{\nu_{x_0}=tr\}} \frac{H^m(Z \cap \mathbf{C}_W(x_0, r) \cap \mathbf{W}(x))}{\alpha(m)r^m} dH^{n-1}(x) \\
&\leq r \int_0^1 dL^{-1}(t) \int_{Z \cap \{\nu_{x_0}=tr\}} \frac{H^m(Z \cap \mathbf{B}(x, r(1+t) + 8m\Lambda r^2) \cap \mathbf{W}(x))}{\alpha(m)r^m} dH^{n-1}(x) \\
&\text{(by 8.3(2))} \\
&\leq r \int_0^1 dL^{-1}(t) \int_{Z \cap \{\nu_{x_0}=tr\}} (1 - \varepsilon_1) \frac{\eta(n, m)}{(2 + \varepsilon_1)^m} (1 + t + 8m\Lambda r)^m dH^{n-1} \\
&\text{(by (35))} \\
&\leq (1 - \varepsilon_1) \eta(n, m) \int_0^r dL^{-1}(\rho) \int_{Z \cap \{\nu_{x_0}=\rho\}} dH^{n-1} \\
&= (1 - \varepsilon_1) \eta(n, m) \int_{Z \cap \{\nu_{x_0} \leq r\}} |\nabla \nu_{x_0}(x)| dL^n(x) \\
&\text{(by [5, 3.4.3])} \\
&= (1 - \varepsilon_1) \eta(n, m) L^n(Z \cap \mathbf{C}_W(x_0, r)) \leq (1 - \varepsilon_1) \eta(n, m) L^n(\mathbf{C}_W(x_0, r)) \\
&\text{(by 8.3(1)).}
\end{aligned}$$

Plugging (40) into (39), we obtain

$$\begin{aligned}
(1 - c_{7.6}(n)\varepsilon_3) L^n(\mathbf{C}_W(x_0, r)) &\leq \left( \frac{1 - \varepsilon_1}{1 - \varepsilon_2} \right) L^n(\mathbf{C}_W(x_0, r)) \\
&< (1 - c_{7.6}(n)\varepsilon_3) L^n(\mathbf{C}_W(x_0, r))
\end{aligned}$$

(by (36) and (37)), a contradiction.

## REFERENCES

- [1] J. BOURGAIN – “A remark on the maximal function associated to an analytic vector field”, in *Analysis at Urbana, Vol. I (Urbana, IL, 1986–1987)*, London Math. Soc. Lecture Note Ser., vol. 137, Cambridge Univ. Press, Cambridge, 1989, p. 111–132.
- [2] A. M. BRUCKNER – “Differentiation of integrals”, *Amer. Math. Monthly* **78** (1971), no. 9, part 2, p. 1–51.
- [3] D. L. COHN – *Measure theory*, second ed., Birkhäuser Advanced Texts: Basler Lehrbücher, Birkhäuser/Springer, New York, 2013.
- [4] TH. DE PAUW – “Concentrated, nearly monotonic, epiperimetric measures in Euclidean space”, *J. Differential Geom.* **77** (2007), no. 1, p. 77–134.
- [5] L. C. EVANS & R. F. GARIEPY – *Measure theory and fine properties of functions*, revised ed., Textbooks in Math., CRC Press, Boca Raton, FL, 2015.
- [6] K. J. FALCONER – “Sets with prescribed projections and Nikodým sets”, *Proc. London Math. Soc.* (3) **53** (1986), no. 1, p. 48–64.
- [7] H. FEDERER – “Curvature measures”, *Trans. Amer. Math. Soc.* **93** (1959), p. 418–491.



- [8] ———, *Geometric measure theory*, Grundlehren Math. Wiss., vol. 153, Springer-Verlag New York, Inc., New York, 1969.
- [9] M. DE GUZMÁN – *Real variable methods in Fourier analysis*, Notas de Matemática, vol. 75, North-Holland Publishing Co., Amsterdam-New York, 1981.
- [10] A. B. KHARAZISHVILI – *Nonmeasurable sets and functions*, North-Holland Math. Studies, vol. 195, Elsevier Science B.V., Amsterdam, 2004.
- [11] M. LACEY & X. LI – *On a conjecture of E. M. Stein on the Hilbert transform on vector fields*, Mem. Amer. Math. Soc., vol. 205, no.965, American Mathematical Society, Providence, RI, 2010.
- [12] O. NIKODÝM – “Sur la mesure des ensembles plans dont tous les points sont rectilinéairement accessibles”, *Fund. Math.* **10** (1927), p. 116–168.
- [13] S. M. SRIVASTAVA – *A course on Borel sets*, Graduate Texts in Math., vol. 180, Springer-Verlag, New York, 1998.

Manuscript received 21st April 2022

accepted 3rd August 2022

THIERRY DE PAUW, Université Paris Cité and Sorbonne Université, CNRS, IMJ-PRG  
F-75013 Paris, France  
E-mail : depauw@imj-prg.fr  
Url : <https://www.thierrydepauw.org/>