



Journal de l'École polytechnique

Mathématiques

VÍCTOR ARNAIZ & GABRIEL RIVIÈRE

Quantum limits of perturbed sub-Riemannian contact Laplacians in dimension 3

Tome II (2024), p. 909-956.

<https://doi.org/10.5802/jep.269>

© Les auteurs, 2024.



Cet article est mis à disposition selon les termes de la licence LICENCE INTERNATIONALE D'ATTRIBUTION CREATIVE COMMONS BY 4.0.

<https://creativecommons.org/licenses/by/4.0/>

Publié avec le soutien
du Centre National de la Recherche Scientifique



Publication membre du
Centre Mersenne pour l'édition scientifique ouverte

www.centre-mersenne.org

e-ISSN : 2270-518X

QUANTUM LIMITS OF PERTURBED SUB-RIEMANNIAN CONTACT LAPLACIANS IN DIMENSION 3

BY VÍCTOR ARNAIZ & GABRIEL RIVIÈRE

ABSTRACT. — On the unit tangent bundle of a compact Riemannian surface, we consider a natural sub-Riemannian Laplacian associated with the canonical contact structure. In the large eigenvalue limit, we study the escape of mass at infinity in the cotangent space of eigenfunctions for hypoelliptic selfadjoint perturbations of this operator. Using semiclassical methods, we show that, in this subelliptic regime, eigenfunctions concentrate on certain quantized level sets along the geodesic flow direction and that they verify invariance properties involving both the geodesic vector field and the perturbation term.

RÉSUMÉ (Limites quantiques pour les laplaciens sous-riemanniens de contact en dimension 3)

Sur le fibré tangent unitaire d'une surface compacte riemannienne, nous considérons un sous-laplacien riemannien naturel associé à la structure de contact canonique. Dans la limite des grandes valeurs propres et pour des perturbations hypoelliptiques auto-adjointes de cet opérateur, nous étudions la façon dont la masse des fonctions propres s'échappe à l'infini dans l'espace co-tangent. En utilisant des méthodes semi-classiques, nous montrons que, dans ce régime sous-elliptique, les fonctions propres se concentrent sur certains ensembles de niveaux quantifiés le long de la direction du flot géodésique et qu'elles vérifient des propriétés d'invariance impliquant à la fois le champ de vecteurs géodésique et le terme de perturbation.

CONTENTS

1. Introduction.....	910
2. Semiclassical conventions.....	918
3. Reduction to the subelliptic regime.....	921
4. Support of the limit measure.....	929
5. Normal form reduction.....	936
6. Invariance properties.....	941
7. Summary of the properties of μ_∞	944
8. The case of the flat torus.....	947
Appendix A. Spectral properties of \widehat{P}_h	951
Appendix B. Reminder on semiclassical analysis on $\mathbb{R}^2 \times \mathbb{S}^1$	954
References.....	955

MATHEMATICAL SUBJECT CLASSIFICATION (2020). — 58J50, 35H10, 53C17.

KEYWORDS. — Hypoelliptic operators, semiclassical analysis, contact flows.

Both authors are supported by the Agence Nationale de la Recherche through the PRC grant ADYCT (ANR-20-CE40-0017), the first author is partially supported by the projects MTM2017-85934-C3-3-P and PID2021-124195NB-C31 (MINECO, Spain) and the second author acknowledges the support of the Institut Universitaire de France.

1. INTRODUCTION

Let (M, g) be a smooth, compact, oriented, and boundaryless Riemannian *surface* and denote by $K(m)$ its sectional curvature at a given point $m \in M$. The unit tangent bundle of M is defined by

$$\mathcal{M} := SM = \{q = (m, v) \in TM : \|v\|_{g(m)} = 1\}.$$

There are two natural vector fields on SM : the geodesic vector field X and the vertical vector field V , i.e., the vector field corresponding to the action by rotation in the fibers of SM . One can then define $X_\perp := [X, V]$ and these vector fields verify the following commutation relations [PSU23, §3.5.1]:

$$(1) \quad [X, X_\perp] = -KV, \quad [X, V] = X_\perp, \quad \text{and} \quad [X_\perp, V] = -X,$$

where K is understood as a function on SM (via pullback). The manifold \mathcal{M} is naturally endowed with a Riemannian metric g_S (the Sasaki metric) which makes (X, X_\perp, V) into an orthonormal basis. The corresponding volume form that we will denote by $d\mu_L$ makes these three vector fields divergence free and we can define the sub-Riemannian Laplacian associated with this geodesic frame:

$$-\Delta_{\text{sR}} := X_\perp^* X_\perp + V^* V = -X_\perp^2 - V^2.$$

More precisely, we consider the Friedrichs extension of this formally selfadjoint operator (see Appendix A for a brief reminder) which is hypoelliptic by Hörmander's Theorem [Hör67, Th. 1.1]. See Section 8 for a concrete description of these operators and their spectrum in the case of the flat torus. In the context of contact geometry, $-\Delta_{\text{sR}}$ is referred as the Rumin Laplacian for the Sasaki metric [Rum94]. See also Section 1.4 for a discussion on the case of general Hörmander (contact) operators in dimension 3.

We now let $Q, W \in \mathcal{C}^\infty(\mathcal{M}, \mathbb{R})$. Our goal is to study, in the semiclassical limit $h \rightarrow 0^+$, the eigenfunctions of the following (formally selfadjoint) operator:

$$(2) \quad \widehat{P}_h := -h^2 \Delta_{\text{sR}} - ih^2 QX - \frac{ih^2 X(Q)}{2} + W, \quad h \in (0, 1].$$

Recall from [RS76, §1] that such operators appear naturally when studying the boundary analogue of the $\bar{\partial}$ operator in complex analysis [FS74] whose local expression is exactly of this form and which is sometimes referred as the Kohn Laplacian. In the present work, the relevance of considering subprincipal terms of the form $QX + (QX)^* = Q[V, X_\perp] + X(Q)/2$ is also the richer dynamical structure displayed by the eigenfunctions of \widehat{P}_h in the semiclassical limit (see Remark 1.2 after Theorem 1.1).

Under the assumption $\|Q\|_{\mathcal{C}^0} < 1$, one can both consider the Friedrichs extension of this operator and apply the Rothschild-Stein theorem [RS76, Th. 16]. In some sense, these operators are also among the simplest example of selfadjoint *hypoelliptic* operator that cannot be written as a sum of squares. Combining this last theorem with classical tools from spectral theory [RS72, RS75], one can find $h_0 > 0$ such that,

for all $0 < h \leq h_0$, there exists a nondecreasing sequence

$$\min W + \mathcal{O}_Q(h) \leq \Lambda_h(0) \leq \lambda_h(1) \leq \dots \leq \Lambda_h(j) \dots \longrightarrow +\infty, \quad \text{as } j \longrightarrow +\infty,$$

and an orthonormal basis $(\psi_h^j)_{j \geq 0}$ of $L^2(\mathcal{M})$ verifying, for all $j \geq 0$,

$$\widehat{P}_h \psi_h^j = \Lambda_h(j) \psi_h^j.$$

We refer to Lemma A.4 in Appendix A for details. Moreover, any solution ψ_h^j to this eigenvalue problem belongs to the space $\mathcal{C}^\infty(\mathcal{M})$ and, thanks to Lemma A.5, it satisfies the a priori estimate for $h > 0$ small enough:

$$\|hX_\perp \psi_h^j\|_{L^2}^2 + \|hV \psi_h^j\|_{L^2}^2 + \|h^2 X \psi_h^j\|_{L^2}^2 \leq C_{Q,W}(1 + |\Lambda_h(j)|)^2,$$

where $C_{Q,W} > 0$ is a constant depending only on (Q, W) . The fact that the first two terms on the left hand side of this inequality are bounded follows by standard energy estimates together with the facts that $\|Q\|_{\mathcal{C}^0} < 1$ and that $X = [V, X_\perp]$. The fact that $\|h^2 X \psi_h^j\|$ is bounded is much more subtle and it follows from the classical Rothschild–Stein theorem. As we shall discuss it later on, it is a manifestation of the lack of ellipticity of the operator along the X direction. In the following, we aim precisely at analyzing the structure of the eigenfunctions in the *subelliptic regime* where formally speaking one has $h^{-1} \ll |X| \lesssim h^{-2}$.

1.1. QUANTUM LIMITS AND SEMICLASSICAL MEASURES. — We are interested in describing the asymptotic properties of the semiclassical eigenmodes satisfying:⁽¹⁾

$$(3) \quad \widehat{P}_h \psi_h = \Lambda_h \psi_h, \quad \|\psi_h\|_{L^2} = 1, \quad \Lambda_h \longrightarrow \Lambda_0 \in \mathbb{R}, \quad \text{as } h \longrightarrow 0^+.$$

When $W \equiv 0$, a natural choice is to pick $\Lambda_h = 1$ that would correspond to studying the large eigenvalue limit for the hypoelliptic operator $\mathcal{L} = \Delta_{\text{sR}} + iQX + iX(Q)/2$. Yet, as we want to emphasize the semiclassical nature of this spectral problem, we keep a general W and thus some general value $\lambda_0 \geq \min W$. Still from Lemma A.5, one finds that, for any sequence $\Lambda_h \rightarrow \Lambda_0$, there exists some $h_0 > 0$ such that, for all $0 < h \leq h_0$ and for any solution to (3),

$$(4) \quad \|hX_\perp \psi_h\|_{L^2}^2 + \|hV \psi_h\|_{L^2}^2 + \|h^2 X \psi_h\|_{L^2}^2 \leq C_{Q,W}(1 + 2|\Lambda_0|)^2.$$

One says that a probability measure ν is a *quantum limit* for this spectral problem if, for every $a \in \mathcal{C}^0(\mathcal{M})$,

$$\lim_{h \rightarrow 0^+} \int_{\mathcal{M}} a |\psi_h|^2 d\mu_L = \int_{\mathcal{M}} a d\nu,$$

where $(\psi_h)_{h \rightarrow 0^+}$ is a sequence verifying (3). Up to extraction of a subsequence, one can always find such an accumulation point. Given $\Lambda_0 \geq \min W$, we denote by \mathcal{N}_{Λ_0} the set of quantum limits associated with the spectral parameter Λ_0 . The relevance of these measures from the point of view of quantum mechanics is that they describe the probability of finding a particle in the quantum state ψ_h . From the mathematical

⁽¹⁾All along the article, we use the standard conventions from semiclassical analysis to write $h \rightarrow 0^+$ instead of writing a sequence $h_n \rightarrow 0$ as $n \rightarrow \infty$.

perspective, they allow to give some information on the regularity of eigenfunctions in the large eigenvalue limit.

In view of describing the regularity properties of ν , one lifts the problem to the cotangent bundle $T^*\mathcal{M}$ by introducing

$$w_h : a \in \mathcal{C}_c^\infty(T^*\mathcal{M}) \longmapsto \langle \text{Op}_h(a)\psi_h, \psi_h \rangle_{L^2},$$

where $\text{Op}_h(a)$ is a h -pseudodifferential operator with principal symbol a [Zwo12, Th. 14.1] and $(\psi_h)_{h \rightarrow 0^+}$ is the sequence used to generate ν . Thanks to the Calderón-Vaillancourt theorem [Zwo12, Th. 5.1], $(w_h)_{h \rightarrow 0^+}$ is a bounded sequence in $\mathcal{D}'(T^*\mathcal{M})$. Hence, up to extraction, it converges to some limit w which is referred as a semiclassical measure for the sequence $(\psi_h)_{h \rightarrow 0^+}$. The theory of semiclassical pseudodifferential operators allows to prove that any such w is a *finite nonnegative measure* on $T^*\mathcal{M}$ that is supported on

$$\mathcal{E}^{-1}(\Lambda_0) := \{(q, p) \in T^*\mathcal{M} : \mathcal{E}(q, p) := H_2(q, p)^2 + H_3(q, p)^2 + W(q) = \Lambda_0\},$$

and that satisfies the following invariance property

$$\{H_2^2 + H_3^2 + W, w\} = 0,$$

where

$$H_2(q, p) := p(X_\perp), \quad \text{and} \quad H_3(q, p) := p(V).$$

See for instance [Zwo12, §5.2] for proofs of these classical facts in the case of \mathbb{R}^{2d} . We emphasize that, contrary to the case of eigenvalue problems of elliptic nature, the energy layer $\mathcal{E}^{-1}(\Lambda_0)$ is not compact and there may be some escape of mass at infinity. In particular, w could be equal to 0. See for instance Section 8 for concrete examples in the case of the flat torus. Due to this escape of mass at infinity, it is natural to study the measure

$$\nu_\infty := \nu - \pi_* w,$$

where $\pi : (q, p) \in T^*\mathcal{M} \rightarrow q \in \mathcal{M}$, and this is the main purpose of the present work.

1.2. DECOMPOSITION OF THE MEASURE ν_∞ AND INVARIANCE PROPERTIES

Colin de Verdière, Hillairet and Trélat proved in [CHT18, Th. B] that $X(\nu_\infty) \equiv 0$ when $Q \equiv 0$ and $W \equiv 0$. Our results generalize this theorem in two directions. First, we will provide a refined description of ν_∞ , showing that the measure ν_∞ decomposes into a discrete sum of non-negative Radon measures covering different asymptotic regimes $h^{-1} \ll |X| \lesssim h^{-2}$ across the non-compact part of $\mathcal{E}^{-1}(\Lambda_0)$. Second, we will prove that each of these measures satisfies a new invariance property, different from each-other, as soon as $\nabla(Q)$ does not vanish. In view of formulating our results, we associate to each smooth function f on \mathcal{M} a natural vector field lying in the contact plane $D := \text{Span}(X_\perp, V)$ given by

$$\Omega_f := V(f)X_\perp - X_\perp(f)V.$$

Our main theorem then reads:

THEOREM 1.1. — *Let $Q, W \in \mathcal{C}^\infty(\mathcal{M}, \mathbb{R})$ such that $\|Q\|_{\mathcal{C}^0} < 1$, let $\Lambda_0 > \max_{q \in \mathcal{M}} W(q)$ and set*

$$(5) \quad Y_W := X + \Omega_{\ln(\Lambda_0 - W)}.$$

Then, for every $\nu \in \mathcal{N}_{\Lambda_0}$, the measure ν_∞ decomposes as

$$(6) \quad \nu_\infty = \bar{\nu}_\infty + \sum_{k=0}^\infty (\nu_{k,\infty}^+ + \nu_{k,\infty}^-),$$

where $\bar{\nu}_\infty$ and $\nu_{k,\infty}^\pm$ are non-negative Radon measures on \mathcal{M} verifying, for all $a \in \mathcal{C}^1(\mathcal{M})$ and for all $k \in \mathbb{Z}_+$,

$$\int_{\mathcal{M}} Y_W(a) d\bar{\nu}_\infty = 0, \quad \text{and} \quad \int_{\mathcal{M}} Y_{W,Q,k}^\pm(a) d\nu_{k,\infty}^\pm = 0,$$

with

$$Y_{W,Q,k}^\pm := (\pm(2k+1) + Q)Y_W - \Omega_Q.$$

REMARK 1.2. — We emphasize the importance for Theorem 1.1 of considering sub-principal terms of the form $-ih^2QX - ih^2X(Q)/2$ in the definition of \widehat{P}_h . Indeed, the resulting quantum limits $\bar{\nu}_\infty$ and ν_k^\pm each satisfy a different invariance property while, for $Q \equiv 0$, only one invariance property for ν_∞ occurs.

REMARK 1.3. — Condition $\Lambda_0 > \max_{q \in \mathcal{M}} W(q)$ ensures that the classical forbidden region is empty. In the case $\min W \leq \Lambda_0 \leq \max W$, the support of $\bar{\nu}_\infty$ becomes confined inside the compact set $\mathcal{M}_{\Lambda_0,W} := \{q \in \mathcal{M} : \Lambda_0 - W \geq 0\}$, while the support of $\nu_{k,\infty}^\pm$ is contained in the open subset $\mathcal{U}_{\Lambda_0,W} := \{q \in \mathcal{M} : \Lambda_0 - W > 0\}$. This more general situation is covered by the more precise description of semiclassical measures stated in Theorem 7.1.

REMARK 1.4. — In Section 8, by working on the flat torus $M = \mathbb{T}^2$, we show examples of sequences (ψ_h, Λ_h) satisfying (3) for which the measures $\bar{\nu}_\infty$ or $\nu_{k,\infty}^\pm$ we construct carry the total mass of ν .

Decomposition (6) reflects the stratification of the asymptotic phase-space distribution of a given sequence (ψ_h, Λ_h) satisfying (3). Indeed, in Section 7 below, we will provide a more general description of ν_∞ by lifting the analysis to the phase-space via introducing an adapted semiclassical measure μ_∞ on $\mathcal{M} \times \mathbb{R}$ such that, by projection:⁽²⁾

$$\nu_\infty(q) = \int_{\mathbb{R}} \mu_\infty(q, dE).$$

The extra variable $E \in \mathbb{R}$ parameterizes the phase-space escape of mass along the degenerate direction of X as $h \rightarrow 0^+$. We refer to (28) below for the explicit construction of the measure μ_∞ using semiclassical tools and we just give here some informal

⁽²⁾Letting μ be a finite Radon measure on $\mathcal{M} \times \mathbb{R}$, the measure $\nu(q) = \int_{\mathbb{R}} \mu(q, dE)$ is defined by

$$\langle \nu, a \rangle := \int_{\mathcal{M} \times \mathbb{R}} a(q) d\mu(q, E), \quad \text{for all } a \in \mathcal{C}^0(\mathcal{M}).$$

explanation. Letting $H_1(q, p) = p(X)$, we will study precisely two different asymptotic regimes generating a splitting of the measure μ_∞ into two parts

$$\mu_\infty = \mathbf{1}_{E \neq 0} \mu_\infty + \mathbf{1}_{E=0} \mu_\infty$$

of qualitatively different nature:

– The *critical subelliptic regime* $h|H_1| \asymp 1$, captured by $\mathbf{1}_{E \neq 0} \mu_\infty$, displays a quantum behavior which manifests as a decomposition of this measure into a discrete sum of Radon measures $(\mu_{k,\infty}^\pm)_{k \in \mathbb{N}}$ supported on quantized level sets $\mathcal{H}_\pm^{-1}(2k+1) \subset \mathcal{M} \times \mathbb{R}_\pm^*$ for the energy functions

$$(7) \quad \mathcal{H}_\pm(q, E) := \pm \left(\frac{\Lambda_0 - W(q)}{E} - Q(q) \right), \quad (q, E) \in \mathcal{M} \times \mathbb{R}_\pm^*.$$

These measures project on the manifold \mathcal{M} and give the measures $\nu_{k,\infty}^\pm$:

$$\nu_{k,\infty}^\pm(q) = \int_{\mathbb{R}} \mu_{k,\infty}^\pm(q, dE).$$

– The *subcritical subelliptic regime* $1 \ll |H_1| \ll h^{-1}$, captured by the measure $\bar{\mu}_\infty = \mathbf{1}_{E=0} \mu_\infty$, which is supported on $\mathcal{M} \times \{0\}$. This measure projects on \mathcal{M} so that:

$$\bar{\nu}_\infty(q) = \int_{\mathbb{R}} \bar{\mu}_\infty(q, dE).$$

Besides this distinction between the different oscillation regimes, our analysis will show the influence of the hypoelliptic perturbations of $-\Delta_{\text{sR}}$ given by (2) in the previous description by obtaining new invariance properties of μ_∞ in terms of Q and W . Among other things, it illustrates that the introduction of the new variable $E = hH_1$ becomes essential for this description even in the non-semiclassical set-up where $W \equiv 0$.

1.3. MORE RELATED RESULTS AND QUESTIONS

The fine analysis of these regimes $h \ll |E| = h|H_1| \lesssim 1$ in the subelliptic region of $T^*\mathcal{M}$ is reminiscent of the analysis made by Burq and Sun for the semiclassical measures of Baouendi-Grushin operators in [BS22] (see also [LS23, AS23] for related works). More precisely, Theorem 7.1 below can be compared with the results obtained by the first author and Sun in [AS23] where a detailed study of semiclassical measures in the subelliptic regime for quasimodes of the Baouendi-Grushin operator was performed. In these references, the model operator is $\Delta_G := \partial_x^2 + a(x)^2 \partial_y^2$, where $(x, y) \in \mathbb{T}^2$ and $a(x)$ is a smooth function that may vanish at finitely many isolated points (with non-vanishing derivative). In particular, this operator can be written as a sum of squares and one has $[\partial_x, a(x)\partial_y] = a'(x)\partial_y$. Thanks to Hörmander's bracket condition [Hör67], it defines an hypoelliptic operator and similar questions can be raised on the asymptotic properties of its eigenfunctions. These are exactly the questions raised in [BS22] and the role of H_1 is then played by the cotangent variable η that is dual to y . As our Theorem 1.1, [AS23, §3] gives a full description of the eigenmodes in the regime $1 \ll |\eta| \lesssim h^{-1}$ through their semiclassical measures. In particular, the

invariance of these measures through the vector field ∂_y is shown and it replaces the geodesic vector field X in that context.

In [CHT18], the hypoelliptic model is closer to ours but this extra variable hH_1 did not appear in the description of ν_∞ because of the use of microlocal methods. The reason for introducing this new variable $E = hH_1$ is that, in the regime $E \neq 0$, the term h^2QX is not negligible anymore compared with $-h^2\Delta_{\text{sR}}$ and it has to be taken into account in the description of the quantum limit. It results in new invariance properties as in Theorem 1.1. The fact that the eigenfunctions are localized on specific levels is a manifestation of the fact that our hypoelliptic operators are modeled locally on the 3-dimensional Heisenberg group (and thus related to the 1-dimensional harmonic oscillator). More specifically, our proof of this support property will only rely on the fact that the sub-Riemannian Laplacian can be written as

$$(8) \quad \Delta_{\text{sR}} = Z^*Z - iX = ZZ^* + iX, \quad [Z, Z^*] = 2iX, \quad \text{where } Z = X_\perp + iV.$$

This quantization of the level sets can be thought as an analogue in our (non-algebraic) set-up of the decomposition appearing in the works of Fermanian-Kammerer and Fischer [FKF21, Th. 1.1, Th. 2.10]. See also [FKL21] for related results in the compact setting. In these references, the decomposition of the semiclassical measures along these quantized levels shows up because there is a natural way to diagonalize the sub-Riemannian Laplacian along the elliptic variables. This is exactly where the harmonic oscillator appears in these references and the subelliptic variable H_1 corresponds exactly to the center direction of the Lie algebra setting from [FKF21, FKL21]. Yet, this algebraic decomposition does not distinguish the various scales of oscillations $h \ll h|H_1| \lesssim 1$ as we are doing in the present work or as it was the case in [BS22, AS23]. In [FKF21, FKL21], the proof of this decomposition required the introduction of operator-valued semiclassical measures. In the case of more general contact manifolds, we can also mention the works of Taylor [Tay20] regarding the question of microlocal Weyl laws for operator-valued symbols. Here, our proof of these support properties will not rely on the introduction of such analytical objects. It will simply follow from a careful use of the relation (8) where Z and Z^* will play the role of ladder operators, in a similar way to the proof that the eigenvalues of the harmonic oscillator are given by $\{2k + 1, k \geq 0\}$.

REMARK 1.5. — In the general 3-dimensional contact case treated in [CHT18], the quantum normal form as formulated in [Col23, §6.2] should in principle allow to get as in [FKF21] a natural decomposition of the measure ν_∞ using the spectral decomposition of the harmonic oscillator. Yet, due to its microlocal nature, it would again not distinguish the various subelliptic regimes $1 \ll |H_1| \lesssim h^{-1}$ involved in our problem as we are doing here.

REMARK 1.6. — The related works of Boil and Vu Ngoc [BVN21] are also relevant towards this kind of decomposition. Indeed, in this reference the long time dynamics of the semiclassical magnetic Schrödinger equation is studied in details in dimension 2. At the time scale $1/h$, it is in fact shown that coherent states split according to the

low Landau levels of the operator with an effective dynamics given by the magnetic field which is the analogue of our Reeb vector field. In other words, it is the exact analogue in this context of our splitting of quantum limits through the measures $\nu_{k,\infty}^\pm$. We also refer to [AS23, §5] where a similar coherent state decomposition is described for the Baouendi-Grushin operator Δ_G .

As our semiclassical analysis of eigenfunctions for hypoelliptic operators is inspired by the fine analysis performed for the Baouendi-Grushin operator in [BS22, AS23], it is natural to expect that such results would remain true for similar hypoelliptic perturbations of the Baouendi-Grushin operator. Similarly, the analysis presented here should in principle allow to deal with the controllability of the Schrödinger equation as in [BS22] and with the stabilization of the wave equation as in [AS23]. Yet, this would require more work that is beyond the scope of the present article. Another natural question would be to study more precisely the regularity of quantum limits when the geodesic vector field X enjoys specific dynamical structure, e.g. on Zoll surfaces, on flat tori or on negatively curved surfaces. Among the natural questions to explore is whether one can always find sequences of eigenfunctions concentrating on a given level set $\mathcal{H}_\pm^{-1}(2k+1)$. Related to this, it would be interesting to describe semiclassical Weyl laws for symbols involving the extra variable $E = hH_1$. In that direction, we refer one more time to [Tay20] for microlocal Weyl laws with operator-valued symbols (including the case where Q is not identically equal to 0) on contact manifolds of dimension ≥ 3 . Finally, these hypoelliptic models are naturally related to semiclassical magnetic Schrödinger operators. For instance, in view of the works [RV15, HKRV16, Mor22, Mor24], it would be natural to compare how the fine structure of eigenfunctions of these models could be understood following the lines of the present work. Recall that rather precise descriptions of the low-energy eigenfunctions were already given via WKB and normal form methods in [BR20, GRV21, GNRV21].

1.4. A FEW WORDS ON MORE GENERAL SUB-RIEMANNIAN CONTACT LAPLACIANS IN DIMENSION 3. — The simple geometric model considered in this article ensures that we have globally defined vector fields (X_\perp, V) generating the contact structure and that $[X_\perp, X] = KV$ and $[V, X] = -X_\perp$. It makes some aspects of the exposition somewhat lighter (e.g. regarding the normal form procedure) but it is not essential in our analysis. In fact, one would only need to have two locally defined generating vector fields (X_2, X_3) on a 3-dimensional manifold \mathcal{N} so that the operator Δ_{sR} writes down locally as $X_2^*X_2 + X_3^*X_3$ (modulo some order 0 operator) where the adjoint is taken with respect to a smooth (nonvanishing) volume form and where one has (locally)

$$(9) \quad T_q\mathcal{N} = \text{Span}(X_2(q), X_3(q), [X_2, X_3](q)).$$

This last condition ensures that $D = \text{Span}(X_2, X_3)$ is non-integrable and thus a contact structure. The Hörmander type condition (9) is in fact the only ingredient needed to perform our normal form procedure in Section 5. For the sake of exposition and as geodesic vector fields form a natural and rich family of Reeb vector fields, we do

not deal with the most general case and we focus on the somehow simplest example of contact structure⁽³⁾ that is not already in normal form. In fact, we emphasize that, contrary to the flat Heisenberg case [CHT18, FKF21, FKL21], the brackets $[X_\perp, X]$ and $[V, X]$ do not identically vanish. This implies that we do not have a nice algebraic structure at hand and that we have to take these nonvanishing brackets into account in our analysis as it is the case in the general contact set-up treated in [CHT18]. In fact, as we shall see below, the way we deal with the normal form procedure slightly differs from the one in [CHT18] by avoiding an “explicit” construction of symplectic coordinates and thus the use of Fourier integral operators. Yet this simplified method does not rely on the specific form of our operator. It would work as well for more general sub-Riemannian contact Laplacians in dimension 3 modulo dealing with more cumbersome cohomological equations and modifying conveniently the various functions and vector fields in the subelliptic direction. In particular, if we write $-ihX_2$ and $-ihX_3$ as $\text{Op}_h^w(H_2)$ and $\text{Op}_h^w(H_3)$ using locally the semiclassical Weyl quantization (modulo terms of order 0), then we could set $H_1 := \{H_3, H_2\}$ to measure the escape of mass at infinity. When studying the measure ν_∞ (i.e., the regime $1 \ll |H_1| \lesssim h^{-1}$), the geodesic vector field X would be replaced as in [CHT18] by the Reeb vector field $X_1 + \alpha X_2 + \beta X_3$ with

$$X_1 := [X_3, X_2], \quad [X_2, X_1] = \beta X_1 \pmod{D}, \quad \text{and} \quad [X_1, X_3] = \alpha X_1 \pmod{D}.$$

1.5. ORGANIZATION OF THE ARTICLE. — In Section 2, we fix the geometric and semiclassical conventions that are used all along the article. In particular, we reduce the analysis to a local chart which allows us to use the Weyl quantization. Then, in Section 3, we microlocalize our eigenfunctions in the region $|H_1| \gg 1$ and we introduce microlocal lifts of our measures that capture the escape of mass at infinity. This is where we introduce the new variable $E = hH_1$ in order to blow up the direction H_1 where the eigenmodes concentrate at infinity. Equivalently this amounts to perform a second microlocalization along this direction. In Section 4, we show that these microlocal lifts are concentrated on certain quantized layers along the E -variable using the commutation relations (1) and the a priori estimates (4) together with standard rules of pseudodifferential calculus. In Section 5, we introduce a simple normal form procedure that is well adapted to the geometry of our problem and we implement it in Section 6 to derive the invariance properties of our lifted measure. Section 7 summarizes the main results of the article and show how they can be used to derive Theorem 1.1 from the introduction. Section 8 treats the simple example of the flat torus in view of illustrating our analysis in a concrete example. Finally, the article contains two appendices: one devoted to the spectral properties of our hypoelliptic operators (Appendix A) and another one collecting a few standard facts from semiclassical analysis (Appendix B).

⁽³⁾Another nice class of examples would be given by the operator $X^2 + X_\perp^2$ on negatively/positively curved surfaces.

Acknowledgements. — We warmly thank G. Carron, B. Chantraine, C. Fermanian-Kammerer, B. Helffer, L. Hillairet, F. Macià and C. Sun for discussions related to various aspects of this work. We are also grateful to an anonymous referee for a careful and detailed report.

2. SEMICLASSICAL CONVENTIONS

In this preliminary section, we introduce the conventions from differential geometry and semiclassical analysis required for our study and used all along the article. In Section 2.1, we recall the existence of local isothermal coordinates to write down the differential objects appearing in our framework in a simple and concrete manner. Then, in Section 2.2, we introduce the principal symbols of these operators together with their commutation expressions inherited from (1). With these expressions at hand, we rewrite in Section 2.3 our main differential operators and the eigenvalue equation (3) using the Weyl quantization in the local chart. We conclude this section by discussing in Section 2.4 the symbolic properties of certain class of polynomials expressions that will appear in the normal form procedure of Section 5.

2.1. LOCAL ISOTHERMAL COORDINATES. — Near any given point $m_0 \in M$, one can find a system of local coordinates $(x, y) \in U_0 \subset \mathbb{R}^2$ (with $(0, 0)$ being the image of m_0) such that the metric g writes down in a conformal way [PSU23, Th. 3.4.8]:

$$g(x, y) := e^{2\lambda(x, y)}(dx^2 + dy^2).$$

We denote this neighborhood inside M by U in the sequel.

To write down the geometrical objects involved in the problem in terms of local isothermal coordinates, we follow the presentation of [PSU23, §3.5] and we refer to it for more details. If we denote by z the angle between a unit vector $p \in S_q U_0$ and $\partial/\partial x$, then we have the following expressions for our vector fields [PSU23, Lem. 3.5.6]:

$$X := e^{-\lambda}(\cos z \partial_x + \sin z \partial_y) + e^{-\lambda}(-\partial_x \lambda \sin z + \partial_y \lambda \cos z) \partial_z,$$

$$X_\perp := e^{-\lambda}(\sin z \partial_x - \cos z \partial_y) + e^{-\lambda}(\partial_x \lambda \cos z + \partial_y \lambda \sin z) \partial_z,$$

and $V := \partial_z$.

These expressions are obtained by solving the Hamilton-Jacobi equation for the Hamiltonian function $e^{2\lambda(x, y)}(\xi^2 + \eta^2)$ on $T^*\mathbb{R}^2$. The Sasaki metric is not conformal in the system of coordinates (x, y, z) . Yet, the volume form has a simple expression

$$d\mu_L(x, y, z) = e^{2\lambda(x, y)} dx dy dz.$$

REMARK 2.1. — In order to prove Theorem 1.1, it is sufficient to prove the result in a local chart and to work locally with a standard Euclidean quantization procedure. We use this isothermal chart for the sake of concreteness and for simplicity of exposition. Among several advantages, it will allow us to write very concretely our operators in this chart through the Weyl quantization and to have concrete expressions for the subprincipal symbols. It will also be convenient to write rather simple expressions in

the normal form expansion of Section 5. Yet, our dynamical and semiclassical arguments would work as well for more general contact flows for which such a nice chart does not exist; thus with slightly more cumbersome expressions for our operators.

REMARK 2.2. — Without loss of generality, we can extend these operators on $\mathcal{U}_0 := SU_0 = U_0 \times \mathbb{S}^1$ to operators on $\mathbb{R}^2 \times \mathbb{S}^1$ by extending λ into a smooth compactly supported function on \mathbb{R}^2 .

2.2. HAMILTONIAN FORMULATION. — In the following, we will make use of different tools of semiclassical pseudodifferential calculus. This leads to define the symbols corresponding to the operators of interest. To this aim, we introduce the Hamiltonian functions associated with the orthonormal frame (X, X_\perp, V) . Namely, we define the following symbols on $T^*(\mathbb{R}^2 \times \mathbb{S}^1)$:

$$H_1(x, y, z, \xi, \eta, \zeta) := e^{-\lambda(x,y)} (\xi \cos z + \eta \sin z + \zeta(-\partial_x \lambda \sin z + \partial_y \lambda \cos z)),$$

$$H_2(x, y, z, \xi, \eta, \zeta) := e^{-\lambda(x,y)} (\xi \sin z - \eta \cos z + \zeta(\partial_x \lambda \cos z + \partial_y \lambda \sin z)),$$

and $H_3(x, y, z, \xi, \eta, \zeta) := \zeta$.

Notice, in particular, that there exists a positive constant C_0 (depending only on our local isothermal coordinates and on our extension of λ to \mathbb{R}^2) verifying

$$(10) \quad C_0^{-1}(\xi^2 + \eta^2 + \zeta^2) \leq H_1^2 + H_2^2 + H_3^2 \leq C_0(\xi^2 + \eta^2 + \zeta^2).$$

The commutator relations (1) can then be translated into the following Poisson bracket commutation formulas:

$$(11) \quad \{H_1, H_3\} = H_2, \quad \{H_1, H_2\} = -KH_3, \quad \text{and} \quad \{H_2, H_3\} = -H_1,$$

where we recall that $K(x, y)$ is the scalar curvature. We also collect a few useful relations involving H_1, H_2, H_3 , and λ in the next lemma, whose proof is immediate:

LEMMA 2.3. — *The following identities hold:*

$$(12) \quad \partial_x H_3 = \partial_y H_3 = \partial_z H_3 = 0,$$

$$(13) \quad \partial_z H_2 = H_1, \quad \partial_z H_1 = -H_2,$$

$$(14) \quad \begin{pmatrix} \partial_x H_1 \\ \partial_y H_1 \end{pmatrix} = - \begin{pmatrix} \partial_x \lambda & e^{-\lambda} (\partial_x^2 \lambda \cos z + \partial_{xy}^2 \lambda \sin z) \\ \partial_y \lambda & e^{-\lambda} (\partial_{xy}^2 \lambda \cos z + \partial_y^2 \lambda \sin z) \end{pmatrix} \begin{pmatrix} H_1 \\ H_3 \end{pmatrix},$$

$$(15) \quad \begin{pmatrix} \partial_x H_2 \\ \partial_y H_2 \end{pmatrix} = - \begin{pmatrix} \partial_x \lambda & e^{-\lambda} (\partial_x^2 \lambda \sin z - \partial_{xy}^2 \lambda \cos z) \\ \partial_y \lambda & e^{-\lambda} (\partial_{xy}^2 \lambda \sin z - \partial_y^2 \lambda \cos z) \end{pmatrix} \begin{pmatrix} H_2 \\ H_3 \end{pmatrix}.$$

2.3. SEMICLASSICAL WEYL QUANTIZATION. — With the above conventions, we can next rewrite the geometrical objects introduced in Section 2.1 in terms of pseudodifferential operators by making use of the Hamiltonian formulation of Section 2.2. Precisely, we have:

$$\frac{\hbar}{i}X = \text{Op}_\hbar^w(H_1 - ihX(\lambda)), \quad \frac{\hbar}{i}X_\perp = \text{Op}_\hbar^w(H_2 - ihX_\perp(\lambda)), \quad \text{and} \quad \frac{\hbar}{i}V = \text{Op}_\hbar^w(H_3),$$

where Op_h^w stands for the semiclassical Weyl quantization on $\mathbb{R}^2 \times \mathbb{S}^1$ (see Appendix B). In particular,

$$-h^2 \Delta_{\text{sR}} = \text{Op}_h^w(H_2^2 + H_3^2 - 2ihX_\perp(\lambda)H_2 - h^2 r_\lambda),$$

where $r_\lambda(x, y, z)$ is a smooth compactly supported function that is independent of h (but depending on the choice of local coordinates). It will be slightly more convenient to work in the local chart with the operator $-h^2 e^\lambda \Delta_{\text{sR}} e^{-\lambda}$ due to the following conjugation formula:

LEMMA 2.4. — *The following holds on \mathcal{U}_0 :*

$$-h^2 e^\lambda \Delta_{\text{sR}} e^{-\lambda} = \text{Op}_h^w(H_2^2 + H_3^2 + h^2 \mathbf{r}_\lambda),$$

where $\mathbf{r}_\lambda(x, y, z)$ is a smooth compactly supported function that is independent of h (but depending on the choice of local coordinates).

Proof. — One has

$$e^\lambda \frac{h}{i} X_\perp e^{-\lambda} = \frac{h}{i} X_\perp + ihX_\perp(\lambda) = \text{Op}_h^w(H_2),$$

from which we infer that

$$-h^2 e^\lambda \Delta_{\text{sR}} e^{-\lambda} = \text{Op}_h^w(H_2)^2 + \text{Op}_h^w(H_3)^2.$$

The conclusion of the lemma follows then from the composition rule (64) for pseudodifferential operators. \square

We are now in position to obtain the expression of the full operator \widehat{P}_h . To do that, we first use the Weyl pseudodifferential calculus to write

$$-ih^2 QX - \frac{ih^2 X(Q)}{2} = \text{Op}_h^w(hQH_1 + h^2 W_1),$$

where $W_1 \in \mathcal{C}_c^\infty(\mathbb{R}^2 \times \mathbb{S}^1, \mathbb{C})$ is independent of h . Using the composition rules for the Weyl quantization, we obtain

$$(16) \quad \widehat{P}_h = e^{-\lambda} \text{Op}_h^w(H_2^2 + H_3^2 + hQH_1 + W + h^2 W_{1,\lambda}) e^\lambda,$$

where $W_{1,\lambda} \in \mathcal{C}_c^\infty(\mathbb{R}^2 \times \mathbb{S}^1, \mathbb{C})$ is independent of h . Regarding this expression, it is natural to set

$$(17) \quad u_h = e^\lambda \psi_h,$$

and thanks to (16), u_h solves locally in \mathcal{U}_0 the eigenvalue equation

$$(18) \quad \widehat{P}_{h,\lambda} u_h = \Lambda_h u_h,$$

where

$$\widehat{P}_{h,\lambda} := \text{Op}_h^w(H_2^2 + H_3^2 + hQH_1 + W + h^2 W_{1,\lambda}).$$

The a priori estimate (4) from the introduction then reads

$$(19) \quad \|\text{Op}_h^w(H_2)u_h\|_{L^2(\mathcal{X})} + \|\text{Op}_h^w(H_3)u_h\|_{L^2(\mathcal{X})} + \|\text{Op}_h^w(hH_1)u_h\|_{L^2(\mathcal{X})} \leq 2C_{Q,W,\mathcal{X}}(1 + |\Lambda_0|),$$

where \mathcal{K} is any compact subset of \mathcal{U}_0 and the L^2 norm is now taken with respect to the standard Lebesgue measure $dx dy dz$ on $\mathcal{K} \subset \mathbb{R}^2 \times \mathbb{S}^1$.

2.4. CLASS OF SYMBOLS IN THE REGION $|H_1| \gg \sqrt{H_2^2 + H_3^2}$. — In Section 5 below, we will describe a normal form procedure that will naturally involve functions in the spaces:

$$\mathcal{P}_N(\mathbb{R}^2 \times \mathbb{S}^1) := \left\{ \sum_{\substack{\alpha=(\alpha_2, \alpha_3) \in \mathbb{Z}_+^2 \\ |\alpha| \leq N}} a_\alpha (H_2/H_1)^{\alpha_2} (H_3/H_1)^{\alpha_3} : \forall \alpha, a_\alpha \in \mathcal{C}^\infty(\mathbb{R}^2 \times \mathbb{S}^1) \right\}.$$

Notice that as a consequence of (12), (13), (14) and (15), one can verify the following:

LEMMA 2.5. — *Let P be an element in $\mathcal{P}(\mathbb{R}^2 \times \mathbb{S}^1) := \bigcup_N \mathcal{P}_N(\mathbb{R}^2 \times \mathbb{S}^1)$. Then, $\partial_x P$, $\partial_y P$ and $\partial_z P$ belong to $\mathcal{P}(\mathbb{R}^2 \times \mathbb{S}^1)$. Similarly, letting $p = (\xi, \eta, \zeta)$, one has that, for every $\gamma \in \mathbb{Z}_+^3$, $H_1^{|\gamma|} \partial_p^\gamma P$ belongs to $\mathcal{P}(\mathbb{R}^2 \times \mathbb{S}^1)$.*

Proof. — The first part of the lemma is direct consequence of (12), (13), (14) and (15). For the second part, we proceed by induction on $|\gamma|$ and use the fact that H_1, H_2 and H_3 are linear functions in (ξ, η, ζ) . □

To make the necessary estimates in the sub-elliptic regime arising from our problem, we will be naturally led to work in the “conic” region

$$C_\varepsilon(\mathcal{K}) := \left\{ (q, p) \in T^*(\mathcal{K}) : \varepsilon |H_1(q, p)| \geq \sqrt{1 + H_2^2(q, p) + H_3^2(q, p)} \right\},$$

where \mathcal{K} is a compact subset of $\mathbb{R}^2 \times \mathbb{S}^1$ and where $\varepsilon \in (0, 1]$ is some small parameter that is intended to tend to 0 in the end. We record the following corollary of Lemma 2.5:

COROLLARY 2.6. — *Let P in $\mathcal{P}(\mathbb{R}^2 \times \mathbb{S}^1)$ and let \mathcal{K} be a compact subset of $\mathbb{R}^2 \times \mathbb{S}^1$. For every $0 < \varepsilon \leq 1$ and for every $(\alpha, \beta) \in \mathbb{Z}_+^6$, one can find a constant $C_{\varepsilon, P, \mathcal{K}, \alpha, \beta}$ such that, for every $(q, p) = (x, y, z, \xi, \eta, \zeta)$ in $C_\varepsilon(\mathcal{K})$, one has*

$$|\partial_q^\alpha \partial_p^\beta P| \leq C_{\varepsilon, P, \mathcal{K}, \alpha, \beta} \langle p \rangle^{-|\beta|},$$

where $\langle p \rangle := (1 + \xi^2 + \eta^2 + \zeta^2)^{1/2}$.

In particular, elements in $\mathcal{P}(\mathbb{R}^2 \times \mathbb{S}^2)$ satisfy the properties of the class of (Kohn-Nirenberg) symbols $S_{\text{cl}}^0(T^*(\mathbb{R}^2 \times \mathbb{S}^1))$ defined in Appendix B inside $C_\varepsilon(\mathcal{K})$ for any compact subset \mathcal{K} of $\mathbb{R}^2 \times \mathbb{S}^1$.

3. REDUCTION TO THE SUBELLIPTIC REGIME

Since our results on quantum limits and semiclassical measures in the subelliptic regime are essentially local, we will restrict ourselves to study the following measures on \mathcal{U}_0 :

$$\nu_h : a \in \mathcal{C}_c^\infty(\mathcal{U}_0) \longmapsto \int_{\mathcal{U}_0} a(x, y, z) |\psi_h(x, y, z)|^2 e^{2\lambda(x, y)} dx dy dz,$$

where \mathcal{U}_0 is a bounded open subset of $\mathbb{R}^2 \times \mathbb{S}^1$ given by local isothermal coordinates and where (ψ_h, Λ_h) is a sequence satisfying (3). As the sequence (ψ_h) is normalized,

this defines a sequence of measures on \mathcal{U}_0 that are of finite mass ≤ 1 . In fact, up to an extraction, one can suppose that $\nu_h \rightharpoonup \nu$ as $h \rightarrow 0^+$ and the limit measure is supported in \mathcal{U}_0 with total mass ≤ 1 . We fix this converging subsequence for the rest of the article.

Using the convention from (18), this can be rewritten as

$$\nu_h : a \in \mathcal{C}_c^\infty(\mathcal{U}_0) \mapsto \int_{\mathcal{U}_0} a(x, y, z) |u_h(x, y, z)|^2 dx dy dz,$$

which allows us to work with the standard Lebesgue measure. Before trying to prove our main theorem, we will first show in this Section how to reduce these integrals to the region of the phase space where $1 \ll |H_1| \lesssim h^{-1}$ and thus how to define the measure μ_∞ from the introduction. We remark that the measures ν_h can be rewritten as

$$(20) \quad \langle \nu_h, a \rangle = \langle \text{Op}_h^w(a) u_h, u_h \rangle_{L^2}.$$

More generally, as anticipated in the introduction, we consider the associated Wigner distribution

$$\forall a \in \mathcal{C}_c^\infty(T^*\mathcal{U}_0), \quad \langle w_h, a \rangle := \langle \text{Op}_h^w(a) u_h, u_h \rangle_{L^2},$$

and, up to another extraction, we can suppose that it converges to some (finite) limit measure w on $T^*\mathcal{U}_0$.

REMARK 3.1. — As usual when working with coordinate charts, we make a small abuse of notations and write ψ_h for the image of ψ_h in the coordinate system. As we always suppose a to be compactly supported in the chart, this causes no difficulties (up to $\mathcal{O}(h^\infty)$ remainders) and we may view ψ_h as a smooth compactly supported function on $\mathbb{R}^2 \times \mathbb{S}^1$.

In the rest of the article, we fix a smooth function $\chi : \mathbb{R} \rightarrow [0, 1]$ which is equal to 1 on $[-1, 1]$ and to 0 outside $[-2, 2]$. Moreover, we make the assumption that $\chi' \geq 0$ on \mathbb{R}_- and $\chi' \leq 0$ on \mathbb{R}_+ . For such a function, we also set

$$\tilde{\chi} = 1 - \chi.$$

For the description of the limit measure $\nu_\infty = \nu - \pi_* w$ and for the definition of μ_∞ , we reduce the analysis of the sequence ν_h to the subelliptic regime $1 \ll |H_1| \lesssim h^{-1}$. To do so, we proceed in three steps:

– In Section 3.1, we microlocalize our measures on the sphere at infinity in T^*M through the use of appropriate cut-off functions.

– In Section 3.2, we show that eigenmodes are in fact microlocalized on two points of this sphere at infinity (namely the two points corresponding to the direction H_1) using one more time appropriate cutoff functions.

– In Section 3.3 and Section 3.4, we perform a kind of blow-up of these two points at infinity by introducing a new variable $E := hH_1$ and we show that eigenfunctions are microlocalized in the region $|E| < \infty$.

3.1. REDUCTION TO THE REGION AT INFINITY. — First, we split the measure ν_h into two parts corresponding to the compact and non-compact distribution of the sequence $(u_h)_{h \rightarrow 0^+}$ in phase space. It leads respectively to the definition of the weak limits $\pi_* w$ and ν_∞ . Let $R > 1$, we introduce the cut-off functions

$$(21) \quad \chi_R^B := \chi\left(\frac{H_1^2 + H_2^2 + H_3^2}{R}\right), \quad \tilde{\chi}_R^B = 1 - \chi_R^B.$$

These cut-offs allow us to split $\nu_h = \nu_{h,R} + \nu_h^R$ where

$$\forall a \in \mathcal{C}_c^\infty(\mathcal{U}_0), \quad \langle \nu_{h,R}, a \rangle = \langle w_h, a \chi_R^B \rangle, \quad \text{and} \quad \langle \nu_h^R, a \rangle = \langle w_h, a \tilde{\chi}_R^B \rangle.$$

Notice moreover that the cut-offs χ_R^B and $\tilde{\chi}_R^B$ belong to the admissible class of symbols $S_{\text{cl}}^0(T^*(\mathbb{R}^2 \times \mathbb{S}^1))$ defined in Appendix B. Letting $h \rightarrow 0^+$ and $R \rightarrow +\infty$ (in this order), one finds

$$\lim_{R \rightarrow +\infty} \lim_{h \rightarrow 0^+} \langle \nu_{h,R}, a \rangle = \langle \pi_* w, a \rangle,$$

where $\pi : T^*\mathcal{U}_0 \ni (q, p) \mapsto q \in \mathcal{U}_0$, and

$$\langle \nu_\infty, a \rangle = \lim_{R \rightarrow +\infty} \lim_{h \rightarrow 0^+} \langle \nu_h^R, a \rangle.$$

REMARK 3.2. — Again we implicitly consider sequences $R_n \rightarrow +\infty$ (say 2^n) but we just write $R \rightarrow +\infty$ for simplicity.

3.2. REDUCTION TO THE CONES $C_\varepsilon(\bar{\mathcal{U}}_0)$. — We next introduce a further cut-off restricting the measure ν_h to a conic region containing the semiclassical wave-front set of the sequence $(u_h)_{h \rightarrow 0^+}$ in the subelliptic regime $1 \ll |H_1| \lesssim h^{-1}$. We set, for $0 < \varepsilon < 1$,

$$\chi_\varepsilon^C := \chi\left(\frac{\varepsilon H_1}{\sqrt{H_2^2 + H_3^2 + 1}}\right), \quad \tilde{\chi}_\varepsilon^C = 1 - \chi_\varepsilon^C.$$

Before including these cut-offs in our analysis of the sequence ν_h^R , we show the following result:

LEMMA 3.3. — *For every $0 < \varepsilon < 1$, the symbols χ_ε^C and $\tilde{\chi}_\varepsilon^C$ belong to the admissible class of symbols $S_{\text{cl}}^0(T^*(\mathbb{R}^2 \times \mathbb{S}^1))$ defined in Appendix B. Moreover, for every $a \in \mathcal{C}_c^\infty(\mathcal{U}_0)$, the function*

$$\frac{a \chi_\varepsilon^C \tilde{\chi}_R^B}{1 + H_2^2 + H_3^2}$$

belongs to the class of symbols $S_{\text{cl}}^{-2}(T^(\mathbb{R}^2 \times \mathbb{S}^1))$.*

REMARK 3.4. — The corresponding seminorms of χ_ε^C and $\tilde{\chi}_\varepsilon^C$ in $S_{\text{cl}}^0(T^*(\mathbb{R}^2 \times \mathbb{S}^1))$ depend on ε .

Proof. — From the definition of χ_ε^C and $\tilde{\chi}_\varepsilon^C$, it is sufficient to verify that all the derivatives of

$$g := \frac{H_1}{\sqrt{H_2^2 + H_3^2 + 1}}$$

are bounded (with some further decay for the derivatives with respect to (ξ, η, ζ)) in the region where $\varepsilon g \in \text{supp}(\chi')$, i.e.,

$$(22) \quad \frac{1}{\varepsilon} \sqrt{1 + H_2^2 + H_3^2} \leq |H_1| \leq \frac{2}{\varepsilon} \sqrt{1 + H_2^2 + H_3^2}.$$

Thanks to (12), (13), (14) and (15), we can verify by induction that, for every $\alpha \in \mathbb{Z}_+^3$,

$$\partial_{xyz}^\alpha g = \sum_{j=0}^{|\alpha|} \frac{P_{j,\alpha}(H_1, H_2, H_3)}{(1 + H_2^2 + H_3^2)^{j+1/2}},$$

where, for every $0 \leq j \leq |\alpha|$, $(u, v) \mapsto P_{j,\alpha}(u, v)$ is a polynomial of degree $\leq 2j + 1$. In particular, using (22), all these derivatives are bounded (with a constant depending on ε). It now remains to deal with the derivatives with respect to (ξ, η, ζ) . To this aim, we recall that H_1, H_2 and H_3 are linear functions in these variables. Arguing by induction as for the derivatives with respect to (x, y, z) , we can then conclude that, for every $(\alpha, \beta) \in \mathbb{Z}_+^6$, $\partial_{xyz}^\alpha \partial_{\xi\eta\zeta}^\beta g$ is uniformly bounded by $C_\varepsilon (1 + H_2^2 + H_3^2)^{-|\beta|/2}$ under the assumption (22). Using the upper bound in (22), we deduce that this is bounded by $C_\varepsilon (1 + H_1^2 + H_2^2 + H_3^2)^{-|\alpha|/2}$, for some slightly larger constant $C_\varepsilon > 0$. This concludes the first part of the proof of the lemma thanks to (10). For the second part, we can remark that $a\chi_\varepsilon^C \tilde{\chi}_R^B$ belongs to S_{cl}^0 and that $\varepsilon^2 H_1^2 \leq 4(H_2^2 + H_3^2 + 1)$ on the support of χ_ε^C . Hence, we can argue as for g above to deduce that $(1 + H_2^2 + H_3^2)^{-1}$ belongs to S^{-2} on the support of $a\chi_\varepsilon^C \tilde{\chi}_R^B$ which concludes the second part of the lemma. \square

The next step of our analysis consists of inserting these two cutoffs in the construction of ν_h^R . This produces the splitting:

$$\langle \nu_h^R, a \rangle = \langle \text{Op}_h^w(a\tilde{\chi}_R^B \chi_\varepsilon^C) u_h, u_h \rangle_{L^2} + \langle \text{Op}_h^w(a\tilde{\chi}_R^B \tilde{\chi}_\varepsilon^C) u_h, u_h \rangle_{L^2},$$

and we can introduce the object of interest for our analysis:

$$(23) \quad \langle \nu_h^{R,\varepsilon}, a \rangle := \langle \text{Op}_h^w(a\tilde{\chi}_R^B \tilde{\chi}_\varepsilon^C) u_h, u_h \rangle_{L^2}.$$

We are in position to prove the following lemma:

LEMMA 3.5. — *With the above conventions, one has, for every $0 < \varepsilon < 1$,*

$$\langle \nu_\infty, a \rangle = \lim_{R \rightarrow +\infty} \lim_{h \rightarrow 0^+} \langle \nu_h^{R,\varepsilon}, a \rangle.$$

Proof. — From Lemma 3.3, the composition rules for pseudodifferential operators and the Calderón-Vaillancourt theorem, one has

$$\text{Op}_h^w(a\chi_\varepsilon^C \tilde{\chi}_R^B) = \text{Op}_h^w\left(\frac{a\chi_\varepsilon^C \tilde{\chi}_R^B}{1 + H_2^2 + H_3^2}\right) \text{Op}_h^w(1 + H_2^2 + H_3^2) + \mathcal{O}_{L^2 \rightarrow L^2}(h).$$

Combining this composition rule with (18) and (19), one obtains the estimate

$$|\langle \text{Op}_h^w(a\tilde{\chi}_R^B \chi_\varepsilon^C) u_h, u_h \rangle_{L^2}| \leq \left\| \text{Op}_h^w\left(\frac{a\chi_\varepsilon^C \tilde{\chi}_R^B}{1 + H_2^2 + H_3^2}\right) \right\|_{L^2 \rightarrow L^2} + \mathcal{O}_{\varepsilon,R}(h).$$

Moreover, by construction of our cutoff functions and by using Calderón-Vaillancourt theorem one more time, one gets

$$|\langle \text{Op}_h^w(a \tilde{\chi}_R^B \tilde{\chi}_\varepsilon^C) u_h, u_h \rangle_{L^2}| \leq \frac{C_{M,g,a}}{R\varepsilon^2} + \mathcal{O}_{\varepsilon,R}(h^{1/2}).$$

Indeed, the Calderón-Vaillancourt theorem states the existence of a constant $C > 0$ such that $\|\text{Op}_h^w(b)\|_{L^2 \rightarrow L^2} \leq C \|b\|_{L^\infty} + \mathcal{O}_b(h^{1/2})$ and this can be applied to

$$|b| = \left| \frac{a \chi_\varepsilon^C \tilde{\chi}_R^B}{1 + H_2^2 + H_3^2} \right| \leq \frac{2\|a\|_{L^\infty}}{(\varepsilon^2 H_1^2/4) + H_2^2 + H_3^2} \leq \frac{8\|a\|_{L^\infty}}{R\varepsilon^2}.$$

Hence, one ends up with

$$\langle \nu_h^R, a \rangle = \langle \text{Op}_h^w(a \tilde{\chi}_R^B \tilde{\chi}_\varepsilon^C) u_h, u_h \rangle_{L^2} + \mathcal{O}((R\varepsilon^2)^{-1}) + \mathcal{O}_{\varepsilon,R}(h^{1/2}),$$

which concludes the proof. □

REMARK 3.6. — Note that, so far, the parameter $\varepsilon > 0$ does not play any particular role. However, it will become important when analyzing the invariance and support properties of ν_∞ where we will need to take the limit $\varepsilon \rightarrow 0$.

Finally, we record the following useful lemma that follows from the proof of Lemma 3.3:

LEMMA 3.7. — *Let \mathcal{K} be a compact subset of $\mathbb{R}^2 \times \mathbb{S}^1$. For every $0 < \varepsilon \leq 1$, for every $N_0 \geq 2$, and for every $(\alpha, \beta) \in \mathbb{Z}_+^6$, one can find a positive constant $C_{\varepsilon, N_0, \mathcal{K}, \alpha, \beta}$ such that, for every $(q, p) = (x, y, z, \xi, \eta, \zeta)$ in $C_{2N_0\varepsilon}(\mathcal{K}) \setminus C_{2^{-N_0}\varepsilon}(\mathcal{K})$, one has*

$$\left| \partial_q^\alpha \partial_p^\beta \left(\frac{H_1}{\sqrt{1 + H_2^2 + H_3^2}} \right) \right| \leq C_{\varepsilon, N_0, \mathcal{K}, \alpha, \beta} \langle p \rangle^{-|\beta|},$$

where $\langle p \rangle := (1 + \xi^2 + \eta^2 + \zeta^2)^{1/2}$.

In other words, the function $H_1/\sqrt{1 + H_2^2 + H_3^2}$ belongs to an amenable class of symbols inside the “cone” $C_{2N_0\varepsilon}(\mathcal{K}) \setminus C_{2^{-N_0}\varepsilon}(\mathcal{K})$.

3.3. REDUCTION TO THE REGION $1 \ll |H_1| \lesssim h^{-1}$. — We will now localize the phase-space distribution of the sequence (u_h) in the sub-elliptic region $1 \ll |H_1| \lesssim h^{-1}$. To do that, we introduce the cutoff functions, for $R_1 > 1$,

$$\rho_{R_1} := \tilde{\chi}(hH_1/R_1), \quad \text{and} \quad \tilde{\rho}_{R_1} := 1 - \rho_{R_1}.$$

The cut-off $\tilde{\rho}_{R_1}$ localizes the sequence (u_h) in the region of interest to us. Let us first show that this last localization keeps the analysis in the admissible symbol class.

LEMMA 3.8. — *Let $a \in \mathcal{C}_c^\infty(\mathcal{U}_0)$. The functions $a \tilde{\chi}_R^B \tilde{\chi}_\varepsilon^C \rho_{R_1}$ and $a \tilde{\chi}_R^B \tilde{\chi}_\varepsilon^C \tilde{\rho}_{R_1}$ belong to the admissible symbol class $S_{\text{cl}}^0(T^*(\mathbb{R}^2 \times \mathbb{S}^1))$ with seminorms that are uniformly bounded for $0 < h \leq 1$.*

Proof. — We already know from Lemma 3.3 that $a\tilde{\chi}_\varepsilon^C$ belongs to $S_{\text{cl}}^0(T^*(\mathbb{R}^2 \times \mathbb{S}^1))$ and we have observed that $\tilde{\chi}_R^B$ belongs to $S_{\text{cl}}^0(T^*(\mathbb{R}^2 \times \mathbb{S}^1))$. Thus, we need to show that $\rho_{R_1}(hH_1)$ belongs to this class when restricted to the region of phase space given by the support of $a\tilde{\chi}_\varepsilon^C \tilde{\chi}_R^B$. Among other constraints, in this region we have that (x, y, z) in \mathcal{U}_0 and

$$(24) \quad \frac{\varepsilon|H_1|}{\sqrt{1 + H_2^2 + H_3^2}} \geq 1.$$

In other words, we need to show that the derivatives of hH_1 verify the properties of the class $S_{\text{cl}}^0(\mathbb{R}^2 \times \mathbb{S}^1)$ under these support properties and the additional assumption that $\rho'_{R_1}(hH_1) \neq 0$, which leads to the additional constraint

$$(25) \quad R_1 \leq h|H_1| \leq 2R_1.$$

In view of (12), (13), (14) and (15), one can verify that all the derivatives of ρ_{R_1} of order ℓ with respect (x, y, z) are linear combinations of functions of the form

$$P(hH_1, hH_2, hH_3)\chi^{(k)}(hH_1/R_1),$$

where P is a polynomial of degree k with coefficients in $\mathcal{C}^\infty(\mathbb{R}^2 \times \mathbb{S}^1)$ and with $k \leq \ell$. In particular, thanks to the support properties (24) and (25), these quantities are bounded as expected. It now remains to differentiate these quantities with respect to (ξ, η, ζ) . As H_1, H_2, H_3 are polynomials of degree 1 in (ξ, η, ζ) , it has the effect to lower the degree of the polynomial and to get a bound of order $h^{\ell'}$ where ℓ' is the number of derivatives with respect to these variables. Using (24) and (25) one more time together with (10), this yields the expected decaying properties of the class $S_{\text{cl}}^0(T^*(\mathbb{R}^2 \times \mathbb{S}^1))$ with constants that are independent of $0 < h \leq 1$. \square

We next include these cutoff functions in (23). The goal is to verify that the contribution of the term

$$\langle \text{Op}_h^w(a\tilde{\chi}_\varepsilon^C \tilde{\chi}_R^B \tilde{\chi}(hH_1/R_1)) u_h, u_h \rangle_{L^2}$$

is small as $R_1 \rightarrow +\infty$. Notice, to this aim, that on the support of $a\tilde{\chi}_\varepsilon^C \tilde{\chi}_R^B$, the function $1/H_1$ belongs to the class of symbols $S_{\text{cl}}^0(T^*(\mathbb{R}^2 \times \mathbb{S}^1))$. Hence, by the composition rules for pseudodifferential operators, one has

$$\begin{aligned} & \langle \text{Op}_h^w(a\tilde{\chi}_\varepsilon^C \tilde{\chi}_R^B \tilde{\chi}(hH_1/R_1)) u_h, u_h \rangle_{L^2} \\ &= \left\langle \text{Op}_h^w\left(\frac{a\tilde{\chi}_\varepsilon^C \tilde{\chi}_R^B \tilde{\chi}(hH_1/R_1)}{hH_1}\right) \text{Op}_h^w(hH_1) u_h, u_h \right\rangle_{L^2} + \mathcal{O}_{R, R_1, \varepsilon}(h). \end{aligned}$$

Using then the a priori estimate (19) together with the Calderón-Vaillancourt theorem, we find

$$\langle \text{Op}_h^w(a\tilde{\chi}_\varepsilon^C \tilde{\chi}_R^B \tilde{\chi}(hH_1/R_1)) u_h, u_h \rangle_{L^2} = \mathcal{O}(R_1^{-1}) + \mathcal{O}_{R, R_1, \varepsilon}(h).$$

Therefore, by another application of pseudodifferential calculus rules, we get finally that

$$(26) \quad \langle \nu_h^{R, \varepsilon}, a \rangle = \langle \text{Op}_h^w(a\tilde{\rho}_{R_1}(hH_1)\tilde{\chi}_\varepsilon^C \tilde{\chi}_R^B) u_h, u_h \rangle_{L^2(\mathcal{U}_0)} + \mathcal{O}(R_1^{-1}) + \mathcal{O}_{R, R_1, \varepsilon}(h).$$

3.4. **ADDING A NEW VARIABLE $E = hH_1$.** — To complete the preliminaries concerning the phase-space localization of the measure $\nu_h^{R,\varepsilon}$, we lift slightly our analysis by introducing more general distributions in terms of a new variable hH_1 for the symbols a considered above. Namely, we set

$$(27) \quad \mu_h^{R,\varepsilon} : b \in \mathcal{C}_c^\infty(\mathcal{U}_0 \times \mathbb{R}) \longmapsto \langle \text{Op}_h^w(b(x, y, z, hH_1)) \tilde{\chi}_R^B \tilde{\chi}_\varepsilon^C u_h \rangle_{L^2(\mathcal{U}_0)}.$$

LEMMA 3.9. — *The sequence $(\mu_h^{R,\varepsilon})_{h \rightarrow 0^+}$ is bounded in $\mathcal{D}'(\mathcal{U}_0 \times \mathbb{R})$. Moreover, any accumulation point as $h \rightarrow 0^+$ is a finite nonnegative measure and there exists a constant $C > 0$ (independent of R, ε) such that any accumulation point $\mu^{R,\varepsilon}$ satisfies*

$$\forall b \in \mathcal{C}_c^\infty(\mathcal{U}_0 \times \mathbb{R}), \quad |\langle \mu^{R,\varepsilon}, b \rangle| \leq C \|b\|_{\mathcal{C}^0}.$$

There exists $\varepsilon_0 > 0$ such that, for every $0 < \varepsilon < \varepsilon_0$, any accumulation point μ^ε of the sequence $(\mu^{R,\varepsilon})_{R \rightarrow +\infty}$ verifies

$$\forall a \in \mathcal{C}_c^\infty(\mathcal{U}_0), \quad \langle \nu_\infty, a \rangle = \langle \mu^\varepsilon, a \rangle.$$

From this lemma, we can infer that our analysis reduces to the description of the measure μ^ε . Finally, up to another extraction as $\varepsilon \rightarrow 0^+$, we can suppose that μ^ε converges to some (finite) Radon measure on $\mathcal{U}_0 \times \mathbb{R}$, i.e.,

$$(28) \quad \forall b \in \mathcal{C}_c^\infty(\mathcal{U}_0 \times \mathbb{R}), \quad \langle \mu_\infty, b \rangle = \lim_{\varepsilon \rightarrow 0^+} \lim_{R \rightarrow +\infty} \lim_{h \rightarrow 0^+} \langle \nu_h^{R,\varepsilon}, b \rangle.$$

From (29), one has

$$\forall a \in \mathcal{C}_c^0(\mathcal{U}_0), \quad \int_{\mathcal{U}_0} a(q) d\nu_\infty(q) = \int_{\mathcal{U}_0 \times \mathbb{R}} a(q) d\mu_\infty(q, E),$$

and our analysis thus boils down to the properties of the extended measure μ_∞ .

REMARK 3.10. — As explained in Remark 3.6, the fact that we take $\varepsilon \rightarrow 0^+$ is not important so far but will turn to be later on.

Proof of Lemma 3.9. — The same argument as in the proof of Lemma 3.8 shows that the symbol $b(x, y, z, hH_1) \tilde{\chi}_\varepsilon^C \tilde{\chi}_R^B$ belongs to $S_{\text{cl}}^0(T^*(\mathbb{R}^2 \times \mathbb{S}^1))$. In particular, from the Calderón-Vaillancourt theorem, we find that

$$\forall b \in \mathcal{C}_c^\infty(\mathcal{U}_0 \times \mathbb{R}), \quad |\langle \mu_h^{R,\varepsilon}, b \rangle| \leq C \|b\|_{\mathcal{C}^0} + \mathcal{O}_{R,\varepsilon}(h^{1/2}).$$

Hence, this defines a bounded sequence in $\mathcal{D}'(\mathcal{U}_0 \times \mathbb{R})$. Thus, up to another extraction, we may suppose that $\mu_h^{R,\varepsilon}$ converges (for the weak- \star topology) to some distribution $\mu^{R,\varepsilon}$. Thanks to the Gårding inequality (65), this is a positive distribution, thus a finite measure. Moreover, since the sequence (ψ_h) is normalized (and hence (u_h) is bounded), this defines a finite measure. Up to another extraction, we can suppose that $\mu^{R,\varepsilon}$ weakly converges to some limit measure μ^ε as $R \rightarrow +\infty$. Coming back to (26), we find that

$$(29) \quad \langle \nu_\infty, a \rangle = \lim_{R \rightarrow +\infty} \lim_{h \rightarrow 0^+} \langle \nu_h^{R,\varepsilon}, a \rangle = \langle \mu^\varepsilon, a \tilde{\rho}_{R_1}(E) \rangle + \mathcal{O}(R_1^{-1}).$$

Applying the dominated convergence theorem, one finds that, for all $\varepsilon > 0$ (small enough),

$$\forall a \in \mathcal{C}_c^\infty(\mathcal{U}_0), \quad \langle \nu_\infty, a \rangle = \langle \mu^\varepsilon, a \rangle. \quad \square$$

3.5. RELATION WITH 2-MICROLOCAL DEFECT MEASURES. — Another way to understand the reductions in this paragraph would have been to start with the larger class of test functions $a \in S_{\text{cl}}^0(T^*(\mathbb{R}^2 \times \mathbb{S}^1))$ and to define the generalized Wigner distribution

$$W_h : a \in S_{\text{cl}}^0(T^*(\mathbb{R}^2 \times \mathbb{S}^1)) \longmapsto \langle \text{Op}_h^w(a)u_h, u_h \rangle_{L^2}.$$

Compared with w_h which only consider test functions in $\mathcal{C}_c^\infty(T^*(\mathbb{R}^2 \times \mathbb{S}^1))$, this point of view would permit to deal with the escape of mass at infinity and to define a limit measure W supported on the compactified space

$$\bar{T}^*(\mathbb{R}^2 \times \mathbb{S}^1) = (T^*(\mathbb{R}^2 \times \mathbb{S}^1)) \cup (\mathbb{R}^2 \times \mathbb{S}^1 \times \mathbb{S}^3) \simeq (T^*(\mathbb{R}^2 \times \mathbb{S}^1)) \cup (S^*(\mathbb{R}^2 \times \mathbb{S}^1)),$$

where the 3-sphere corresponds to the sphere at infinity in the cotangent variable. This would result into a semiclassical defect measure carried on

$$\{H_2^2 + H_3^2 = 1\} \cup (\mathbb{R}^2 \times \mathbb{S}^1 \times \{\pm(1, 0, 0)\}),$$

where the points $\pm(1, 0, 0)$ would correspond to the direction $\pm H_1$ on the sphere at infinity. The measure at infinity can thus be identified with a measure on the configuration space $\mathbb{R}^2 \times \mathbb{S}^1$ (up to the two connected components) and this is the reason why we only worked with test functions in $\mathcal{C}_c^\infty(\mathcal{U}_0)$. The role of Lemma 3.5 and of the various cutoff functions therein is exactly to isolate this part of the limit measure W without introducing test functions $a \in S_{\text{cl}}^0(T^*(\mathbb{R}^2 \times \mathbb{S}^1))$ while in the end only functions in $\mathcal{C}_c^\infty(\mathcal{U}_0)$ will be needed.

Once this reduction to this part of phase space is done, Section 3.4 is intended to analyze in more depth this part of the measure formally carried by these two “points” at infinity by making some kind of blow-up procedure of these points through the introduction of the rescaled variable $E = hH_1$. This last part of our analysis is reminiscent of what is done when defining two-microlocal defect measures as in [FK95, Mil96, Nie96, FK00, AM14]. Yet, we emphasize that, compared with these references, one major simplification occurs as we did not introduce the full cotangent variables. This results into the fact that the corresponding test functions, namely $b(x, y, z, hH_1)\tilde{\chi}_R^B\tilde{\chi}_\varepsilon^C$, lies in a nice class of symbols amenable to the standard arguments on scalar semiclassical defect measures. Introducing the full cotangent variables would in principle require to deal with more exotic class of symbols and with operator valued symbols as in the above works. Such a point of view was for instance taken in the Euclidean setting of Grushin operators [AS23]. In this reference, the authors introduced operator-valued measures lifting the analogue of the measure ν_∞ and described completely these lifted objects involving the full cotangent variables. Here, we only focus on the behaviour along the variables $(q, hH_1(q, p))$. Introducing the full cotangent variables in our Riemannian setting would require some extra and delicate work (especially when dealing with coordinate charts in the critical regime $h|H_1| \asymp 1$) that is not necessary to prove the results we are aiming at.

4. SUPPORT OF THE LIMIT MEASURE

Before describing the propagation and invariance properties of ν_∞ , we discuss first the support properties of μ_∞ along the new variable $E \in \mathbb{R}$. More precisely, the goal of this section is to prove the following:

PROPOSITION 4.1. — *The measure μ_∞ defined in (28) decomposes as*

$$\mu_\infty(q, E) = \bar{\mu}_\infty(q, E) + \sum_{k=0}^{\infty} (\mu_{k,\infty}^+(q, E) + \mu_{k,\infty}^-(q, E)),$$

where $\bar{\mu}_\infty$ and $(\mu_{k,\infty}^\pm)_{k \geq 0}$ are finite non-negative Radon measures on $\mathcal{M} \times \mathbb{R}$ satisfying the following concentration properties:

(S.1) $\text{supp } \bar{\mu}_\infty \subset \mathcal{M}_{\Lambda_0, W} \times \{0\}$;

(S.2) for every $k \in \mathbb{Z}_+$, $\text{supp } \mu_{k,\infty}^\pm \subset \mathcal{H}_\pm^{-1}(2k + 1) \subset \mathcal{U}_{\Lambda_0, W} \times \mathbb{R}_\pm^*$.

Recall that $\mathcal{M}_{\Lambda_0, W} := \{\Lambda_0 - W \geq 0\}$ is the classical allowed region, $\mathcal{U}_{\Lambda_0, W} = \{\Lambda_0 - W > 0\}$, and that \mathcal{H}_\pm was defined in (7). As we shall see in the proof, our arguments are reminiscent of what one does when computing the spectrum of the 1-dimensional harmonic oscillator [Zwo12, Ch. 6], i.e., by using creation and annihilation type operators and by playing with their commutation properties. Recall that, given an hypoelliptic operator that can be written as a sum of squares, there is always a local underlying Lie algebra model [RS76]. Here, due to the commutation relations (1), this underlying structure is the one of the 3-dimensional Heisenberg group for which the harmonic oscillator naturally appears when performing Fourier analysis on this group [FKF21]. We do not explicitly use this local algebraic structure but this is in some sense the mechanism at work in the upcoming proof of Proposition 4.1.

4.1. PRELIMINARY LEMMAS. — We first define the following creation and annihilation type operators (ladder operators):

$$A_h := \text{Op}_h^w(H_2 + iH_3) \quad \text{and} \quad A_h^* := \text{Op}_h^w(H_2 - iH_3),$$

so that

$$(30) \quad \text{Op}_h^w(H_2^2 + H_3^2) = A_h^* A_h + h \text{Op}_h^w(H_1) + h^2 c_0 = A_h A_h^* - h \text{Op}_h^w(H_1) + h^2 c_0,$$

where c_0 is a smooth compactly supported and real-valued function on $\mathbb{R}^2 \times \mathbb{S}^1$ (recall that $\lambda \equiv 0$ outside a compact set containing the isothermal neighborhood U_0) which is independent of h . Notice that, by (11) and by the composition rule for the Weyl quantization,

$$(31) \quad [A_h, A_h^*] = 2h \text{Op}_h^w(H_1).$$

We begin with the following lemma:

LEMMA 4.2. — *Let $k \geq 1$. Then*

$$A_h^k = \text{Op}_h^w \left((H_2 + iH_3)^k + \sum_{j=1}^k P_{j,k,h}(hH_1, hH_2, hH_3) (H_2 + iH_3)^{k-j} \right),$$

where $P_{j,k,h}(u, v, w)$ is a polynomial with coefficients depending polynomially on h and smoothly on $(x, y, z) \in \mathbb{R}^2 \times \mathbb{S}^1$ (with uniformly bounded derivatives).

In fact, modulo some extra work, we could be slightly more precise on the nature of the polynomials as we know that the full symbol is a polynomial of degree k in the cotangent variables (ξ, η, ζ) . Yet, as it is not necessary for our analysis, we do not try to be more precise and we just keep track of the information that is relevant for our proofs.

Proof. — Recall that

$$A_h = \text{Op}_h^w(H_2 + iH_3) = \frac{h}{i}X_\perp + hV + ihX_\perp(\lambda).$$

In particular, A_h^k is a differential operator of order k for every $k \geq 1$, and its symbol is polynomial of degree $\leq k$ in the cotangent variables (ξ, η, ζ) . We now proceed by induction and suppose that the lemma is true for a given $k \geq 1$. Using the composition rule from Theorem B.1, we can write

$$\begin{aligned} A_h^{k+1} &= \text{Op}_h^w\left((H_2 + iH_3)^k + \sum_{j=1}^k P_{j,k,h}(hH_1, hH_2, hH_3)(H_2 + iH_3)^{k-j}\right)A_h \\ &= \text{Op}_h^w\left((H_2 + iH_3)^{k+1} + \sum_{j=1}^k P_{j,k,h}(hH_1, hH_2, hH_3)(H_2 + iH_3)^{k+1-j}\right) \\ &\quad + \text{Op}_h^w(R_k(h)), \end{aligned}$$

where

$$R_k(h) = \sum_{\ell=1}^{k+1} \sum_{j=1}^k \frac{h^\ell}{\ell!} A(D)^\ell \left((P_{j,k,h}(hH_1, hH_2, hH_3)(H_2 + iH_3)^{k-j})(H_2 + iH_3) \right).$$

Here the sum stops at $\ell = k + 1$ as each symbol is a polynomial of respective degree k and 1 in the (ξ, η, ζ) variables. Recall that $A(D) = \frac{1}{2i}(\partial_{p_1} \cdot \partial_{q_2} - \partial_{q_1} \cdot \partial_{p_2})$ so that the symbols of interest for the remainder take the form

$$A_\ell(D) \left((P_{j,k,h}(hH_1, hH_2, hH_3)(H_2 + iH_3)^{k-j})(H_2 + iH_3) \right),$$

with $A_\ell(D) = (\partial_{p_1} \cdot \partial_{q_2})^\ell - \ell(\partial_{q_1} \cdot \partial_{p_2})(\partial_{p_1} \cdot \partial_{q_2})^{\ell-1}$. To see this, we observe that $H_2 + iH_3$ is of degree 1 in (ξ, η, ζ) so that ∂_{p_2} can occur at most once in the expansion of $A(D)^\ell$. By induction, we get the expected expression for the terms of order h^ℓ in the asymptotic expansion. \square

As a corollary of this lemma and of the composition rule for pseudodifferential operators, we also find:

COROLLARY 4.3. — *Let $k \geq 1$. Then*

$$\text{Op}_h^w((H_2 + iH_3)^k) = A_h^k + \sum_{j=1}^k \text{Op}_h^w(\tilde{P}_{j,k,h}(hH_1, hH_2, hH_3))A_h^{k-j},$$

where $\tilde{P}_{j,k,h}(u, v, w)$ is a polynomial with coefficients depending polynomially on h and smoothly on $(x, y, z) \in \mathbb{R}^2 \times \mathbb{S}^1$ (with uniformly bounded derivatives).

We now turn to the commutation properties of A_h^k with the operators of interest for our analysis:

LEMMA 4.4. — *Let Q and W be two smooth functions on $\mathbb{R}^2 \times \mathbb{S}^1$ whose derivatives are uniformly bounded. Then, for every $k \geq 1$,*

$$[A_h^k, \text{Op}_h^w(hQH_1)] = h \sum_{j=0}^{k-1} \text{Op}_h^w(P_{j,k,h}(hH_1, hH_2, hH_3))A_h^{k-1-j},$$

and

$$[A_h^k, \text{Op}_h^w(W)] = h \sum_{j=0}^{k-1} \text{Op}_h^w(\tilde{P}_{j,k,h}(hH_1, hH_2, hH_3))A_h^{k-1-j},$$

where $P_{j,k,h}(u, v, w)$ and $\tilde{P}_{j,k,h}(u, v, w)$ are polynomials whose coefficients depend polynomially on h and smoothly on $(x, y, z) \in \mathbb{R}^2 \times \mathbb{S}^1$ (with derivatives that are uniformly bounded).

Proof. — First, we observe that, thanks to Lemma 4.2, one has

$$A_h^k = \text{Op}_h^w\left((H_2 + iH_3)^k + \sum_{j=1}^k P_{j,k,h}(hH_1, hH_2, hH_3)(H_2 + iH_3)^{k-j}\right),$$

where $P_{j,k,h}$ are polynomials verifying the properties of the present lemma. The second bracket formula is then a direct consequence of the composition rule for pseudodifferential operators (see Theorem B.1) together with Corollary 4.3. In fact, since W depends only on (x, y, z) , the terms in the asymptotic expansion will only involves derivatives of the symbol of A_h^k with respect to the variables (ξ, η, ζ) .

We now turn to the first bracket which can be rewritten as

$$[A_h^k, \text{Op}_h^w(hQH_1)] = \sum_{j=0}^k [\text{Op}_h^w(P_{j,k,h}(hH_1, hH_2, hH_3)(H_2 + iH_3)^{k-j}), \text{Op}_h^w(QhH_1)],$$

with $P_{0,k,h} = 1$. Given $0 \leq j \leq k$, one can apply the composition rule from Theorem B.1 to each term, i.e.,

$$\begin{aligned} & [\text{Op}_h^w(P_{j,k,h}(hH_1, hH_2, hH_3)(H_2 + iH_3)^{k-j}), \text{Op}_h^w(QhH_1)] \\ &= 2 \sum_{0 \leq 2\ell \leq k} \frac{h^{2\ell+1}}{(2\ell + 1)!} \text{Op}_h^w(A(D)^{2\ell+1}(*)), \end{aligned}$$

with $(*)$ being given by $(P_{j,k,h}(hH_1, hH_2, hH_3)(H_2 + iH_3)^{k-j})(QhH_1)$. Here, the sum over ℓ is bounded as we are only considering polynomials symbols in the variables (ξ, η, ζ) (with the total degree being bounded by $k + 1$). Recalling the exact expression of $A(D)$ from Theorem B.1 and Corollary 4.3 (together with several applications of the composition formula), we find the expected result. \square

REMARK 4.5. — Similar statements as those of Lemmas 4.2, 4.4 and Corollary 4.3 hold for A_h^* replacing A_h .

4.2. **INDUCTIVE ARGUMENT: PROOF OF PROPOSITION 4.1.** — In order to prove the localization properties (S.1) and (S.2) for the measure μ_∞ , we start from the following semiclassical estimates. For every $k \geq 0$,

$$(32) \quad \left\langle \text{Op}_h^w(b(x, y, z, hH_1) \widetilde{\chi}_R^B \widetilde{\chi}_\varepsilon^C)^2 A_h^k (\widehat{P}_{h,\lambda} - \Lambda_h) u_h, A_h^k u_h \right\rangle = \mathcal{O}(h^\infty),$$

$$(33) \quad \left\langle \text{Op}_h^w(b(x, y, z, hH_1) \widetilde{\chi}_R^B \widetilde{\chi}_\varepsilon^C)^2 (A_h^*)^k (\widehat{P}_{h,\lambda} - \Lambda_h) u_h, (A_h^*)^k u_h \right\rangle = \mathcal{O}(h^\infty).$$

We note that the remainder is $\mathcal{O}(h^\infty)$ (and not 0) because the eigenvalue equation (18) is only local. Using estimates (32) and (33) together with (30) and the symbolic calculus developed in Section 4.1, we aim at deriving inductively suitable concentration properties for the distributions $\varrho_{k,\varepsilon,R,h}^\pm$ defined by

$$\begin{aligned} \varrho_{k,\varepsilon,R,h}^+ : b &\longmapsto \left\langle \text{Op}_h^w(b(x, y, z, hH_1) \widetilde{\chi}_R^B \widetilde{\chi}_\varepsilon^C) A_h^k u_h, A_h^k u_h \right\rangle, \\ \varrho_{k,\varepsilon,R,h}^- : b &\longmapsto \left\langle \text{Op}_h^w(b(x, y, z, hH_1) \widetilde{\chi}_R^B \widetilde{\chi}_\varepsilon^C) (A_h^*)^k u_h, (A_h^*)^k u_h \right\rangle. \end{aligned}$$

REMARK 4.6. — Note that, in order to make sense of the limit measures for $k \geq 1$, one needs to have an a priori upper bound on

$$\left\| \text{Op}_h^w(b(x, y, z, hH_1) \widetilde{\chi}_R^B \widetilde{\chi}_\varepsilon^C) A_h^k u_h \right\|,$$

which will be part of the argument below. For $k = 1$, such an upper bound follows for instance from the a priori estimate (19) but, for $k \geq 2$, this does not longer work immediately.

We will show in Lemma 4.9 that, up to additional extractions, the weak limits of these distributions are well defined as non-negative Radon measures:

$$\langle \varrho_k^\pm, b \rangle := \lim_{\varepsilon \rightarrow 0^+} \lim_{R \rightarrow +\infty} \lim_{h \rightarrow 0} \langle \varrho_{k,\varepsilon,R,h}^\pm, b \rangle,$$

and we will deduce from these measures the desired support properties of μ_∞ . For the sake of exposition, we start with the first step $k = 0$ which is slightly easier to handle:

LEMMA 4.7. — *The measure $\mu_\infty = \varrho_0^\pm$ satisfies:*

$$\text{supp}(\mu_\infty) \subset \left\{ (q, E) \in \mathcal{M} \times \mathbb{R} : -\frac{\Lambda_0 - W}{1 - Q} \leq E \leq \frac{\Lambda_0 - W}{1 + Q} \right\}.$$

In particular, the support of the measure μ_∞ is compact in the E variable and disjoint with the classical forbidden region $\{W > \Lambda_0\}$. Moreover,

$$\text{supp}(\mu_\infty) \setminus \mathcal{H}_-^{-1}(1) = \text{supp} \varrho_1^- \quad \text{and} \quad \text{supp}(\mu_\infty) \setminus \mathcal{H}_+^{-1}(1) = \text{supp} \varrho_1^+$$

where we recall that $\mathcal{H}_\pm(q, E) = \pm E^{-1}(\Lambda_0 - W - EQ)$.

REMARK 4.8. — Recall also from Appendix A that condition $\|Q\|_{\mathcal{C}^0} < 1$ was initially imposed to ensure the hypoellipticity (and the semiboundedness) of the operator \widehat{P}_h .

Proof. — Given $b \in \mathcal{C}_c^\infty(\mathcal{U}_0 \times \mathbb{R})$, one has

$$(34) \quad \langle \text{Op}_h^w(b(x, y, z, hH_1)) \widetilde{\chi}_R^B \widetilde{\chi}_\varepsilon^C (\widehat{P}_{h,\lambda} - \Lambda_h) u_h, u_h \rangle = \mathcal{O}(h^\infty).$$

Recalling that

$$(35) \quad \text{Op}_h^w(H_2^2 + H_3^2) = A_h^* A_h + h \text{Op}_h^w(H_1) + h^2 c_0,$$

one can use the composition rule for pseudodifferential operators together with the a priori estimate (19). This yields

$$\langle \text{Op}_h^w(b(x, y, z, hH_1))(hH_1 + hQH_1 + W - \Lambda_h) \widetilde{\chi}_R^B \widetilde{\chi}_\varepsilon^C u_h, u_h \rangle = -\langle \varrho_{1,\varepsilon,R,h}^+, b \rangle + \mathcal{O}(h).$$

Thanks to (19) and to the Calderón-Vaillancourt theorem, the right-hand side defines a bounded sequence in $\mathcal{D}'(\mathcal{U}_0 \times \mathbb{R})$. Moreover, the Gårding inequality (65) ensures that the limit distribution is a nonnegative Radon measure. Hence, letting $h \rightarrow 0^+$ and $R \rightarrow +\infty$ (in this order), one finds that, for every b compactly supported in $\mathcal{U}_0 \times \mathbb{R}$,

$$\mu^\varepsilon(b(x, y, z, E)(E(1 + Q) + W - \Lambda_0)) = -\varrho_{1,\varepsilon}^+(b).$$

From this, one infers that, on the support of μ^ε , $(1 + Q)E + W - \Lambda_0 \leq 0$, and moreover that

$$(36) \quad \text{supp}(\mu^\varepsilon) \setminus \mathcal{H}_+^{-1}(1) = \text{supp} \varrho_{1,\varepsilon}^+.$$

Similarly, using now the identity

$$\text{Op}_h^w(H_2^2 + H_3^2) = A_h A_h^* - h \text{Op}_h^w(H_1) + h^2 c_0$$

instead of (35), and using again (34), one finds

$$\langle \text{Op}_h^w(b(x, y, z, hH_1))(-hH_1 + hQH_1 + W - \Lambda_h) \widetilde{\chi}_R^B \widetilde{\chi}_\varepsilon^C u_h, u_h \rangle = -\langle \varrho_{1,\varepsilon,h}^-, b \rangle + \mathcal{O}(h),$$

and thus

$$\mu^\varepsilon(b(x, y, z, E)(-E(1 - Q) + W - \Lambda_0)) = -\varrho_{1,\varepsilon}^-(b).$$

This implies that $-E(1 - Q) + W - \Lambda_0 \leq 0$ on the support of μ^ε and moreover that

$$(37) \quad \text{supp}(\mu^\varepsilon) \setminus \mathcal{H}_-^{-1}(1) = \text{supp} \varrho_{1,\varepsilon}^-.$$

Putting together (36) and (37) and, letting $\varepsilon \rightarrow 0^+$ this concludes the proof. \square

We now turn to the general case for which we cannot make use of the a priori estimate (19) directly:

LEMMA 4.9. — *For every $k \geq 0$ and for every $R > 1$ and $\varepsilon > 0$, the family*

$$(\varrho_{k,R,\varepsilon,h}^\pm)_{0 < h \leq h_0}$$

is bounded in $\mathcal{D}'(\mathcal{U}_0 \times \mathbb{R})$ and any accumulation point (as $h \rightarrow 0^+$) $\varrho_{k,R,\varepsilon}^\pm$ is a finite nonnegative Radon measure. Moreover, $(\varrho_{k,R,\varepsilon}^\pm)_{R,\varepsilon}$ is bounded and any accumulation point as $R \rightarrow +\infty$ and $\varepsilon \rightarrow 0^+$ (in this order) is a finite nonnegative Radon measure ϱ_k^\pm verifying for every $k \geq 0$

$$(\pm E(2k + 1 \pm Q) + W - \Lambda_0) \varrho_k^\pm = -\varrho_{k+1}^\pm.$$

In particular, one can deduce from this lemma that

$$\text{supp } \varrho_k^\pm \subset \{(q, E) \in \mathcal{M} \times \mathbb{R} : \pm E(2k + 1 \pm Q) + W - \Lambda_0 \leq 0\},$$

that, for every $k \geq 0$, $\text{supp } \varrho_{k+1}^\pm \subset \text{supp } \varrho_k^\pm$ and that

$$\text{supp } \varrho_k^\pm \setminus \text{supp } \varrho_{k+1}^\pm \subset \mathcal{H}_\pm^{-1}(2k + 1).$$

Proof. — Recall first that the limits $R \rightarrow \infty$ and $\varepsilon \rightarrow 0^+$ are understood along subsequences $(R_n)_{n \geq 1}$ and $(\varepsilon_m)_{m \geq 1}$. For simplicity of exposition, we now fix them to be of the form⁽⁴⁾ $R_n = 2^n$ and $\varepsilon_m = 2^{-m}$. The case $k = 0$ follows by Lemma 4.7. Assume that the claim holds for every $0 \leq j \leq k$ and moreover that the following a priori estimates hold, for any $b \in \mathcal{C}_c^\infty(\mathcal{U}_0 \times \mathbb{R})$ and for any $h > 0$ small enough,

$$(38) \quad \left\| \text{Op}_h^w(b(x, y, z, hH_1) \tilde{\chi}_{2^j}^B \tilde{\chi}_{2^{k-j} 2^{-m}}^C) A_h^j u_h \right\|_{L^2} \leq C(\|b\|_\infty + \mathcal{O}_{n,m,b}(h^{1/2})),$$

for $0 \leq j \leq k$, for $n, m \geq k + 1 - j$, and for some constant $C > 0$ that is independent of n and m and that depends on the support of b (and also on W and Q). Moreover, the constant in the remainder depends on a finite number of derivatives of b . Let us prove the claim together with (38) for $k + 1$. One more time, we will keep the notations $R \rightarrow +\infty$ and $\varepsilon \rightarrow 0^+$ to alleviate notations.

We begin by proving the same a priori estimate on

$$\left\| \text{Op}_h^w(b(x, y, z, hH_1) \tilde{\chi}_R^B \tilde{\chi}_\varepsilon^C) A_h^{k+1} u_h \right\|_{L^2},$$

which is the main technical point of the analysis. To do that, we begin with equality (32) which can be expanded as follows

$$\begin{aligned} & \left\langle \text{Op}_h^w(b(x, y, z, hH_1) \tilde{\chi}_R^B \tilde{\chi}_\varepsilon^C)^2 A_h^k \text{Op}_h^w(r_h) u_h, A_h^k u_h \right\rangle \\ &= - \left\langle \text{Op}_h^w(b(x, y, z, hH_1) \tilde{\chi}_R^B \tilde{\chi}_\varepsilon^C)^2 A_h^k A_h^* A_h u_h, A_h^k u_h \right\rangle + \mathcal{O}(h^\infty), \end{aligned}$$

with $r_h := hH_1 + hQH_1 + W + h^2(W_{1,\lambda} + c_0) - \Lambda_h$. Applying Lemma 4.4 together with (31), we find that the right-hand side can be rewritten as

$$\begin{aligned} & \left\langle \text{Op}_h^w(b(x, y, z, hH_1) \tilde{\chi}_R^B \tilde{\chi}_\varepsilon^C)^2 A_h^k A_h^* A_h u_h, A_h^k u_h \right\rangle \\ &= \left\langle \text{Op}_h^w(b(x, y, z, hH_1) \tilde{\chi}_R^B \tilde{\chi}_\varepsilon^C)^2 A_h^{k-1} A_h^* A_h^2 u_h, A_h^k u_h \right\rangle \\ & \quad + \left\langle \text{Op}_h^w(b(x, y, z, hH_1) \tilde{\chi}_R^B \tilde{\chi}_\varepsilon^C)^2 \text{Op}_h^w(2hH_1) A_h^k u_h, A_h^k u_h \right\rangle \\ & \quad + h \sum_{j=0}^{k-2} \left\langle \text{Op}_h^w(b(x, y, z, hH_1) \tilde{\chi}_R^B \tilde{\chi}_\varepsilon^C)^2 \text{Op}_h^w(*) A_h^{k-1-j} u_h, A_h^k u_h \right\rangle, \end{aligned}$$

with $(*)$ given by $\mathbf{P}_{j,k-1,h}(hH_1, hH_2, hH_3)$. Applying the composition formula for pseudodifferential operators together with the Calderón-Vaillancourt theorem, one finds that each term in the sum over $0 \leq j \leq k - 2$ is of the form

$$\left\langle \text{Op}_h^w(b_{h,R,\varepsilon} \tilde{\chi}_{R/2}^B \tilde{\chi}_{2\varepsilon}^C) A_h^{k-1-j} u_h, A_h^k u_h \right\rangle + \mathcal{O}(h),$$

⁽⁴⁾Other sequences can be dealt along similar lines.

where $b_{h,R,\varepsilon}$ depends on j and belongs to $S_{\text{cl}}^0(T^*(\mathbb{R}^2 \times \mathbb{S}^1))$ with all its seminorms uniformly bounded in terms of $0 < h \leq 1$ (but not R, ε a priori) and with compact support in \mathcal{U}_0 . This follows from the fact that b is initially compactly supported in the variable $E = hH_1$ and that $H_2^2 + H_3^2 \ll H_1^2$ thanks to our cutoff functions. Hence, combining this observation with (38) and with elliptic estimates for pseudodifferential operators as in [DZ19, Th. E.33], we can deduce that each term in the sum over $0 \leq j \leq k - 2$ is uniformly bounded in terms of $0 < h \leq 1$ so that we get

$$\begin{aligned} & \langle \text{Op}_h^w(b(x, y, z, hH_1)\tilde{\chi}_R^B\tilde{\chi}_\varepsilon^C)^2 A_h^k \text{Op}_h^w(\tilde{r}_h)u_h, A_h^k u_h \rangle \\ &= - \langle \text{Op}_h^w(b(x, y, z, hH_1)\tilde{\chi}_R^B\tilde{\chi}_\varepsilon^C)^2 A_h^{k-1} A_h^* A_h^2 u_h, A_h^k u_h \rangle + \mathcal{O}_{R,\varepsilon}(h), \end{aligned}$$

with $\tilde{r}_h := 3hH_1 + hQH_1 + W + h^2(W_{1,\lambda} + c_0) - \Lambda_h$. Iterating this commutation argument k times, we find that

$$\begin{aligned} & \langle \text{Op}_h^w(b(x, y, z, hH_1)\tilde{\chi}_R^B\tilde{\chi}_\varepsilon^C)^2 \text{Op}_h^w((2k+1)hH_1 + hQH_1 + W - \Lambda_h)A_h^k u_h, A_h^k u_h \rangle \\ &= - \langle \text{Op}_h^w(b(x, y, z, hH_1)\tilde{\chi}_R^B\tilde{\chi}_\varepsilon^C)^2 A_h^* A_h^{k+1} u_h, A_h^k u_h \rangle + \mathcal{O}_{R,\varepsilon}(h). \end{aligned}$$

Applying the composition rule for pseudodifferential operators together with the induction hypothesis, we find that

$$\begin{aligned} & \langle \text{Op}_h^w((b(x, y, z, hH_1)\tilde{\chi}_R^B\tilde{\chi}_\varepsilon^C)^2((2k+1)hH_1 + hQH_1 + W - \Lambda_h)A_h^k u_h, A_h^k u_h) \rangle \\ &= - \|\text{Op}_h^w(b(x, y, z, hH_1)\tilde{\chi}_R^B\tilde{\chi}_\varepsilon^C)A_h^{k+1} u_h\|^2 + \mathcal{O}_{R,\varepsilon}(h). \end{aligned}$$

By induction, the left-hand side is bounded from which we deduce the expected upper bound at step $k + 1$.

We note that the upper bound (38) allows to verify by induction that any accumulation point $\varrho_{k,\varepsilon,R}^\pm$ (as $h \rightarrow 0^+$) is a finite nonnegative Radon measure whose total mass is independent of ε and R . We can thus take the limit $R \rightarrow +\infty$ and $\varepsilon \rightarrow 0^+$ in this order. We obtain the expected limit measure ϱ_{k+1}^\pm and we can now derive its support properties. Repeating the same argument with $\text{Op}_h^w(b(x, y, z, hH_1)\tilde{\chi}_R^B\tilde{\chi}_\varepsilon^C)$ instead of $\text{Op}_h^w(b(x, y, z, hH_1)\tilde{\chi}_R^B\tilde{\chi}_\varepsilon^C)^2$ (and with $(A_h^*)^k$ instead of A_h^k) and taking the limits $h \rightarrow 0^+$ and $R \rightarrow +\infty$ (in this order), we can prove that

$$(39) \quad \varrho_k^\pm(b(x, y, z, E)(\pm E(2k+1 \pm Q) + W - \Lambda_0)) = -\varrho_{k+1}^\pm(b).$$

Finally, from (39), we get the first statement of the lemma and moreover:

$$\text{supp } \varrho_k^\pm \setminus \mathcal{H}_\pm^{-1}(2k+1) = \text{supp } \varrho_{k+1}^\pm.$$

This concludes the proof. □

Finally, Lemma 4.9 implies that:

$$\text{supp } \mu_\infty \subset (\mathcal{M}_{\Lambda_0, W} \times \{0\}) \cup \bigcup_{k=0}^\infty (\mathcal{H}_+^{-1}(2k+1) \cup \mathcal{H}_-^{-1}(2k+1)).$$

Defining, for $k \in \mathbb{Z}_+$,

$$\mu_{k,\infty}^\pm := \mathbf{1}_{\mathcal{H}_\pm^{-1}(2k+1)} \mu_\infty, \quad \text{and} \quad \bar{\mu}_\infty := \mathbf{1}_{\mathcal{M}_{\Lambda_0, W} \times \{0\}} \mu_\infty,$$

we obtain the proof of Proposition 4.1.

5. NORMAL FORM REDUCTION

In order to study the invariance properties of the measure μ_∞ , we will require a normal form procedure in the subelliptic regime $1 \ll |H_1| \lesssim h^{-1}$. This will allow us to work with functions that are adapted to the geometry of the problem. Roughly speaking, it amounts to work in a system of asymptotically symplectic coordinates as $|H_1| \rightarrow \infty$. This normal form approach was pioneered in Melrose’s works [Mel85] and recently revisited in the context of sub-Riemannian Laplacians associated with 3-dimensional contact flows [CHT18, CHT21] as we are dealing here. See also [RV15] for earlier related normal forms procedure in the context of 2-dimensional magnetic semiclassical Schrödinger operators and [HKRV16, Mor22, Mor24] regarding higher dimensional normal forms for magnetic operators. More precisely, our normal form expansion reads

THEOREM 5.1. — *One can find $P \in \mathcal{P}_3(\mathbb{R}^2 \times \mathbb{S}^1)/\mathcal{P}_1(\mathbb{R}^2 \times \mathbb{S}^1)$ and $(R_j)_{1 \leq j \leq 3}$ in $\mathcal{P}(\mathbb{R}^2 \times \mathbb{S}^1)/\mathcal{P}_1(\mathbb{R}^2 \times \mathbb{S}^1)$ such that, in the local chart \mathcal{U}_0 , one has, for $H_1 \neq 0$,*

$$\{H_2^2 + H_3^2, H_1(1 + P)\} = H_2^2 R_1 + H_3^2 R_2 + H_2 H_3 R_3.$$

Moreover, for every $a \in \mathcal{C}_c^\infty(\mathcal{U}_0)$, one can find $P_a \in \mathcal{P}_2(\mathbb{R}^2 \times \mathbb{S}^1)/\mathcal{P}_0(\mathbb{R}^2 \times \mathbb{S}^1)$ and $R_a \in \mathcal{P}(\mathbb{R}^2 \times \mathbb{S}^1)/\mathcal{P}_1(\mathbb{R}^2 \times \mathbb{S}^1)$ such that, in the local chart \mathcal{U}_0 , one has, for $H_1 \neq 0$,

$$\{H_2^2 + H_3^2, a + P_a\} = \frac{H_2^2 + H_3^2}{H_1} X(a) + \frac{R_a}{H_1}.$$

Recall that $\mathcal{P}_N(\mathbb{R}^2 \times \mathbb{S}^1)$ and $\mathcal{P}(\mathbb{R}^2 \times \mathbb{S}^1)$ were defined in Lemma 2.5 and consist of functions of the form $\sum_{|\alpha| \leq N} b_\alpha (H_2/H_1)^{\alpha_2} (H_3/H_1)^{\alpha_3}$ with b_α depending only in the (x, y, z) variables. The first part of this result states that $\mathbf{H}_1 = H_1(1 + P)$ is a small deformation H_1 (in the sense that $P = \mathcal{O}(\varepsilon^2)$ in the region $|H_2| |H_3| \lesssim \varepsilon |H_1|$ where eigenmodes are microlocalized) whose Poisson bracket with $H_2^2 + H_3^2$ is of order $\mathcal{O}(\varepsilon^2)$ in the region of interest. This error is modulo quadratic terms in H_2 and H_3 but these should be understood as bounded terms in the region where our eigenmodes are microlocalized thanks to (19). Similarly $\mathbf{a} = a + P_a$ is a small deformation of a whose Poisson bracket with $H_2^2 + H_3^2$ has a nice asymptotic expansion.

As already alluded, such a theorem is obtained through a normal form procedure. Compared with [CHT18], we will simplify some aspects of this normal form procedure. Rather than making a symplectic change of variables at each step of the iterative scheme and using the machinery of Fourier integral operators, we will just encode a slightly simpler change of variables (close to but not necessarily symplectic) into a small deformation of the test function

$$\mathbf{b} = b + \mathcal{O}\left(\frac{|H_2| + |H_3|}{|H_1|}\right).$$

As already explained, this deformation makes the remainder term in $\{H_2^2 + H_3^2, \mathbf{b}\}$ as small as possible in the regime $H_2^2 + H_3^2 \ll H_1^2$. This simplified version of the normal form method is in fact sufficient to obtain the desired invariance properties

of the semiclassical measure $\mu_\infty = \lim_{\varepsilon \rightarrow 0} \mu^\varepsilon$. We emphasize that Wigner type distributions enjoy somehow more flexibility regarding change of variables than Fourier integral operators (since many negligible terms disappear in the limit $h \rightarrow 0$), and we will crucially exploit this fact to avoid the use of cumbersome symplectic changes of coordinates that in the end match with ours in the semiclassical limit (see [AS23, §3.1] for a related construction involving two-microlocal semiclassical measures).

REMARK 5.2. — In this section, the fact that we have simple bracket formulas as (11) will make the proof slightly simpler and more explicit regarding the terms appearing in the normal form. Yet, the strategy would remain the same if $\{H_1, H_2\}$ and $\{H_1, H_3\}$ were more general linear combinations of H_1, H_2 and H_3 (as in the general contact case). In any case, the fact that these brackets do not identically vanish is exactly where the situation is more involved than in the flat Heisenberg case [FKF21, FKL21] like when considering varying magnetic fields rather than constant ones.

5.1. APPROXIMATE SOLUTIONS OF COHOMOLOGICAL EQUATIONS. — We first recall the algebraic relations given by (11) and producing the subelliptic structure of our problem:

$$(40) \quad \{H_1, H_2\} = -KH_3, \quad \{H_1, H_3\} = H_2, \quad \{H_2, H_3\} = -H_1.$$

It is convenient to introduce complex notations:

$$Z := H_2 + iH_3 \quad \text{and} \quad \bar{Z} := H_2 - iH_3,$$

so that

$$H_2^2 + H_3^2 = |Z|^2.$$

The relations (40) now become

$$\{H_1, Z\} = iK_+Z + iK_-\bar{Z}, \quad \{H_1, \bar{Z}\} = -iK_-Z - iK_+\bar{Z}, \quad \{Z, \bar{Z}\} = 2iH_1,$$

where

$$K_+ := \frac{1 + K}{2} \quad \text{and} \quad K_- = \frac{1 - K}{2}.$$

In particular, one has

$$(41) \quad \{|Z|^2, H_1\} = iK_-(Z^2 - \bar{Z}^2).$$

In view of proving Theorem 5.1, we will have to modify the function of interest (either a or H_1) through an inductive scheme that is achieved by solving *cohomological equations* of the form

$$(42) \quad \{|Z|^2, f\} = \frac{b_\alpha(x, y, z)}{H_1^{\alpha_1}} Z^{\alpha_2} \bar{Z}^{\alpha_3},$$

where $(\alpha_1, \alpha_2, \alpha_3) \in \mathbb{Z}_+^3$ and the unknown of the equation is f . The key observation to solve (approximately) this kind of equation is that, for every $k \neq \ell$ in \mathbb{Z}_+ , one has

$$(43) \quad \left\{ |Z|^2, Z^k \bar{Z}^\ell / 2i(\ell - k) \right\} = H_1 Z^k \bar{Z}^\ell.$$

In particular, by a direct computation, one has the following approximate solution to the cohomological equation (42):

LEMMA 5.3. — For $\alpha_2 \neq \alpha_3$, set

$$f := \frac{b_\alpha}{2i(\alpha_3 - \alpha_2)H_1^{\alpha_1+1}} Z^{\alpha_2} \bar{Z}^{\alpha_3}.$$

Then, one has

$$\begin{aligned} \{|Z|^2, f\} &= \frac{b_\alpha(x, y, z)}{H_1^{\alpha_1}} Z^{\alpha_2} \bar{Z}^{\alpha_3} - iK_- \frac{b_\alpha(x, y, z)}{H_1^{\alpha_1+2}} Z^{\alpha_2} \bar{Z}^{\alpha_3} (Z^2 - \bar{Z}^2) \\ &\quad + \frac{\{\bar{Z}, b_\alpha\}}{H_1^{\alpha_1+1}} Z^{\alpha_2+1} \bar{Z}^{\alpha_3} + \frac{\{Z, b_\alpha\}}{H_1^{\alpha_1+1}} Z^{\alpha_2} \bar{Z}^{\alpha_3+1}. \end{aligned}$$

Here, we have not exactly solved (42) but only up to smaller order term in the region where our eigenmodes are microlocalized (roughly speaking $|Z| \lesssim 1$ and $|H_1| \rightarrow +\infty$). Such an observation will be enough to construct inductively the functions P and P_a appearing in Theorem 5.1.

REMARK 5.4. — In the rest of this Section, we will always suppose that we work in the region $H_1 \neq 0$. In the end, the formulas will only be used in the regime $|H_2|, |H_3| \ll |H_1|$.

5.2. A SMALL DEFORMATION OF H_1 . — We start with the deformation of the variable H_1 and prove the first part of Theorem 5.1.

5.2.1. Normal form procedure. — In view of (41) and of Lemma 5.3, we set

$$P_2 := \frac{K_-}{2} ((H_2/H_1)^2 - (H_3/H_1)^2).$$

By construction, one has

$$\{|Z|^2, H_1(1 + P_2)\} = \frac{(Z^2 + \bar{Z}^2)}{4} \{|Z|^2, K_-/H_1\}.$$

Iterating this procedure by applying Lemma 5.3 one more time, one can find P_3 in $\mathcal{P}(\mathbb{R}^2 \times \mathbb{S}^1)$ of the form

$$P_3 = \sum_{|\alpha|=3} P_{3,\alpha}(x, y, z) (H_2/H_1)^{\alpha_2} (H_3/H_1)^{\alpha_3}$$

and such that, if we set

$$(44) \quad \mathbf{H}_1 := H_1 (1 + P_2 + P_3),$$

then one has

$$(45) \quad \{H_2^2 + H_3^2, \mathbf{H}_1\} = H_2^2 R_1 + H_3^2 R_2 + H_2 H_3 R_3,$$

with R_j belonging in $\mathcal{P}(\mathbb{R}^2 \times \mathbb{S}^1)$ and being of the form

$$R_j = \sum_{|\alpha| \geq 2} R_{j,\alpha}(x, y, z) (H_2/H_1)^{\alpha_2} (H_3/H_1)^{\alpha_3}.$$

Hence, in the regime $H_2^2 + H_3^2 \ll H_1^2$, this Poisson bracket is somehow of smaller order than the one appearing in (41). This concludes the first part of the proof of Theorem 5.1.

We finally state the following analogue of Lemma 3.7:

LEMMA 5.5. — *Let \mathcal{K} be a compact subset of $\mathbb{R}^2 \times \mathbb{S}^1$. For every $0 < \varepsilon \leq 1$, for every $N_0 \geq 2$ and for every $(\alpha, \beta) \in \mathbb{Z}_+^6$, one can find a constant $C_{\varepsilon, N_0, \mathcal{K}, \alpha, \beta}$ such that, for every $(q, p) = (x, y, z, \xi, \eta, \zeta)$ in $C_{2N_0\varepsilon}(\mathcal{K}) \setminus C_{2-N_0\varepsilon}(\mathcal{K})$, one has, for every $j \in \{2, 3\}$,*

$$\left| \partial_q^\alpha \partial_p^\beta \left(\frac{\mathbf{H}_1}{\sqrt{1 + H_2^2 + H_3^2}} \right) \right| \leq C_{\varepsilon, N_0, \mathcal{K}, \alpha, \beta} \langle p \rangle^{-|\beta|},$$

where $\langle p \rangle := (1 + \xi^2 + \eta^2 + \zeta^2)^{1/2}$.

Proof. — This is a direct consequence of Corollary 2.6 and Lemma 3.7. □

5.2.2. *Rewriting $\mu_h^{R, \varepsilon}$ using \mathbf{H}_1 .* — Before proving the last part of Theorem 5.1, we briefly come back to the distribution $\mu_h^{R, \varepsilon}$ that was introduced in (27) and defined using H_1 . We next show that we can replace H_1 by \mathbf{H}_1 modulo small remainders in h and ε .

LEMMA 5.6. — *For b in $\mathcal{C}_c^\infty(\mathcal{U}_0 \times \mathbb{R})$, we set*

$$(46) \quad \langle \mu_h^{R, \varepsilon}, b \rangle := \langle \text{Op}_h(b(x, y, z, h\mathbf{H}_1) \tilde{\chi}_\varepsilon^C \tilde{\chi}_\varepsilon^C \tilde{\chi}_R^B \tilde{\chi}_R^B) u_h, u_h \rangle_{L^2},$$

where

$$\tilde{\chi}_R^B := \tilde{\chi} \left(\frac{\mathbf{H}_1^2 + H_2^2 + H_3^2}{4R} \right), \quad \text{and} \quad \tilde{\chi}_\varepsilon^C := \tilde{\chi} \left(\frac{\varepsilon \mathbf{H}_1}{4\sqrt{H_2^2 + H_3^2 + 1}} \right).$$

Then, one has, as $h \rightarrow 0^+$, $R \rightarrow +\infty$ and $\varepsilon \rightarrow 0$ (in this order),

$$\langle \mu_h^{R, \varepsilon}, b \rangle = \langle \mu_h^{R, \varepsilon}, b \rangle + \mathcal{O}_{b, \varepsilon, R}(h) + \mathcal{O}_{b, \varepsilon}(R^{-1}) + \mathcal{O}_b(\varepsilon^2).$$

REMARK 5.7. — Note that, on the support of $\tilde{\chi}_\varepsilon^C$, one has $\mathbf{H}_1 = H_1(1 + \mathcal{O}(\varepsilon^2))$, so that $\tilde{\chi}_\varepsilon^C = 1$ on the support of $\tilde{\chi}_\varepsilon^C \tilde{\chi}_\varepsilon^C$ if $\varepsilon > 0$ is chosen small enough. We just keep the function $\tilde{\chi}_\varepsilon^C$ to ensure that \mathbf{H}_1 is well defined. The same holds for $\tilde{\chi}_R^B \tilde{\chi}_R^B$.

Proof. — Let $b(x, y, z, E)$ be an element in $\mathcal{C}_c^\infty(\mathcal{U}_0 \times \mathbb{R}^*)$. One has

$$\langle \mu_h^{R, \varepsilon}, b \rangle = \langle \text{Op}_h(b(x, y, z, h\mathbf{H}_1) \tilde{\chi}_\varepsilon^C \tilde{\chi}_R^B) u_h, u_h \rangle_{L^2},$$

where $\tilde{\chi}_\varepsilon^C$ was defined in (21) as

$$\tilde{\chi}_\varepsilon^C := \tilde{\chi} \left(\frac{\varepsilon H_1}{\sqrt{H_2^2 + H_3^2 + 1}} \right).$$

Arguing as in the proof of Lemma 3.3, one knows that

$$\tilde{\chi}_\varepsilon^C \tilde{\chi} \left(\frac{\varepsilon \mathbf{H}_1}{4\sqrt{H_2^2 + H_3^2 + 1}} \right) b(x, y, z, h\mathbf{H}_1)$$

belongs to $S_{\text{cl}}^0(T^*(\mathbb{R}^2 \times \mathbb{S}^1))$ with all seminorms uniformly bounded in terms of h such that $0 < h \leq 1$ (but not on ε a priori).

Set now $\chi_\varepsilon^C = 1 - \tilde{\chi}_\varepsilon^C$. On the support of $b(x, y, z, h\mathbf{H}_1) \tilde{\chi}_\varepsilon^C \chi_\varepsilon^C$, one has

$$1 \leq \frac{\varepsilon |H_1|}{\sqrt{1 + H_2^2 + H_3^2}} \leq 10,$$

for small enough $\varepsilon > 0$. In particular, we can argue as in (20) and write

$$b(x, y, z, hH_1) \tilde{\chi}_R^B \tilde{\chi}_\varepsilon^C \chi_\varepsilon^C = \frac{b(x, y, z, hH_1) \tilde{\chi}_R^B \tilde{\chi}_\varepsilon^C \chi_\varepsilon^C}{1 + H_2^2 + H_3^2} (1 + H_2^2 + H_3^2),$$

where the first term of the product on the right-hand side belongs to S_{cl}^{-2} thanks to the above support properties. Using the Calderón-Vaillancourt theorem together with (18) and (19), we find that

$$\langle \mu_h^{R,\varepsilon}, b \rangle = \langle \text{Op}_h(b(x, y, z, hH_1) \tilde{\chi}_\varepsilon^C \tilde{\chi}_\varepsilon^C \tilde{\chi}_R^B) u_h, u_h \rangle_{L^2} + \mathcal{O}_{R,\varepsilon}(h) + \mathcal{O}_\varepsilon(R^{-1}).$$

Similarly, we can insert the cutoff function $\tilde{\chi}_R^B$ if we note that, on the support of $\chi_R^B \tilde{\chi}_R^B$, one has

$$\frac{1}{10}R \leq H_1^2 + H_2^2 + H_3^2 \leq 10R,$$

hence the involved symbol is compactly supported. We can then apply the exact same argument and show that

$$\langle \mu_h^{R,\varepsilon}, b \rangle = \langle \text{Op}_h(b(x, y, z, hH_1) \tilde{\chi}_\varepsilon^C \tilde{\chi}_\varepsilon^C \tilde{\chi}_R^B \tilde{\chi}_R^B) u_h, u_h \rangle_{L^2} + \mathcal{O}_{R,\varepsilon}(h) + \mathcal{O}_\varepsilon(R^{-1}).$$

We are now left with replacing H_1 by \mathbf{H}_1 in the last component of b . We write

$$b(x, y, z, hH_1) = b(x, y, z, h\mathbf{H}_1) + h(H_1 - \mathbf{H}_1) \int_0^1 \partial_E b(x, y, z, h\mathbf{H}_1 + th(H_1 - \mathbf{H}_1)) dt.$$

Hence, applying Corollary 2.6 together with (19) and (44), the composition rule for pseudodifferential operators and the Calderón-Vaillancourt theorem, one finds the expected conclusion. \square

5.3. END OF THE PROOF OF THEOREM 5.1. — We now proceed similarly and introduce a small deformation of a whose Poisson bracket with $H_2^2 + H_3^2$ is small in the regime $H_2^2 + H_3^2 \ll H_1^2$. Let a be a smooth function compactly supported on \mathcal{U}_0 , we write

$$\{|Z|^2, a\} = Z\{\bar{Z}, a\} + \bar{Z}\{Z, a\}.$$

In the region $H_1 \neq 0$ and in view of (43), we can set

$$a_1 := \frac{Z}{2iH_1} \{\bar{Z}, a\} - \frac{\bar{Z}}{2iH_1} \{Z, a\}.$$

We then find

$$\{|Z|^2, a + a_1\} = \frac{Z}{2i} \left\{ |Z|^2, \{\bar{Z}, a\} / H_1 \right\} - \frac{\bar{Z}}{2i} \left\{ |Z|^2, \{Z, a\} / H_1 \right\}.$$

Recalling that a is a function on \mathcal{U}_0 , one has

$$\begin{aligned} \{Z, \{\bar{Z}, a\}\} - \{\bar{Z}, \{Z, a\}\} &= (X_\perp + iV)(X_\perp - iV)(a) - (X_\perp - iV)(X_\perp + iV)(a) \\ &= 2i[V, X_\perp](a) = 2iX(a). \end{aligned}$$

Hence, letting X_Z (resp. $X_{\bar{Z}}$) be the (complex) vector field generated by Z (resp. \bar{Z}) the above expression can be simplified as

$$\{|Z|^2, a + a_1\} = \frac{|Z|^2}{H_1} X(a) + \frac{(Z^2 X_{\bar{Z}}^2 - \bar{Z}^2 X_Z^2)(a)}{2iH_1} - \frac{Z^2 X_{\bar{Z}}(a)}{2iH_1^2} \{\bar{Z}, H_1\} + \frac{\bar{Z}^2 X_Z(a)}{2iH_1^2} \{Z, H_1\}.$$

We would now like to eliminate the terms of magnitude $1/H_1$ using Lemma 5.3. Namely, we set

$$a_2 = -\frac{(Z^2 X_{\bar{Z}}^2 + \bar{Z}^2 X_Z^2)(a)}{8H_1^2}.$$

This allows us, after defining $\mathbf{a} := a + a_1 + a_2$, to get

$$(47) \quad \{H_2^2 + H_3^2, \mathbf{a}\} = \frac{H_2^2 + H_3^2}{H_1} X(a) + \frac{1}{H_1} \mathbf{R}_a$$

where \mathbf{R}_a is an element of the form

$$\mathbf{R}_a = \sum_{|\alpha| \geq 2} R_{a,\alpha}(x, y, z) (H_2/H_1)^{\alpha_2} (H_3/H_1)^{\alpha_3},$$

with $R_{a,\alpha}$ that is compactly supported in \mathcal{U}_0 .

For simplicity of exposition, we will use more compact notations for \mathbf{a} :

$$(48) \quad \mathbf{a} = \sum_{|\alpha| \leq 2} a_\alpha (H_2/H_1)^{\alpha_2} (H_3/H_1)^{\alpha_3},$$

where the functions a_α are also defined on \mathcal{U}_0 and explicitly given by

$$(49) \quad a_{(0,0)} := a, \quad a_{(1,0)} := -V(a), \quad a_{(0,1)} := X_\perp(a),$$

and

$$(50) \quad a_{(2,0)} := -a_{(0,2)} := -\frac{(X_\perp^2 - V^2)(a)}{4}, \quad a_{(1,1)} := -\frac{(X_\perp V + V X_\perp)(a)}{2}.$$

6. INVARIANCE PROPERTIES

In view of Lemma 5.6, we are left to study the Wigner type distribution $\mu_h^{R,\varepsilon}$ given by (46). Our goal is to prove that any accumulation point μ_∞ of this sequence as $h \rightarrow 0$, $R \rightarrow +\infty$ and $\varepsilon \rightarrow 0$ (in this order) verifies certain invariance properties:

PROPOSITION 6.1. — *Let (ψ_h, Λ_h) be a sequence satisfying (3) and set*

$$\mathcal{H}_1(q, E) := \Lambda_0 - W(q) - EQ(q),$$

and

$$(51) \quad \mathcal{X}_{W,Q} := (\Lambda_0 - W)X + \Omega_{\mathcal{H}_1} + EX(\mathcal{H}_1)\partial_E.$$

Let μ_∞ be any semiclassical measure obtained as a weak limit for the sequence of distributions $\mu_h^{R,\varepsilon}$ defined from the sequence (ψ_h, Λ_h) . Then, for every $b \in \mathcal{C}_c^1(\mathcal{U}_0 \times \mathbb{R})$,

$$\int_{\mathcal{U}_0 \times \mathbb{R}} (\mathcal{X}_{W,Q}(b) + X(\mathcal{H}_1)b) d\mu_\infty = 0.$$

In order to prove Proposition 6.1, we start by fixing an element b in $\mathcal{C}_c^\infty(\mathcal{U}_0 \times \mathbb{R})$. Rather than looking at b directly, we will consider test functions based on the normal form \mathbf{b} from (48), that is,

$$(52) \quad \mathbf{b}(x, y, z, E) = \sum_{|\alpha| \leq 2} b_\alpha(x, y, z, E) (H_2/H_1)^{\alpha_2} (H_3/H_1)^{\alpha_3},$$

where b_α is given by (49) and (50) with b instead of a (the variable E plays the role of a parameter). One then has

$$\left\langle \left[\widehat{P}_{h,\lambda}, \text{Op}_h^w(\mathbf{H}_1 \mathbf{b}(\cdot, h\mathbf{H}_1) \widetilde{\chi}_\varepsilon^C \widetilde{\chi}_\varepsilon^C \widetilde{\chi}_R^B \widetilde{\chi}_R^B) \right] u_h, u_h \right\rangle = \mathcal{O}(h^\infty).$$

On the other hand, we know that $\mathbf{H}_1 \mathbf{b}(\cdot, \cdot, \cdot, h\mathbf{H}_1) \widetilde{\chi}_\varepsilon^C \widetilde{\chi}_\varepsilon^C \widetilde{\chi}_R^B \widetilde{\chi}_R^B$ belongs to S_{cl}^1 while the symbol of $\widehat{P}_{h,\lambda}$ lies in S_{cl}^2 . Then, by the composition rules for the Weyl quantization (see Appendix B), one finds

$$(53) \quad \left\langle \text{Op}_h^w(\{P_h, \mathbf{H}_1 \mathbf{b}(\cdot, h\mathbf{H}_1) \widetilde{\chi}_\varepsilon^C \widetilde{\chi}_\varepsilon^C \widetilde{\chi}_R^B \widetilde{\chi}_R^B\}) u_h, u_h \right\rangle = \mathcal{O}(h^2),$$

where $P_h = H_2^2 + H_3^2 + hQH_1 + W$. Recall that, as before, the measure in the L^2 scalar product is the standard Lebesgue measure thanks to our conventions for $\widehat{P}_{h,\lambda}$ and u_h (see (16), (17) and (18)).

6.1. REMOVING THE DERIVATIVES OF THE CUTOFF FUNCTION. — Before exploiting the normal form procedure, we start with the following Lemma in view of removing the cutoff functions $\widetilde{\chi}_\varepsilon^C \widetilde{\chi}_\varepsilon^C \widetilde{\chi}_R^B \widetilde{\chi}_R^B$ from the Poisson bracket:

LEMMA 6.2. — *With the above conventions, one has:*

$$\left\langle \text{Op}_h^w(\{P_h, \mathbf{H}_1 \mathbf{b}(\cdot, h\mathbf{H}_1)\} \chi_{R,\varepsilon}) u_h, u_h \right\rangle_{L^2(\text{Leb})} = \mathcal{O}(\varepsilon^2) + \mathcal{O}_{R,\varepsilon}(h^{1/2}),$$

where $\chi_{R,\varepsilon} := \widetilde{\chi}_\varepsilon^C \widetilde{\chi}_\varepsilon^C \widetilde{\chi}_R^B \widetilde{\chi}_R^B$.

Proof. — In light of (53), proving this equality amounts to show that

$$\left\langle \text{Op}_h^w(\mathbf{H}_1 \mathbf{b}(\cdot, h\mathbf{H}_1) \{H_2^2 + H_3^2, \chi_{R,\varepsilon}\}) u_h, u_h \right\rangle_{L^2(\text{Leb})} = \mathcal{O}(\varepsilon^2) + \mathcal{O}_{R,\varepsilon}(h^{1/2}).$$

To do that, we first write

$$\begin{aligned} \mathbf{H}_1 \mathbf{b}(\cdot, h\mathbf{H}_1) \{H_2^2 + H_3^2, \chi_{R,\varepsilon}\} &= \mathbf{H}_1 \mathbf{b}(\cdot, h\mathbf{H}_1) \{H_2^2 + H_3^2, \mathbf{H}_1\} \widetilde{\chi}_\varepsilon^C \widetilde{\chi}_R^B \\ &\times \left(\frac{\mathbf{H}_1 \widetilde{\chi}_\varepsilon^C}{2R} \widetilde{\chi}' \left(\frac{H_1^2 + H_2^2 + H_3^2}{4R} \right) + \frac{\varepsilon \widetilde{\chi}_R^B}{4\sqrt{1 + H_2^2 + H_3^2}} \widetilde{\chi}' \left(\frac{\varepsilon \mathbf{H}_1}{4\sqrt{1 + H_2^2 + H_3^2}} \right) \right). \end{aligned}$$

The first observation is that, from Theorem 5.1, the Poisson bracket $\{H_2^2 + H_3^2, \chi_{R,\varepsilon}\}$ is of the form $R_1 H_2^2 + R_2 H_3^2 + R_3 H_2 H_3$ where R_j are elements of $\mathcal{P}(\mathbb{R}^2 \times \mathbb{S}^1)$. Thus each R_j is of order $\mathcal{O}(\varepsilon^2)$ in the region where the eigenmodes are microlocalized thanks to the cutoff function $\widetilde{\chi}_\varepsilon^C \widetilde{\chi}_R^B$. Now, we can consider the first term in the right hand side which is compactly supported in (ξ, η, ζ) and which is of the form $b_1 H_2^2 + b_2 H_3^2 + b_3 H_2 H_3$ where each b_j belongs to S_{cl}^0 and has its supremum of order $\mathcal{O}(\varepsilon^2)$ (recall that $\mathbf{H}_1^2 \leq 8R$ thanks to the support properties of $\widetilde{\chi}$). Hence, applying (19) together with the Calderón-Vaillancourt theorem, we get that the contribution of this first term to our semiclassical quantities is of size $\mathcal{O}(\varepsilon^2) + \mathcal{O}_{R,\varepsilon}(h^{1/2})$.

It now remains to deal with the contribution of the second term where one differentiates $\tilde{\chi}_R^C$. In that case, our symbol is supported in the region where

$$\frac{1}{10}\sqrt{1 + H_2^2 + H_3^2} \leq \varepsilon|H_1| \leq 10\sqrt{1 + H_2^2 + H_3^2}.$$

In that case, we can remark that we end up with terms that are of the form

$$\frac{\varepsilon \mathbf{b}(\cdot, h\mathbf{H}_1)\mathbf{H}_1}{\sqrt{1 + H_2^2 + H_3^2}} (R_1H_2^2 + R_2H_3^2 + R_3H_2H_3).$$

Using Lemma 3.7, the first term lies in the class of symbol S_{cl}^0 inside the support of our symbol (recall that we have differentiated $\tilde{\chi}_\varepsilon^C$) and is uniformly bounded in terms of ε and R (but not small a priori). Using one more time the semiclassical a priori estimates (19), we find using the composition rule together with the Calderón-Vaillancourt theorem that, up to remainders of size $\mathcal{O}_{R,\varepsilon}(h^{1/2})$, the contribution of this second term is given by terms of the form

$$\left\langle \text{Op}_h^w \left(\tilde{\chi}_\varepsilon^C \tilde{\chi}_R^B \tilde{\chi}_R^B \tilde{\chi}' \left(\frac{\varepsilon \mathbf{H}_1}{4\sqrt{1 + H_2^2 + H_3^2}} \right) \frac{\varepsilon \mathbf{b}(\cdot, h\mathbf{H}_1)\mathbf{H}_1}{\sqrt{1 + H_2^2 + H_3^2}} R_j \right) u_h, u_h \right\rangle.$$

Using the Calderón-Vaillancourt theorem one last time, we get the expected result thanks to the fact that $R_j = \mathcal{O}(\varepsilon^2)$ according to the localization properties of the function $\tilde{\chi}$. □

6.2. PROOF OF PROPOSITION 6.1. — Let $b \in \mathcal{C}_c^\infty(\mathcal{U}_0 \times \mathbb{R})$, we consider as before the small deformation $\mathbf{b}(x, y, x, E)$ of b given by (52). Thanks to Lemma 6.2, one has

$$(54) \quad \left\langle \text{Op}_h^w \left(\{P_h, \mathbf{H}_1 \mathbf{b}(\cdot, h\mathbf{H}_1)\} \tilde{\chi}_\varepsilon^C \tilde{\chi}_\varepsilon^C \tilde{\chi}_R^B \tilde{\chi}_R^B \right) u_h, u_h \right\rangle = \mathcal{O}(\varepsilon^2) + \mathcal{O}_{R,\varepsilon}(h^{1/2}).$$

Hence, we need to understand

$$\{P_h, \mathbf{H}_1 \mathbf{b}\} = \{H_2^2 + H_3^2, \mathbf{H}_1 \mathbf{b}\} + \{W, \mathbf{H}_1 \mathbf{b}\} + \{hH_1Q, \mathbf{H}_1 \mathbf{b}\}.$$

Recalling (45) and (47), one has

$$\{H_2^2 + H_3^2, \mathbf{H}_1 \mathbf{b}\} = R_1H_2^2 + R_2H_3^2 + R_3H_2H_3,$$

and

$$\begin{aligned} \{H_2^2 + H_3^2, \mathbf{b}\} &= \frac{H_2^2 + H_3^2}{H_1} X(\mathbf{b}) + \frac{1}{H_1} \mathbf{R}_b \\ &\quad + h\{H_2^2 + H_3^2, \mathbf{H}_1 \mathbf{b}\} \sum_{|\alpha| \leq 2} \partial_E b_\alpha (H_2/H_1)^{\alpha_2} (H_3/H_1)^{\alpha_3}, \end{aligned}$$

where \mathbf{R}_b and $(R_j)_{j \geq 3}$ belongs to the class of symbol S_{cl}^0 inside the support of our cutoff functions with supremum that is of order $\mathcal{O}(\varepsilon^2)$. Hence, using the eigenvalue equation (18) and the semiclassical a priori estimates (19) together with the composition rule for pseudodifferential operators and the expressions for \mathbf{b} and \mathbf{H}_1 given

in (52) and (44), equation (54) becomes

$$(55) \quad \langle \text{Op}_h^w(X(b)(\Lambda_h - W - hQH_1) \tilde{\chi}_\varepsilon^C \tilde{\chi}_\varepsilon^C \tilde{\chi}_R^B \tilde{\chi}_R^B) u_h, u_h \rangle \\ + \langle \text{Op}_h^w(\{W + hH_1Q, \mathbf{H}_1 \mathbf{b}(\cdot, h\mathbf{H}_1)\} \tilde{\chi}_\varepsilon^C \tilde{\chi}_\varepsilon^C \tilde{\chi}_R^B \tilde{\chi}_R^B) u_h, u_h \rangle = \mathcal{O}(\varepsilon^2) + \mathcal{O}_{R,\varepsilon}(h^{1/2}).$$

Hence, it remains to analyze the terms

$$\{W + hH_1Q, \mathbf{H}_1 \mathbf{b}(\cdot, h\mathbf{H}_1)\}.$$

To do that, we recall that, from the exact expressions for \mathbf{H}_1 and \mathbf{b} given in (44), (49) and (52), one has

$$\mathbf{H}_1 = H_1 \left(1 + \sum_{|\alpha| \in \{2,3\}} P_\alpha(x, y, z) (H_2/H_1)^{\alpha_2} (H_3/H_1)^{\alpha_3} \right),$$

$$\text{and } \mathbf{b}(\cdot, h\mathbf{H}_1) \mathbf{H}_1 = b(\cdot, h\mathbf{H}_1) H_1 + X_\perp(b)(\cdot, h\mathbf{H}_1) H_3 - V(b)(\cdot, h\mathbf{H}_1) H_2 \\ + \sum_{|\alpha| \geq 2} Q_\alpha(x, y, z, h\mathbf{H}_1) H_2^{\alpha_2} H_3^{\alpha_3} / H_1^{|\alpha|-1},$$

where Q_α are smooth compactly supported functions. We also observe that, in the support of our cutoff functions, one has

$$\{W + hH_1Q, h\mathbf{H}_1\} = \{W + hH_1Q, hH_1\} + \mathcal{O}(\varepsilon).$$

For similar reasons, the terms of order $|\alpha| \geq 2$ in the expression of $\mathbf{b}(\cdot, h\mathbf{H}_1) \mathbf{H}_1$ yields a contribution of order $\mathcal{O}(\varepsilon)$ and (55) can be rewritten as

$$(56) \quad \langle \text{Op}_h^w(X(b)(\Lambda_h - W - hQH_1) \tilde{\chi}_\varepsilon^C \tilde{\chi}_\varepsilon^C \tilde{\chi}_R^B \tilde{\chi}_R^B) u_h, u_h \rangle + \mathcal{O}(\varepsilon) + \mathcal{O}_{R,\varepsilon}(h^{1/2}) \\ = - \langle \text{Op}_h^w(\{W + hH_1Q, H_1b + X_\perp(b)H_3 - V(b)H_2\} \tilde{\chi}_\varepsilon^C \tilde{\chi}_\varepsilon^C \tilde{\chi}_R^B \tilde{\chi}_R^B) u_h, u_h \rangle.$$

As one has, on the support of our functions,

$$\{W, H_1b + X_\perp(b)H_3 - V(b)H_2\} \\ = -X(W)b - hH_1X(W)\partial_E b - X_\perp(b)V(W) + V(b)X_\perp(W) + \mathcal{O}(\varepsilon)$$

and

$$\{hH_1Q, H_1b + X_\perp(b)H_3 - V(b)H_2\} = -hH_1X(Q)b - (hH_1)^2X(Q)\partial_E b + hH_1QX(b) \\ - hH_1X_\perp(b)V(Q) + hH_1V(b)X_\perp(Q) + \mathcal{O}(\varepsilon),$$

we finally obtain the expected result by letting $h \rightarrow 0^+$, $R \rightarrow +\infty$ and $\varepsilon \rightarrow 0^+$ (in this order) in (56).

7. SUMMARY OF THE PROPERTIES OF μ_∞

In this short section, we summarize our description of the semiclassical measures μ_∞ obtained as weak limits of the Wigner distributions $\mu_h^{R,\varepsilon}$ given by (46) (or equivalently (28)). More precisely, as a consequence of Propositions 4.1 and 6.1, one has the following theorem:

THEOREM 7.1. — *Let $Q, W \in \mathcal{C}^\infty(\mathcal{M}, \mathbb{R})$ such that $\|Q\|_{\mathcal{C}^0} < 1$ and let $\Lambda_0 \geq \min W$. Given a sequence (ψ_h, Λ_h) satisfying (3) then any measure μ_∞ obtained from the sequence (28) decomposes as:*

$$\mu_\infty(q, E) = \bar{\mu}_\infty(q, E) + \sum_{k=0}^\infty (\mu_{k,\infty}^+(q, E) + \mu_{k,\infty}^-(q, E)),$$

where $\bar{\mu}_\infty$ and $(\mu_{k,\infty}^\pm)_{k \geq 0}$ are finite non-negative Radon measures satisfying the following concentration properties:

(S.1) $\text{supp } \bar{\mu}_\infty \subset \mathcal{M}_{\Lambda_0, W} \times \{0\}$, with $\mathcal{M}_{\Lambda_0, W} := \{q \in \mathcal{M} : \Lambda_0 - W(q) \geq 0\}$,

(S.2) for every $k \in \mathbb{Z}_+$,

$$\text{supp } \mu_{k,\infty}^\pm \subset \mathcal{H}_\pm^{-1}(2k + 1) \subset \mathcal{U}_{\Lambda_0, W} \times \mathbb{R}_\pm^*.$$

Moreover, they verify the following invariance properties:

(P.1) for every $a \in \mathcal{C}_c^1(\mathcal{U}_{\Lambda_0, W})$,

$$\int_{\mathcal{M} \times \{0\}} Y_W(a) d\bar{\mu}_\infty = 0,$$

with Y_W being defined in (5),

(P.2) for every $k \in \mathbb{N}$ and every $a \in \mathcal{C}_c^1(\mathcal{M} \times \mathbb{R}^*)$,

$$\int_{\mathcal{M} \times \mathbb{R}_\pm^*} \frac{\mathcal{X}_{W,Q}(a)}{E} d\mu_{k,\infty}^\pm = 0,$$

with $\mathcal{X}_{W,Q}$ being defined in (51),

(P.3) for every $a \in \mathcal{C}^1(\mathcal{M})$,

$$\int_{(\mathcal{M}_{\Lambda_0, W} \setminus \mathcal{U}_{\Lambda_0, W}) \times \{0\}} (\Omega_W(a) + X(W)a) d\bar{\mu}_\infty = 0.$$

Notice that, in the last item, the vector field Ω_W is “tangential” to the set $\mathcal{M}_{\Lambda_0, W} \setminus \mathcal{U}_{\Lambda_0, W}$.

Proof. — The only remaining point compared with Proposition 6.1 is to verify that the invariance properties restrict to each layer $\mathcal{H}_\pm^{-1}(2k + 1)$ and $\mathcal{M} \times \{0\}$. To see this, we first work inside $\mathcal{U}_{\Lambda_0, W} \times \mathbb{R}$ and prove properties (P.1) and (P.2). We let $k \in \mathbb{Z}_+$ and $a \in \mathcal{C}_c^\infty(\mathcal{U}_{\Lambda_0, W} \times \mathbb{R})$ whose support does not intersect $\text{supp}(\mu_\infty) \setminus \mathcal{H}_\pm^{-1}(2k + 1)$. For such a function, we deduce the expected property (P.2). If we now consider a to be an element in $\mathcal{C}_c^1(\mathcal{M} \times \mathbb{R}^*)$, then it can be split as a sum of a function of the previous form and a function that is supported away from $\mathcal{H}_\pm^{-1}(2k + 1)$. Thus, we obtain property (P.2) for the expected class of functions. Combining this with Proposition 6.1, we also find that, for every $a \in \mathcal{C}_c^1(\mathcal{U}_{\Lambda_0, W} \times \mathbb{R})$,

$$\int_{\mathcal{M} \times \{0\}} Y_W((\Lambda_0 - W)a) d\bar{\mu}_\infty = 0,$$

from which we infer (P.1). It now remains to discuss what happens on the critical set

$$\mathcal{M}_{\Lambda_0, W} \setminus \mathcal{U}_{\Lambda_0, W} := \{q \in \mathcal{M} : W(q) = \Lambda_0\}.$$

To do this, we rewrite the conclusion of Proposition 6.1 slightly more explicitly: for every $a \in \mathcal{C}_c^\infty(\mathcal{M} \times \mathbb{R})$,

$$\int_{\mathcal{M} \times \mathbb{R}} ((\Lambda_0 - W)X(a) + \Omega_{\Lambda_0 - W}(a) - E\Omega_Q(a) + EX(\mathcal{H}_1)\partial_E a + X(\mathcal{H}_1)a) d\mu_\infty = 0.$$

This expression can be rewritten as

$$\begin{aligned} & \int_{\mathcal{M} \times \{0\}} ((\Lambda_0 - W)X(a) + \Omega_{\Lambda_0 - W}(a) - X(W)a) d\mu_\infty \\ &= - \int_{\mathcal{M} \times \{E \neq 0\}} ((\Lambda_0 - W)X(a) + \Omega_{\Lambda_0 - W}(a) - E\Omega_Q(a) + EX(\mathcal{H}_1)\partial_E a + X(\mathcal{H}_1)a) d\mu_\infty. \end{aligned}$$

If we take a to be of the form $\chi(E/\delta)b(q)$ where $b \in \mathcal{C}^\infty(\mathcal{M})$ and where χ is the same cutoff function as in the previous sections (recall that $\chi(x) = x\chi'(x) = 0$ for $|x| \geq 2$), we find using the dominated convergence theorem that the right-hand-side converges to 0. Hence, for every $b \in \mathcal{C}^\infty(\mathcal{M})$,

$$\int_{\mathcal{M} \times \{0\}} ((\Lambda_0 - W)X(b) - \Omega_W(b) - X(W)b) d\bar{\mu}_\infty = 0.$$

We now take the test function b to be of the form $\chi((W(q) - \Lambda_0)/\delta)\tilde{b}(q)$ with χ a smooth cutoff function (near 0) as above. Letting $\delta \rightarrow 0$ in the previous equality, we find thanks to the dominated convergence theorem

$$\int_{(\mathcal{M}_{\Lambda_0, W} \setminus \cup_{\Lambda_0, W}) \times \{0\}} (\Omega_W(\tilde{b}) + X(W)\tilde{b}) d\bar{\mu}_\infty = 0. \quad \square$$

Note that, compared with Theorem 1.1 from the introduction, this result holds without any assumption on Λ_0 . It also involves the more general measure μ_∞ which describes precisely how H_1 escape at infinity.

Let us now explain that it directly implies Theorem 1.1. We also remark that, if $\Lambda_0 > \max W$, then (P.1) reads equivalently as $\int_{\mathcal{M} \times \{0\}} Y_W(a) d\bar{\mu}_\infty = 0$ for every $a \in \mathcal{C}^1(\mathcal{M})$. This implies the property of $\bar{\nu}_\infty$ in Theorem 1.1 by letting

$$\bar{\nu}_\infty(q) = \int_{\mathbb{R}} \bar{\mu}_\infty(q, dE).$$

Notice, since $\Omega_{\mathcal{H}_1}(\mathcal{H}_1) = 0$, that $\mathcal{X}_{W,Q}(\mathcal{H}_\pm) = 0$. This implies in particular that the vector field $E^{-1}\mathcal{X}_{W,Q}$ is tangent to the level sets $\mathcal{H}_\pm^{-1}(2k+1)$ and thus induces a well-defined flow on these layers. Finally, we can derive from (S.2) and (P.2) that

$$\int_{\mathcal{M} \times \mathbb{R}_\pm^*} ((\pm(2k+1) + Q)Y_W(a) - \Omega_Q(a) + X(\mathcal{H}_1)\partial_E a) d\mu_{k,\infty}^\pm = 0,$$

which implies the last part of Theorem 1.1 by letting

$$\nu_{k,\infty}^\pm(q) := \int_{\mathbb{R}} \mu_{k,\infty}^\pm(q, dE).$$

8. THE CASE OF THE FLAT TORUS

In this section, we briefly discuss the case where $M = \mathbb{T}^2 = \mathbb{R}^2/2\pi\mathbb{Z}^2$, $Q = W = 0$ and $g = dx^2 + dy^2$ is the canonical Euclidean metric. Our aim is to show examples of different sequences of eigenfunctions for the operator $-h^2\Delta_{\text{sR}}$ which select any given choice among the semiclassical measures $\bar{\mu}_\infty$ and $\mu_{k,\infty}^\pm$ by putting their total mass on them as $h \rightarrow 0^+$.

In this particular example, the operators X , X_\perp and V can be written by global formulas in the canonical coordinates $(x, y, z) \in \mathbb{T}^3 \simeq S\mathbb{T}^2$. Precisely:

$$X = \cos z\partial_x + \sin z\partial_y, \quad X_\perp = \sin z\partial_x - \cos z\partial_y, \quad \text{and } V = \partial_z,$$

so that

$$\Delta_{\text{sR}} = (\sin z\partial_x - \cos z\partial_y)^2 + \partial_z^2.$$

We restrict ourselves to search for solutions to (3) of the particular form

$$\psi_h(x, y, z) = u_h(z)e^{in \cdot (x,y)}, \quad \mathbf{n} = (n_1, n_2) \in \mathbb{Z}^2.$$

As we impose that ψ_h solves our eigenvalue problem, then u_h must satisfy

$$h^2(n_1 \sin z - n_2 \cos z)^2 u_h(z) - h^2 u_h''(z) = u_h(z),$$

or equivalently

$$-\frac{1}{\|\mathbf{n}\|^2} u_h''(z) + \sin^2(z - z_{\mathbf{n}}) u_h(z) = \frac{1}{h^2 \|\mathbf{n}\|^2} u_h(z),$$

with $(\cos z_{\mathbf{n}}, \sin z_{\mathbf{n}}) = \mathbf{n}/\|\mathbf{n}\|$. We recognize in this expression the semiclassical Mathieu operator

$$\widehat{M}_{\mathbf{n}} := -\frac{1}{\|\mathbf{n}\|^2} \partial_z^2 + \sin^2(z - z_{\mathbf{n}}),$$

on the circle \mathbb{S}^1 whose spectral analysis is a classical topic. Indeed, it is a one-dimensional Schrödinger operator with a double well potential. Hence, we are led to the equation

$$(57) \quad \widehat{M}_{\mathbf{n}} u_{\mathbf{n}} = \lambda(\mathbf{n}) u_{\mathbf{n}}, \quad \lambda(\mathbf{n}) = \frac{1}{(h_{\mathbf{n}} \|\mathbf{n}\|)^2}.$$

Since $\widehat{M}_{\mathbf{n}}$ has compact resolvent on $L^2(\mathbb{S}^1)$, for every $\mathbf{n} \in \mathbb{Z}^2$ there is an increasing sequence of eigenvalues $(\Lambda_k(\mathbf{n}))_{k \geq 0}$ with corresponding (normalized) eigenfunctions $(u_{\mathbf{n},k}(z))_{k \geq 0}$ of $\widehat{M}_{\mathbf{n}}$.

REMARK 8.1. — Note that

$$\text{Sp}(\Delta_{\text{sR}}) = \bigcup_{\mathbf{n} \in \mathbb{Z}^2} \text{Sp}(\|\mathbf{n}\|^2 \widehat{M}_{\mathbf{n}}).$$

Note that, up to translation by $z_{\mathbf{n}}$, we can restrict ourselves to the case where $z_{\mathbf{n}} = 0$ which amounts to take a lattice point of the form $\mathbf{n} = (n, 0)$ with say $n > 0$. Under this assumption, we first show the following standard fact:

LEMMA 8.2. — For every $k \in \mathbb{N}$, there exists $\Lambda_k(n, 0) := \Lambda_k(n) \in \text{Sp}_{L^2(S^1)}(\widehat{M}_{(n,0)})$ such that:

$$(58) \quad \Lambda_k(n) = \frac{(2k + 1)}{n} (1 + \mathcal{O}_k(1/\sqrt{n})), \quad \text{as } n \rightarrow +\infty.$$

REMARK 8.3. — In principle, there could be other sequences of eigenvalues verifying different asymptotic formulas, say $\Lambda_\alpha(n) = \alpha/n + o(n^{-1})$. Yet, one could show that this is not the case by comparing eigenfunctions of \widehat{M}_n with quasimodes of the harmonic oscillator

$$\widehat{H}_n := -\frac{1}{n^2} \partial_z^2 + z^2,$$

on $L^2(\mathbb{R})$, whose spectrum is given explicitly by

$$\text{Sp}_{L^2(\mathbb{R})}(\widehat{H}_n) = \left\{ \frac{2k + 1}{n} : k \in \mathbb{N} \right\}.$$

However, since we are only interested in showing the existence of sequences of eigenfunctions of $-h_n^2 \Delta_{\text{sR}}$ which put positive mass on the semiclassical measures $\bar{\mu}_\infty$ and $\mu_{k,\infty}^\pm$, for which we already know the concentration properties (S.1) and (S.2) of Theorem 7.1, we omit this discussion.

Proof. — The proof of this lemma is classical and we just briefly recall it for the sake of completeness. Recall that is sufficient to construct a sequence of (almost) normalized quasimodes $(v_{k,n}, E_k(n))_{n \rightarrow \infty}$ satisfying

$$\widehat{M}_n v_{k,n} = E_k(n) v_{k,n} + \mathcal{O}_k(1/n^{3/2}); \quad E_k(n) = \frac{2k + 1}{n}.$$

To this aim, let $\delta > 0$ small, and set $v_{k,n}(z) := \chi(z/\delta) \varphi_{k,n}$ where χ is a cutoff function supported in a neighborhood of 0 and where

$$\varphi_{k,n}(z) = n^{1/4} \varphi_k(\sqrt{n}z),$$

and φ_k is the normalized Hermite function of degree k [Zwo12, Th. 6.2]. Observe that, for any $N \geq 1$ and for any $\ell \geq 0$,

$$(59) \quad \int_{|z| \geq \delta} |\varphi_{k,n}^{(\ell)}(z)|^2 dz = \mathcal{O}_{\delta,\ell,k,N}(1/n^{N/2}), \quad \text{as } \|n\| \rightarrow \infty.$$

In particular, we get that $\|v_{k,n}\|_{L^2(\mathbb{T})} = 1 + \mathcal{O}_k(n^{-1})$ as $n \rightarrow \infty$ as expected. Moreover, using the Taylor expansion

$$\sin^2 z = z^2 + \mathcal{O}(|z|^3), \quad \text{as } z \rightarrow 0,$$

and the fact that

$$\left(\int_{\mathbb{R}} |z^3 \varphi_{k,n}(z)|^2 dz \right)^{1/2} = \mathcal{O}_k(1/n^{3/2}), \quad \text{as } n \rightarrow \infty,$$

the claim holds by using the eigenvalue equation combined with (59) and

$$\widehat{H}_n \varphi_{k,n} = E_k(n) \varphi_{k,n}, \quad z \in \mathbb{R}. \quad \square$$

Let us now fix, for every $n > 0$,

$$(60) \quad h_n := \frac{1}{\sqrt{(2k+1+o_k(1))n}}, \quad \text{as } n \rightarrow \infty,$$

so that (57) and (58) hold. For this sequence, take a sequence of solutions to (3) that are of the form

$$(61) \quad \psi_n(x, y, z) = u_n(z)e^{inx}.$$

PROPOSITION 8.4. — *Let $(\psi_n)_{n \geq 1}$ be a normalized sequence of the form (61). Let us assume that $(\psi_n)_{n \geq 1}$ satisfy (3) with $(h_n)_{n \geq 1}$ given by (60). Then the total mass of $\mu_{k,\infty}^+ + \mu_{k,\infty}^-$ is equal to one.*

Again, we will just make a rough analysis and a more careful work would show that the sequences of eigenmodes put equal mass on $\mu_{k,\infty}^\pm$ due to symmetry of our double well potential. If we were looking for quasimodes, then we could ensure that the full mass is put either on $\mu_{k,\infty}^+$ or $\mu_{k,\infty}^-$.

Proof. — Let us consider $\delta > 0$ to be chosen sufficiently small along the proof. Let $\chi_\delta = \chi(\cdot/\delta)$ where χ is still a small cutoff function near 0. We have, by the functional analysis of pseudodifferential operators [Zwo12, Th. 14.9], the localization property:

$$\begin{aligned} \text{Op}_{1/\|n\|}^{\mathbb{S}^1,w}(\chi_\delta(\sin^2(z) + \zeta^2))u_n &= \chi_\delta(\widehat{M}_n)u_n + O_\delta(1/n) \\ &= \chi_\delta\left(\frac{2k+1}{n}(1+o_k(1))\right)u_n + O_\delta(1/n) \\ &= u_n + O_{\delta,k}(1/n). \end{aligned}$$

On the other hand, we have:

$$\langle \text{Op}_{h_n}^w(\chi(h_n H_1)\widetilde{\chi}_\varepsilon^C \widetilde{\chi}_R^B)\psi_n, \psi_n \rangle_{L^2(\mathbb{T}^3)} = \langle \text{Op}_{h_n}^{\mathbb{S}^1,w}(\kappa_n^{\varepsilon,R})u_n, u_n \rangle_{L^2(\mathbb{S}^1)},$$

where

$$\kappa_n^{\varepsilon,R}(z, \zeta) := \chi(h_n^2 n \cos(z))\widetilde{\chi}\left(\frac{\varepsilon h_n n \cos(z)}{\sqrt{1+(h_n n)^2 \sin^2(z) + \zeta^2}}\right)\widetilde{\chi}\left(\frac{(h_n n)^2 + \zeta^2}{R}\right).$$

Arguing as in Section 3, one finds that this symbol belongs to the class of symbols $S_{\text{cl}}^0(T^*\mathbb{S}^1)$ amenable to pseudodifferential calculus on the circle. Observe also that, for n large enough, the last function in this product is identically equal to 1. Notice also that

$$\text{Op}_{1/\|n\|}^{\mathbb{S}^1,w}(\chi_\delta((\sin^2(z) + \zeta^2))) = \text{Op}_{h_n}^{\mathbb{S}^1,w}(\chi_\delta(\sin^2(z) + (\zeta/h_n n)^2)) =: \text{Op}_{h_n}^{\mathbb{S}^1,w}(\sigma_n^\delta).$$

Thus, by using the previous localization property for u_n and the semiclassical pseudodifferential calculus, we have the composition formula:

$$\begin{aligned} \langle \text{Op}_{h_n}^{\mathbb{S}^1,w}(\kappa_n^{\varepsilon,R})u_n, u_n \rangle_{L^2(\mathbb{S}^1)} &= \langle \text{Op}_{h_n}^{\mathbb{S}^1,w}(\kappa_n^{\varepsilon,R}) \text{Op}_{h_n}^{\mathbb{S}^1,w}(\sigma_n^\delta)u_n, u_n \rangle_{L^2(\mathbb{S}^1)} + \mathcal{O}_{\delta,k}(1/n) \\ &= \langle \text{Op}_{h_n}^{\mathbb{S}^1,w}(\kappa_n^{\varepsilon,R} \sigma_n^\delta)u_n, u_n \rangle_{L^2(\mathbb{S}^1)} + \mathcal{O}_{\delta,\varepsilon,R}(h_n). \end{aligned}$$

In this expression, if we take δ sufficiently small (i.e., so that $\delta \ll \varepsilon^2$), we have

$$\kappa_n^{\varepsilon,R}(z, \zeta) \sigma_n^\delta(z, \zeta) = \chi(h_n^2 n \cos(z)) \sigma_n^\delta(z, \zeta),$$

since $\tilde{\chi}(x) = 1$ for $|x| \geq 2$. Moreover, for $\delta > 0$ sufficiently small, we can also decompose σ_n^δ as the sum of two functions $\sigma_n^{1,\delta}$ and $\sigma_n^{2,\delta}$ compactly supported respectively near $z = 0$ and $z = \pi$, that is:

$$\sigma_n^\delta(z, \zeta) = \sigma_n^{1,\delta}(z, \zeta) + \sigma_n^{2,\delta}(z, \zeta),$$

with $\text{supp } \sigma_n^{1,\delta} \cap \text{supp } \sigma_n^{2,\delta} = \emptyset$. Using next that

$$h_n^2 n \cos(z) = \frac{(-1)^{j-1}}{2k+1} + \mathcal{O}(\delta), \quad \text{as } \delta \rightarrow 0, \quad j = 1, 2,$$

respectively on the support of $\sigma_n^{j,\delta}(z, \zeta)$, we get thanks to the Calderón-Vaillancourt theorem:

$$\begin{aligned} & \langle \text{Op}_{h_n}^{\mathbb{S}^1, w}(\kappa_n^{\varepsilon,R}) u_n, u_n \rangle_{L^2(\mathbb{S}^1)} \\ &= \sum_{j \in \{1,2\}} \chi\left(\frac{(-1)^{j-1}}{2k+1}\right) \langle \text{Op}_{h_n}^{\mathbb{S}^1, w}(\sigma_n^{j,\delta}) u_n, u_n \rangle_{L^2(\mathbb{S}^1)} + \mathcal{O}(\delta) + \mathcal{O}_{\delta, \varepsilon, R}(h_n). \end{aligned}$$

Therefore, taking limits in $n \rightarrow +\infty$ through a subsequence, we obtain, for $\delta \ll \varepsilon^2$,

$$\lim_{h_n \rightarrow 0} \langle \text{Op}_{h_n}^w(\chi(h_n H_1) \tilde{\chi}_\varepsilon^C \tilde{\chi}_R^B) \psi_n, \psi_n \rangle_{L^2(\mathcal{M})} = \alpha_1^\delta \chi\left(\frac{1}{2k+1}\right) + \alpha_2^\delta \chi\left(\frac{-1}{2k+1}\right) + \mathcal{O}(\delta),$$

where $\alpha_j^\delta = \lim_{h_n \rightarrow 0^+} \langle \text{Op}_{h_n}^{\mathbb{T}^1, w}(\sigma_n^{j,\delta}) u_n, u_n \rangle_{L^2(\mathbb{T}^1)}$, and $\alpha_1^\delta + \alpha_2^\delta = 1$ (using one more time the localization property of the sequence $(u_n)_{n \geq 1}$). Finally, in view of the fact that

$$\lim_{\varepsilon \rightarrow 0} \lim_{R \rightarrow \infty} \lim_{h_n \rightarrow 0} \langle \text{Op}_{h_n}^w(\chi(h_n H_1) \tilde{\chi}_\varepsilon^C \tilde{\chi}_R^B) \psi_n, \psi_n \rangle_{L^2(\mathbb{T}^3)} = \int_{\mathbb{T}^3 \times \mathbb{R}} \chi(E) d\mu_\infty(q, E),$$

we can take $\delta \rightarrow 0$ and use Theorem 7.1 (property (S.2)) to conclude the proof. \square

We finally show the existence of sequences of eigenfunctions (ψ_n) satisfying (3) which put positive mass on the semiclassical measure $\bar{\mu}_\infty$. To this aim, we note that, thanks to Lemma 8.2 and for every $k \geq 1$, we can find some $n_k \geq 1$ such that, for every $n \geq n_k$, there is an eigenvalue $E_k(n)$ of $\widehat{M}_{(n,0)}$ verifying

$$\frac{1}{2\sqrt{(2k+1)n}} \leq E_k(n) = \frac{1}{(h_n n)^2} \leq \frac{\sqrt{2}}{\sqrt{(2k+1)n}} \leq \frac{1}{\sqrt{k}}.$$

Hence, we can take $n = n_k$ and pick a sequence $(K_{n_k})_{k \geq 1}$ such that $K_{n_k} \rightarrow +\infty$ and thus the sequence $(h_{n_k})_{k \geq 1}$ satisfying now

$$(62) \quad \frac{1}{n_k} \ll h_{n_k} = \frac{1}{\sqrt{K_{n_k} n_k}} \ll \frac{1}{\sqrt{n_k}}, \quad \text{as } k \rightarrow \infty.$$

Adapting the proof of Proposition 8.4 and using property (S.1) of Theorem 7.1, we obtain:

COROLLARY 8.5. — *Let $(\psi_{n_k})_{k \geq 1}$ be a normalized sequence of the form (61). Let us assume that ψ_{n_k} satisfy (3) with h_{n_k} given by (62). Then the total mass of $\bar{\mu}_\infty$ is equal to one.*

APPENDIX A. SPECTRAL PROPERTIES OF \widehat{P}_h

In this appendix, we briefly review the spectral properties of \widehat{P}_h . A key ingredient of the analysis is the following standard result [RS76, Cor. 17.14]:

THEOREM A.1 (Rothschild-Stein). — *Let $Q \in \mathcal{C}^\infty(\mathcal{M}, \mathbb{R})$ such that $\|Q\|_{\mathcal{C}^0} < 1$. Set*

$$\mathcal{L} = -\Delta_{sR} - iQX = -X_\perp^2 - V^2 - iQ[V, X_\perp].$$

Then, for every $N \geq 1$, one can find continuous maps $P_N : H^s \rightarrow H^{1+s}$ and $S_N : H^s \rightarrow H^{s+N/2}$ (for all $s \geq 0$) such that

$$P_N \mathcal{L} = \text{Id} + S_N.$$

In particular, there exists a constant $C_{M,g} > 0$ such that

$$(63) \quad \forall \psi \in \mathcal{C}^\infty(\mathcal{M}), \quad \|\psi\|_{H^1} \leq C_{M,g} (\|\mathcal{L}\psi\|_{L^2} + \|\psi\|_{L^2}),$$

and, for every $s \geq 0$,

$$\mathcal{L}(\psi) \in H^s, \quad \psi \in L^2 \implies \psi \in H^{s+1}.$$

Let us now discuss the spectral properties of \widehat{P}_h . For an introduction on the spectral properties of unbounded operators, the reader is referred to [RS72, Ch. VIII] and [RS75, Ch. X] that we closely follow for the terminology. For any $\psi \in H^2(\mathcal{M})$, we define

$$\widetilde{P}_h(\psi) := \left(-h^2 \Delta_{sR} + \frac{h^2}{2i} (QX - (QX)^*) + W \right) \psi,$$

which induces an unbounded operator

$$\widetilde{P}_h : D(\widetilde{P}_h) := H^2(\mathcal{M}) \subset L^2(\mathcal{M}) \longrightarrow L^2(\mathcal{M}).$$

One can define its adjoint \widetilde{P}_h^* by defining the domain

$$D(\widetilde{P}_h^*) := \left\{ \psi \in L^2(\mathcal{M}) : \exists u \in L^2(\mathcal{M}) \text{ such that } \forall \varphi \in H^2(\mathcal{M}), \langle \psi, \widetilde{P}_h \varphi \rangle = \langle u, \varphi \rangle \right\},$$

or equivalently

$$D(\widetilde{P}_h^*) := \{ \psi \in L^2(\mathcal{M}) : \widetilde{P}_h \psi \in L^2(\mathcal{M}) \}$$

The operator $\widetilde{P}_h^* : \psi \in D(\widetilde{P}_h^*) \rightarrow \widetilde{P}_h \psi \in L^2(\mathcal{M})$ is closed and it is densely defined. Hence, according to [RS72, Th. VIII.1], \widetilde{P}_h is closable and we denote its closure by $\overline{\widetilde{P}_h}$ whose domain is denoted by $D(\overline{\widetilde{P}_h})$ and equal to the set of $\psi \in L^2(\mathcal{M})$ such that

$$\exists \psi_j \in H^2(\mathcal{M}), \exists v \in L^2(\mathcal{M}) \text{ such that } \|\psi_j - \psi\|_{L^2} + \|\widetilde{P}_h \psi_j - v\|_{L^2} \longrightarrow 0.$$

In general, one only has $D(\overline{\widetilde{P}_h^*}) = D(\widetilde{P}_h^*) \subset D(\overline{\widetilde{P}_h})$ so that $\overline{\widetilde{P}_h}$ is not necessarily selfadjoint. In order to fix this problem, we can make some assumptions on the size of Q and use positivity arguments.

More precisely, \tilde{P}_h is associated with the real quadratic form

$$\tilde{B}(\psi) := \int_{\mathcal{M}} (\tilde{P}_h \psi) \bar{\psi} d\mu_L, \quad \psi \in H^2(\mathcal{M}),$$

which, thanks to (1), is bounded from below by

$$\begin{aligned} \tilde{B}(\psi) &\geq \|hX_{\perp}\psi\|_{L^2}^2 + \|hV\psi\|_{L^2}^2 - 2\|Q\|_{\mathcal{C}^0} \|hX_{\perp}\psi\|_{L^2} \|hV\psi\|_{L^2} \\ &\quad + (\min W - h^2\|Q\|_{\mathcal{C}^1}) \|\psi\|_{L^2}^2 \\ &\quad - h\|Q\|_{\mathcal{C}^1} \|\psi\|_{L^2} (\|hX_{\perp}\psi\|_{L^2} + \|hV\psi\|_{L^2}) \\ &\geq \left(1 - \|Q\|_{\mathcal{C}^0} - \frac{h\|Q\|_{\mathcal{C}^1}}{2}\right) (\|hX_{\perp}\psi\|_{L^2}^2 + \|hV\psi\|_{L^2}^2) \\ &\quad + (\min W - (h^2 + h)\|Q\|_{\mathcal{C}^1}) \|\psi\|_{L^2}^2. \end{aligned}$$

Hence, if $\|Q\|_{\mathcal{C}^0} < 1$ (and $h > 0$ is small enough in a way that depends on Q), it follows from [RS75, Th. X.23] that \tilde{B} is a closable form whose closure B corresponds to a unique selfadjoint operator \hat{P}_h referred as the *Friedrichs extension* of \tilde{P}_h . Moreover, the spectrum of this selfadjoint extension is bounded from below by $\min W + \mathcal{O}_Q(h)$ and its domain verifies

$$D(\hat{P}_h) \subset H_{\text{sR}}^1(\mathcal{M}) := \{\psi \in \mathcal{D}'(\mathcal{M}) : \|\psi\|_{L^2}^2 + \|X_{\perp}\psi\|_{L^2}^2 + \|V\psi\|_{L^2}^2 < \infty\}.$$

In particular, $\hat{P}_h : D(\hat{P}_h) \subset L^2(\mathcal{M}) \rightarrow L^2(\mathcal{M})$ is a closed selfadjoint operator and thus $(\hat{P}_h + C)$ has a bounded inverse for $C > 0$ large enough:

$$(\hat{P}_h + C)^{-1} : L^2(\mathcal{M}) \longrightarrow (D(\hat{P}_h), \|\cdot\|_{L^2}) \subset L^2(\mathcal{M}).$$

We would like to show that this defines a compact operator. To see this, recall from (63) that

$$\forall \psi \in \mathcal{C}^{\infty}(\mathcal{M}), \quad \|(\hat{P}_h + C)^{-1}\psi\|_{H^1(\mathcal{M})} \leq c_h (\|\psi\|_{L^2} + \|(\hat{P}_h + C)^{-1}\psi\|_{L^2}),$$

so that, if $(\psi_j)_{j \geq 0}$ is a bounded sequence in $L^2(\mathcal{M})$, then $((\hat{P}_h + C)^{-1}\psi_j)_{j \geq 0}$ is also bounded in $H^1(\mathcal{M})$.

REMARK A.2. — Along the way, this discussion shows that $H^2(\mathcal{M}) \subset D(\hat{P}_h) \subset H^1(\mathcal{M})$ (with continuous inclusions).

As the inclusion $H^1(\mathcal{M}) \subset L^2(\mathcal{M})$ is compact, $(\hat{P}_h + C)^{-1} : L^2(\mathcal{M}) \rightarrow L^2(\mathcal{M})$ is indeed a compact operator. As \hat{P}_h is selfadjoint, there exists an orthonormal basis of $L^2(\mathcal{M})$ made of eigenmodes of \hat{P}_h . Moreover, if one has $\hat{P}_h \psi_h = \Lambda_h \psi_h$ with $\psi_h \in D(\hat{P}_h)$, then $\mathcal{L}\psi_h \in H^1(\mathcal{M})$ and, according to Theorem A.1, one finds that $\psi_h \in H^2(\mathcal{M})$. By induction, we get that these eigenmodes are smooth.

REMARK A.3. — If we let $C > 0$ be a large enough constant, then $\tilde{P}_h + C$ is a positive symmetric operator and its adjoint is given by $\tilde{P}_h^* + C$ with domain $D(\tilde{P}_h^*)$. In particular, if ψ belongs to the kernel of $\tilde{P}_h^* + C$, then, by the Rothschild-Stein theorem, ψ belongs to $H^1(\mathcal{M})$ (and by induction to $\mathcal{C}^{\infty}(\mathcal{M})$). Hence, it lies in the domain of \hat{P}_h and we can deduce that $\psi = 0$. According to [RS75, Th. X.26], it implies

that the Friedrichs extension is the only (semibounded) selfadjoint extension of $\tilde{P}_h + C$ (hence of \hat{P}_h).

The spectral properties of \hat{P}_h that we have proved so far are summarized by the next statement:

LEMMA A.4. — *Suppose that $\|Q\|_{\mathcal{C}^0} < 1$. Then, there exists $h_0 > 0$ such that, for every $0 < h < h_0$,*

$$\hat{P}_h : D(\hat{P}_h) \longrightarrow L^2(\mathcal{M})$$

is a selfadjoint operator whose spectrum consists in a discrete sequence of eigenvalues

$$\min W + \mathcal{O}_Q(h) \leq \Lambda_h(0) \leq \Lambda_h(1) \leq \dots \leq \Lambda_h(j) \dots \longrightarrow +\infty.$$

Moreover,

$$\hat{P}_h \psi_h = \Lambda_h \psi_h, \text{ with } \psi_h \in D(-\hat{P}_h) \implies \psi_h \in \mathcal{C}^\infty(\mathcal{M}).$$

We conclude this appendix with the following a priori estimates that are used all along the article:

LEMMA A.5. — *Suppose that $\|Q\|_{\mathcal{C}^0} < 1$. Then, one can find $C_{Q,W} > 0$ and $0 < h_Q \leq 1$, such that, for all $0 < h \leq h_Q$,*

$$\begin{aligned} \hat{P}_h \psi_h &= \Lambda_h \psi_h, \text{ with } \psi_h \in D(\hat{P}_h) \\ \implies \|hX_\perp \psi_h\|_{L^2}^2 + \|hV \psi_h\|_{L^2}^2 + \|h^2 X \psi_h\|_{L^2}^2 &\leq C_{Y,W} (1 + |\Lambda_h|)^2 \|\psi_h\|_{L^2}^2. \end{aligned}$$

Proof. — Let $\psi_h \in D(\hat{P}_h)$ such that $\hat{P}_h = \Lambda_h \psi_h$. One has then

$$\|hX_\perp \psi_h\|_{L^2}^2 + \|hV \psi_h\|_{L^2}^2 = \Lambda_h \|\psi_h\|_{L^2}^2 - \langle W \psi_h, \psi_h \rangle - \frac{h^2}{i} \langle (QX + 1/2X(Q)) \psi_h, \psi_h \rangle.$$

Hence, one has

$$\|hX_\perp \psi_h\|_{L^2}^2 + \|hV \psi_h\|_{L^2}^2 \leq (\|W\|_{\mathcal{C}^0} + |\Lambda_h|) \|\psi_h\|_{L^2}^2 + h^2 |\langle (QX + \frac{1}{2}X(Q)) \psi_h, \psi_h \rangle|.$$

Recall that $X = [V, X_\perp]$ from which we infer

$$\begin{aligned} |\langle (QX + \frac{1}{2}X(Q)) \psi_h, \psi_h \rangle| &\leq 2\|Q\|_{\mathcal{C}^0} \|hX_\perp \psi_h\|_{L^2} \|hV \psi_h\|_{L^2} \\ &\quad + h\|Q\|_{\mathcal{C}^1} (\|hX_\perp \psi_h\|_{L^2} + \|hV \psi_h\|_{L^2}) \|\psi_h\|_{L^2} + \frac{h^2}{2} \|Q\|_{\mathcal{C}^1} \|\psi_h\|_{L^2}^2. \end{aligned}$$

Then, we get

$$\|hX_\perp \psi_h\|_{L^2}^2 + \|hV \psi_h\|_{L^2}^2 \leq \frac{1}{1 - \|Q\|_{\mathcal{C}^0} - h\|Q\|_{\mathcal{C}^1}/2} (\|W\|_{\mathcal{C}^0} + |\Lambda_h| + 2h\|Q\|_{\mathcal{C}^1}) \|\psi_h\|_{L^2}^2.$$

Hence, under the assumption that $\|Q\|_{\mathcal{C}^0} < 1$, there exists a constant $C_{Q,W} > 0$ (depending only Q and W) and $0 < h_Q \leq 1$ (depending only on Q) such that, for every $0 < h \leq h_Q$,

$$\|hX_\perp \psi_h\|_{L^2}^2 + \|hV \psi_h\|_{L^2}^2 \leq C_{Q,W} (1 + |\Lambda_h|) \|\psi_h\|_{L^2}^2.$$

Finally, using the Rothschild-Stein theorem one more time, one finds that there exists a constant $C_{M,g} > 0$ such that

$$\|X\psi_h\|_{L^2} \leq \|\psi_h\|_{H^1} \leq C_{M,g} (\|\mathcal{L}\psi_h\|_{L^2} + \|\psi_h\|_{L^2}).$$

Multiplying this inequality by h^2 and using the fact that $\widehat{P}_h\psi_h = \Lambda_h\psi_h$ to control the upper bound in terms of $\|\psi_h\|_{L^2}$, we obtain the expected upper bound. \square

APPENDIX B. REMINDER ON SEMICLASSICAL ANALYSIS ON $\mathbb{R}^2 \times \mathbb{S}^1$

In this appendix, we review a few facts about semiclassical analysis on $T^*(\mathbb{R}^2 \times \mathbb{S}^1)$ that are used all along our analysis of the measure at infinity. A standard textbook is [Zwo12] which treats the case of $T^*\mathbb{R}^3$ in great details in Chapter 4. The case of $T^*(\mathbb{R}^2 \times \mathbb{S}^1)$ can be handled similarly by proper use of Fourier series along the z -variable rather than Fourier transform. See for instance [Zwo12, §5.3] for a detailed discussion in the case of $T^*\mathbb{T}^3$.

For a nice enough smooth function a on $T^*(\mathbb{R}^2 \times \mathbb{S}^1)$ (say compactly supported) and for every $h > 0$, the Weyl (semiclassical) quantization of a is defined, for all u in $\mathcal{C}_c^\infty(\mathbb{R}^3)$, by

$$\text{Op}_h^w(a)(u)(q) := \frac{1}{(2\pi h)^3} \int_{\mathbb{R}^6} e^{(i/h)(q-q') \cdot p} a\left(\frac{q+q'}{2}, p\right) u(q') dq' dp.$$

Using the periodicity along the \mathbb{S}^1 -variable, one can verify that this definition extends to smooth test functions $u \in \mathcal{C}_c^\infty(\mathbb{R}^2 \times \mathbb{S}^1)$ [Zwo12, §5.3.1].

Regarding the regularity needed for a , this definition still makes sense when working with smooth functions a belonging to the class of (Kohn-Nirenberg) symbols [Zwo12, §9.3]:

$$S_{\text{cl}}^m(T^*(\mathbb{R}^2 \times \mathbb{S}^1)) = \{a \in \mathcal{C}^\infty(T^*(\mathbb{R}^2 \times \mathbb{S}^1)) : \forall(\alpha, \beta) \in \mathbb{Z}_+^6, P_{m,\alpha,\beta}(a) < +\infty\},$$

where $m \in \mathbb{R}$, and

$$P_{m,\alpha,\beta}(a) := \sup_{(q,p)} \{ \langle p \rangle^{-m+|\beta|} |\partial_q^\alpha \partial_p^\beta a(x, \xi)| \}.$$

In other words, we gain some decay in p when differentiating in the p -variable. Even if such a decay is not necessary to work in an Euclidean set-up, it is of crucial importance in our analysis to have this extra decay in view of dealing with the escape at infinity in the fibers.

A nice property of the Weyl quantization is that, for a real-valued a , $\text{Op}_h^w(a)$ is a (formally) selfadjoint operator [Zwo12, Th. 4.1]. Another property that we extensively use all along this article is the composition rule for pseudodifferential operators⁽⁵⁾ [Zwo12, Th. 9.5, Th. 4.12].

⁽⁵⁾Technically speaking, this reference deals with the Weyl quantization on $T^*\mathbb{R}^3$ but the proof works as well in our set-up.

THEOREM B.1. — *Let $a \in S_{\text{cl}}^{m_1}(T^*(\mathbb{R}^2 \times \mathbb{S}^1))$ and $b \in S_{\text{cl}}^{m_2}(T^*(\mathbb{R}^2 \times \mathbb{S}^1))$. Then, there exists $c \in S_{\text{cl}}^{m_1+m_2}(T^*(\mathbb{R}^2 \times \mathbb{S}^1))$ (depending on h) such that*

$$(64) \quad \text{Op}_h^w(a) \circ \text{Op}_h^w(b) = \text{Op}_h^w(c).$$

Moreover,

$$c(q, p) = \sum_{k=0}^N \frac{h^k}{k!} (A(D))^k (a(q_1, p_1)b(q_2, p_2))|_{q_1=q_2=q, p_1=p_2=p} + \mathcal{O}_{S^{m_1+m_2-N-1}}(h^{N+1}),$$

where the constant in the remainder depends on a finite number of seminorms of a and b (depending on N and on the semi-norm in $S^{m_1+m_2-N-1}$), and where

$$A(D) := \frac{1}{2i} (\partial_{p_1} \cdot \partial_{q_2} - \partial_{p_2} \cdot \partial_{q_1}).$$

In particular, we can see from this result that $c = \mathcal{O}_{S^{m_1+m_2-N-1}}(h^{N+1})$ if a and b have disjoint supports. We can also verify that, all the even powers in h in the asymptotic expansion of $[\text{Op}_h^w(a), \text{Op}_h^w(b)]$ cancels out and that the first term is given by $(h/i)\{a, b\}$.

Another key property for us is the Calderón-Vaillancourt theorem [Zwo12, Ch. 5] that states the existence of constants C_0, N_0 such that, for every $a \in S_{\text{cl}}^0(T^*(\mathbb{R}^2 \times \mathbb{S}^1))$,

$$\|\text{Op}_h^w(a)\|_{L^2 \rightarrow L^2} \leq C_0 \sum_{|\alpha| \leq N_0} h^{|\alpha|/2} \|\partial^\alpha a\|_\infty.$$

Recall also the Gårding property that is valid for elements in $S_{\text{cl}}^0(T^*(\mathbb{R}^2 \times \mathbb{S}^1))$. Given any a in that class satisfying $a \geq 0$, it ensures the existence of a constant $C_a > 0$ [Zwo12, Th. 4.32] such that

$$(65) \quad \forall u \in L^2(\mathbb{R}^2 \times \mathbb{S}^1), \quad \langle \text{Op}_h^w(a)u, u \rangle \geq -C_a h \|u\|_{L^2}^2.$$

REFERENCES

- [AM14] N. ANANTHARAMAN & F. MACIÀ – “Semiclassical measures for the Schrödinger equation on the torus”, *J. Eur. Math. Soc. (JEMS)* **16** (2014), no. 6, p. 1253–1288.
- [AS23] V. ARNAIZ & C. SUN – “Sharp resolvent estimate for the Baouendi-Grushin operator and applications”, *Comm. Math. Phys.* **400** (2023), no. 1, p. 541–637.
- [BVN21] G. BOIL & S. VU NGOC – “Long-time dynamics of coherent states in strong magnetic fields”, *Amer. J. Math.* **143** (2021), no. 6, p. 1747–1789.
- [BR20] Y. BONTHONNEAU & N. RAYMOND – “WKB constructions in bidimensional magnetic wells”, *Math. Res. Lett.* **27** (2020), no. 3, p. 647–663.
- [BS22] N. BURQ & C. SUN – “Time optimal observability for Grushin Schrödinger equation”, *Anal. PDE* **15** (2022), no. 6, p. 1487–1530.
- [Col23] Y. COLIN DE VERDIÈRE – “A proof of a trace formula by Richard Melrose”, *Adv. Nonlinear Stud.* **23** (2023), no. 1, article no. 20220054 (13 pages).
- [CHT18] Y. COLIN DE VERDIÈRE, L. HILLAIRET & E. TRÉLAT – “Spectral asymptotics for sub-Riemannian Laplacians, I: Quantum ergodicity and quantum limits in the 3-dimensional contact case”, *Duke Math. J.* **167** (2018), no. 1, p. 109–174.
- [CHT21] Y. COLIN DE VERDIÈRE, L. HILLAIRET & E. TRÉLAT – “Spiraling of sub-Riemannian geodesics around the Reeb flow in the 3d contact case”, 2021, [arXiv:2102.12741](https://arxiv.org/abs/2102.12741).
- [DZ19] S. DYATLOV & M. ZWORSKI – *Mathematical theory of scattering resonances*, Graduate Studies in Math., vol. 200, American Mathematical Society, Providence, RI, 2019.
- [FK95] C. FERMANIAN-KAMMERER – “Mesures semi-classiques et équation de la chaleur”, PhD Thesis, Université Paris-Sud Orsay, 1995.
- [FK00] ———, “Mesures semi-classiques 2-microlocales”, *C. R. Acad. Sci. Paris Sér. I Math.* **331** (2000), no. 7, p. 515–518.

- [FKF21] C. FERMANIAN-KAMMERER & V. FISCHER – “Quantum evolution and sub-Laplacian operators on groups of Heisenberg type”, *J. Spectral Theory* **11** (2021), no. 3, p. 1313–1367.
- [FKL21] C. FERMANIAN-KAMMERER & C. LETROUIT – “Observability and controllability for the Schrödinger equation on quotients of groups of Heisenberg type”, *J. Éc. polytech. Math.* **8** (2021), p. 1459–1513.
- [FS74] G. B. FOLLAND & E. M. STEIN – “Estimates for the $\bar{\partial}_b$ complex and analysis on the Heisenberg group”, *Comm. Pure Appl. Math.* **27** (1974), p. 429–522.
- [GNRV21] Y. GUEDES BONTHONNEAU, D. NGUYEN, N. RAYMOND & S. VU NGOC – “Magnetic WKB constructions on surfaces”, *Rev. Math. Phys.* **33** (2021), no. 7, article no. 2150022 (41 pages).
- [GRV21] Y. GUEDES BONTHONNEAU, N. RAYMOND & S. VU NGOC – “Exponential localization in 2d pure magnetic wells”, *Ark. Mat.* **59** (2021), no. 1, p. 53–85.
- [HKRV16] B. HELFFER, Y. KORDYUKOV, N. RAYMOND & S. VU NGOC – “Magnetic wells in dimension three”, *Anal. PDE* **9** (2016), no. 7, p. 1575–1608.
- [Hör67] L. HÖRMANDER – “Hypoelliptic second order differential equations”, *Acta Math.* **119** (1967), p. 147–171.
- [LS23] C. LETROUIT & C. SUN – “Observability of Baouendi-Grushin-type equations through resolvent estimates”, *J. Inst. Math. Jussieu* **22** (2023), no. 2, p. 541–579.
- [Mel85] R. B. MELROSE – “The wave equation for a hypoelliptic operator with symplectic characteristics of codimension two”, *J. Analyse Math.* **44** (1984/85), p. 134–182.
- [Mil96] L. MILLER – “Propagation d’ondes semi-classiques à travers une interface et mesures 2-microlocales”, PhD Thesis, École polytechnique, 1996.
- [Mor22] L. MORIN – “A semiclassical Birkhoff normal form for symplectic magnetic wells”, *J. Spectral Theory* **12** (2022), no. 2, p. 459–496.
- [Mor24] ———, “A semiclassical Birkhoff normal form for constant-rank magnetic fields”, *Anal. PDE* **17** (2024), no. 5, p. 1593–1632.
- [Nie96] F. NIER – “A semi-classical picture of quantum scattering”, *Ann. Sci. École Norm. Sup. (4)* **29** (1996), no. 2, p. 149–183.
- [PSU23] G. P. PATERNAIN, M. SALO & G. UHLMANN – *Geometric inverse problems—with emphasis on two dimensions*, Cambridge Studies in Advanced Math., vol. 204, Cambridge University Press, Cambridge, 2023.
- [RV15] N. RAYMOND & S. VU NGOC – “Geometry and spectrum in 2D magnetic wells”, *Ann. Inst. Fourier (Grenoble)* **65** (2015), no. 1, p. 137–169.
- [RS72] M. REED & B. SIMON – *Methods of modern mathematical physics. I. Functional analysis*, Academic Press, New York-London, 1972.
- [RS75] ———, *Methods of modern mathematical physics. II. Fourier analysis, self-adjointness*, Academic Press, New York-London, 1975.
- [RS76] L. P. ROTHSCHILD & E. M. STEIN – “Hypoelliptic differential operators and nilpotent groups”, *Acta Math.* **137** (1976), no. 3-4, p. 247–320.
- [Rum94] M. RUMIN – “Differential forms on contact manifolds”, *J. Differential Geom.* **39** (1994), no. 2, p. 281–330.
- [Tay20] M. TAYLOR – “Microlocal Weyl formula on contact manifolds”, *Comm. Partial Differential Equations* **45** (2020), no. 5, p. 392–413.
- [Zwo12] M. ZWORSKI – *Semiclassical analysis*, Graduate Studies in Math., vol. 138, American Mathematical Society, Providence, RI, 2012.

Manuscript received 30th June 2023

accepted 30th July 2024

VICTOR ARNAIZ, Laboratoire de Mathématiques Jean Leray, Nantes Université, UMR CNRS 6629,
2 rue de la Houssinière, 44322 Nantes Cedex 03, France

E-mail : victor.arnaiz@univ-nantes.fr

E-mail : victor.arnaiz-solorzano@math.u-bordeaux.fr

GABRIEL RIVIÈRE, Laboratoire de Mathématiques Jean Leray, Nantes Université, UMR CNRS 6629,
2 rue de la Houssinière, 44322 Nantes Cedex 03, France

& Institut Universitaire de France, Paris, France

E-mail : gabriel.riviere@univ-nantes.fr

Url : <https://www.math.sciences.univ-nantes.fr/~riviere-g>