



INSTITUT  
POLYTECHNIQUE  
DE PARIS

# *J*ournal de l'École polytechnique *Mathématiques*


Gustave BILLON

**Moduli spaces of marked branched projective structures on surfaces**

Tome II (2024), p. 1373-1410.

<https://doi.org/10.5802/jep.279>

© L'auteur, 2024.



Cet article est mis à disposition selon les termes de la licence  
LICENCE INTERNATIONALE D'ATTRIBUTION CREATIVE COMMONS BY 4.0.  
<https://creativecommons.org/licenses/by/4.0/>

Publié avec le soutien  
du Centre National de la Recherche Scientifique



Publication membre du  
Centre Mersenne pour l'édition scientifique ouverte  
[www.centre-mersenne.org](http://www.centre-mersenne.org)  
e-ISSN : 2270-518X

# MODULI SPACES OF MARKED BRANCHED PROJECTIVE STRUCTURES ON SURFACES

BY GUSTAVE BILLON

**ABSTRACT.** — We show that the moduli space  $\mathcal{P}_g(n)$  of marked branched projective structures of genus  $g$  and branching degree  $n$  is a complex analytic space. In the case  $g \geq 2$ , we show that  $\mathcal{P}_g(n)$  is of dimension  $6g - 6 + n$  and we characterize its singular points in terms of their monodromy. We introduce a notion of branching class, that is an infinitesimal description of branched projective structures at the branched points. We show that the space  $\mathcal{A}_g(n)$  of marked branching classes of genus  $g$  and branching degree  $n$  is a complex manifold. We show that if  $n < 2g - 2$  the space  $\mathcal{P}_g(n)$  is an affine bundle over  $\mathcal{A}_g(n)$ , while if  $n > 4g - 4$ ,  $\mathcal{P}_g(n)$  is an analytic subspace of  $\mathcal{A}_g(n)$ .

**RÉSUMÉ** (Espaces de modules de structures projectives branchées sur les surfaces)

On montre que l'espace de modules  $\mathcal{P}_g(n)$  des structures projectives branchées de genre  $g$  et de degré de branchement  $n$  est un espace analytique complexe. Dans le cas où  $g \geq 2$ , on montre que  $\mathcal{P}_g(n)$  est de dimension  $6g - 6 + n$  et on caractérise ses points singuliers en termes de leur monodromie. On introduit une notion de classe de branchement, qui est une description infinitésimale des structures projectives branchées aux points de branchement. On montre que l'espace  $\mathcal{A}_g(n)$  des classes de branchement marquées de genre  $g$  et de degré de branchement  $n$  est une variété différentielle complexe. On montre que si  $n < 2g - 2$ , l'espace  $\mathcal{P}_g(n)$  est un fibré affine sur  $\mathcal{A}_g(n)$ , tandis que si  $n > 4g - 4$ ,  $\mathcal{P}_g(n)$  est un sous-espace analytique de  $\mathcal{A}_g(n)$ .

## CONTENTS

Introduction.....	1373
1. Branching classes.....	1378
2. Families of branched projective structures over a family of curves.....	1382
3. Holonomy and singularities of the moduli space of marked branched projective structures in genus at least two.....	1396
References.....	1409

## INTRODUCTION

**DEFINITION AND EXAMPLES.** — Holomorphic projective structures were introduced at the end of the nineteenth century in relation to linear differential equations of order 2. They were used in particular by Poincaré, as an analogous for curves of genus at least 2 of elliptic functions, in his proof of the uniformization theorem of Riemann surfaces.

**MATHEMATICAL SUBJECT CLASSIFICATION (2020).** — 57M50, 14H15, 32G15, 14H30.

**KEYWORDS.** — Complex projective structures, moduli spaces, families of Riemann surfaces.

They were then thoroughly studied throughout the twentieth century, in particular by Gunning (see for instance [Gun66]).

In [Man72], Mandelbaum introduced a notion of *branched* projective structures, that can be seen as a generic generalization of the concept of projective structure.

Let us give a precise definition. Fix an oriented surface  $S$ .

**DEFINITION.** — A *branched projective structure* on  $S$  is the datum of

- an open cover  $(U_i)_{i \in I}$  of  $S$ ,
  - for each  $i \in I$ , an open subset  $V_i \subset \mathbb{CP}^1$  and a ramified covering  $f_i : U_i \rightarrow V_i$  such that on the intersections  $U_i \cap U_j$ , one has  $f_i = g_{ij} \circ f_j$ , where  $g_{ij} \in \mathrm{PGL}(2, \mathbb{C})$ .
- The number of ramification points of the maps  $f_i$ , counted with multiplicity, is called the *branching degree* of the branched projective structure.

In the case where the branching degree is zero, one gets *projective structures*, see [Dum09] and [LMP09] for overview papers. Here are a few examples of branched projective structures:

*Tautological projective structure.* — The Riemann sphere  $\mathbb{CP}^1$  is endowed with a trivial projective structure, of branching degree zero, given by the identity map.

*Uniformizing projective structure.* — Suppose the surface  $S$  is closed, and let  $X$  be a Riemann surface with underlying differential surface  $S$ . Let  $\pi : \tilde{X} \rightarrow X$  be the universal covering of  $X$ . By the uniformization theorem, there are inclusions  $\tilde{X} \subset \mathbb{CP}^1$  and  $\mathrm{Aut}(\pi) \subset \mathrm{PGL}(2, \mathbb{C})$ . As a consequence, the local inverses of the map  $\pi$  are charts of a projective structure on  $X$ , without ramification point, called the *uniformizing projective structure* of  $X$ .

*Pullback by a ramified covering.* — Let  $S'$  be another oriented surface endowed with a branched projective structure  $(V_i, h_i)_{i \in I}$ , and let  $\phi : S \rightarrow S'$  be a ramified covering. Then  $S$  inherits a *pullback* projective structure: the open covering is  $(\phi^{-1}(V_i))_{i \in I}$  and the family of charts is  $(h_i \circ \phi)_{i \in I}$ . In particular, a ramified covering of the Riemann sphere can be seen as a branched projective structure.

*Curve not preserved by a foliation.* — There is a notion of projective structure *transverse to a foliation*, see [Blu79], [God91], [Scá97]. Precisely, let  $M$  be a complex manifold and  $\mathcal{F}$  a codimension 1 holomorphic foliation on  $M$ . A projective structure transverse to  $\mathcal{F}$  is an open cover  $(\mathcal{U}_i)_{i \in I}$  of  $M$  and holomorphic submersions  $s_i : \mathcal{U}_i \rightarrow \mathbb{CP}^1$ , constant on the leaves of  $\mathcal{F}$ , such that on the intersections  $\mathcal{U}_i \cap \mathcal{U}_j$  one has  $s_i = g_{ij} \circ s_j$  with  $g_{ij} \in \mathrm{PGL}(2, \mathbb{C})$ . Now if  $X \subset M$  is a complex curve that is not invariant by  $\mathcal{F}$ , then  $X$  inherits a branched projective structure, whose branching degree is the number of tangencies between  $X$  and  $\mathcal{F}$ .

*Linear differential equations of order 2.* — Projective structures with branching degree zero can be obtained from equations of the form  $a(x)y''(x) + b(x)y'(x) + c(x)y(x) = 0$  where  $x$  is a local coordinate,  $y$  is an unknown holomorphic function and  $a, b, c$  are holomorphic functions, with  $a$  nonvanishing. Given a basis  $(y_1, y_2)$  of local solutions, the quotient  $y_1/y_2$  is a local biholomorphism from an open subset of  $X$  to  $\mathbb{CP}^1$ .

If  $(w_1, w_2)$  is another basis of local solutions, then there exists  $\alpha, \beta, \gamma, \delta \in \mathbb{C}$  with  $\alpha\delta - \beta\gamma \neq 0$  such that  $w_1 = \alpha y_1 + \beta y_2$  and  $w_2 = \gamma y_1 + \delta y_2$ . Thus there exists a Möbius transformation  $g = \begin{pmatrix} \alpha & \beta \\ \gamma & \delta \end{pmatrix}$  such that  $w_1/w_2 = g \circ (y_1/y_2)$ . In other words, the equation defines a projective structure on  $X$ .

This description of projective structures gives rise to a notion of *meromorphic projective structure*, corresponding to order 2 differential equations with meromorphic coefficients. This notion was studied in particular in [AB20], [GM21], [GM20] and [Sér22]. Branched projective structures are very special meromorphic projective structures, that are usually excluded from the studies of general meromorphic projective structures.

Note that the notion of linear differential equation on Riemann surfaces is formalized in the notion of *oper*, see [BD05]. In this language, projective structures are  $\mathrm{PGL}(2, \mathbb{C})$ -opers. See [Fre07] for the role of opers in the Langlands program. This article thus deals with *branched*  $\mathrm{PGL}(2, \mathbb{C})$ -opers. The notion of branched oper has been investigated in [FG10] and [BDH22]. See also [BDG19] for the link between branched projective structures and logarithmic connections.

*Topological constructions.* — Branched projective structures can also be modified by cut-and-paste techniques, some preserving the branching degree, such as *grafting* or moving branch points, others changing the branching degree, such as *bubbling*. See for instance [Dum09], [GKM00] or [CDF14]. These cut-and-paste techniques allow to construct branched projective structures of any degree.

*Main result.* — Suppose the surface  $S$  is closed, of genus  $g$ . Fix a nonnegative integer  $n \in \mathbb{N}$ . There is a notion of *marked* branched projective structure, similar to the notion of marked complex structure on  $S$  (see 2.3 for precise definitions). We are interested in the following space:

$$\mathcal{P}_g(n) = \frac{\{\text{marked branched projective structures on } S\}}{\{\text{isomorphisms}\}}.$$

Given a marked branched projective structure  $(U_i, f_i)_{i \in I}$  with  $f_i = g_{ij} \circ f_j$ ,  $g_{ij} \in \mathrm{PSL}(2, \mathbb{C})$ , the changes of charts  $g_{ij}$  are holomorphic maps, thus the (ramified) atlas  $(U_i, f_i)_{i \in I}$  defines a marked complex structure on  $S$ . Denote by  $\mathcal{T}_g$  the Teichmüller space for genus  $g$ , i.e., the space of isomorphism classes of marked Riemann surfaces of genus  $g$ . One has a forgetful map

$$\mathrm{Teich}_g(n) : \mathcal{P}_g(n) \longrightarrow \mathcal{T}_g.$$

Moreover, the family  $(g_{ij})_{i,j \in I}$  is a constant cocycle with values in  $\mathrm{PSL}(2, \mathbb{C})$ , and thus defines a flat  $\mathbb{C}P^1$ -bundle on  $X$ , to which is associated a holonomy representation  $\rho \in \mathrm{Hom}(\pi_1(S), \mathrm{PSL}(2, \mathbb{C}))$ , defined up to conjugation with an element of  $\mathrm{PSL}(2, \mathbb{C})$ . Thus there is a holonomy map

$$\mathrm{Hol}_g(n) : \mathcal{P}_g(n) \longrightarrow \mathrm{Hom}(\pi_1(S), \mathrm{PSL}(2, \mathbb{C}))/\mathrm{PSL}(2, \mathbb{C}).$$

The holonomy of branched projective structures has been extensively studied, see [GKM00], [CDF14], [LF23].

In the unbranched case, i.e., in the case  $n = 0$ , if  $g \geq 2$ , it is well-known ([Gun66], [Hej75], [Hub81]) that the space  $\mathcal{P}_g(0)$  is a smooth analytic space. Moreover the fiber of  $\text{Teich}_g(0)$  at a marked complex curve  $X \in \mathcal{T}_g$  is an affine space, directed by the space  $H^0(X, K_X^{\otimes 2})$  of global holomorphic quadratic differentials on  $X$ , and the holonomy map  $\text{Hol}_g(0)$  is a local biholomorphism. This very nice structure of the moduli space of projective structures is one of the main reasons why they are extensively used.

The main result of this paper is an analog of these properties in the branched case (see Theorems 2.3.7, 3.0.1 and 3.5.1 for more precise statements).

**THEOREM A.** — *The space  $\mathcal{P}_g(n)$  is an analytic space of dimension  $6g - 6 + n$ . Moreover, if  $g \geq 2$ , one has:*

- (i) *A point  $p \in \mathcal{P}_g(n)$  is singular if and only if its holonomy representations  $\text{Hol}_g(n)(p)$  are abelian and fix a point in  $\mathbb{CP}^1$ .*
- (ii) *The map  $\text{Hol}_g(n)$  is a holomorphic submersion over branched projective structures with non-elementary holonomy.*

Recall that a representation  $\rho \in \text{Hom}(\pi_1(S), \text{PSL}(2, \mathbb{C}))$  is said to be *elementary* if its image is an elementary subgroup of  $\text{PSL}(2, \mathbb{C})$ . A subgroup of  $\text{PSL}(2, \mathbb{C})$  is elementary if its action on the hyperbolic space  $\mathbb{H}^3$  admits a finite orbit. Namely, elementary subgroups of  $\text{PSL}(2, \mathbb{C})$  are the ones that fix a point in  $\mathbb{CP}^1$ , the ones that fix a pair of points in  $\mathbb{CP}^1$  (conjugated to a subgroup of

$$\{z \mapsto az \mid a \in \mathbb{C}^*\} \cup \{z \mapsto a/z \mid a \in \mathbb{C}^*\},$$

and the ones that fix a point in the interior of  $\mathbb{H}^3$  (spherical subgroups). Note that the condition of having non-elementary holonomy in point (ii) is not optimal, see Theorem 3.5.1.

Theorem A answers a question asked in the appendix of [CDF14], where it is shown that the fibers of  $\text{Hol}_g(n)$  over non-elementary representations are smooth analytic spaces.

Let us recall the results obtained by Mandelbaum in his papers introducing branched projective structures, that are related to Theorem A. In [Man72], Mandelbaum fixes a Riemann surface  $X$  of genus  $g$  as well as an effective divisor  $D = \sum_{i=0}^r k_i x_i$  on  $X$ , where the  $x_i$  are distinct points of  $X$ . He considers the set  $P_X(D)$  of *holomorphic* branched projective structures (i.e., branched projective structures with holomorphic charts) on  $X$  with ramifications at the points of  $D$ . He shows that  $P_X(D)$  is an analytic space, that is of dimension  $3g - 3$  in the case where  $\deg D \leq 2g - 2$  or when the divisor  $D$  is supported on a single point. In [Man73], he shows that, when the complex structure on  $X$  and the distinct points  $x_i$  of  $D$  vary (but not the coefficients  $k_i$ ), the analytic spaces  $P_X(D)$  can be glued together, so that their union is itself an analytic space. Note that these results can be seen as consequences of the results in this paper, except the computations of the dimension of  $P_X(D)$  when  $\deg D = 2g - 2$  or when the divisor  $D$  is supported on a single point. When  $\deg D < 2g - 2$ , Proposition 2.4.2 shows that  $P_X(D)$  is a *smooth* analytic space of dimension  $3g - 3$ . Finally, note that Theorem A shows that the spaces  $P_X(D)$  glue

together even when the coefficients  $k_i$  vary, the only invariants being the genus  $g$  and the degree  $\deg D$ . See also [Man75] about branched projective structures.

The key point in the proof of Theorem A is what we call *branching classes*. Their existence is due to the fact that, unlike local biholomorphisms, local branched coverings are not all obtained from one another by postcomposition with a local biholomorphism. See Section 1 for details.

*Structure of the paper.* — In Section 1, we introduce the notion of branching class over a curve with a divisor. We investigate the reasons why branching classes are a useful tool for our purpose. These reasons are the very nice structure of the space of branching classes over a fixed curve with divisor, and the very nice structure of the space of branching classes with fixed branching class.

In Section 2, we define the moduli spaces of branched projective structures  $\mathcal{P}_g(n)$ , and we exhibit its analytic structure. We consider in turn the analytic structures of spaces of marked curves (the Teichmüller spaces), the spaces of marked curves with divisors and the spaces of marked branching classes, before being able to endow  $\mathcal{P}_g(n)$  with an analytic structure. We also have to introduce a technical notion of *restricted* branched projective structures, that we use as an intermediate between branching classes and branched projective structures. We emphasize in this section the universal properties of the moduli spaces we consider, working with *relative* branched projective structures, or equivalently families of such structures, in the spirit of [Hub81].

In Section 3 we study  $\mathcal{P}_g(n)$  when the genus  $g$  is at least 2. Most of this section is dedicated to the characterization of the singular points of  $\mathcal{P}_g(n)$ , and follows the same structure as Section 2, considering in turn the moduli spaces of curves, divisors, branching classes, restricted branched projective structures and finally branched projective structures. The last part of Section 3 is dedicated to the study of the holonomy map. The main tool for this section, in the spirit of [Hub81], is the Kodaira-Spencer machinery that allows to identify the tangent space of a moduli space at a point with the first cohomology group of infinitesimal automorphisms of this point.

*Notations.* — If  $X$  is a Riemann surface,  $T_X$  denotes the tangent bundle of  $X$  and  $K_X$  denotes its cotangent bundle, which is also the canonical bundle of  $X$ .

If  $E$  is a holomorphic vector bundle on a complex manifold, we denote also by  $E$  its sheaf of holomorphic sections.

Let  $z$  be a local coordinate on an open subset  $U$  of a Riemann surface, and  $\phi = \varphi(z)$  a non constant holomorphic function on  $U$ . The two following differential operators are studied in [Gun66]:

- (i) We denote by  $[\phi, z]$  the *affine distortion* of  $\phi$  in the coordinate  $z$ :  $[\phi, z] = \varphi''/\varphi'$ .
- (ii) We denote by  $\{\phi, z\}$  the *Schwarzian derivative* of  $\phi$  in the coordinate  $z$ :  $\{\phi, z\} = (\varphi''/\varphi')' - \frac{1}{2}(\varphi''/\varphi')^2$ .

*Acknowledgements.* — I wish to thank my adviser Sorin Dumitrescu for his help and support. I also thank Bertrand Deroin and Frank Loray for fruitful conversations. I thank the referee for helpful comments.

## 1. BRANCHING CLASSES

**1.1. THE SPACE OF BRANCHING CLASSES ON A CURVE WITH DIVISOR.** — Let  $X$  be a Riemann surface,  $x \in X$  and  $n \in \mathbb{N}$ . Denote by  $(X, x)$  the germ of  $X$  at  $x$ , and by  $R_x^n$  the complex manifold of  $2(n+1)$ -jets of  $(n+1)$ -sheeted ramified coverings from  $(X, x)$  to  $\mathbb{CP}^1$ :

$$(1) \quad R_x^n = \left\{ j_x^{2(n+1)} \phi \left| \begin{array}{l} \phi \text{ germ at } x \text{ of holomorphic } (n+1)\text{-fold} \\ \text{ramified covering with values in } \mathbb{CP}^1 \end{array} \right. \right\}.$$

The group of Möbius transformations  $\mathrm{PSL}(2, \mathbb{C})$  acts on  $R_x^n$  by postcomposition.

**PROPOSITION 1.1.1.** — *The group  $\mathrm{PSL}(2, \mathbb{C})$  acts freely on  $R_x^n$ . Moreover, the quotient  $\mathrm{PSL}(2, \mathbb{C}) \backslash R_x^n$  is a complex manifold of dimension  $n$ , isomorphic to  $\mathbb{C}^n$ .*

*Proof.* — Take  $z$  a local coordinate on  $X$  centered at  $x$ . In an affine coordinate  $w$  of  $\mathbb{CP}^1 = \mathbb{C} \cup \{\infty\}$ , an element  $j_x^{2(n+1)} \phi \in R_x^n$  can be written:

$$j_x^{2(n+1)} \phi = a_0 + a_{n+1}z^{n+1} + a_{n+2}z^{n+2} + \cdots + a_{2(n+1)}z^{2(n+1)}$$

with  $a_{n+1} \neq 0$ . Recall that the datum of a Möbius transformation  $g \in \mathrm{PSL}(2, \mathbb{C})$  is equivalent to the datum of its 2-jet at  $a_0$ . If

$$g = \alpha_0 + \alpha_1(w - a_0) + \alpha_2(w - a_0)^2 + O((w - a_0)^3),$$

the action of  $g$  on  $j_x^{2(n+1)} \phi$  writes

$$\begin{aligned} g \cdot j_x^{2(n+1)} \phi &= j_x^{2(n+1)} g \circ \phi \\ &= (\alpha_0 + a_0) + \alpha_1 a_{n+1} z^{n+1} + \alpha_1 a_{n+2} z^{n+2} + \cdots + \alpha_1 a_{2n+1} z^{2n+1} \\ &\quad + (\alpha_1 a_{2(n+1)} + \alpha_2 a_{n+1}^2) z^{2(n+1)}. \end{aligned}$$

Note that the coordinate  $w$  can be chosen so that neither  $\phi(x)$  nor  $g(a_0)$  is  $\infty$  in the coordinate  $w$ . Since  $a_{n+1} \neq 0$ , the coefficients of degree 0,  $n+1$  and  $2(n+1)$  of  $g \cdot j_x^{2(n+1)} \phi$  determine the 2-jet of  $g \in \mathrm{PSL}(2, \mathbb{C})$  at  $a_0$ , thus  $g$  itself, showing that  $\mathrm{PSL}(2, \mathbb{C})$  acts freely on  $R_x^n$ . Moreover the orbit of  $j_x^{2(n+1)} \phi$  under the action of  $\mathrm{PSL}(2, \mathbb{C})$  is given by the complex parameters

$$(a_{n+2}/a_{n+1}, a_{n+3}/a_{n+1}, \dots, a_{2n+1}/a_{n+1}),$$

showing that the quotient  $\mathrm{PSL}(2, \mathbb{C}) \backslash R_x^n$  is isomorphic to  $\mathbb{C}^n$ .  $\square$

Consider now  $D = \sum_{i \in I} n_i x_i$  an effective divisor of degree  $n$  on  $X$ . A *branching class* on  $X$  of divisor  $D$  is the choice for each point  $x_i$  of  $D$  of an orbit for the action of  $\mathrm{PSL}(2, \mathbb{C})$  on  $R_{x_i}^{n_i}$ . The space  $A_X^D$  of branching classes over  $(X, D)$  is thus given by

$$A_X^D := \prod_{i \in I} \mathrm{PSL}(2, \mathbb{C}) \backslash R_{x_i}^{n_i}.$$

If  $D$  has finite degree, for instance if  $X$  is compact,  $A_X^D$  is a complex manifold of dimension  $n = \sum_{i=1}^r n_i$ .

**PROPOSITION 1.1.2.** — *The space  $A_X^D$  of branching classes over a Riemann surface with divisor  $(X, D)$  is an affine space directed by the vector space  $H^0(X, K_X|_D)$  of jets of holomorphic differentials at the points of  $D$ .*

*Proof.* — Let  $z$  be a local coordinate defined on an open subset  $U \subset X$ . Let  $x_0 \in U$  and let  $\phi_1 = \varphi_1(z), \phi_2 = \varphi_2(z)$  be non constant holomorphic functions on  $U$  such that  $\phi_1, \phi_2$  are both ramified at  $x_0$  with  $(n+1)$  branches and have no other ramification points. Recall that  $[\phi_1, z]$  denotes the affine distortion  $\varphi_1''/\varphi_1'$ . A straightforward calculation shows the following (see also [Gun66]):

(i) The 1-form  $[\phi_1, z]dz$  is meromorphic on  $U$ , holomorphic away from  $x_0$  and with a simple pole at  $x_0$  with residue  $n$ . In particular, the difference  $[\phi_2, z]dz - [\phi_1, z]dz$  is a holomorphic 1-form on  $U$ .

(ii) The holomorphic 1-form  $[\phi_2, z]dz - [\phi_1, z]dz$  does not depend on the choice of the holomorphic coordinate  $z$  on  $U$ .

(iii) If  $\omega$  is a holomorphic 1-form on  $U$ , there exists a neighborhood  $V$  of  $x_0$  in  $U$  and a non constant holomorphic function  $\phi$  on  $V$  whose only ramification point is  $x_0$ , where it has  $(n+1)$  branches, such that  $\omega = [\phi, z]dz - [\phi_1, z]dz$ .

(iv) The  $n$ -jet  $j_{x_0}^n([\phi_2, z]dz - [\phi_1, z]dz)$  only depends on the  $2n+1$ -jet  $j_{x_0}^{2n+1}\phi_1$ .

(v) One has  $j_{x_0}^n([\phi_2, z]dz - [\phi_1, z]dz) = 0$  if and only if there exists a local biholomorphism  $\gamma$  such that  $j_{x_0}^{2n+1}\phi_2 = j_{x_0}^{2n+1}\gamma \circ \phi_1$ .

These five properties imply that the space  $A_U^{n, x_0}$  is an affine space, directed by the vector space  $H^0(X, K_U|_{n, x_0})$  of  $n$ -jets of holomorphic 1-forms at  $x_0$ . The difference between the class of  $j_{x_0}^{2(n+1)}\phi_2$  and the class of  $j_{x_0}^{2(n+1)}\phi_1$  is given by

$$j_{x_0}^n([\phi_2, z]dz - [\phi_1, z]dz) \in H^0(X, K_U|_{n, x_0}).$$

Write  $D = \sum_{i \in I} n_i x_i$ , and take a family of open sets  $(U_i)_{i \in I}$  such that  $x_i \in U_i$  and if  $i \neq j$ ,  $U_i \cap U_j = \emptyset$ . Then  $A_X^D = \prod_{i \in I} A_{U_i}^{n_i, x_i}$  is an affine space directed by the vector space

$$\bigoplus_{i \in I}^r H^0(U_i, K_{U_i}|_{n_i, x_i}) = H^0(X, K_X|_D). \quad \square$$

Let us end this paragraph by fixing some notations. Let  $D = \sum_{x \in X} n_x x$  be a divisor on a Riemann surface  $X$ ,

$$a = (a_x)_{x \in X} \in \prod_{x \in X} \mathrm{PSL}(2, \mathbb{C}) \backslash R_x^{n_x} = A_X^D$$

a branching class over  $(X, D)$  and  $U \subset X$  an open subset. In what follows, we denote by  $D|_U = \sum_{x \in U} n_x x$  the divisor  $D$  restricted to  $U$  and by

$$a|_U = (a_x)_{x \in U} \in \prod_{x \in U} \mathrm{PSL}(2, \mathbb{C}) \backslash R_x^{n_x} = A_U^{D|_U}$$

the branching class  $a$  restricted to  $U$ .



1.2. BRANCHED PROJECTIVE STRUCTURES AND BRANCHING CLASSES. — Let us first recall the definition of a branched projective structure.

DEFINITION 1.2.1. — A *branched projective atlas* on a Riemann surface  $X$  is given by an open cover  $(U_i)_{i \in I}$  of  $X$ , and for each  $i \in I$  a non constant holomorphic map  $f_i : U_i \rightarrow \mathbb{CP}^1$  such that for any  $i, j \in I$  with  $U_i \cap U_j \neq \emptyset$  there is a Möbius transformation  $g_{ij} \in \mathrm{PSL}(2, \mathbb{C})$  such that  $f_i|_{U_i \cap U_j} = g_{ij} \circ f_j|_{U_i \cap U_j}$ .

Two branched projective atlases are said to be *equivalent* if their union is also a branched projective atlas.

A *branched projective structure* on  $X$  is an equivalence class of branched projective atlases.

Let  $X$  be a Riemann surface, let  $p$  be a branched projective structure on  $X$  and  $(U_i, f_i)_{i \in I}$  an atlas for  $p$ . On the intersections  $U_i \cap U_j$ ,  $f_i$  is obtained from  $f_j$  by postcomposition with a Möbius transformation, so the vanishing orders of the differentials  $df_i$  and  $df_j$  are the same at any point of  $U_i \cap U_j$ . Moreover at any point  $x \in U_i \cap U_j$ , if  $n$  is the (possibly zero) vanishing order of  $df_i$  (and thus of  $df_j$ ) at  $x$ , then  $f_i$  and  $f_j$  both determine elements of  $R_x^n$  in (1) that are in the same orbit under the action of  $\mathrm{PSL}(2, \mathbb{C})$ .

The branched projective structure  $p$  thus determines two additional objects on  $X$ :

- (i) An effective divisor  $D$ , called the *branching divisor* of  $p$ , given by the vanishing locus of the differentials of the charts of  $p$ :  $D = \sum_{x \in X} n_x x$  where  $n_x$  is the vanishing order at  $x$  of the charts of  $p$  defined at  $x$ .
- (ii) A *branching class*  $a \in A_X^D$  on  $X$  of divisor  $D$ , given by the orbits of the jets of charts of  $p$  at the points of  $D$ .

For a branching class  $a$  over  $(X, D)$ , let us denote by  $P_X^a$  the set of branched projective structures on  $X$  with branching class  $a$  (thus with branching divisor  $D$ ).

PROPOSITION 1.2.2. — Let  $D$  be an effective divisor on a Riemann surface  $X$ . Let  $a \in A_X^D$  be a branching class over  $(X, D)$ .

- (i) There exists an obstruction  $c(a) \in H^1(X, K_X^{\otimes 2}(-D))$  such that  $c(a) = 0$  if and only if  $P_X^a \neq \emptyset$ .
- (ii) The set  $P_X^a$ , if nonempty, is an affine space directed by the vector space  $H^0(X, K_X^{\otimes 2}(-D))$  of holomorphic quadratic differentials on  $X$  vanishing over the divisor  $D$ .

Let us state point (i) in a slightly more general framework. Let  $\mathcal{S}$  be a sheaf of abelian groups on  $X$ . By a *sheaf of affine spaces directed by  $\mathcal{S}$*  we mean a sheaf  $\mathfrak{S}$  such that if  $U \subset X$  is an open subset, the space of sections  $\Gamma(U, \mathfrak{S})$  is either empty or an affine space directed by  $\Gamma(U, \mathcal{S})$ . We say moreover that  $\mathfrak{S}$  is *locally nonempty* if for any  $x \in X$  there exists an open neighborhood  $U$  of  $x$  such that  $\Gamma(U, \mathfrak{S}) \neq \emptyset$ .

LEMMA 1.2.3. — Let  $\mathcal{S}$  be a sheaf of abelian groups over the Riemann surface  $X$  and  $\mathfrak{S}$  a locally nonempty sheaf of affine spaces directed by  $\mathcal{S}$ . There exists a cohomology class

$c(\mathfrak{S}) \in H^1(X, \mathbb{S})$  that is an obstruction to the existence of a global section of  $\mathfrak{S}$ . More explicitly,  $c(\mathfrak{S}) = 0$  if and only if  $H^0(X, \mathfrak{S}) \neq 0$ .

*Proof.* — Since  $\mathfrak{S}$  is locally nonempty, there exists an open cover  $(U_i)_{i \in I}$  of  $X$  such that for each  $i \in I$  there is a local section  $s_i \in \Gamma(U_i, \mathfrak{S})$ . For any  $i, j \in I$ , write  $c_{ij} = s_j - s_i \in \Gamma(U_{ij}, \mathbb{S})$ , where  $U_{ij} = U_i \cap U_j$ . The family  $(c_{ij})_{i, j \in I}$  is clearly a 1-cocycle for  $\mathbb{S}$  and thus defines a cohomology class  $c(\mathfrak{S}) \in H^1(X, \mathbb{S})$ .

The class  $c(\mathfrak{S})$  does not depend on the choice of the family  $(s_i)_{i \in I}$ . Indeed, if  $(s'_i)_{i \in I}$  is another such family and  $c'_{ij} = s'_j - s'_i$  the corresponding cocycle, then one has  $c'_{ij} - c_{ij} = (s'_j - s_j) - (s'_i - s_i)$ , so that the cohomology class of the cocycle  $(c'_{ij} - c_{ij})_{i, j \in I}$  is 0. In particular, if  $H^0(X, \mathfrak{S}) \neq 0$  and  $s$  is a global section of  $\mathfrak{S}$ , then one can take  $s_i = s|_{U_i}$  in which case  $c_{ij} = 0$  thus  $c(\mathfrak{S}) = 0$ .

Conversely, if  $c(\mathfrak{S}) = 0$ , then up to refining the open cover  $(U_i)_{i \in I}$ , there is a family  $(t_i)_{i \in I}$  such that  $t_i \in \Gamma(U_i, \mathbb{S})$  and  $c_{ij} = t_j - t_i$ . In that case, write  $\tilde{s}_i = s_i - t_i$ . The sections  $\tilde{s}_i$  coincide on the intersections  $U_{ij}$  and thus glue together to provide a global section of  $\mathfrak{S}$ .  $\square$

*Proof of Proposition 1.2.2.* — Let us first focus on the affine structure of  $P_X^a$  in case it is nonempty. Let  $z$  be a local coordinate defined on an open subset  $U \subset X$ . Let  $x_0 \in U$  and let  $a_0 \in A_X^{n, x_0}$  be a branching class of order  $n$  at  $x_0$ . Let also  $\phi_1 = \varphi_1(z), \phi_2 = \varphi_2(z)$  be non constant holomorphic functions on  $U$  such that  $\phi_1, \phi_2$  are both ramified at  $x_0$  with  $(n+1)$  branches and branching class  $a_0$ . Suppose  $\phi_1$  and  $\phi_2$  have no other ramification point. Recall that  $\{\phi_1, z\}$  denotes the Schwarzian derivative  $(\varphi_1''/\varphi_1')' - \frac{1}{2}(\varphi_1''/\varphi_1')^2$ . It is well-known (see for instance [Gun66]) that the difference  $\{\phi_2, z\} - \{\phi_1, z\}$  does not depend on the choice of the coordinate  $z$ . Moreover, this difference vanishes uniformly on  $U$  if and only if there is a Möbius transformation  $g \in \text{PSL}(2, \mathbb{C})$  such that  $\phi_2 = g \circ \phi_1$ . A straightforward calculation shows the following:

- (i) The quadratic differential  $\{\phi_1, z\} dz^{\otimes 2}$  is meromorphic on  $U$ , holomorphic away from  $x_0$  and admits a pole of order 2 at  $x_0$ .
- (ii) The coefficient of order  $-2$  at  $x_0$  of  $\{\phi_1, z\} dz^{\otimes 2}$  is  $(1 - (n+1)^2)/2$ .
- (iii) The coefficients of order  $-1, 0, 1, \dots, n-1$  at  $x_0$  of  $\{\phi_1, z\} dz^{\otimes 2}$  determine and are determined by the branching class of  $\phi_1$  at  $x_0$ . In particular, the difference  $\{\phi_2, z\} dz^{\otimes 2} - \{\phi_1, z\} dz^{\otimes 2}$  is a holomorphic quadratic differential on  $U$  with a zero of order  $n$  at  $x_0$ .

Moreover, it is a consequence of Fuchs's local theory (see for instance [Man72] or [dSG16, §IX.1]) that if  $\omega$  is a holomorphic quadratic differential on  $U$  with a zero of order  $n$  at  $x_0$ , then on an open neighborhood  $V$  of  $x_0$  in  $U$ , there exists a function  $\phi_3$  such that  $\omega = \{\phi_3, z\} dz^{\otimes 2} - \{\phi_1, z\} dz^{\otimes 2}$ . The function  $\phi_3$  is ramified at  $x_0$ , with  $(n+1)$  branches and branching class  $a_0$ .

Now, suppose  $P^a \neq \emptyset$  and let  $p_1, p_2 \in P_X^a$ . Let respectively  $(U_i, \phi_i)$  and  $(U_i, \psi_i)$  be atlases for  $p_1$  and  $p_2$  and for each  $i \in I$  let  $z_i$  be a coordinate on  $U_i$ . The above facts about Schwarzian derivatives imply that the local holomorphic quadratic differentials

$\{\psi_i, z_i\}dz_i^{\otimes 2} - \{\phi_i, z_i\}dz_i^{\otimes 2}$  coincide on the intersections  $U_i \cap U_j$  and the global quadratic differential  $\omega$  obtained by gluing them is a section of  $H^0(X, K_X^{\otimes 2}(-D))$ . We write  $p_2 - p_1 := \omega$ . Fuchs's local theory ensures that any quadratic differential in  $H^0(X, K_X^{\otimes 2}(-D))$  is obtained as the difference of two branched projective structures in  $P_X^a$ . This shows that when  $P_X^a \neq \emptyset$ , it is an affine space directed by  $H^0(X, K_X^{\otimes 2}(-D))$ .

Now we do not suppose  $P_X^a \neq \emptyset$  anymore. Denote by  $\mathfrak{P}_X^a$  the *sheaf of branched projective structures* on  $X$  of divisor  $D$  and branching class  $a$ : if  $U \subset X$  is an open subset, the space of local sections  $\Gamma(U, \mathfrak{P}_X^a)$  is the space  $P_U^{a|U}$  of branched projective structures on  $U$  of divisor  $D|_U$  and branching class  $a|_U$ . As we have just proved, the sheaf  $\mathfrak{P}_X^a$  is an affine sheaf directed by  $K_X^{\otimes 2}(-D)$ . The sheaf  $\mathfrak{P}_X^a$  is moreover locally nonempty: if  $x \in X$ , any local holomorphic function around  $x$  with the right branching order and branching class at  $x$  provides a local section of  $\mathfrak{P}_X^a$ . Thus by Lemma 1.2.3, there is a canonical cohomology class  $c(\mathfrak{P}_X^a) \in H^1(X, K_X^{\otimes 2}(-D))$  that vanishes if and only if  $H^0(X, \mathfrak{P}_X^a) \neq 0$ . We write  $c(a) := c(\mathfrak{P}_X^a)$ .  $\square$

## 2. FAMILIES OF BRANCHED PROJECTIVE STRUCTURES OVER A FAMILY OF CURVES

By a *family of complex curves*, we mean a triple  $(S, \mathcal{X}, \pi)$  where  $\mathcal{X}$  and  $S$  are connected, reduced analytic spaces and  $\pi : \mathcal{X} \rightarrow S$  is a proper and smooth morphism. The analytic space  $S$  is called the *base* of  $\mathcal{F}$ . It is equivalent to require that  $\pi$  is proper and locally trivial on  $\mathcal{X}$ : each point of  $\mathcal{X}$  admits a neighborhood  $U \subset \mathcal{X}$  such that  $V = \pi(U) \subset S$  is open and there exists an isomorphism of analytic spaces  $\phi : U \simeq V \times \Omega$  where  $\Omega$  is an open subset of  $\mathbb{C}$  and  $\pi = \text{pr}_1 \circ \phi$ , where  $\text{pr}_1$  is the projection on the first factor. In particular, the fibers of  $\pi$  are smooth and in the case where  $S$  is smooth,  $\pi$  is a holomorphic submersion. According to a theorem by Ehresmann (see for instance [Voi07]),  $\pi : \mathcal{X} \rightarrow S$  is topologically a locally trivial bundle.

Given a family of complex curves  $\mathcal{F} = (S, \mathcal{X}, \pi)$ , an analytic space  $S'$  and a morphism of analytic spaces  $f : S' \rightarrow S$ , we denote by  $f^*\mathcal{F} = (S', f^*\mathcal{X}, f^*\pi)$  the *pullback family* of  $\mathcal{F}$  by  $\pi$ :  $f^*\mathcal{X}$  is the fiber product  $S' \times_S \mathcal{X}$  and  $f^*\pi$  is the projection on the first factor.

**2.1. FAMILIES OF DIVISORS OVER A FAMILY OF CURVES.** — By a *family of complex curves with divisors*, or more shortly *family of divisors*, we mean a family  $\mathcal{F} = (S, \mathcal{X}, \mathcal{D}, \pi)$ , where  $(S, \mathcal{X}, \pi)$  is a family of complex curves, and  $\mathcal{D}$  is an effective Cartier divisor on  $\mathcal{X}$  that does not contain any fiber of  $\pi$ . Writing  $X_s = \pi^{-1}(\{s\})$  ( $s \in S$ ), Rouché's theorem implies that the degree of  $D_s = \mathcal{D}|_{X_s}$  does not depend on  $s$ . The *degree* of the family  $\mathcal{F}$  is the degree of the divisors  $D_s$ .

As in the case of families of curves, if  $f : S' \rightarrow S$  is a morphism of analytic spaces and  $\mathcal{F}$  a family of divisors, one can define the *pullback*  $f^*\mathcal{F}$  of  $\mathcal{F}$  by  $f$ : the base of  $f^*\mathcal{F}$  is  $S'$  and the fiber of  $f^*\mathcal{F}$  over  $s' \in S'$  is isomorphic to the fiber of  $\mathcal{F}$  over  $f(s') \in S$ .

Let  $\mathcal{G}$  be a family of complex curves, with base an analytic space  $T$ . A family of divisors *over*  $\mathcal{G}$  is a family of divisors  $\mathcal{F} = (S, \mathcal{X}, \mathcal{D}, \pi)$  along with a morphism

$f : S \rightarrow T$ , such that the underlying family of curves  $(S, \mathcal{X}, \pi)$  is the pullback of  $\mathcal{G}$  by  $f$ :  $(S, \mathcal{X}, \pi) = f^*\mathcal{G}$ . If  $g : S' \rightarrow S$  is a morphism of analytic spaces, the pullback of  $\mathcal{F}$  by  $g$  is still a family of divisors over  $\mathcal{G}$ , the associated morphism being  $f \circ g : S' \rightarrow T$ . Take  $n \in \mathbb{N}$ . The family  $\mathcal{F}$  is a *universal family of divisors of degree  $n$  over  $\mathcal{G}$*  if for any analytic space  $S'$  and any family  $\mathcal{F}'$  of divisors of degree  $n$  over  $\mathcal{G}$  with base  $S'$ , there exists a unique morphism  $g : S' \rightarrow S$  such that  $\mathcal{F}' = g^*\mathcal{F}$ . A universal family of divisors of degree  $n$  over  $\mathcal{G}$ , if it exists, is clearly unique up to a unique isomorphism.

It is a classical fact that for any family  $\mathcal{G}$  of complex curves, there exists a universal family  $\mathcal{F}$  of divisors of degree  $n$  over  $\mathcal{G}$ . Let us recall briefly the construction of  $\mathcal{F}$ . Write  $\mathcal{G} = (T, \mathcal{X}, \pi)$ . We set  $S = \text{Sym}_T^n \mathcal{X}$  the  $n$ -th symmetric power of  $\mathcal{X}$  relatively to  $T$ , that comes with a map  $\gamma : S \rightarrow T$ . The fiber of  $\gamma$  over  $t \in T$  is the  $n$ -th symmetric power of the fiber  $X_t$  of  $\pi$  over  $t$ . Note that

$$S = \{(t, D) \mid t \in T, D \text{ effective divisor of degree } n \text{ on } X_t\}.$$

The universal family  $\mathcal{F}$  is defined as  $(S, \gamma^*\mathcal{X}, \gamma^*\pi, \mathcal{D})$  where  $\mathcal{D}$  is the Cartier divisor on  $\gamma^*\mathcal{X}$  such that for any  $s = (t, D) \in S$ ,  $\mathcal{D}|_{X_s} = D$ , where  $X_s = \gamma^*\pi^{-1}(\{s\})$ . If  $\mathcal{F}' = (S', \mathcal{X}', \pi', \mathcal{D}')$ , along with a morphism  $f' : S' \rightarrow T$ , is another family of divisors of degree  $n$  over  $\mathcal{G}$ ,  $\mathcal{F}' = g^*\mathcal{F}$ , where  $g : S' \rightarrow S$  is the analytic morphism defined for any  $s' \in S'$  by  $g(s') = (f'(s'), D'_{s'})$  where  $X'_{s'}$  and  $X_{f'(s')}$  are identified (here  $X_t = \pi^{-1}(\{t\})$ ,  $X'_{s'} = \pi'^{-1}(\{s'\})$  and  $D'_{s'} = \mathcal{D}'|_{X'_{s'}}$ ). The morphism  $g$  is clearly the only one such that  $\mathcal{F}' = g^*\mathcal{F}$ . See [BM14] for more details on spaces of effective divisors.

**2.2. FAMILIES OF BRANCHING CLASSES OVER A FAMILY OF DIVISORS.** — The following definition gives a meaning to the notion of a holomorphic deformation of a branching class.

**DEFINITION 2.2.1**

(i) Let  $X$  be a Riemann surface and  $D$  an effective divisor on  $X$ . A *branched atlas* on the complex curve  $X$  with branching divisor  $D$  is the datum of an open covering  $(U_i)_{i \in I}$  of  $X$ , along with holomorphic functions  $f_i : U_i \rightarrow \mathbb{C}$  whose branching divisors are  $D|_{U_i}$ , such that the branching class of  $f_i$  on  $U_{ij} = U_i \cap U_j$  is the same as the branching class of  $f_j$  on  $U_{ij}$ .

(ii) Let  $(S, \mathcal{X}, \pi, \mathcal{D})$  be a family of divisors. A *branched atlas on  $\mathcal{X}$  relative to  $S$ , with branching divisor  $\mathcal{D}$*  is the datum of an open covering  $(U_i)_{i \in I}$  of  $\mathcal{X}$ , along with holomorphic functions  $f_i : U_i \rightarrow \mathbb{C}$  such that for any  $s \in S$ , the family  $(U_i \cap X_s, f_i|_{U_i \cap X_s})_{i \in I}$  is a branched atlas on  $X_s$  of divisor  $D_s$ . Two branched atlases are *equivalent* if their union is still a branched atlas. A *branching class on  $\mathcal{X}$  relative to  $S$ , with branching divisor  $\mathcal{D}$*  is an equivalence class of branched atlases on  $\mathcal{X}$  relative to  $S$ , with branching divisor  $\mathcal{D}$ .

We can now use a terminology similar to that of Section 2.1.

## DEFINITION 2.2.2

(i) Let  $n \in \mathbb{N}$ . A *family of complex curves with divisors (of degree  $n$ ) and branching classes*, or more shortly a *family of branching classes (of degree  $n$ )* is given by a family  $\mathcal{F} = (S, \mathcal{X}, \pi, \mathcal{D}, \mathfrak{a})$  where  $(S, \mathcal{X}, \pi, \mathcal{D})$  is a family of divisors (of degree  $n$ ) and  $\mathfrak{a}$  is a branching class on  $\mathcal{X}$  relative to  $S$ , with branching divisor  $\mathcal{D}$ .

(ii) Let  $\mathcal{G}$  be a family of divisors (respectively of complex curves) of basis  $T$ . A *family of branching classes over  $\mathcal{G}$*  is a family of branching classes  $\mathcal{F}$  of basis  $S$ , along with a morphism  $f : S \rightarrow T$  such that the underlying family of divisors (respectively of complex curves) is  $f^*\mathcal{G}$ .

(iii) Let  $S$  be an analytic space,  $\mathcal{F}$  a family of branching classes over  $\mathcal{G}$  with basis  $S$ . The family  $\mathcal{F}$  is said to be *universal* if any family  $\mathcal{F}'$  of branching classes over  $\mathcal{G}$  is uniquely obtained as the pullback of  $\mathcal{F}$  by some morphism  $g : S' \rightarrow S$ , where  $S'$  is the basis of  $\mathcal{F}'$ .

The discussion that will follow in this paragraph can be summarized in the following proposition:

PROPOSITION 2.2.3. — *Let  $\mathcal{G} = (T, \mathcal{X}, \pi, \mathcal{D})$  be a family of divisors. There exists a universal family  $\mathcal{F}$  of branching classes over  $\mathcal{G}$ .*

*Moreover, the basis of  $\mathcal{F}$  is an affine bundle over  $T$ , directed by the vector bundle  $\pi_* K_{\mathcal{X}/T}|_{\mathcal{D}}$  of relative differentials on  $\mathcal{X}$  restricted to  $\mathcal{D}$  with respect to  $T$ .*

## REMARK 2.2.4

(i) With the notations of Proposition 2.2.3, the fiber over  $t \in T$  of the vector bundle  $\pi_* K_{\mathcal{X}/T}|_{\mathcal{D}}$  is  $H^0(X_t, K_{X_t}|_{D_t})$

(ii) It is clear from Definition 2.2.2 that the universal family in 2.2.3 is unique up to a unique isomorphism.

(iii) It follows from Proposition 2.2.3 that for any family of curves  $\mathcal{G}$ , there exists a universal family of branching classes over  $\mathcal{G}$ . This is shown by applying Proposition 2.2.3 to the universal family of divisors over  $\mathcal{G}$ .

For the rest of the paragraph, fix a family of divisors  $\mathcal{G} = (T, \mathcal{X}, \pi, \mathcal{D})$ . Let us denote by

$$(2) \quad \mathcal{A}_{\mathcal{X}}^{\mathcal{D}} = \{(t, a) \mid t \in T, a \in A_{X_t}^{\mathcal{D}_t}\}$$

the union of all the spaces of branching classes associated to one of the curves with divisor of the family  $\mathcal{G}$ . There is an obvious projection

$$(3) \quad \delta : \mathcal{A}_{\mathcal{X}}^{\mathcal{D}} \twoheadrightarrow T.$$

For each  $t \in T$ , according to Proposition 1.1.2, the fiber  $\delta^{-1}(s) = A_{X_s}^{\mathcal{D}_s}$  is an affine space directed by the vector space  $H^0(X_t, K_{X_t}|_{D_t}) = (\pi_*(K_{\mathcal{X}/T}|_{\mathcal{D}}))_t$ . In order to get on  $\mathcal{A}_{\mathcal{X}}^{\mathcal{D}}$  the structure of an affine bundle on  $T$  directed by the vector bundle  $\pi_*(K_{\mathcal{X}/S}|_{\mathcal{D}})$ , it is enough to specify local holomorphic sections of  $\delta$ .

Remark that, up to restricting  $T$ , there exists a branched atlas on  $\mathcal{X}$  relative to  $T$  with branching divisor  $\mathcal{D}$ . Indeed, take  $t \in T$  and to any point  $y_k$  of  $D_t = \sum_{k=1}^r n_k \cdot y_k$ ,

associate a neighborhood  $U_k$  of  $y_k$  in  $\mathcal{X}$  and a function  $g_k : U_k \rightarrow \mathbb{C}$  such that  $\mathcal{D} \cap U_k$  is the zero locus of  $g_k$ . Write  $V = \pi(U_1) \cap \cdots \cap \pi(U_r) \subset T$ . It is a neighborhood of  $t$ . Let also  $(W_i, f_i)_{i \in I}$  be a relative atlas (i.e., a branched relative atlas with divisor zero) for the family  $\pi^{-1}(V) \setminus \mathcal{D}$ . Such a relative atlas exists because the family  $(T, \mathcal{X}, \pi)$  is locally trivial on  $\mathcal{X}$ . The union of the families  $(W_i, f_i)_{i \in I}$  and  $(U_k, g_k)_{1 \leq k \leq r}$  is a branched relative atlas for the family  $\pi^{-1}(V)$  with branching divisor  $\mathcal{D}|_{\pi^{-1}(V)}$ .

The datum of a branching class  $\mathfrak{a}$  on  $\mathcal{X}$  relative to  $T$  with branching divisor  $\mathcal{D}$  defines a section  $\sigma_{\mathfrak{a}} : T \rightarrow \mathcal{A}_{\mathcal{X}}^{\mathcal{D}}$  of the projection  $\delta$ :  $\sigma_{\mathfrak{a}}(t)$  is the branching class of the branched atlas  $\mathfrak{a}$  restricted to  $X_t$ .

LEMMA 2.2.5. — *Let  $\mathfrak{a}_1$  and  $\mathfrak{a}_2$  be two branching classes on  $\mathcal{X}$  relative to  $T$  with divisor  $\mathcal{D}$ . For each  $t \in T$ , write  $\sigma_{\mathfrak{a}_2}(t) = \sigma_{\mathfrak{a}_1}(t) + \phi(t) \in A_{X_t}^{D_t}$ , with  $\phi(t) \in H^0(X_t, K_{X_t}|_{D_t})$ . Then  $\phi$  is a holomorphic section of the vector bundle  $\pi_*(K_{\mathcal{X}/T}|_{\mathcal{D}})$ .*

*Proof.* — Take  $t_0 \in T$ , and let  $y \in X_{t_0}$  be a point of  $D_{t_0}$ . The local triviality on  $\mathcal{X}$  of  $(T, \mathcal{X}, \pi)$  implies that there exists a local neighborhood  $V$  of  $t_0$  in  $T$ , an open neighborhood  $U \subset \mathcal{X}$  of  $y$  with  $\pi(U) = V$ , and local coordinates  $(z, w) : U \rightarrow V \times \mathbb{C}$  with  $\pi(z, w) = z$ . Without loss of generality, suppose  $U$  is the domain of a chart both in  $\mathfrak{a}_1$  and  $\mathfrak{a}_2$ . Take charts  $f, g : U \rightarrow \mathbb{C}$  of  $\mathfrak{a}_1$  and  $\mathfrak{a}_2$  respectively. According to the proof of Proposition 1.1.2, it is enough to show that the function

$$\frac{\partial_w^2 g}{\partial_w g} - \frac{\partial_w^2 f}{\partial_w f} : U \longrightarrow \mathbb{C}$$

is holomorphic.

The functions  $f$  and  $g$  have the same branching divisor when restricted to each fiber of  $X$ , i.e., the functions  $\partial_w g$  and  $\partial_w f$  have the same vanishing locus. Thus there exists a nonvanishing holomorphic function  $h : U \rightarrow \mathbb{C}^*$  such that  $\partial_w g = h \cdot \partial_w f$ . Therefore, one has  $\partial_w^2 g / \partial_w g - \partial_w^2 f / \partial_w f = \partial_w h / h$ , which is holomorphic.  $\square$

As a consequence of Lemma 2.2.5,  $\mathcal{A}_{\mathcal{X}}^{\mathcal{D}}$  admits a structure of affine bundle for which the sections coming from branched relative atlases on  $\mathcal{G}$  are holomorphic.

LEMMA 2.2.6. — *Let  $\sigma : T \rightarrow \mathcal{A}_{\mathcal{X}}^{\mathcal{D}}$  be a holomorphic section. There exists a branching class  $\mathfrak{a}$  on  $\mathcal{X}$ , relative to  $T$  and of divisor  $\mathcal{D}$  such that  $\sigma = \sigma_{\mathfrak{a}}$ .*

*Proof.* — The union of two branched relative atlases with the same branching classes given by  $\sigma$  is still a branched relative atlas with relative branching classes given by  $\sigma$ . As a consequence, it is enough to prove the lemma locally on  $T$ .

Take  $t \in T$ . Up to restricting  $T$  to an open neighborhood of  $t$ , there exists a relative branching class  $\mathfrak{a}_0$  on  $\mathcal{G}$ , with associated section  $\sigma_{\mathfrak{a}_0} : T \rightarrow \mathcal{A}_{\mathcal{X}}^{\mathcal{D}}$ . Let  $y \in X_t$  be a point of  $D_t$ . Let  $(U, f)$  be a chart of an atlas in  $\mathfrak{a}_0$  with  $y \in U$ . Since  $(T, \mathcal{X}, \pi)$  is locally trivial on  $\mathcal{X}$ , up to restricting  $U$ , one has a local biholomorphism  $(z, w) : U \rightarrow V \times \mathbb{C}$ , where  $V = \pi(U)$  and  $\pi(z, w) = z$ . The difference  $\sigma - \sigma_{\mathfrak{a}_0}$  is a section of the vector bundle  $\pi_*(K_{\mathcal{X}/T}|_{\mathcal{D}})$  on  $T$ , that can be seen as a section of the vector bundle  $K_{\mathcal{X}/T}|_{\mathcal{D}}$  on the analytic space  $\mathcal{D}$ . With this point of view, up to restricting  $U$ ,  $(\sigma - \sigma_{\mathfrak{a}_0})|_U$  is given by  $\phi|_{\mathcal{D}}$ , where  $\phi$  is a section on  $U$  of  $K_{\mathcal{X}/T}$ , given by  $\phi = \varphi(z, w)dw$ . Up to

restricting  $U$  again (in particular so that it is contractible), there exists a holomorphic function that never vanishes  $h : U \rightarrow \mathbb{C}^*$  such that  $\varphi = \partial_w h / h$ . By restricting  $U$  once more, one gets a function  $g : U \rightarrow \mathbb{C}$  such that  $\partial_w g = h \cdot \partial_w f$ . The branching divisor of  $g$  is then  $\mathcal{D}|_U$  and its branching classes on the fibers of  $\pi$  are given by  $\sigma$ .

By doing so at each point of  $D_t$ , by restricting  $T$  to the intersection of the obtained open sets  $\pi(U)$ , and then by completing with a relative atlas on  $\mathcal{X} \setminus \mathcal{D}$ , one gets a branched relative atlas  $\mathfrak{a}$  on  $\mathcal{G}$  with  $\sigma = \sigma_{\mathfrak{a}}$ .  $\square$

The pullback family of divisors  $\delta^*\mathcal{G} = (\mathcal{A}_{\mathcal{X}}^{\mathcal{D}}, \delta^*\mathcal{X}, \delta^*\pi, \delta^*\mathcal{D})$  comes with a tautological relative branching class  $\mathfrak{a}_0$ : if  $(t, a) \in \mathcal{A}_{\mathcal{X}}^{\mathcal{D}}$ , the branching class defined by  $\mathfrak{a}_0$  on  $(\delta^*\pi)^{-1}(t, a) = X_t$  with divisor  $D_t$  is  $a$ . Let us show that  $\mathfrak{a}_0$  is indeed a relative branching class. The pullback by  $\delta$  of a relative branching class on  $\mathcal{X}$  relative to  $T$  of divisor  $\mathcal{D}$  is clearly a branching class on  $\delta^*\mathcal{X}$ , relative to  $\mathcal{A}_{\mathcal{X}}^{\mathcal{D}}$  of divisor  $\delta^*\mathcal{D}$ . Thus the analytic space  $\mathcal{A}_{\delta^*\mathcal{X}}^{\delta^*\mathcal{D}}$  of branching classes on the fibers of  $\delta^*\mathcal{G}$  is the analytic fiber product  $\mathcal{A}_{\mathcal{X}}^{\mathcal{D}} \times_T \mathcal{A}_{\mathcal{X}}^{\mathcal{D}}$ . The tautological relative branching class  $\mathfrak{a}_0$  comes from the diagonal section of  $\mathcal{A}_{\delta^*\mathcal{X}}^{\delta^*\mathcal{D}}$ , that is holomorphic.

The family  $\mathcal{F} := (\mathcal{A}_{\mathcal{X}}^{\mathcal{D}}, \delta^*\mathcal{X}, \delta^*\pi, \delta^*\mathcal{D}, \mathfrak{a}_0)$ , along with the morphism  $\delta : \mathcal{A}_{\mathcal{X}}^{\mathcal{D}} \rightarrow T$  is a family of branching classes over  $\mathcal{G}$ .

**LEMMA 2.2.7.** — *The family  $\mathcal{F}$  above, along with the map  $\delta$ , is a universal family of branching classes over  $\mathcal{G}$ .*

*Proof.* — Let  $\mathcal{F}' = (S', \mathcal{X}', \pi', \mathcal{D}', \mathfrak{a}')$ , along with a morphism  $f' : S' \rightarrow T$ , be a family of branching classes over  $\mathcal{G}$ . For any  $s' \in S'$ , the relative branched atlas  $\mathfrak{a}'$  defines a branching class  $a_{s'}$  over  $(X'_{s'}, D'_{s'}) \simeq (X_{f'(s')}, D_{f'(s')})$ . Thus we get a map  $g : S' \rightarrow \mathcal{A}_{\mathcal{X}}^{\mathcal{D}}$  given by  $g(s') = (f'(s'), a_{s'})$ . One has  $f' = \delta \circ g$ , for any  $s' \in S'$ , the fiber of  $\mathcal{F}'$  over  $s'$  is identified to the fiber of  $\mathcal{F}$  over  $g(s')$ , and  $g$  is the only map from  $S'$  to  $\mathcal{A}_{\mathcal{X}}^{\mathcal{D}}$  that fulfills these two conditions.

To complete the proof of the lemma, we only need to show that  $g$  is analytic. Clearly,  $\mathcal{A}_{\mathcal{X}'}^{\mathcal{D}'}$  is the analytic fiber product  $\mathcal{A}_{\mathcal{X}'}^{\mathcal{D}'} = S' \times_T \mathcal{A}_{\mathcal{X}}^{\mathcal{D}}$ . By Lemma 2.2.6, the relative branching class  $\mathfrak{a}'$  gives an analytic section  $\sigma_{\mathfrak{a}'}$  of  $\mathcal{A}_{\mathcal{X}'}^{\mathcal{D}'}$ , that can be viewed as an analytic morphism  $\sigma_{\mathfrak{a}'} : S' \rightarrow S' \times_T \mathcal{A}_{\mathcal{X}}^{\mathcal{D}}$ . The map  $g$  is this analytic morphism composed with the projection on  $\mathcal{A}_{\mathcal{X}}^{\mathcal{D}}$ , which is analytic.  $\square$

**2.3. FAMILIES OF BRANCHED PROJECTIVE STRUCTURES.** — The following definition gives a meaning to the notion of a holomorphic deformation of a branched projective structure.

**DEFINITION 2.3.1.** — Let  $(S, \mathcal{X}, \pi, \mathcal{D}, \mathfrak{a})$  be a family of branching classes. A *branched projective atlas on  $\mathcal{X}$  relative to  $S$  with branching divisor  $\mathcal{D}$  and relative branching class  $\mathfrak{a}$*  is the datum of an atlas on  $\mathcal{X}$ , relative to  $S$ , of divisor  $D$  and relative branching class  $\mathfrak{a}$  such that the branched atlas induced on each  $X_s$  is a branched projective atlas. Two relative branched projective atlases are *equivalent* if their union is still a relative branched projective atlas. A *branched projective structure* relative to  $S$  with branching

divisor  $\mathcal{D}$  and relative branching class  $\mathfrak{a}$  is an equivalence class of branched projective atlases on  $\mathcal{X}$  relative to  $S$  with branching divisor  $\mathcal{D}$  and relative branching class  $\mathfrak{a}$ .

We can now introduce a terminology similar to that of Sections 2.1 and 2.2.

**DEFINITION 2.3.2**

(i) Let  $n \in \mathbb{N}$ . A *family of complex curves with divisors (of degree  $n$ ), branching classes and branched projective structures*, or more shortly a *family of branched projective structures (of degree  $n$ )* is given by a family  $\mathcal{F} = (S, \mathcal{X}, \pi, \mathcal{D}, \mathfrak{a}, \mathfrak{p})$ , where  $(S, \mathcal{X}, \pi, \mathcal{D}, \mathfrak{a})$  is a family of branching classes (of degree  $n$ ) and  $\mathfrak{p}$  is a branched projective structure on  $\mathcal{X}$  relative to  $S$ , with branching divisor  $\mathcal{D}$  and branching class  $\mathfrak{a}$ .

(ii) Let  $\mathcal{G}$  be a family of branching classes (respectively of complex curves, respectively of divisors) of basis  $T$ . A *family of branched projective structures over  $\mathcal{G}$*  is a family of branched projective structures  $\mathcal{F}$  of basis  $S$ , along with a morphism  $f : S \rightarrow T$  such that the underlying family of branching classes (respectively of complex curves, respectively of divisors) is  $f^*\mathcal{G}$ .

(iii) Let  $S$  be an analytic space,  $\mathcal{F}$  a family of branched projective structures over  $\mathcal{G}$  with basis  $S$ . The family  $\mathcal{F}$  is said to be *universal* if any family  $\mathcal{F}'$  of branched projective structures over  $\mathcal{G}$  is uniquely obtained as the pullback of  $\mathcal{F}$  by some morphism  $g : S' \rightarrow S$ , where  $S'$  is the basis of  $\mathcal{F}'$ .

The aim of this section is to prove the following proposition:

**PROPOSITION 2.3.3.** — *Let  $\mathcal{G}$  be a family of branching classes. There exists a universal family of branched projective structures over  $\mathcal{G}$ .*

**REMARK 2.3.4.** — It is clear from Definition 2.2.2 that the universal family in Proposition 2.3.3 is unique up to a unique isomorphism.

Before diving into the proof of Proposition 2.3.3 in the next paragraphs, let us state a corollary that is one of the main results of this paper. Let us first recall the definition of a *marked* Riemann surface.

**DEFINITION 2.3.5.** — Let  $X$  be a compact Riemann surface. Let  $\mathcal{F} = (T, \mathcal{X}, \pi)$  be a family of Riemann surfaces that are homeomorphic to  $X$ .

(i) If  $X$  has genus 0, a *marking* on  $X$  is the datum of 3 distinct points on  $X$ . A marking on  $\mathcal{F}$  is the datum of three holomorphic sections of  $\pi$  that do not intersect with each other.

(ii) If  $X$  has genus 1 and  $S$  is a differential surface of genus 1, a *marking on  $X$  with reference  $S$*  is the datum of a point in  $X$  and an isotopy class of diffeomorphisms from  $S$  to  $X$ . A marking on  $\mathcal{F}$  with reference  $S$  is the datum of a holomorphic section of  $\pi$ , and an isotopy class of diffeomorphisms from  $T \times S$  to  $\mathcal{X}$  that commute with  $\pi$ .

(iii) If  $X$  has genus  $g \geq 2$  and  $S$  is a differential surface of genus  $g$ , a *marking on  $X$  with reference  $S$*  is the datum of an isotopy class of diffeomorphisms from  $S$  to  $X$ . A marking on  $\mathcal{F}$  with reference  $S$  is the datum of an isotopy class of diffeomorphisms from  $T \times S$  to  $\mathcal{X}$  that commute with  $\pi$ .



A *marked compact Riemann surface* is a compact Riemann surface with a marking, where a differential surface  $S$  is supposed fixed.

In particular, there is at most one analytic isomorphism between two marked Riemann surfaces.

Marked (families of) divisors (respectively branching classes, respectively branched projective structures) are (families of) divisors (respectively branching classes, respectively branched projective structures) along with a marking on the underlying (family of) curves. A marked family of divisors (respectively branching classes, respectively branched projective structures)  $\mathcal{F}$  is said to be *over* a marked family of curves (respectively divisors, respectively branching classes)  $\mathcal{G}$  if  $\mathcal{F}$  is given as the pullback of  $\mathcal{G}$  by an analytic morphism.

Propositions 2.2.3 and 2.3.3 remain true if we consider marked families: the marking on the universal family is obtained by pulling back the marking of the base family.

Let us give a definition of a *universal family*, that is not over another family:

**DEFINITION 2.3.6.** — Take  $n, g \in \mathbb{N}$ . A family  $\mathcal{F}$  of (marked) Riemann surfaces (respectively divisors of degree  $n$ , respectively branching classes of degree  $n$ , respectively branched projective structures of degree  $n$ ) of genus  $g$  is said to be *universal* if any other such family  $\mathcal{F}'$  can be realized in a unique way as a pullback of  $\mathcal{F}$ .

Let  $g \in \mathbb{N}$ . There is no universal family of curves of genus  $g$ . In particular, the tautological family over the moduli space  $\mathcal{M}_g$  of genus  $g$  curves is not universal, since nontrivial isotrivial families are not pullbacks of it. There exists however a universal family of *marked* Riemann surfaces of genus  $g$ , given by  $\mathcal{U}_g = (\mathcal{T}_g, \mathcal{X}_g, \pi_g, m_g)$ , where  $\mathcal{T}_g$  is the Teichmüller space for curves of genus  $g$ ,  $\mathcal{X}_g$  is the tautological family over  $\mathcal{T}_g$ ,  $\pi_g$  is the projection and  $m_g$  is the tautological marking. For  $n \in \mathbb{N}$ , denote by  $\mathcal{U}_g^{\text{div}}(n)$  the marked family of divisors of degree  $n$  obtained as the universal family of divisors of degree  $n$  over  $\mathcal{U}_g$ , along with the pullback marking.

Let  $\mathcal{U}_g^{\text{bc}}(n)$  be the universal marked family of branching classes over  $\mathcal{U}_g^{\text{div}}(n)$ , given by Proposition 2.2.3, and let  $\mathcal{U}_g^{\text{bps}}(n)$  be the universal marked family of branched projective structures over  $\mathcal{U}_g^{\text{bc}}(n)$ , given by Proposition 2.3.3. The family  $\mathcal{U}_g^{\text{bps}}(n)$  is clearly the universal family of marked branched projective structures of genus  $g$  and degree  $n$ . Denote by  $\mathcal{P}_g(n)$  the basis of the family  $\mathcal{U}_g^{\text{bps}}(n)$ . Now let  $p$  be a marked branched projective structure of degree  $n$  and genus  $g$ . The structure  $p$  can be seen as a family over a single-point analytic space  $\{x\}$ . By universality of  $\mathcal{U}_g^{\text{bps}}(n)$ , there exists a unique morphism  $f : \{x\} \rightarrow \mathcal{P}_g(n)$  such that the family  $p$  is the pullback of  $\mathcal{U}_g^{\text{bps}}(n)$  by  $f$ . To the structure  $p$  is thus associated a unique point in  $\mathcal{P}_g(n)$ . In particular, the structure  $p$  is isomorphic to the fiber of  $\mathcal{U}_g^{\text{bps}}(n)$  over the associated point. Clearly, two marked branched projective structures of genus  $g$  and degree  $n$  have the same associated point if and only if they are isomorphic, and any point in  $\mathcal{P}_g(n)$  is associated to the fiber of  $\mathcal{U}_g^{\text{bps}}(n)$  over itself. In this way we get a bijection between the analytic space  $\mathcal{P}_g(n)$  and the set of isomorphism classes of

marked branched projective structures of degree  $n$  and genus  $g$ . We thus have the following corollary of Proposition 2.3.3:

**THEOREM 2.3.7.** — *Let  $n, g \in \mathbb{N}$ . The moduli space of isomorphism classes of marked branched projective structures of genus  $g$  and degree  $n$  is an analytic space  $\mathcal{P}_g(n)$ , and is the basis of a universal family  $\mathcal{U}_g^{\text{bps}}(n)$  of marked branched projective structures of genus  $g$  and degree  $n$ . Moreover, the fiber of  $\mathcal{U}_g^{\text{bps}}(n)$  over a branched projective structure  $p \in \mathcal{P}_g(n)$  is  $p$  itself.*

Let us now give a proof for Proposition 2.3.3.

**2.4. SCHWARZIAN PARAMETRIZATION IN FAMILY.** — Let  $g, n \in \mathbb{N}$ . Fix a family of branching classes  $\mathcal{G} = (T, \mathcal{X}, \pi, \mathcal{D}, \mathbf{a})$  of genus  $g$  and degree  $n$ .

**LEMMA 2.4.1.** — *Suppose given  $\mathbf{p}_0$  a branched projective structure on  $\mathcal{X}$  relative to  $T$  and of relative branching class  $\mathbf{a}$ .*

(i) *If  $\mathbf{p}$  is a branched projective structure on  $\mathcal{X}$  relative to  $T$  and of relative branching class  $\mathbf{a}$ , then the difference  $\mathbf{p} - \mathbf{p}_0$  is a holomorphic section of the vector bundle  $K_{\mathcal{X}/T}^{\otimes 2}(-\mathcal{D})$  over  $\mathcal{X}$ :  $\mathbf{p} - \mathbf{p}_0 \in H^0(\mathcal{X}, K_{\mathcal{X}/T}^{\otimes 2}(-\mathcal{D}))$ .*

(ii) *If  $q \in H^0(\mathcal{X}, K_{\mathcal{X}/S}^{\otimes 2}(-\mathcal{D}))$ , then there exists a projective structure  $\mathbf{p}$  on  $\mathcal{X}$  relative to  $T$  and of relative branching class  $\mathbf{a}$  such that  $\mathbf{p} - \mathbf{p}_0 = q$ .*

*Proof*

(i) Let  $(z, w)$  be local coordinates on an open subset  $U \subset \mathcal{X}$  adapted to the locally trivial family on  $\mathcal{X}$ ,  $(T, \mathcal{X}, \pi)$ , i.e., with  $z = \tilde{z} \circ \pi$ , where  $\tilde{z}$  is a local coordinate on  $T$ . Let  $\phi_0 : U \rightarrow \mathbb{CP}^1$  be a local chart of  $\mathbf{p}_0$  on  $U$  and  $\phi : U \rightarrow \mathbb{CP}^1$  a local chart of  $\mathbf{p}$  on  $U$ . The difference  $\mathbf{p} - \mathbf{p}_0$  is the meromorphic relative quadratic differential

$$q = \left( (\partial_w(\partial_w^2 \phi / \partial_w \phi) - 1/2(\partial_w^2 \phi / \partial_w \phi)^2) - (\partial_w(\partial_w^2 \phi_0 / \partial_w \phi_0) - 1/2(\partial_w^2 \phi_0 / \partial_w \phi_0)^2) \right) dw^{\otimes 2}.$$

But  $\mathbf{p}_0$  and  $\mathbf{p}$ , when restricted to the fibers of  $\pi$ , define branched projective structures with the same branching class, thus for any  $t \in T$ ,  $q|_{X_t}$  is a local holomorphic section of  $K_{X_t}^{\otimes 2}(-D_t)$ . As a consequence, the relative quadratic differential  $q$  is a local holomorphic section of  $K_{\mathcal{X}/T}^{\otimes 2}(-\mathcal{D})$ .

(ii) Take  $x \in \mathcal{X}$ ,  $U$  an open neighborhood of  $x$ ,  $(U, \phi_0)$  a chart of  $\mathbf{p}_0$ , and  $(z, w)$  coordinates on  $U$  adapted to the locally trivial family  $(T, \mathcal{X}, \pi)$ . Write  $q|_U = u dw^{\otimes 2}$ . Up to restricting  $U$ , according to Cauchy-Lipschitz's theorem, there exists a holomorphic solution  $\phi : U \rightarrow \mathbb{CP}^1$  of the partial differential equation

$$\partial_w(\partial_w^2 \phi / \partial_w \phi) - 1/2(\partial_w^2 \phi / \partial_w \phi)^2 = u + (\partial_w(\partial_w^2 \phi_0 / \partial_w \phi_0) - 1/2(\partial_w^2 \phi_0 / \partial_w \phi_0)^2).$$

As a consequence of Proposition 1.2.2,  $\phi$  defines on  $U$  a relative branched projective structure with relative branching class  $\mathbf{a}$ . By doing so around each point of  $\mathcal{X}$ , one gets a branched projective atlas on  $\mathcal{X}$  relative to  $S$  of relative branching class  $\mathbf{a}$ . This atlas defines a relative branched projective structure  $\mathbf{p}$ , and one has  $\mathbf{p} - \mathbf{p}_0 = q$ .  $\square$

For any  $t \in T$ , write  $a_t$  the branching class defined by  $\mathbf{a}$  on the fiber of  $\mathcal{G}$  over  $t$ . Consider the union

$$(4) \quad \mathcal{P}_{\mathcal{X}}^{\mathbf{a}} = \{(t, p) \mid t \in T, p \in P_{X_t}^{a_t}\}$$

of the spaces of branched projective structures on the fibers of  $\mathcal{G}$ . There is an obvious map

$$(5) \quad \alpha : \mathcal{P}_{\mathcal{X}}^{\mathbf{a}} \longrightarrow T.$$

According to Proposition 1.2.2, for any  $t \in T$ , the fiber  $\alpha^{-1}(t)$  is either empty or an affine space directed by  $H^0(X_t, K_{X_t}^{\otimes 2}(-D_t))$ , according to whether the cohomology class  $c(a_t) \in H^1(X_t, K_{X_t}^{\otimes 2}(-D_t))$  vanishes or not. Thus studying the structure of  $\mathcal{P}_{\mathcal{X}}^{\mathbf{a}}$  is essentially studying the variation with  $t$  of the space  $H^0(X_t, K_{X_t}^{\otimes 2}(-D_t))$  and the cohomology class  $c(a_t)$ .

Write  $\mathfrak{BR}_{\mathcal{X}}^{\mathbf{a}}$  the sheaf on  $\mathcal{X}$  of branched projective structures on  $\mathcal{X}$  relative to  $T$  of relative branching class  $\mathbf{a}$ : the space of sections  $\Gamma(U, \mathfrak{BR}_{\mathcal{X}}^{\mathbf{a}})$  is the set of branched projective structures on  $U$  relative to  $\pi(U)$  of relative branching class  $\mathbf{a}|_{\pi(U)}$ . Lemma 2.4.1 implies that  $\mathfrak{BR}_{\mathcal{X}}^{\mathbf{a}}$  is a sheaf of affine spaces directed by the line bundle  $K_{\mathcal{X}/T}^{\otimes 2}(-\mathcal{D})$ . Moreover  $\mathfrak{BR}_{\mathcal{X}}^{\mathbf{a}}$  is clearly locally nonempty. In particular, it comes with a cohomology class  $\mathfrak{C}(\mathbf{a}) \in H^1(\mathcal{X}, K_{\mathcal{X}/T}^{\otimes 2}(-\mathcal{D}))$  with  $\mathfrak{C}(\mathbf{a}) = 0$  if and only if  $H^0(\mathcal{X}, \mathfrak{BR}_{\mathcal{X}}^{\mathbf{a}}) \neq \emptyset$ , i.e., if and only if there exists a branched projective structure on  $\mathcal{X}$ , relative to  $T$  of relative branching class  $\mathbf{a}$ .

The Leray spectral sequence gives an exact sequence (see [Voi07]):

$$(6) \quad H^1(T, \pi_* K_{\mathcal{X}/T}^{\otimes 2}(-\mathcal{D})) \longrightarrow H^1(\mathcal{X}, K_{\mathcal{X}/T}^{\otimes 2}(-\mathcal{D})) \xrightarrow{\chi_{\mathcal{G}}} H^0(T, R^1 \pi_* K_{\mathcal{X}/T}^{\otimes 2}(-\mathcal{D})).$$

Write

$$(7) \quad \mathfrak{c}(\mathbf{a}) = \chi_{\mathcal{G}}(\mathfrak{C}(\mathbf{a})).$$

Recall the Riemann-Roch formula for  $K_{X_t}^{\otimes 2}(-D_t)$ :

$$h^0(X_t, K_{X_t}^{\otimes 2}(-D_t)) - h^1(X_t, K_{X_t}^{\otimes 2}(-D_t)) = 3g - 3 - n.$$

Let us finish this paragraph by showing Proposition 2.3.3 in the special cases where the branching degree  $n$  is either small enough or large enough when compared to the genus  $g$ .

Suppose  $h^0(X, K_{X_t}^{\otimes 2}(-D_t))$ , thus  $h^1(X, K_{X_t}^{\otimes 2}(-D_t))$ , does not depend on  $t \in T$ . Then the sheaves  $\pi_* K_{\mathcal{X}/T}^{\otimes 2}(-\mathcal{D})$  and  $R^1 \pi_* K_{\mathcal{X}/T}^{\otimes 2}(-\mathcal{D})$  are vector bundles, and the section  $\mathfrak{c}(\mathbf{a})$  in (7) satisfies for any  $t \in T$ :  $\mathfrak{c}(\mathbf{a})(s) = c(a_t)$ .

Denote by  $Z \subset T$  the zero locus of the section  $\mathfrak{c}(\mathbf{a})$ , which is an analytic subspace. Let  $\iota : Z \rightarrow T$  be the holomorphic immersion. In other words, there exists a branched projective structure of divisor  $D_t$  and branching class  $a_t$  on  $X_t$  if and only if  $t \in Z$ . Let  $\mathcal{G}' = (Z, \iota^* \mathcal{X}, \iota^* \pi, \iota^* \mathcal{D}, \iota^* \mathbf{a})$ . The space  $Z$  is exactly the image of the map  $\alpha$  in (5) and the space  $\mathcal{P}_{\mathcal{X}}^{\mathbf{a}}$  in (4) satisfies

$$\mathcal{P}_{\mathcal{X}}^{\mathbf{a}} = \{(s, p) \mid s \in Z, p \in P_{X_t}^{a_t}\} = \mathcal{P}_{\iota^* \mathcal{X}}^{\iota^* \mathbf{a}}.$$

The map  $\chi_{\mathcal{G}'}$ , defined as in (6), is the zero map. Moreover, if  $Z$  is Stein, the exact sequence (6) applied to the family  $\mathcal{G}'$  has vanishing first term, which implies that the

second term also vanishes. As a consequence, up to restricting  $Z$  to an open subset, there are branched projective structures on  $\iota^*\mathcal{X}$  relative to  $Z$  and of branching class  $\iota^*\mathfrak{a}$  (such structures do not necessarily exist globally on  $Z$ , see [DG23]).

By Lemma 2.4.1, the map  $\alpha : \mathcal{P}_{\mathcal{X}}^{\mathfrak{a}} \rightarrow Z$  is an affine bundle directed by the vector bundle

$$(\iota^*\pi)_* K_{\iota^*\mathcal{X}/Z}^{\otimes 2}(-\iota^*\mathcal{D}) = \pi_* K_{\mathcal{X}/T}^{\otimes 2}(-\mathcal{D})|_Z,$$

and whose local holomorphic sections are given by the (local on  $Z$ ) branched projective structures on  $\iota^*\mathcal{X}$  relative to  $Z$  and with branching class  $\iota^*\mathfrak{a}$ . In particular, the map  $\alpha$  is holomorphic.

The pullback family of branching classes  $\alpha^*\mathcal{G} = (\mathcal{P}_{\mathcal{X}}^{\mathfrak{a}}, \alpha^*\mathcal{X}, \alpha^*\pi, \alpha^*\mathcal{D}, \alpha^*\mathfrak{a})$  comes with a tautological relative branched projective structure  $\mathfrak{p}_0$ : if  $(t, p) \in \mathcal{P}_{\mathcal{X}}^{\mathfrak{a}}$ , the branched projective structure defined by  $\mathfrak{p}_0$  on  $(\alpha^*\pi)^{-1}(t, p) = X_t$  with branching class  $a_t$  is  $p$ .

The discussion around Lemma 2.2.7 applies *mutatis mutandis*, to show that the family  $(\mathcal{P}_{\mathcal{X}}^{\mathfrak{a}}, \alpha^*\mathcal{X}, \alpha^*\pi, \alpha^*\mathcal{D}, \alpha^*\mathfrak{a}, \mathfrak{p}_0)$ , along with the morphism  $\alpha : \mathcal{P}_{\mathcal{X}}^{\mathfrak{a}} \rightarrow T$  is a universal family of branched projective structures over  $\mathcal{G}$ .

If  $n < 2g - 2$ , by Serre duality, for any  $t \in T$ , we have

$$h^1(X_t, K_{X_t}^{\otimes 2}(-D_t)) = h^0(X_t, T_{X_t}(D_t)) = 0,$$

because  $\deg(T_{X_t}(D_t)) < 0$ . In particular,  $h^1(X_t, K_{X_t}^{\otimes 2}(-D_t))$  and  $h^0(X_t, K_{X_t}^{\otimes 2}(-D_t))$  do not depend on  $t \in T$ . In that case, the section  $c(\mathfrak{a})$  takes values in the zero vector bundle, thus  $Z = T$  and  $\mathcal{P}_{\mathcal{X}}^{\mathfrak{a}}$  is an affine bundle over  $T$ .

On the other extreme, if  $n > 4g - 4$ , then  $\deg(K_{X_t}^{\otimes 2}(-D_t)) < 0$ , so

$$h^0(X_t, K_{X_t}^{\otimes 2}(-D_t)) = 0.$$

Again  $h^0(X_t, K_{X_t}^{\otimes 2}(-D_t))$  and  $h^1(X_t, K_{X_t}^{\otimes 2}(-D_t))$  are both independent of  $t \in T$ . Moreover the fibers of  $\alpha : \mathcal{P}_{\mathcal{X}}^{\mathfrak{a}} \rightarrow T$  contain at most one point. In that case,  $\mathcal{P}_{\mathcal{X}}^{\mathfrak{a}}$  is an analytic subspace of  $T$ .

One has the following proposition, which is a particular case of Proposition 2.3.3:

**PROPOSITION 2.4.2.** — *Recall  $n, g \in \mathbb{N}$ ,  $\mathcal{G} = (T, \mathcal{X}, \mathcal{D}, \pi, \mathfrak{a})$  is a family of complex curves of genus  $g$  with divisors of degree  $n$  and branching classes and  $\mathcal{P}_{\mathcal{X}}^{\mathfrak{a}}$  is the set of branched projective structures on the fibers of  $\mathcal{G}$ .*

(i) *If  $n < 2g - 2$ , the space  $\mathcal{P}_{\mathcal{X}}^{\mathfrak{a}}$  defined in (4) is an affine bundle on  $T$  directed by the vector bundle  $\pi_*(K_{\mathcal{X}/T}^{\otimes 2}(-\mathcal{D}))$ .*

(ii) *If  $n > 4g - 4$ , the space  $\mathcal{P}_{\mathcal{X}}^{\mathfrak{a}}$  is an analytic subspace of  $T$ .*

*In both cases,  $\mathcal{P}_{\mathcal{X}}^{\mathfrak{a}}$  is the basis of a universal family of branched projective structures over  $\mathcal{G}$ .*

**REMARK 2.4.3.** — In the limit case  $n = 2g - 2$ , the set  $T_0$  of points  $t \in T$  for which  $h^1(X_t, K_{X_t}^{\otimes 2}) \neq 0$  is the set of points for which  $D_t$  is a canonical divisor of  $X_t$ . If  $T' = T \setminus T_0$ ,  $\alpha^{-1}(T') \subset \mathcal{P}_{\mathcal{X}}^{\mathfrak{a}}|_T$  is an affine bundle over  $T'$ . It is striking that, according to [FR21], the branching divisor being canonical for the underlying complex curve

is necessary for a branched projective structure to be a critical point of the holomorphic map  $\mathcal{M}_{\lambda,\rho} \rightarrow \mathcal{T}_g$ . Here  $\mathcal{T}_g$  is the Teichmüller space for genus  $g$ ,  $\lambda$  is a partition of  $k \leq 2g-2$ ,  $\rho$  is a non-elementary representation of  $\pi_1(R)$  in  $\mathrm{PGL}(2, \mathbb{C})$  and  $M_{\lambda,\rho}$  is the complex manifold of branched projective structures on a curve of genus  $g$  with monodromy  $\rho$  and whose branching points follow the partition  $\lambda$ . Moreover, it is proved in [FR21] that the condition is sufficient if the underlying complex curve is hyperelliptic and the divisor is reduced.

**2.5. BRANCHED PROJECTIVE STRUCTURES RESTRICTED TO A DIVISOR.** — To prove Proposition 2.3.3 in its full generality, we need to introduce a technical notion of branched projective structure *restricted* to a divisor. It can be seen as a notion of jet of projective structure. It can also be seen as a way to precise the information contained in a branching class in order to get an analog of Proposition 1.2.2 without the ambiguity in point (ii).

Let  $X$  be a Riemann surface, along with an effective divisor  $D$  and a branching class  $a$  over  $D$ . Let  $D'$  be an effective divisor on  $X$ .

**DEFINITION 2.5.1.** — Let  $U$  and  $U'$  be two open neighborhoods of  $D'$  in  $X$ . Write  $V = U \cap U'$ . Let  $p_1, p_2$  be branched projective structures, defined respectively on  $U$  and  $U'$ , of divisor  $D|_U$  (respectively  $D|_{U'}$ ) and branching class  $a|_U$  (respectively  $a|_{U'}$ ). The projective structures  $p_1$  and  $p_2$  are said to *coincide over the divisor  $D'$*  if the difference  $p_2|_V - p_1|_V \in H^0(V, K_V^{\otimes 2}(-D|_V))$  vanishes over the divisor  $D'$ , i.e.,

$$p_2|_V - p_1|_V \in H^0(V, K_X^{\otimes 2}(-D|_V - D'|_V)).$$

A *branched projective structure restricted to  $D'$*  of divisor  $D$  and branching class  $a$  is an equivalence class for the above equivalence relation. We denote by  $P_{X,D'}^a$  the set of branched projective structures of divisor  $D$  and branching class  $a$  restricted to  $D'$ .

Any local holomorphic map defined around the points of  $D'$  with branching class  $a$  determines an element of  $P_{X,D'}^a$ . Proposition 1.2.2 implies the following:

**PROPOSITION 2.5.2.** — *Let  $D$  be an effective divisor on a Riemann surface  $X$ , let  $a \in A_X^D$  and  $D'$  an effective divisor on  $X$ . The space  $P_{X,D'}^a$  is an affine space directed by  $H^0(X, K_X^{\otimes 2}(-D)|_{D'})$ .*

Now let  $r \in P_{X,D'}^a$  be a branched projective structure on  $X$  with branching class  $a$  restricted to  $D'$ . We denote by  $\mathcal{P}_X^{a,r}$  the space of branched projective structures of branching class  $a$  whose restriction to  $D'$  is  $r$ . The following proposition is a clear consequence of Proposition 1.2.2.

**PROPOSITION 2.5.3.** — *Let  $D$  be an effective divisor on a Riemann surface  $X$ ,  $a \in A_X^D$  and  $D'$  an effective divisor on  $X$ . Let  $r \in \mathcal{P}_{X,D'}^a$  be a restricted branched projective structure.*

(i) *There exists an obstruction  $\bar{c}(a, r) \in H^1(X, K_X^{\otimes 2}(-D - D'))$  such that*

$$\bar{c}(a, r) = 0 \quad \text{if and only if} \quad P_X^{a,r} \neq \emptyset.$$

(ii) The set  $P_X^{a,r}$ , if nonempty, is an affine space directed by the vector space

$$H^0(X, K_X^{\otimes 2}(-D - D')).$$

In particular, if  $D'$  is chosen so that

$$\deg(D) + \deg(D') > 4g - 4,$$

then  $h^0(X, K_X^{\otimes 2}(-D - D')) = 0$ , thus  $P_X^{a,r}$  contains at most one point.

**2.6. FAMILIES OF RESTRICTED BRANCHED PROJECTIVE STRUCTURES.** — In this paragraph we complete the proof of Proposition 2.3.3. By a *family of branching classes with auxiliary divisors of degree  $n'$*  we mean a family  $(T, \mathcal{X}, \pi, \mathcal{D}, \mathbf{a}, \mathcal{D}')$  where  $(T, \mathcal{X}, \pi, \mathcal{D}, \mathbf{a})$  is a family of branching classes and  $\mathcal{D}'$  is an effective divisor on  $\mathcal{X}$  that contains no fiber of  $\pi$ , such that the restriction of  $\mathcal{D}'$  to any fiber of  $\pi$  has degree  $n'$ .

**DEFINITION 2.6.1.** — Let  $\mathcal{G} = (T, \mathcal{X}, \pi, \mathcal{D}, \mathbf{a}, \mathcal{D}')$  be a family of branching classes with auxiliary divisors. Let  $\mathcal{U}$  and  $\mathcal{U}'$  be two open neighborhoods of  $\mathcal{D}'$  in  $\mathcal{X}$ . Set  $\mathcal{V} = \mathcal{U} \cap \mathcal{U}'$ . Let  $\mathbf{p}_1, \mathbf{p}_2$  be relative branched projective structures, defined respectively on  $\mathcal{U}$  and  $\mathcal{U}'$ , of divisor  $\mathcal{D}|_{\mathcal{U}}$  (respectively  $\mathcal{D}|_{\mathcal{U}'}$ ), and of branching class  $\mathbf{a}|_{\mathcal{U}}$  (respectively  $\mathbf{a}|_{\mathcal{U}'}$ ). The projective structures  $\mathbf{p}_1$  and  $\mathbf{p}_2$  are said to *coincide over the divisor  $\mathcal{D}'$*  if the difference  $\mathbf{p}_2|_{\mathcal{V}} - \mathbf{p}_1|_{\mathcal{V}} \in H^0(\mathcal{V}, K_{\mathcal{V}/T}^{\otimes 2}(-\mathcal{D}|_{\mathcal{V}}))$  vanishes over the divisor  $\mathcal{D}'$ , i.e.,  $\mathbf{p}_2|_{\mathcal{V}} - \mathbf{p}_1|_{\mathcal{V}} \in H^0(\mathcal{V}, K_{\mathcal{V}/T}^{\otimes 2}(-\mathcal{D}|_{\mathcal{V}} - \mathcal{D}'))$ .

A *branched projective structure on  $\mathcal{X}$ , relative to  $T$ , restricted to  $\mathcal{D}'$*  of divisor  $\mathcal{D}$  and branching class  $\mathbf{a}$  is an equivalence class for the above equivalence relation.

**DEFINITION 2.6.2**

(i) Let  $n' \in \mathbb{N}$ . A *family of restricted branched projective structures of auxiliary degree  $n'$*  is given by a family  $\mathcal{F} = (S, \mathcal{X}, \pi, \mathcal{D}, \mathbf{a}, \mathcal{D}', \mathbf{r})$ , where  $(S, \mathcal{X}, \pi, \mathcal{D}, \mathbf{a}, \mathcal{D}')$  is a family of branching classes with auxiliary divisors of degree  $n'$  and  $\mathbf{r}$  is a branched projective structure on  $\mathcal{X}$  relative to  $S$ , restricted to  $\mathcal{D}'$ , with branching divisor  $\mathcal{D}$  and branching class  $\mathbf{a}$ .

(ii) Let  $\mathcal{G}$  be a family of branching classes with auxiliary divisors of basis  $T$ . A *family of restricted branched projective structures over  $\mathcal{G}$*  is a family of restricted branched projective structures  $\mathcal{F}$  of basis  $S$ , along with a morphism  $f : S \rightarrow T$  such that the underlying family of branching classes with auxiliary divisors is  $f^*\mathcal{G}$ .

(iii) Let  $S$  be an analytic space,  $\mathcal{F}$  a family of restricted branched projective structures over  $\mathcal{G}$  with basis  $S$ . The family  $\mathcal{F}$  is said to be *universal* if any family  $\mathcal{F}'$  of restricted branched projective structures over  $\mathcal{G}$  is uniquely obtained as the pullback of  $\mathcal{F}$  by some morphism  $g : S' \rightarrow S$ , where  $S'$  is the basis of  $\mathcal{F}'$ .

Fix  $\mathcal{G} = (T, \mathcal{X}, \pi, \mathbf{a})$  a family of branching classes of genus  $g$  and degree  $n$ , as well as an effective divisor  $\mathcal{D}'$  that contains no fiber of  $\mathcal{G}$ . Write  $\mathcal{H}$  the corresponding family of branching classes with auxiliary divisors and  $n'$  the auxiliary degree.

For  $t \in T$ , write respectively  $X_t$ ,  $D_t$ ,  $a_t$  and  $D'_t$  the curve, branching divisor, branching class and auxiliary divisor in the fiber of  $\mathcal{H}$ . Consider the union

$$\mathcal{P}_{\mathcal{X}, \mathcal{D}'}^{\mathbf{a}} = \{(t, r) \mid t \in T, r \in P_{X_t, D'_t}^{a_t}\}$$

of the spaces of restricted projective structures on the fibers of  $\mathcal{H}$ . There is an obvious map

$$\bar{\alpha} : \mathcal{P}_{\mathcal{X}, \mathcal{D}'}^{\mathbf{a}} \longrightarrow T.$$

The following lemma is a direct consequence of Lemma 2.4.1:

LEMMA 2.6.3. — *Let  $\mathbf{r}_0$  be a branched projective structure on  $\mathcal{X}$ , relative to  $T$ , restricted to  $\mathcal{D}'$ , of divisor  $\mathcal{D}$  and relative branching class  $\mathbf{a}$ .*

(i) *If  $\mathbf{r}$  is another such structure, then the difference  $\mathbf{r} - \mathbf{r}_0$  is a holomorphic section of the vector bundle  $K_{\mathcal{X}/T}^{\otimes 2}(-\mathcal{D})|_{\mathcal{D}'}$ .*

(ii) *If  $q \in H^0(\mathcal{D}', K_{\mathcal{X}/T}^{\otimes 2}(-\mathcal{D})|_{\mathcal{D}'})$  is a section, then there exists  $\mathbf{r}$  a branched projective structure on  $\mathcal{X}$ , relative to  $T$ , restricted to  $\mathcal{D}'$ , of relative branching class  $\mathbf{a}$ , such that  $\mathbf{r} - \mathbf{r}_0 = q$ .*

The dimension of  $H^0(D'_t, K_{X_t}^{\otimes 2}(-D_t)|_{D'_t})$  does not depend on  $t \in T$ . Thus the sheaf  $\pi_*(K_{\mathcal{X}/T}^{\otimes 2}(-\mathcal{D})|_{\mathcal{D}'})$  is a vector bundle on  $T$ , and giving a section in  $H^0(\mathcal{D}', K_{\mathcal{X}/T}^{\otimes 2}(-\mathcal{D})|_{\mathcal{D}'})$  is equivalent to giving a section in  $H^0(T, \pi_*(K_{\mathcal{X}/T}^{\otimes 2}(-\mathcal{D})|_{\mathcal{D}'}))$ . As a consequence,  $\mathcal{P}_{\mathcal{X}, \mathcal{D}'}^{\mathbf{a}}$  has the structure of an affine bundle over  $T$ , whose holomorphic (local) sections are the ones defined by (local over  $T$ ) families of branched projective structures on  $\mathcal{G}$  restricted to  $\mathcal{D}'$ .

The pullback family of branching classes

$$\bar{\alpha}^* \mathcal{G} = (\mathcal{P}_{\mathcal{X}, \mathcal{D}'}^{\mathbf{a}}, \bar{\alpha}^* \mathcal{X}, \bar{\alpha}^* \pi, \bar{\alpha}^* \mathcal{D}, \bar{\alpha}^* \mathbf{a})$$

comes with a tautological relative branched projective structure  $\mathbf{r}_0$  restricted to  $\bar{\alpha}^* \mathcal{D}'$ : if  $(t, r) \in \mathcal{P}_{\mathcal{X}, \mathcal{D}'}^{\mathbf{a}}$ , the branched projective structure defined by  $\mathbf{r}_0$  on  $(\bar{\alpha}^* \pi)^{-1}(t, r) = X_t$  with branching class  $a_t$  is  $r$ .

The discussion around Lemma 2.2.7 applies *mutatis mutandis*, to show that the family

$$(\mathcal{P}_{\mathcal{X}, \mathcal{D}'}^{\mathbf{a}}, \bar{\alpha}^* \mathcal{X}, \bar{\alpha}^* \pi, \bar{\alpha}^* \mathcal{D}, \bar{\alpha}^* \mathbf{a}, \bar{\alpha}^* \mathcal{D}', \mathbf{r}_0),$$

along with the morphism  $\bar{\alpha} : \mathcal{P}_{\mathcal{X}}^{\mathbf{a}} \rightarrow T$  is a universal family of branched projective structures over  $\mathcal{H}$ . We thus have the following lemma:

LEMMA 2.6.4. — *If  $\mathcal{G}$  is a family of branching classes with auxiliary divisors, there exists a universal family of restricted branched projective structures over  $\mathcal{G}$ .*

Now let  $\mathcal{J} = (T, \mathcal{X}, \pi, \mathcal{D}, \mathbf{a}, \mathcal{D}', \mathbf{r})$  be a family of restricted branched projective structures with genus  $g$ , branching degree  $n$  and auxiliary degree  $n'$ . Suppose  $n + n' > 4g - 4$ . Denote by  $r_t$  the restricted branched projective structure in  $\mathcal{J}$  over  $t \in T$ . Consider the set

$$\mathcal{P}_{\mathcal{X}}^{\mathbf{a}, \mathbf{r}} = \{(t, p) \mid t \in T, p \in P_{X_t}^{a_t, r_t}\}$$

of branched projective structures whose restrictions are given by the fibers of  $\mathcal{J}$ . There is an obvious map

$$\rho : \mathcal{P}_{\mathcal{X}}^{\mathbf{a}, \mathbf{r}} \longrightarrow T.$$

According to Proposition 2.5.3, the fiber of  $\rho$  over  $t \in T$  is either empty or an affine space directed by  $H^0(X_t, K_{X_t}^{\otimes 2}(-D_t - D'_t))$ . Since moreover  $\deg(D_t) + \deg(D'_t) = n + n' > 4g - 4$ , the line bundle  $K_{X_t}^{\otimes 2}(-D_t - D'_t)$  has negative degree, thus has no

nonzero section. As a consequence, the fibers of  $\alpha$  contain at most one point, and  $\mathcal{P}_{\mathcal{X}}^{\mathbf{a}, \mathbf{r}}$  can be seen as a subset of  $T$ .

Similarly to Section 2.4, consider the sheaf  $\mathcal{BR}_{\mathcal{X}}^{\mathbf{a}, \mathbf{r}}$  of branched projective structures on  $\mathcal{X}$ , relative to  $T$ , branched on  $\mathcal{D}$ , of relative branching class  $\mathbf{a}$ , whose restriction to  $\mathcal{D}'$  is  $\mathbf{r}$ . This is a locally nonempty sheaf of affine spaces directed by  $K_{\mathcal{X}/T}^{\otimes 2}(-\mathcal{D} - \mathcal{D}')$  and thus is associated to a cohomology class  $\bar{\mathcal{C}}(\mathbf{a}, \mathbf{r}) \in H^1(X, K_{\mathcal{X}/T}^{\otimes 2}(-\mathcal{D} - \mathcal{D}'))$  that vanishes if and only if  $\mathcal{BR}_{\mathcal{X}}^{\mathbf{a}, \mathbf{r}}$  has global sections. One also has an exact sequence similar to (6):

$$(8) \quad H^1(T, \pi_* K_{\mathcal{X}/T}^{\otimes 2}(-\mathcal{D} - \mathcal{D}')) \longrightarrow H^1(\mathcal{X}, K_{\mathcal{X}/T}^{\otimes 2}(-\mathcal{D} - \mathcal{D}')) \\ \xrightarrow{\bar{\chi}_{\mathcal{H}}} H^0(T, R^1 \pi_* K_{\mathcal{X}/T}^{\otimes 2}(-\mathcal{D} - \mathcal{D}')).$$

Write

$$\bar{\mathbf{c}}(\mathbf{a}, \mathbf{r}) = \bar{\chi}_{\mathcal{H}}(\bar{\mathcal{C}}(\mathbf{a}, \mathbf{r})).$$

Since  $h^0(K_{\mathcal{X}_t}^{\otimes 2}(-D_t - D'_t))$  vanishes for any  $t \in T$ , Riemann-Roch formula implies that  $h^1(K_{\mathcal{X}_t}^{\otimes 2}(-D_t - D'_t))$  does not depend on  $t \in T$ , so that  $R^1 \pi_* K_{\mathcal{X}/T}^{\otimes 2}(-\mathcal{D} - \mathcal{D}')$  is a vector bundle on  $T$  of which  $\bar{\mathbf{c}}(\mathbf{a}, \mathbf{r})$  is a holomorphic section. For any  $t \in T$ , one has  $\bar{\mathbf{c}}(\mathbf{a}, \mathbf{r}) = \bar{\mathbf{c}}(a_t, r_t)$ . Thus  $\mathcal{P}_{\mathcal{X}}^{\mathbf{a}, \mathbf{r}}$  is the zero locus of the section  $\bar{\mathbf{c}}(\mathbf{a}, \mathbf{r})$ .

Let  $\mathcal{J}' = (\mathcal{P}_{\mathcal{X}}^{\mathbf{a}, \mathbf{r}}, \rho^* \mathcal{X}, \rho^* \pi, \rho^* \mathcal{D}, \rho^* \mathbf{a}, \rho^* \mathcal{D}', \rho^* \mathbf{r})$ . The map  $\chi_{\mathcal{J}'}$ , defined as in (8), is the zero map. Moreover, if  $T$  is Stein, the exact sequence (8) has vanishing first term, which implies that the second term also vanishes as soon as  $\chi_{\mathcal{J}}$  vanishes. As a consequence, up to restricting  $\mathcal{P}_{\mathcal{X}}^{\mathbf{a}, \mathbf{r}}$  to an open subset, there is a branched projective structure on  $\rho^* \mathcal{X}$  relative to  $\mathcal{P}_{\mathcal{X}}^{\mathbf{a}, \mathbf{r}}$  of branching class  $\rho^* \mathbf{a}$  and whose restriction to  $\rho^* \mathcal{D}'$  is  $\mathbf{r}$ . Such a relative branched projective structure is unique since the fibers of  $\rho$  contain at most one element. We denote it by  $\mathbf{p}_0$ .

A family of branched projective structures over the family of restricted branched projective structures  $\mathcal{J} = (T, \mathcal{X}, \pi, \mathcal{D}, \mathbf{a}, \mathcal{D}', \mathbf{r})$  is a family of branched projective structures  $\mathcal{F}_1 = (T_1, \mathcal{X}_1, \pi_1, \mathcal{D}_1, \mathbf{a}_1, \mathbf{p}_1)$  along with a morphism  $f : T \rightarrow T'$  such that the family of branching classes  $(T_1, \mathcal{X}_1, \pi_1, \mathcal{D}_1, \mathbf{a}_1)$  is the pullback by  $f$  of  $(T, \mathcal{X}, \pi, \mathcal{D}, \mathbf{a})$  and the restriction of  $\mathbf{p}_1$  to  $f^* \mathcal{D}'$  is  $f^* \mathbf{r}$ . Such a family  $\mathcal{F}_1$  is *universal* if any other such family is uniquely obtained as a pullback of  $\mathcal{F}_1$ .

Clearly, the family of branched projective structures  $(\mathcal{P}_{\mathcal{X}}^{\mathbf{a}, \mathbf{r}}, \rho^* \mathcal{X}, \rho^* \pi, \rho^* \mathcal{D}, \rho^* \mathbf{a}, \mathbf{p}_0)$ , along with the morphism  $\rho : \mathcal{P}_{\mathcal{X}}^{\mathbf{a}, \mathbf{r}} \rightarrow T$  is a universal family of branched projective structures over  $\mathcal{J}$ . We have shown:

**LEMMA 2.6.5.** — *Let  $\mathcal{J}$  be a family of restricted branched projective structures of genus  $g$ , branching degree  $n$  and auxiliary degree  $n'$ . If  $n + n' > 4g - 4$ , there exists a universal family of branched projective structures over  $\mathcal{J}$ .*

Now we are able to complete the proof of Proposition 2.3.3. Let  $\mathcal{G}$  be a family of branching classes of genus  $g$  and branching degree  $n$ , of basis  $T_{\mathcal{G}}$ . Take  $\mathcal{D}'$  a divisor on the total space of  $\mathcal{G}$  that contains no fiber of  $\mathcal{G}$  and let  $\mathcal{H}$  be the corresponding family of branching classes with auxiliary divisors. Write  $n'$  the auxiliary degree of  $\mathcal{H}$ ,



and choose  $\mathcal{D}'$  such that  $n + n' > 4g - 4$ . Let  $\mathcal{J}$  be the universal family of restricted projective structures over  $\mathcal{H}$ ,  $T_{\mathcal{J}}$  its basis and  $f_{\mathcal{J}\mathcal{J}} : T_{\mathcal{J}} \rightarrow T_{\mathcal{G}}$  the associated morphism. Now, let  $\mathcal{F}$ , of basis  $T_{\mathcal{F}}$ , be the universal family of branched projective structures over  $\mathcal{J}$ . Write  $f_{\mathcal{J}\mathcal{F}} : T_{\mathcal{F}} \rightarrow T_{\mathcal{J}}$  the associated morphism. The family  $\mathcal{F}$ , along with the morphism  $f_{\mathcal{G}\mathcal{F}} = f_{\mathcal{G}\mathcal{J}} \circ f_{\mathcal{J}\mathcal{F}} : T_{\mathcal{F}} \rightarrow T_{\mathcal{G}}$  is a family of branched projective structures over  $\mathcal{G}$ .

Let  $\mathcal{F}_1$ , of basis  $T_{\mathcal{F}_1}$  with a morphism  $f_{\mathcal{G}\mathcal{F}_1} : T_{\mathcal{F}_1} \rightarrow T_{\mathcal{G}}$ , be another family of branched projective structures over  $\mathcal{G}$ . By restricting the relative branched projective structure of  $\mathcal{F}_1$  to the pullback auxiliary divisor  $f_{\mathcal{G}\mathcal{F}_1}^* \mathcal{D}'$ , one gets a family of restricted branched projective structures  $\mathcal{J}_1$ , of basis  $T_{\mathcal{F}_1}$ , over the family  $\mathcal{H}$ , the associated morphism being  $f_{\mathcal{G}\mathcal{F}_1}$ . By universality of  $\mathcal{J}$ , the family  $\mathcal{J}_1$  is the pullback of  $\mathcal{J}$  by some morphism  $f_{\mathcal{J}\mathcal{F}_1}$  verifying  $f_{\mathcal{G}\mathcal{F}_1} = f_{\mathcal{G}\mathcal{J}} \circ f_{\mathcal{J}\mathcal{F}_1}$ . As a consequence,  $\mathcal{F}_1$  is a family of branched projective structures over  $\mathcal{J}$ , the associated morphism being given by  $f_{\mathcal{J}\mathcal{F}_1}$ . By universality of  $\mathcal{F}$ , there exists a morphism  $f_{\mathcal{F}\mathcal{F}_1} : T_{\mathcal{F}_1} \rightarrow T_{\mathcal{F}}$  such that  $\mathcal{F}_1 = f_{\mathcal{F}\mathcal{F}_1}^* \mathcal{F}$  and  $f_{\mathcal{J}\mathcal{F}_1} = f_{\mathcal{J}\mathcal{F}} \circ f_{\mathcal{F}\mathcal{F}_1}$  thus  $f_{\mathcal{G}\mathcal{F}_1} = f_{\mathcal{G}\mathcal{F}} \circ f_{\mathcal{F}\mathcal{F}_1}$ .

It remains to show that  $f_{\mathcal{F}\mathcal{F}_1}$  is the unique such morphism. Suppose given  $f : T_{\mathcal{F}_1} \rightarrow T_{\mathcal{F}}$  such that  $\mathcal{F}_1 = f^* \mathcal{F}$  and  $f_{\mathcal{G}\mathcal{F}_1} = f_{\mathcal{G}\mathcal{F}} \circ f$ . The family of restricted branched projective structures  $\mathcal{J}_1$  is obtained by pulling back  $\mathcal{J}$  by the morphism  $f \circ f_{\mathcal{J}\mathcal{F}}$ . By universality of  $\mathcal{J}$ , one has  $f_{\mathcal{J}\mathcal{F}} \circ f = f_{\mathcal{J}\mathcal{F}_1}$ . Thus  $\mathcal{F}_1$  is the pullback of  $\mathcal{F}$  by  $f$  as a family of branched projective structures over  $\mathcal{J}$ . Since  $\mathcal{F}$  is a universal such family,  $f = f_{\mathcal{F}\mathcal{F}_1}$ . This concludes the proof of Proposition 2.3.3.

### 3. HOLONOMY AND SINGULARITIES OF THE MODULI SPACE OF MARKED BRANCHED PROJECTIVE STRUCTURES IN GENUS AT LEAST TWO

In this section, we prove the following theorem. Recall that if  $n, g \in \mathbb{N}$ ,  $\mathcal{P}_g(n)$  stands for the moduli space of isomorphism classes of branched projective structures of genus  $g$  and branching degree  $n$ .

**THEOREM 3.0.1.** — *Let  $n, g \in \mathbb{N}$ , with  $g \geq 2$ . Let  $p$  be a branched projective structure of genus  $g$  and branching degree  $n$  on a marked Riemann surface  $X$ . Let  $\rho \in \text{Hom}(\pi_1(X), \text{PSL}(2, \mathbb{C}))$  be a monodromy representation for  $g$ . The branched projective structure  $p$  is a singular point of the moduli space  $\mathcal{P}_g(n)$  if and only if the image  $\text{Im } \rho \subset \text{PSL}(2, \mathbb{C})$  of the monodromy representation  $\rho$  has commutative image and the action of  $\text{Im } \rho$  on  $\mathbb{CP}^1$  admits a fixed point.*

#### REMARK 3.0.2

- (i) It is equivalent to say that  $p$  is a singular point of  $\mathcal{P}_g(n)$  if and only if, up to conjugation,  $\text{Im } \rho$  is contained either in  $\{z \mapsto \alpha z \mid \alpha \in \mathbb{C}^*\}$  or in  $\{z \mapsto z + \alpha \mid \alpha \in \mathbb{C}\}$ .
- (ii) In particular, branched projective structures with non-elementary holonomy are smooth points of  $\mathcal{P}_g(n)$ . The holonomy of a branched projective structure whose degree is either odd or smaller than  $2g - 2$  is non-elementary (see [GKM00]). Thus if  $n$  is odd or  $n < 2g - 2$ , then  $\mathcal{P}_g(n)$  is smooth.

(iii) A branched projective structure in  $\mathcal{P}_g(n)$  is a smooth point if and only if its holonomy is a smooth point of the analytic space  $\text{Hom}(\pi_1(X), \text{PGL}(2, \mathbb{C}))$  (see [Hub81] for singularities of the latter), where  $X$  is the underlying Riemann surface.

(iv) The holonomy  $\rho \in \text{Hom}(\pi_1(X), \text{PGL}(2, \mathbb{C}))$  of a branched projective structure  $p$  is only defined up to conjugation with an element of  $\text{PGL}(2, \mathbb{C})$ . However, the criterion on  $\rho$  for  $p$  to be singular does not depend on the choice of  $\rho$ .

In order to study the singularities of  $\mathcal{P}_g(n)$ , we need to have an insight in its tangent spaces. We achieve this by studying the infinitesimal automorphisms of branched projective structures.

### 3.1. INFINITESIMAL AUTOMORPHISMS OF A BRANCHED PROJECTIVE STRUCTURE

Let  $X$  be a compact Riemann surface of genus  $g \geq 2$  and  $p$  a branched projective structure on  $X$  of branching divisor  $D$  and branching class  $a \in A_X^D$ . Denote by  $\Lambda_p$  the sheaf of *infinitesimal automorphisms* of the branched projective structure  $p$ . A local vector field  $V$  defined on an open subset  $U \subset X$  is a local section of  $\Lambda_p$  if the flow of  $V$  (defined at small times on relatively compact subsets of  $U$ ) preserves the charts of  $p$ . More precisely, if  $\phi_V^t$  is the flow of  $V$  at time  $t \in \mathbb{C}$ , then  $V \in \Gamma(U, \Lambda_p)$  if and only if for any relatively compact open subset  $U' \subset U$ , any chart  $f : U' \rightarrow \mathbb{CP}^1$  of  $p$  and any small enough  $t \in \mathbb{C}$ , the map  $f \circ \phi_V^t$  is again a chart of  $p$ .

The sheaf  $\Lambda_p$  is a subsheaf of  $T_X(-D^{\text{red}})$ , where  $D^{\text{red}}$  is the reduced divisor associated to the branching divisor  $D$ . A germ of holomorphic vector field  $V_{(x)}$  at  $x \in X$  is in the stalk of  $\Lambda_p$  at  $x$  if and only if  $V_{(x)}$  is the pullback of a germ of projective vector field by some chart of  $p$ .

Let  $\rho \in \text{Hom}(\pi_1(X), \text{PSL}(2, \mathbb{C}))$  be a monodromy representation of  $p$ . Let  $\text{Ad}_p$  be the sheaf on  $X$  whose stalk at  $x \in X$  contains germs of meromorphic vector fields at  $x$  that are obtained by pulling back a germ of projective vector field by a chart of the branched projective structure  $p$ . In particular,  $\text{Ad}_p$  is a subsheaf of the sheaf of meromorphic functions on  $X$  and  $\Lambda_p$  is a subsheaf of  $\text{Ad}_p$ . The sheaf  $\text{Ad}_p$  is a local system whose stalks are 3-dimensional Lie algebras modeled on  $\mathfrak{sl}(2, \mathbb{C})$ , and whose holonomy is given by  $\rho$ , where  $\text{PSL}(2, \mathbb{C})$  acts on its Lie algebra by the adjoint action.

Let  $x \in X$  be a branched point of  $p$  of degree  $n$  and  $z$  a local coordinate of  $X$  centered at  $x$  such that  $f(z) = z^{n+1} \in \mathbb{CP}^1 = \mathbb{C} \cup \{\infty\}$  is a chart of  $p$ . The pullback by  $f$  of the vector field  $W = (a_0 + a_1\zeta + a_2\zeta^2) \partial_\zeta$ , where  $\zeta$  is the classical coordinate on  $\mathbb{CP}^1 = \mathbb{C} \cup \{\infty\}$ , is given by

$$f^*W = \left( a_0 \frac{1}{z^n} + a_1 z + a_2 z^{n+2} \right) \partial_z.$$

This implies that the quotient  $\text{Ad}_p / \Lambda_p$  is supported on the branched points of  $p$  and the stalk of  $\text{Ad}_p / \Lambda_p$  at any branched point of  $p$  is a complex line, whatever the branching degree. The short exact sequence

$$0 \longrightarrow \Lambda_p \longrightarrow \text{Ad}_p \longrightarrow \text{Ad}_p / \Lambda_p \longrightarrow 0$$

gives in cohomology the long exact sequence (recall that  $X$  has no infinitesimal automorphism, so  $H^0(X, \Lambda_p) = 0$ )

$$(9) \quad 0 \longrightarrow H^0(X, \text{Ad}_p) \longrightarrow H^0(X, \text{Ad}_p / \Lambda_p) \longrightarrow H^1(X, \Lambda_p) \longrightarrow H^1(X, \text{Ad}_p) \\ \longrightarrow 0 \longrightarrow H^2(X, \Lambda_p) \longrightarrow H^2(X, \text{Ad}_p) \longrightarrow 0.$$

In particular, the injection  $\Lambda_p \rightarrow \text{Ad}_p$  induces an isomorphism

$$H^2(X, \Lambda_p) \simeq H^2(X, \text{Ad}_p).$$

The sheaf cohomology of the local system  $\text{Ad}_p$  is the same as its cohomology with local coefficients, see [Ste43]. Moreover, Poincaré duality provides an isomorphism  $H^2(X, \text{Ad}_p) \simeq H_0(X, \text{Ad}_p)$ , where the latter group is the homology with local coefficients of  $\text{Ad}_p$ , see [Sun17] and references therein. We have the following lemma, see [Whi78]:

LEMMA 3.1.1. — *Let  $L_0 \subset \mathfrak{g}$  be the linear subspace generated by the family*

$$(\rho(\gamma) \cdot W - W)_{\substack{\gamma \in \pi_1(X) \\ W \in \mathfrak{g}}}.$$

*Let  $L_1 \subset \mathfrak{g}$  be the linear subspace of fixed points of the action on  $\pi_1(X)$  on  $\mathfrak{g}$ :*

$$L_1 = \{W \in \mathfrak{g} \mid \forall \gamma \in \pi_1(X), \rho(\gamma) \cdot W = W\}.$$

*One has*

- (i)  $H_0(X, \text{Ad}_p) = \mathfrak{g}/L_0$ ,
- (ii)  $H^0(X, \text{Ad}_p) = L_1$ .

Take  $W \in \mathfrak{g}$ . Recall that  $\mathfrak{g}$  is the Lie algebra of projective vector fields, so  $W$  can be written  $W = (a_0 + a_1\zeta + a_2\zeta^2) \partial_\zeta$ . A computation shows that if  $g \in G$  is the Möbius transformation  $g : \zeta \mapsto (\alpha\zeta + \beta)/(\gamma\zeta + \delta)$ , then

- (i) if  $\alpha = \delta = 1$ ,  $\gamma = 0$ , i.e.,  $g$  is parabolic fixing  $\infty \in \mathbb{CP}^1$ , then

$$g \cdot W = W + ((a_2\beta^2 - a_1\beta) - 2a_2\beta\zeta) \partial_\zeta;$$

- (ii) if  $\beta = \gamma = 0$ ,  $\delta = 1$ , i.e.,  $g$  is either elliptic or loxodromic fixing 0 and  $\infty$  in  $\mathbb{CP}^1$ , then

$$g \cdot W = W + (a_0(\alpha - 1) + a_2((1/\alpha) - 1)\zeta^2) \partial_\zeta.$$

As a consequence, there are three cases for the dimensions of the spaces in Equation (9). Here we denote by  $r$  the number of branched points of  $p$ , without taking the multiplicity into account, so that  $h^0(X, \text{Ad}_p / \Lambda_p) = r$ . Recall that the Euler characteristic of  $X$  is given by the alternate sum of the Betti numbers of the cohomology of any local system on  $X$ , so in particular  $3(2-2g) = h^0(X, \text{Ad}_p) - h^1(X, \text{Ad}_p) + h^2(X, \text{Ad}_p)$  (here the factor 3 stands for the dimension of the model space for coefficients  $\mathfrak{g}$ ).

- (i) If the image of  $\rho$  is the trivial subgroup of  $G$ ,  $L_0 = \{0\}$ ,  $L_1 = \mathfrak{g}$ , so

$$h^0(X, \text{Ad}_p) = 3 \quad \text{and} \quad h^2(X, \Lambda_p) = h^2(X, \text{Ad}_p) = h_0(X, \text{Ad}_p) = 3.$$

As a consequence  $h^1(X, \text{Ad}_p) = 6g$ , and  $h^1(X, \Lambda_p) = 6g - 3 + r$ .

(ii) Suppose the image  $\text{Im}(\rho) \subset G$  is not the trivial subgroup but all the nontrivial elements of  $\text{Im}(\rho)$  have the same set of fixed points. More precisely, the nontrivial elements of  $\text{Im}(\rho)$  are either all parabolic transformations with the same fixed point (conjugate to  $z \mapsto z + \beta$ ), or all loxodromic or elliptic transformations with the same two fixed points (conjugate to  $z \mapsto \alpha z$ ). Then  $h^0(X, \text{Ad}_\rho) = 1$ , and  $h^2(X, \Lambda_\rho) = h^2(X, \text{Ad}_\rho) = h_0(X, \text{Ad}_\rho) = 1$ . As a consequence  $h^1(X, \text{Ad}_\rho) = 6g - 4$ , and  $h^1(X, \Lambda_\rho) = 6g - 5 + r$ .

(iii) In the other cases, i.e., if the image  $\text{Im}(\rho) \subset G$  contains at least two nontrivial transformations  $g_1$  and  $g_2$  with  $g_1$  not fixing a fixed point of  $g_2$ , then  $h^0(X, \text{Ad}_\rho) = 0$ , and  $h^2(X, \Lambda_\rho) = h^2(X, \text{Ad}_\rho) = h_0(X, \text{Ad}_\rho) = 0$ . As a consequence  $h^1(X, \text{Ad}_\rho) = 6g - 6$ , and  $h^1(X, \Lambda_\rho) = 6g - 6 + r$ .

REMARK 3.1.2. — The group  $h^2(X, \Lambda_\rho)$  is nonzero if and only if the monodromy representation  $\rho$  is a singular point of  $\text{Hom}(\pi_1(X), G)$  (see [Hub81]).

3.2. TANGENT SPACES OF THE UNIVERSAL SPACES OF DIVISORS. — Let  $n, g \in \mathbb{N}$  with  $g \geq 2$ . Following Section 2.3, let  $\mathcal{U}_g = (\mathcal{T}_g, \mathcal{X}_g, \pi_g, m_g)$  be the universal family of marked Riemann surfaces of genus  $g$  ( $m_g$  is the marking). In particular  $\mathcal{T}_g$  is the Teichmüller space. Let

$$\mathcal{U}_g^{\text{div}}(n) = (\mathcal{T}_g(n), \mathcal{X}_g^{\text{div}}(n), \pi_g^{\text{div}}(n), m_g^{\text{div}}(n), \mathcal{D}_g^{\text{div}}(n))$$

be the universal family of marked curves of genus  $g$  with divisors of degree  $n$ , and let  $\gamma_g(n) : \mathcal{T}_g(n) \rightarrow \mathcal{T}_g$  be the forgetful morphism. In particular,  $\mathcal{T}_g(n)$  is the moduli space of marked Riemann surfaces with divisors.

A *partition* of  $n$  is an ordered finite sequence of strictly positive integers  $(k_1 \leq k_2 \leq \dots \leq k_r)$  with  $k_1 + \dots + k_r = n$ . Denote by  $\text{Part}(n)$  the set of partitions of  $n$ . Let  $(X, D, m) \in \mathcal{T}_g(n)$ :  $X$  is a Riemann surface,  $m$  is a marking on  $X$ , and  $D$  is a divisor of degree  $n$  on  $X$ . The divisor  $D$  is given by  $D = k_1 \cdot x_1 + \dots + k_r \cdot x_r$  with  $k_i \in \mathbb{N}^*$ ,  $x_i \in X$  pairwise distinct and  $k_1 + \dots + k_r = n$ . The  $k_i$ 's can be chosen to be ordered, so that the sequence  $(k_1, \dots, k_r)$  is a partition of  $n$ , uniquely determined by  $D$ .

In this way we define a map  $\varpi : \mathcal{T}_g(n) \rightarrow \text{Part}(n)$ . For any partition  $\kappa = (k_1 \leq \dots \leq k_r)$  of  $n$ ,  $\varpi^{-1}(\kappa)$  is a smooth  $3g - 3 + r$ -dimensional analytic subspace of  $\mathcal{T}_g(n)$  (see [BM14] for details on the structure of the symmetric powers of a smooth curve). Let us denote this space by  $\mathcal{T}_g(\kappa) = \varpi^{-1}(\kappa)$ , and by

$$\mathcal{U}_g(\kappa) = (\mathcal{T}_g(\kappa), \mathcal{X}_g^{\text{div}}(\kappa), \mathcal{D}_g^{\text{div}}(\kappa), m_g^{\text{div}}(\kappa), \pi_g^{\text{div}}(\kappa))$$

the restriction of the family  $\mathcal{U}_g(n)$  to the basis  $\mathcal{T}_g(\kappa)$ . Let us also denote by  $\gamma_g(\kappa) : \mathcal{T}_g(\kappa) \rightarrow \mathcal{T}_g$  the forgetful submersion, restriction of  $\gamma_g(n)$ . Finally, denote by  $\text{Sym}_\kappa(X)$  the submanifold of the symmetric power  $\text{Sym}^n(X)$  containing the divisors whose associated partition is  $\kappa$ .

Let us now focus on the space  $\mathcal{T}_g(\kappa)$  from the point of view of infinitesimal deformations of marked curves with divisor. Let  $(X, D, m) \in \mathcal{T}_g(\kappa)$  be a smooth marked

complex curve with divisor ( $m$  is the marking on  $X$ ). One has a first exact sequence:

$$(10) \quad 0 \longrightarrow (T_{\text{Sym}_\kappa(X)})_D \longrightarrow (T_{\mathcal{T}_g(\kappa)})_{(X,D,m)} \longrightarrow (T_{\mathcal{T}_g})_{(X,m)} \longrightarrow 0$$

and another one, denoting by  $D^{\text{red}}$  the reduced divisor on  $X$  with the same support as  $D$ :

$$0 \longrightarrow T_X(-D^{\text{red}}) \longrightarrow T_X \longrightarrow T_X|_{D^{\text{red}}} \longrightarrow 0.$$

Since  $g \geq 2$ , one has in cohomology:

$$0 \longrightarrow H^0(X, T_X|_{D^{\text{red}}}) \longrightarrow H^1(X, T_X(-D^{\text{red}})) \longrightarrow H^1(X, T_X) \longrightarrow 0.$$

There is an obvious isomorphism  $\iota : (T_{\text{Sym}_\kappa(X)})_D \xrightarrow{\sim} H^0(X, T_X|_{D^{\text{red}}})$ . One also has an isomorphism  $\text{KS} : (T_{\mathcal{T}_g})_{(X,m)} \xrightarrow{\sim} H^1(X, T_X)$ , called the Kodaira-Spencer isomorphism, of which we recall the definition.

Consider the exact sequence

$$(11) \quad 0 \longrightarrow T_{\mathcal{X}_g/\mathcal{T}_g} \longrightarrow T_{\mathcal{X}_g} \xrightarrow{d\pi_g} \pi_g^* T_{\mathcal{T}_g} \longrightarrow 0.$$

By applying the left exact functor  $\pi_{g*}$ , it gives a morphism of vector bundles  $\pi_{g*}\pi_g^* T_{\mathcal{T}_g} = T_{\mathcal{T}_g} \rightarrow R^1\pi_{g*}T_{\mathcal{X}_g/\mathcal{T}_g}$ . The map  $\text{KS}$  is the former isomorphism restricted to the fiber above  $(X, m)$  (the equality  $\pi_{g*}\pi_g^* T_{\mathcal{T}_g} = T_{\mathcal{T}_g}$  stands because  $\pi_g$  has compact fibers). It is a property of the Teichmüller space that the map  $\text{KS}$  is an isomorphism (see for instance [EE69]).

Let us denote by  $\mathcal{D}_g^{\text{div}}(\kappa)$  the restriction of the divisor  $\mathcal{D}_g^{\text{div}}(n)$  to the manifold  $\mathcal{X}_g^{\text{div}}(\kappa)$ . The reduced divisor associated to  $\mathcal{D}_g^{\text{div}}(\kappa)$ , denoted by  $\mathcal{D}_g^{\text{div}}(\kappa)^{\text{red}}$ , is a complex manifold (a smooth analytic subspace), that is transverse to the fibers of  $\pi_g^{\text{div}}(\kappa)$ . As a consequence, when writing the Kodaira-Spencer exact sequence (11) for the family  $\mathcal{U}_g^{\text{div}}(\kappa)$ , we can restrict to the vector fields on  $\mathcal{X}_g^{\text{div}}(\kappa)$  that are tangent to  $\mathcal{D}_g^{\text{div}}(\kappa)^{\text{red}}$ :

$$0 \longrightarrow T_{\mathcal{X}_g^{\text{div}}(\kappa)/\mathcal{T}_g(\kappa)}(-\mathcal{D}_g^{\text{div}}(\kappa)^{\text{red}}) \longrightarrow T_{\mathcal{X}_g^{\text{div}}(\kappa)}(-\log \mathcal{D}_g^{\text{div}}(\kappa)^{\text{red}}) \xrightarrow{d\pi_g^{\text{div}}(\kappa)} \pi_g^{\text{div}}(\kappa)^* T_{\mathcal{T}_g(\kappa)} \longrightarrow 0.$$

Recall that  $T_{\mathcal{X}_g^{\text{div}}(\kappa)}(-\log \mathcal{D}_g^{\text{div}}(\kappa)^{\text{red}})$  is the sheaf of vector fields on  $\mathcal{X}_g^{\text{div}}(\kappa)$  that are tangent to  $\mathcal{D}_g^{\text{div}}(\kappa)^{\text{red}}$ . Note that the reason why, in the above diagram, the arrow  $d\pi_g^{\text{div}}(\kappa)$  is surjective is that  $\mathcal{D}_g^{\text{div}}(\kappa)^{\text{red}}$  is a smooth analytic subspace transverse to the fibers of  $d\pi_g^{\text{div}}(\kappa)$ . By applying the left exact functor  $\pi_g^{\text{div}}(\kappa)_*$ , one obtains in particular a morphism of vector bundles

$$T_{\mathcal{T}_g(\kappa)} \longrightarrow R^1\pi_g^{\text{div}}(\kappa)_* T_{\mathcal{X}_g^{\text{div}}(\kappa)/\mathcal{T}_g(\kappa)}(-\mathcal{D}_g^{\text{div}}(\kappa)^{\text{red}}).$$

By looking at that morphism at the point  $(X, D, m) \in \mathcal{T}_g(\kappa)$ , one obtains a linear map

$$(12) \quad \text{KS}^{\text{div}} : (T_{\mathcal{T}_g(\kappa)})_{(X,D,m)} \longrightarrow H^1(X, T_X(-D^{\text{red}})).$$

It is easy to see that the following diagram commutes:

$$\begin{array}{ccccccc}
 0 & \longrightarrow & (T_{\text{Sym}_\kappa(X)})_D & \longrightarrow & (T_{\mathcal{T}_g(\kappa)})_{(X,D,m)} & \longrightarrow & (T_{\mathcal{T}_g})_{(X,m)} \longrightarrow 0 \\
 & & \downarrow \iota & & \downarrow \text{KS}^{\text{div}} & & \downarrow \text{KS} \\
 0 & \longrightarrow & H^0(X, T_X|_{D^{\text{red}}}) & \longrightarrow & H^1(X, T_X(-D^{\text{red}})) & \longrightarrow & H^1(X, T_X) \longrightarrow 0
 \end{array}$$

By the five lemma, we have:

LEMMA 3.2.1. — *The map  $\text{KS}^{\text{div}}$  in (12) is an isomorphism.*

3.3. TANGENT SPACES OF THE UNIVERSAL SPACES OF BRANCHING CLASSES. — Let  $X$  be a Riemann surface of genus  $g \geq 2$ ,  $D$  a divisor on  $X$  and  $a \in A_X^D$  a branching class. Let  $U$  be an open set in  $X$  and let  $V \in \Gamma(U, T_X(-D^{\text{red}}))$ . For any  $t \in \mathbb{R}$  near 0, denote by  $\phi_V^t$  the flow of  $V$  at time  $t$ . Write

$$\chi_a(V) = \lim_{t \rightarrow 0} \frac{(\phi_V^t)^* a - a}{t} \in \Gamma(U, K_X|_D),$$

where the pullback of branching classes is the pullback of jets of holomorphic maps (thus the pullback of holomorphic maps).

Let  $\zeta$  be a non constant holomorphic map on  $U$ , whose jets at the points of  $D|_U$  are representatives of the branching class  $a|_U$ . In particular,  $\zeta$  has branching divisor  $D|_U$ . Write  $V = f(\zeta)\partial_\zeta$ , where  $f$  is a multivalued holomorphic function. A straightforward computation shows that  $\chi_a(V) = f''(\zeta)d\zeta$ . As a consequence, if  $D|_U = nx$  with  $x \in U$  and if  $z$  is a local coordinate centered at  $x$  and such that  $\zeta = z^{n+1}$ , then if  $V = (a_1z + a_2z^2 + \dots)\partial_z$ , the section  $\chi_a(V) \in \Gamma(U, K_X|_D)$  is given in the coordinate  $z$  by:

$$(13) \quad \chi_a(V) = ((n+2)a_2 + 2(n+3)a_3z + 3(n+4)a_4z^2 + \dots + n(2n+1)a_{n+1}z^{n-1})dz.$$

In particular, the morphism of sheaves  $\chi_a : T_X(-D^{\text{red}}) \rightarrow K_X|_D$  is surjective. Denoting by  $\Gamma_a$  its kernel, one gets an exact sequence:

$$(14) \quad 0 \longrightarrow \Gamma_a \longrightarrow T_X(-D^{\text{red}}) \xrightarrow{\chi_a} K_X|_D \longrightarrow 0$$

and thus in cohomology

$$(15) \quad 0 \longrightarrow H^0(X, K_X|_D) \longrightarrow H^1(X, \Gamma_a) \longrightarrow H^1(X, T_X(-D^{\text{red}})) \longrightarrow 0.$$

Let  $n \in \mathbb{N}$ . Following Section 2.3, let

$$\mathcal{U}_g^{\text{bc}} = (\mathcal{A}_g(n), \mathcal{X}_g^{\text{bc}}(n), \pi_g^{\text{bc}}(n), m_g^{\text{bc}}(n), \mathcal{D}_g^{\text{bc}}(n), \mathfrak{a}_g^{\text{bc}}(n))$$

be the universal family of marked branching classes of genus  $g$  and degree  $n$ . Write  $\delta_g(n) : \mathcal{A}_g(n) \rightarrow \mathcal{T}_g(n)$  the forgetful morphism. According to Section 2.2,  $\mathcal{A}_g(n)$  is an affine bundle over  $\mathcal{T}_g(n)$ .

For any partition  $\kappa$  of the integer  $n$ , denote by  $\mathcal{A}_g(\kappa)$  the bundle  $\mathcal{A}_g(n) \rightarrow \mathcal{T}_g(n)$  restricted to the basis  $\mathcal{T}_g(\kappa) \subset \mathcal{T}_g(n)$ . Let

$$\mathcal{U}_g^{\text{bc}}(\kappa) = (\mathcal{A}_g(\kappa), \mathcal{X}_g^{\text{bc}}(\kappa), \mathcal{D}_g^{\text{bc}}(\kappa), \pi_g^{\text{bc}}(\kappa), \mathfrak{a}_g^{\text{bc}}(\kappa))$$

be the pullback by the inclusion  $\mathcal{A}_g(\kappa) \rightarrow \mathcal{A}_g(n)$  of the family  $\mathcal{U}_g^{\text{bc}}(n)$ . Denote by  $\delta_g(\kappa)$  the restriction of  $\delta_g(n)$  to  $\mathcal{A}_g(\kappa)$ .

Let  $(X, D, a, m) \in \mathcal{A}_g(n)$  be a marked curve with divisor of degree  $n$  and branching class. Denote by  $\kappa$  the partition of  $n$  associated to the divisor  $D$ . It is a consequence of Proposition 1.1.2 that the tangent space of the fiber  $A_X^D = \delta_g(\kappa)^{-1}(X, D, m)$  at  $a$  is  $(T_{A_X^D})_a = H^0(X, K_X|_D)$ . Moreover we saw in Section 3.2 that there is a Kodaira-Spencer isomorphism

$$\text{KS}^{\text{div}} : (T_{\mathcal{T}_g(\kappa)})_{(X, D, m)} \xrightarrow{\sim} H^1(X, T_X(-D^{\text{red}})).$$

As a consequence, similarly to Section 3.2, one has identifications between the kernels and cokernels of the exact sequences (15) and

$$0 \longrightarrow (T_{A_X^D})_a \longrightarrow (T_{\mathcal{A}_g(\kappa)})_{(X, D, a, m)} \longrightarrow (T_{\mathcal{T}_g(\kappa)})_{(X, D, m)} \longrightarrow 0.$$

Let  $\mathcal{K}$  (respectively  $\mathcal{K}^{\text{rel}}$ ) be the subsheaf of  $T_{\mathcal{X}_g^{\text{bc}}(\kappa)}$  (respectively  $T_{\mathcal{X}_g^{\text{bc}}(\kappa)/\mathcal{A}_g(\kappa)}$ ) that contains the vector fields preserving the relative branched atlas  $\mathfrak{a}_g^{\text{bc}}(\kappa)$ . The Kodaira-Spencer exact sequence for the family  $\mathcal{X}_g^{\text{bc}}(\kappa)$ , when restricted to the vector fields preserving  $\mathfrak{a}_g^{\text{bc}}(\kappa)$ , is given by:

$$0 \longrightarrow \mathcal{K}^{\text{rel}} \longrightarrow \mathcal{K} \longrightarrow \pi_g^{\text{bc}}(\kappa)^* T_{\mathcal{A}_g(\kappa)} \longrightarrow 0.$$

Applying the left exact functor  $\pi_g^{\text{bc}}(\kappa)_*$ , one gets a morphism of sheaves  $T_{\mathcal{A}_g(\kappa)} \rightarrow R^1 \pi_g^{\text{bc}}(\kappa)_* \mathcal{K}^{\text{rel}}$ . In particular, at the point  $(X, D, a, m)$ , this morphism gives a linear map:

$$(16) \quad \text{KS}^{\text{bc}} : (T_{\mathcal{A}_g(\kappa)})_{(X, D, a, m)} \longrightarrow H^1(X, \Gamma_a).$$

It is easy to see that the following diagram is commutative:

$$\begin{array}{ccccccc} 0 & \longrightarrow & (T_{A_X^D})_a & \longrightarrow & (T_{\mathcal{A}_g(\kappa)})_{(X, D, a, m)} & \longrightarrow & (T_{\mathcal{T}_g(\kappa)})_{(X, D, m)} \longrightarrow 0 \\ & & \parallel & & \downarrow \text{KS}^{\text{bc}} & & \downarrow \text{KS}^{\text{div}} \\ 0 & \longrightarrow & H^0(X, K_X|_D) & \longrightarrow & H^1(X, \Gamma_a) & \longrightarrow & H^1(X, T_X(-D^{\text{red}})) \longrightarrow 0 \end{array}$$

Since  $\text{KS}^{\text{div}}$  is an isomorphism by Lemma 3.2.1, we have the following lemma:

LEMMA 3.3.1. — *The map  $\text{KS}^{\text{bc}}$  in (16) is an isomorphism.*

### 3.4. TANGENT SPACES OF THE UNIVERSAL SPACES OF BRANCHED PROJECTIVE STRUCTURES

Let  $X$  be a Riemann surface of genus  $g \geq 2$ ,  $D$  a divisor on  $X$ ,  $a \in A_X^D$  a branching class and  $p \in P_X^a$  a branched projective structure. Let  $U$  be an open subset of  $X$  and let  $V \in \Gamma(U, \Gamma_a)$ , where  $\Gamma_a$  is defined in (14). For any  $t \in \mathbb{R}$  that is near 0, the flow of  $V$  at time  $t$  is denoted by  $\phi_V^t$ . Note that, by definition of  $\Gamma_a$ ,  $V \in \Gamma_a$  means that for all  $t$  near 0 one has  $(\phi_V^t)^* a - a = 0$ . Write

$$\xi_p(V) = \lim_{t \rightarrow 0} \frac{(\phi_V^t)^* p - p}{t} \in \Gamma(U, K_X^{\otimes 2}(-D)).$$

Let  $\zeta$  be a chart on  $U$  of the branched projective structure  $p$ . Write  $V = f(\zeta)\partial_\zeta$ , where  $f$  is a multivalued holomorphic function. A straightforward computation shows

that  $\xi_p(V) = f'''(\zeta)d\zeta^{\otimes 2}$ . Suppose  $D|_U = nx$  with  $x \in U$  and let  $z$  be a coordinate centered at  $x$  such that  $\zeta = z^{n+1}$ . Write  $V = (a_1z + a_2z^2 + \cdots)\partial_z$ . Then

$$\xi_p(V) = \left( \sum_{k \geq 1} k(k+n+1)(k-(n+1))a_{k+1}z^{k-2} \right) dz^{\otimes 2}.$$

According to Equation (13),  $V$  is a section of  $\Gamma_a$  if and only if  $a_2 = a_3 = \cdots = a_{n+1} = 0$ . As a consequence:

$$(17) \quad \xi_p(V) = \left( \sum_{\ell \geq n} (\ell+2)(\ell+n+3)(\ell-n+1)a_{\ell+3}z^\ell \right) dz^{\otimes 2}.$$

Since for any  $\ell \geq n$ ,  $(\ell+2)(\ell+n+3)(\ell-n+1) \neq 0$ , the morphism of sheaves  $\xi_p : \Gamma_a \rightarrow K_X^{\otimes 2}(-D)$  is surjective. Denote by  $\Lambda_p$  its kernel. It is the sheaf of infinitesimal automorphisms of  $p$ . There is an exact sequence:

$$0 \longrightarrow \Lambda_p \longrightarrow \Gamma_a \xrightarrow{\xi_p} K_X^{\otimes 2}(-D) \longrightarrow 0$$

which gives, by taking cohomologies, an exact sequence:

$$(18) \quad 0 \longrightarrow H^0(X, K_X^{\otimes 2}(-D)) \longrightarrow H^1(X, \Lambda_p) \longrightarrow H^1(X, \Gamma_a) \\ \longrightarrow H^1(X, K_X^{\otimes 2}(-D)) \longrightarrow H^2(X, \Lambda_p) \longrightarrow 0.$$

For any  $n \in \mathbb{N}$ , according to Section 2.3, let

$$\mathcal{U}_g^{\text{bps}}(n) = (\mathcal{P}_g(n), \mathcal{X}_g^{\text{bps}}(n), \pi_g^{\text{bps}}(n), m_g^{\text{bps}}(n), \mathcal{D}_g^{\text{bps}}(n), \mathfrak{a}_g^{\text{bps}}(n), \mathfrak{p}_g^{\text{bps}}(n))$$

be the universal family of marked branched projective structures of genus  $g$  and degree  $n$ . In particular  $\mathcal{P}_g(n)$  is the moduli space of marked branched projective structures of degree  $n$  and genus  $g$ . Denote by  $\alpha_g(n) : \mathcal{P}_g(n) \rightarrow \mathcal{A}_g(n)$  the forgetful map.

**3.4.1. Case  $n < 2g - 2$  or  $n > 4g - 4$ .** — Let us first handle the case, corresponding to the framework of Proposition 2.4.2, where the branching degree  $n$  and the genus  $g$  verify  $n < 2g - 2$  or  $n > 4g - 4$ .

In this paragraph we prove the following proposition, which, combined with the study of Section 3.1, shows Theorem 3.0.1.

**PROPOSITION 3.4.1.** — *Take  $g \geq 2$  and  $n \in \mathbb{N}$ .*

- (i) *If  $n < 2g - 2$ , the analytic space  $\mathcal{P}_g(n)$  is smooth.*
- (ii) *If  $n > 4g - 4$  and  $p \in \mathcal{P}_g(n)$ ,  $p$  is a smooth point of  $\mathcal{P}_g(n)$  if and only if its sheaf  $\Lambda_p$  of infinitesimal automorphisms satisfies  $H^2(X, \Lambda_p) = 0$ .*

*Proof.* — In the case  $n < 2g - 2$ , according to Proposition 2.4.2,  $\mathcal{P}_g(n)$  is an affine bundle on the smooth variety  $\mathcal{A}_g(n)$ , thus it is smooth. The case  $n > 4g - 4$  is more subtle, it is the reason why we introduced the Kodaira-Spencer machinery.

From now on, suppose  $n > 4g - 4$ . Consider the vector bundle on  $\mathcal{A}_g(n)$  whose fiber over a point  $(X, D, a, m)$  is  $H^1(X, K_X^{\otimes 2}(-D))$ , and denote by  $\mathfrak{c}_g(n) = \mathfrak{c}(\mathfrak{a}_g^{\text{bc}}(n))$  its global section defined in (7). By Proposition 2.4.2, the map  $\alpha_g(n)$  is an embedding and thus identifies  $\mathcal{P}_g(n)$  with an analytic subspace of  $\mathcal{A}_g(n)$ . More precisely, we saw



in the proof of Proposition 2.4.2 that the subspace  $\mathcal{P}_g(n) \subset \mathcal{A}_g(n)$  is given by the zeroes of the section  $\mathfrak{c}_g(n)$ . In particular, if  $(X, D, a, m) \in \mathcal{P}_g(n) \subset \mathcal{A}_g(n)$ , one has  $\mathfrak{c}_g(n)(X, D, a, m) = 0$ . In that case, the differential  $(d\mathfrak{c}_g(n))_{(X, D, a, m)}$  is a linear map from  $(T_{\mathcal{A}_g(n)})_{(X, D, a, m)}$  to  $H^1(X, K_X^{\otimes 2}(-D))$ . We will show that this differential is surjective for any  $p \in \mathcal{A}_g(n)$  such that  $H^2(X, \Lambda_p) = 0$ , which implies smoothness of  $\mathcal{P}_g(n)$  at  $p$ .

For any partition  $\kappa$  of the integer  $n$ , denote by  $\mathcal{P}_g(\kappa)$  the restriction of the bundle  $\mathcal{P}_g(n) \rightarrow \mathcal{A}_g(n)$  to the basis  $\mathcal{A}_g(\kappa) \subset \mathcal{A}_g(n)$ . Let

$$\mathcal{U}_g^{\text{bps}}(\kappa) = (\mathcal{P}_g(\kappa), \mathfrak{X}_g^{\text{bps}}(\kappa), \pi_g^{\text{bps}}(\kappa), m_g^{\text{bps}}(\kappa), \mathcal{D}_g^{\text{bps}}(\kappa), \mathfrak{a}_g^{\text{bps}}(\kappa))$$

be the pullback of the family  $\mathcal{U}_g^{\text{bps}}(n)$  by the inclusion  $\mathcal{P}_g(\kappa) \rightarrow \mathcal{P}_g(n)$ . Denote by  $\alpha_g(\kappa)$  the restriction of  $\alpha_g(n)$  to  $\mathcal{P}_g(\kappa)$ .

Let  $(X, D, a, p, m) \in \mathcal{A}_g(n)$  be a marked curve with divisor of degree  $n$ , branching class and branched projective structure. Let  $\kappa$  be the partition of  $n$  associated to the divisor  $D$ . Proposition 1.2.2 implies that the tangent space of the fiber  $P_X^a = \alpha_g(\kappa)^{-1}(X, D, a, m)$  at  $p$  is  $(T_{P_X^a})_p = H^0(X, K_X^{\otimes 2}(-D))$ . Moreover, as we saw in Section 3.3, there is a Kodaira-Spencer isomorphism

$$\text{KS}^{\text{bc}} : (T_{\mathcal{A}_g(\kappa)})_{(X, D, a, m)} \xrightarrow{\sim} H^1(X, \Gamma_a).$$

Thus the first (respectively the third) terms of the two exact sequences (18) and

$$0 \longrightarrow (T_{P_X^a})_p \longrightarrow (T_{\mathcal{P}_g(\kappa)})_{(X, D, a, p, m)} \longrightarrow (T_{\mathcal{A}_g(\kappa)})_{(X, D, a, m)} \xrightarrow{d\mathfrak{c}_g(\kappa)_{(X, D, a, m)}} H^1(X, K_X^{\otimes 2}(-D))$$

are identified.

Let  $\mathcal{G}$  (respectively  $\mathcal{G}^{\text{rel}}$ ) be the subsheaf of  $T_{\mathcal{X}_g^{\text{bps}}(\kappa)}$  (respectively  $T_{\mathcal{X}_g^{\text{bps}}(\kappa)/\mathcal{P}_g(\kappa)}$ ) of vector fields preserving the relative branched projective structure  $\mathfrak{p}_g(n)$ . The Kodaira-Spencer exact sequence for the family  $\mathcal{X}_g^{\text{bps}}(\kappa)$ , when restricted to vector fields preserving  $\mathfrak{p}_g(n)$ , is given by:

$$0 \longrightarrow \mathcal{G}^{\text{rel}} \longrightarrow \mathcal{G} \longrightarrow \pi_g^{\text{bps}}(\kappa)^* T_{\mathcal{P}_g(\kappa)} \longrightarrow 0.$$

By applying the left exact functor  $\pi_g^{\text{bps}}(\kappa)_*$ , one gets a morphism of sheaves  $T_{\mathcal{P}_g(\kappa)} \rightarrow R^1\pi_g^{\text{bps}}(\kappa)_*\mathcal{G}^{\text{rel}}$ . In particular, at the point  $(X, D, a, p, m)$ , one gets a linear map

$$(19) \quad \text{KS}^{\text{bps}} : (T_{\mathcal{P}_g(\kappa)})_{(X, D, a, p, m)} \longrightarrow H^1(X, \Lambda_p).$$

Since  $n > 4g - 4$ ,  $(T_{P_X^a})_p = H^0(X, K_X^{\otimes 2}(-D)) = 0$ . Thus one has the following diagram, that is easily seen to be commutative:

$$\begin{array}{ccccccc} 0 \rightarrow & (T_{\mathcal{P}_g(\kappa)})_{(X, D, a, p, m)} & \rightarrow & (T_{\mathcal{A}_g(\kappa)})_{(X, D, a, m)} & \xrightarrow{d\mathfrak{c}_g(\kappa)} & H^1(X, K_X^{\otimes 2}(-D)) & \\ & \downarrow \text{KS}^{\text{bps}} & & \downarrow \text{KS}^{\text{bc}} & & \parallel & \\ 0 \longrightarrow & H^1(X, \Lambda_p) & \longrightarrow & H^1(X, \Gamma_a) & \longrightarrow & H^1(X, K_X^{\otimes 2}(-D)) & \rightarrow H^2(X, \Lambda_p) \end{array}$$

Since  $\mathrm{KS}^{\mathrm{bc}}$  is an isomorphism, one has:

LEMMA 3.4.2. — *The map  $\mathrm{KS}^{\mathrm{bps}}$  in (19) is an isomorphism. Moreover the differential  $d\mathbf{c}_g(\kappa)$  is surjective at any point  $p$  of  $\mathcal{P}_g(\kappa)$  such that  $H^2(X, \Lambda_p) = 0$ .*

Let  $p \in \mathcal{P}_g(\kappa)$  be such that  $H^2(X, \Lambda_p) = 0$ . Since the restricted differential at  $d\mathbf{c}_g(\kappa)$  is surjective at  $p$ , the total differential  $d\mathbf{c}_g(n)$  is a fortiori surjective at  $p$ . This implies that  $p$  is a smooth point of  $\mathcal{P}_g(n) \subset \mathcal{A}_g(n)$ . Reciprocally, if  $H^2(X, \Lambda_p) \neq 0$  and  $D$  is reduced, then  $h^1(X, \Lambda_p) = 6g - 6 + n + h^2(X, \Lambda_p) > 6g - 6 + n$ , so  $p$  is singular.

Finally, let  $p \in \mathcal{P}_g(n)$  be any branched projective structure such that  $h^2(X, \Lambda_p) > 0$ . It is shown in [CDF14] that  $p$  is the limit in  $\mathcal{P}_g(n)$  of a sequence  $(p_n)_{n \in \mathbb{N}}$  of branched projective structures with the same holonomy as  $p$  and reduced branching divisor. By Section 3.1, the dimension of  $H^2(X, \Lambda_p)$  depends only on the holonomy of  $\rho$ , thus the points  $p_n$  are all singular points of  $\mathcal{P}_g(n)$ , and since the singular locus is closed, the point  $p$  is itself a singular point.  $\square$

3.4.2. *General case.* — In order to address the case where  $2g - 2 \leq n \leq 4g - 4$ , we need to have a closer look at the point of view on  $\mathcal{P}_g(n)$  given in Section 2.6, and to work with restricted branched projective structures.

Let  $X$  be a Riemann surface of genus  $g \geq 2$ ,  $D$  and  $D'$  divisors on  $X$ ,  $a \in A_X^D$  a branching class and  $r \in P_{X, D'}^a$  a jet of branched projective structure. Let  $U$  be an open subset of  $X$  and let  $V \in \Gamma(U, \Gamma_a)$ , where  $\Gamma_a$  is defined in (14). For any  $t \in \mathbb{R}$  that is near 0, the flow of  $V$  at time  $t$  is still denoted by  $\phi_V^t$ . Write

$$\tilde{\xi}_r(V) = \lim_{t \rightarrow 0} \frac{(\phi_V^t)^* r - r}{t} \in \Gamma(U, K_X^{\otimes 2}(-D)|_{D'}).$$

According to Equation 17, the morphism of sheaves  $\tilde{\xi}_r : \Gamma_a \rightarrow K_X^{\otimes 2}(-D)|_{D'}$  is surjective. Denote by  $\tilde{\Lambda}_r$  its kernel, so that there is an exact sequence:

$$0 \longrightarrow \tilde{\Lambda}_r \longrightarrow \Gamma_a \xrightarrow{\tilde{\xi}_r} K_X^{\otimes 2}(-D)|_{D'} \longrightarrow 0$$

which gives, by taking cohomologies, an exact sequence:

$$(20) \quad 0 \longrightarrow H^0(X, K_X^{\otimes 2}(-D)|_{D'}) \longrightarrow H^1(X, \tilde{\Lambda}_r) \longrightarrow H^1(X, \Gamma_a) \longrightarrow 0.$$

Fix  $n \in \mathbb{N}$ . Let  $\mathfrak{D}$  be a divisor on  $\mathcal{X}_g^{\mathrm{bc}}(n)$  such that  $\mathfrak{D}$  contains no fiber of the map  $\pi_g^{\mathrm{bc}}(n) : \mathcal{X}_g^{\mathrm{bc}}(n) \rightarrow \mathcal{A}_g(n)$ . Let  $\mathfrak{n}$  be the degree of the restriction of  $\mathfrak{D}$  to the fibers of  $\mathcal{X}_g^{\mathrm{bc}}(n)$ . Choose  $\mathfrak{D}$  such that  $n + \mathfrak{n} > 4g - 4$ .

Applying Lemma 2.6.4, let us write

$$\mathcal{U}_g(n, \mathfrak{D}) = (\mathcal{P}_g(n, \mathfrak{D}), \mathcal{X}_g(n, \mathfrak{D}), \pi_g(n, \mathfrak{D}), m_g(n, \mathfrak{D}), \mathcal{D}_g(n, \mathfrak{D}), \mathfrak{a}_g(n, \mathfrak{D}), \mathfrak{D}', \mathfrak{r}_g(n, \mathfrak{D}))$$

the universal family of restricted branched projective structures over the family of branching classes with auxiliary divisors

$$(\mathcal{A}_g(n), \mathcal{X}_g^{\mathrm{bc}}(n), \pi_g^{\mathrm{bc}}(n), m_g^{\mathrm{bc}}(n), \mathcal{D}_g^{\mathrm{bc}}(n), \mathfrak{a}_g^{\mathrm{bc}}(n), \mathfrak{D}).$$

Let  $\bar{\alpha}_g(n, \mathfrak{D}) : \mathcal{P}_g(n, \mathfrak{D}) \rightarrow \mathcal{A}_g(n)$  be the forgetful morphism. It is an affine bundle by Section 2.6.

For any partition  $\kappa$  of the integer  $n$ , denote by  $\mathcal{P}_g(\kappa, \mathfrak{D})$  the restriction of the bundle  $\bar{\alpha}_g(n, \mathfrak{D}) : \mathcal{P}_g(n, \mathfrak{D}) \rightarrow \mathcal{A}_g(n)$  to the basis  $\mathcal{A}_g(\kappa) \subset \mathcal{A}_g(n)$ . Let

$$\mathcal{U}_g(\kappa, \mathfrak{D}) = (\mathcal{P}_g(\kappa, \mathfrak{D}), \mathcal{X}_g(\kappa, \mathfrak{D}), \mathcal{D}_g(\kappa, \mathfrak{D}), \pi_g(\kappa, \mathfrak{D}), \mathfrak{a}_g(\kappa, \mathfrak{D}), \mathfrak{D}'(\kappa), \mathfrak{r}_g(\kappa, \mathfrak{D}))$$

be the pullback by the inclusion  $\mathcal{P}_g(\kappa, \mathfrak{D}) \rightarrow \mathcal{P}_g(n, \mathfrak{D})$  of the family  $\mathcal{U}_g(n, \mathfrak{D})$ . Denote by  $\bar{\alpha}_g(\kappa, \mathfrak{D})$  the restriction of  $\bar{\alpha}_g(n, \mathfrak{D})$  to  $\mathcal{P}_g(\kappa, \mathfrak{D})$ .

Let  $r \in \mathcal{P}_g(n, \mathfrak{D})$ . Let  $(X, D, a, m) \in \mathcal{A}_g(n)$  be the marked curve with divisor of degree  $n$  and branching class, image of  $r$  by the map  $\bar{\alpha}_g(n, \mathfrak{D})$ . Let  $D'$  be the divisor of degree  $\mathfrak{n}$  on  $X$ , restriction of  $\mathfrak{D}$  to the fiber of  $\mathcal{X}_g^{\text{bc}}(n)$  over  $(X, D, a, m)$ . Denote by  $\kappa$  the partition of  $n$  associated to the divisor  $D$ . The tangent space of the fiber  $P_{X, D'}^a = \bar{\alpha}_g(\kappa, \mathfrak{D})^{-1}(X, D, a, m)$  at  $r$  is  $(T_{P_{X, D'}^a}^a)_r = H^0(X, K_X^{\otimes 2}(-D)|_{D'})$ , according to Proposition 2.5.2.

Moreover we saw that there is a Kodaira-Spencer isomorphism

$$\text{KS}^{\text{bc}} : (T_{\mathcal{A}_g(\kappa)})_{(X, D, a, m)} \xrightarrow{\sim} H^1(X, \Gamma_a).$$

As a consequence, one has identifications between the kernels and cokernels of the exact sequences (20) and

$$0 \longrightarrow (T_{P_{X, D'}^a}^a)_r \longrightarrow (T_{\mathcal{P}_g(\kappa, \mathfrak{D})})_r \longrightarrow (T_{\mathcal{A}_g(\kappa)})_{(X, D, a, m)} \longrightarrow 0.$$

Let  $\mathcal{G}_{\mathfrak{D}}$  (respectively  $\mathcal{G}_{\mathfrak{D}}^{\text{rel}}$ ) be the subsheaf of  $\mathcal{T}_{\mathcal{X}_g(\kappa, \mathfrak{D})}$  (respectively  $\mathcal{T}_{\mathcal{X}_g(\kappa, \mathfrak{D})/\mathcal{P}_g(\kappa, \mathfrak{D})}$ ) that contains the vector fields preserving the restricted relative projective structure  $\mathfrak{r}_g(n, \mathfrak{D})$ . The Kodaira-Spencer exact sequence for the family  $\mathcal{X}_g(\kappa, \mathfrak{D})$ , when restricted to the vector fields preserving  $\mathfrak{r}_g(\kappa, \mathfrak{D})$ , is given by:

$$0 \longrightarrow \mathcal{G}_{\mathfrak{D}}^{\text{rel}} \longrightarrow \mathcal{G}_{\mathfrak{D}} \longrightarrow \pi_g(\kappa, \mathfrak{D})^* T_{\mathcal{P}_g(\kappa, \mathfrak{D})} \longrightarrow 0.$$

Applying the left exact functor  $\pi_g(\kappa, \mathfrak{D})_*$ , one gets a morphism of sheaves  $T_{\mathcal{P}_g(\kappa, \mathfrak{D})} \rightarrow R^1 \pi_g(\kappa, \mathfrak{D})_* \mathcal{G}_{\mathfrak{D}}^{\text{rel}}$ . In particular, at the point  $r$ , this morphism gives a linear map

$$(21) \quad \text{KS}_{\mathfrak{D}} : (T_{\mathcal{P}_g(\kappa, \mathfrak{D})})_r \longrightarrow H^1(X, \tilde{\Lambda}_r).$$

It is easy to see that the following diagram is commutative:

$$\begin{array}{ccccccc} 0 & \longrightarrow & (T_{P_{X, D'}^a}^a)_r & \longrightarrow & (T_{\mathcal{P}_g(\kappa, \mathfrak{D})})_r & \longrightarrow & (T_{\mathcal{A}_g(\kappa)})_{(X, D, a, m)} \longrightarrow 0 \\ & & \parallel & & \downarrow \text{KS}_{\mathfrak{D}} & & \downarrow \text{KS}^{\text{bc}} \\ 0 & \longrightarrow & H^0(X, K_X^{\otimes 2}(-D)|_{D'}) & \longrightarrow & H^1(X, \tilde{\Lambda}_r) & \longrightarrow & H^1(X, \Gamma_a) \longrightarrow 0 \end{array}$$

Since  $\text{KS}^{\text{bc}}$  is an isomorphism, we have the following lemma:

LEMMA 3.4.3. — *The map  $\text{KS}_{\mathfrak{D}}$  in (21) is an isomorphism.*

Since  $n + \mathfrak{n} > 4g - 4$ , one has a vector bundle on  $\mathcal{P}_g(n, \mathfrak{D})$  whose fiber over a point  $r \in P_{X, D'}^a$  is  $H^1(X, K_X^{\otimes 2}(-D - D'))$ . Denote by  $\mathfrak{c}_g(n, \mathfrak{D}) = \bar{\mathfrak{c}}(\mathfrak{a}_g(n, \mathfrak{D}), \mathfrak{r}_g(n, \mathfrak{D}))$  its global section defined in Section 2.6. According to Section 2.6, the space  $\mathcal{P}_g(n)$  is

an analytic subspace of  $\mathcal{P}_g(n, \mathfrak{D})$ , namely the vanishing locus of the section  $\mathfrak{c}_g(n, \mathfrak{D})$ . In particular, for any  $p \in \mathcal{P}_g(n)$ , the differential  $(d\mathfrak{c}_g(n, \mathfrak{D}))_p$  is a linear map from  $(T_{\mathcal{P}_g(n, \mathfrak{D})})_p$  to  $H^1(X, K_X^{\otimes 2}(-D - D'))$ , where  $X$  is the curve underlying  $p$ ,  $D$  is the branching divisor of  $p$  and  $D'$  is the auxiliary divisor. We will show that this differential is surjective for any  $p \in \mathcal{P}_g(n)$  such that  $H^2(X, \Lambda_p) = 0$ , which implies smoothness of  $\mathcal{P}_g(n)$  at  $p$ .

For any partition  $\kappa$  of the integer  $n$ ,  $\mathcal{P}_g(\kappa)$  is the intersection of the subspace  $\mathcal{P}_g(n) \subset \mathcal{P}_g(n, \mathfrak{D})$  with  $\mathcal{P}_g(\kappa, \mathfrak{D}) \subset \mathcal{P}_g(n, \mathfrak{D})$ . Recall that the family  $\mathcal{U}_g^{\text{bps}}(\kappa)$  is the pullback of the family  $\mathcal{U}_g^{\text{bps}}(n)$  by the inclusion  $\mathcal{P}_g(\kappa) \rightarrow \mathcal{P}_g(n)$ .

Let  $(X, D, a, p, m) \in \mathcal{P}_g(n)$  be a marked curve with divisor of degree  $n$ , branching class and branched projective structure. Let  $\kappa$  be the partition of  $n$  associated to the divisor  $D$ . Denote by  $r$  the branched projective structure  $p$ , seen as an element of  $\mathcal{P}_g(\kappa, \mathfrak{D})$ .

Denote by  $\mathcal{G}$  (respectively  $\mathcal{G}^{\text{rel}}$ ) the subsheaf of  $T_{\mathcal{X}_g^{\text{bps}}(\kappa)}$  (respectively  $T_{\mathcal{X}_g^{\text{bps}}(\kappa)/\mathcal{P}_g(\kappa)}$ ) containing vector fields preserving the relative branched projective structure  $\mathfrak{p}_g(\kappa)$ . The Kodaira-Spencer exact sequence for the family  $\mathcal{X}_g(\kappa, \mathfrak{D})$ , when restricted to vector fields preserving  $\mathfrak{p}_g(\kappa)$ , is given by:

$$0 \longrightarrow \mathcal{G}^{\text{rel}} \longrightarrow \mathcal{G} \longrightarrow \pi_g^{\text{bps}}(\kappa)^* T_{\mathcal{P}_g(\kappa)} \longrightarrow 0.$$

By applying the left exact functor  $\pi_g^{\text{bps}}(\kappa)_*$ , one gets a morphism of sheaves  $T_{\mathcal{P}_g(\kappa)} \rightarrow R^1 \pi_g^{\text{bps}}(\kappa)_* \mathcal{G}^{\text{rel}}$ . At the point  $(X, D, a, p, m)$ , one gets a linear map

$$(22) \quad \text{KS}^{\text{bps}} : (T_{\mathcal{P}_g(\kappa)})_{(X, D, a, p, m)} \longrightarrow H^1(X, \Lambda_p).$$

It is easy to see that the following diagram is commutative, where  $D'$  is the auxiliary divisor for the branching class  $a$  and  $r$  is  $p$  restricted to  $D'$ :

$$\begin{array}{ccccccc} 0 \longrightarrow & (T_{\mathcal{P}_g(\kappa)})_{(X, D, a, p, m)} & \text{KS}^{\text{bps}} \longrightarrow & (T_{\mathcal{P}_g(\kappa, \mathfrak{D})})_r & \xrightarrow{(d\mathfrak{c}_g(\kappa, \mathfrak{D}))_r} & H^1(X, K_X^{\otimes 2}(-D - D')) & \\ & \downarrow & & \downarrow \text{KS}_{\mathfrak{D}} & & \parallel & \\ 0 \longrightarrow & H^1(X, \Lambda_p) & \longrightarrow & H^1(X, \tilde{\Lambda}_r) & \longrightarrow & H^1(X, K_X^{\otimes 2}(-D - D')) & \\ & & & & & & \longrightarrow H^2(X, \Lambda_p). \end{array}$$

Since  $\text{KS}_{\mathfrak{D}}$  is an isomorphism, one has:

LEMMA 3.4.4. — *The map  $\text{KS}^{\text{bps}}$  in (22) is an isomorphism. Moreover the differential  $d\mathfrak{c}_g(\kappa, \mathfrak{D})$  is surjective at any point  $p$  of  $\mathcal{P}_g(\kappa)$  such that  $H^2(X, \Lambda_p) = 0$ .*

Take  $p \in \mathcal{P}_g(\kappa)$  such that  $H^2(X, \Lambda_p) = 0$ . Since the restricted differential  $d\mathfrak{c}_g(\kappa, \mathfrak{D})$  is surjective at  $p$ , the total differential  $d\mathfrak{c}_g(n, \mathfrak{D})$  is a fortiori surjective at  $p$ . This implies that  $p$  is a smooth point of  $\mathcal{P}_g(n) \subset \mathcal{P}_g(n, \mathfrak{D})$ . Conversely, if  $H^2(X, \Lambda_p) \neq 0$  and  $D$  is reduced, then  $h^1(X, \Lambda_p) = 6g - 6 + n + h^2(X, \Lambda_p) > 6g - 6 + n$ , so  $p$  is singular.

Finally, let  $p \in \mathcal{P}_g(n)$  be any branched projective structure such that  $h^2(X, \Lambda_p) > 0$ . It is shown in [CDF14] that  $p$  is the limit in  $\mathcal{P}_g(n)$  of a sequence  $(p_n)_{n \in \mathbb{N}}$  of branched projective structures with the same holonomy as  $p$  and reduced branching divisor. By Section 3.1, the dimension of  $H^2(X, \Lambda_p)$  depends only on the holonomy of  $\rho$ , thus the points  $p_n$  are all singular points of  $\mathcal{P}_g(n)$ , and since the singular locus is closed, the point  $p$  is itself a singular point.

**3.5. THE HOLONOMY MAP.** — Let  $S$  be a surface of genus  $g$  (with  $g \geq 2$ ). Denote by  $\text{Hom}^*(\pi_1(S), \text{PSL}(2, \mathbb{C}))$  the space of group morphisms  $\rho : \pi_1(S) \rightarrow \text{PSL}(2, \mathbb{C})$  such that  $\text{Im } \rho \subset \text{PSL}(2, \mathbb{C})$  is not abelian and is not conjugated to a subgroup of the group  $\{z \mapsto az \mid a \in \mathbb{C}^*\} \cup \{z \mapsto a/z \mid a \in \mathbb{C}^*\}$ .<sup>(1)</sup> In particular,  $\text{Hom}^*(\pi_1(S), \text{PSL}(2, \mathbb{C}))$  contains all non-elementary representations. The space  $\text{Hom}^*(\pi_1(S), \text{PSL}(2, \mathbb{C}))$  is an open subspace of  $\text{Hom}(\pi_1(S), \text{PSL}(2, \mathbb{C}))$ . The space  $\text{Hom}^*(\pi_1(S), \text{PSL}(2, \mathbb{C}))$  is a smooth analytic space (see [Hub81]) and the group  $\text{PSL}(2, \mathbb{C})$  acts freely and properly by conjugation on  $\text{Hom}^*(\pi_1(S), \text{PSL}(2, \mathbb{C}))$ . As a consequence, the quotient  $\text{Hom}^*(\pi_1(S), \text{PSL}(2, \mathbb{C}))/\text{PSL}(2, \mathbb{C})$  is a smooth analytic space.

Denote by  $\mathcal{P}_g(n)^*$  the subset of  $\mathcal{P}_g(n)$  of branched projective structures whose holonomy lies in  $\text{Hom}^*(\pi_1(S), \text{PSL}(2, \mathbb{C}))$ . There is a map

$$\text{Hol}_g(n)^* : \mathcal{P}_g(n)^* \longrightarrow \text{Hom}^*(\pi_1(S), \text{PSL}(2, \mathbb{C}))/\text{PSL}(2, \mathbb{C})$$

which associates to a branched projective structure the conjugacy class of its holonomy representations. The space  $\mathcal{P}_g(n)^*$  is clearly an open subset of  $\mathcal{P}_g(n)$ , and it is smooth by Theorem 3.0.1. Since  $\mathcal{P}_g(n)^*$  is the basis of a family of branched projective structures, the map  $\text{Hol}_g(n)^*$  is holomorphic.

**THEOREM 3.5.1.** — *The holonomy map*

$$\text{Hol}_g(n)^* : \mathcal{P}_g(n)^* \longrightarrow \text{Hom}^*(\pi_1(S), \text{PSL}(2, \mathbb{C}))/\text{PSL}(2, \mathbb{C})$$

*is a holomorphic submersion.*

Let  $\mathcal{X}_g$  be the  $\text{PSL}(2, \mathbb{C})$ -character variety associated to  $S$ , i.e., the GIT quotient  $\text{Hom}(\pi_1(S), \text{PSL}(2, \mathbb{C}))/\text{PSL}(2, \mathbb{C})$ , that is an analytic space. There is an analytic morphism  $\text{Hol}_g(n) : \mathcal{P}_g(n) \rightarrow \mathcal{X}_g$ , that to a branched projective structure associates its holonomy. There is an obvious map

$$\iota : \text{Hom}^*(\pi_1(S), \text{PSL}(2, \mathbb{C}))/\text{PSL}(2, \mathbb{C}) \longrightarrow \mathcal{X}_g,$$

that verifies in particular  $\text{Hol}_g(n) = \iota \circ \text{Hol}_g(n)^*$ . In restriction to the preimage of the smooth locus of  $\mathcal{X}_g$ , the map  $\iota$  is a biholomorphism. As a consequence, Theorem 3.5.1 implies:

**COROLLARY 3.5.2.** — *In restriction to the preimage of the smooth locus of the character variety  $\mathcal{X}_g$ , the holonomy map  $\text{Hol}_g(n) : \mathcal{P}_g(n) \rightarrow \mathcal{X}_g$  is a holomorphic submersion.*

---

<sup>(1)</sup>This second condition is necessary for the action by conjugation of  $\text{PSL}(2, \mathbb{C})$  to be free.

*Proof of Theorem 3.5.1.* — Take  $p \in \mathcal{P}_g(n)^*$ , with underlying Riemann surface  $X$ . Let  $\rho \in \text{Hom}^*(\pi_1(S), \text{PSL}(2, \mathbb{C}))$  be a holonomy representation of  $p$ , and write  $[\rho] = \text{Hol}_g(n)^*(p)$ . It is proved in [Hub81] that the tangent space of

$$\text{Hom}^*(\pi_1(S), \text{PSL}(2, \mathbb{C})) / \text{PSL}(2, \mathbb{C})$$

at  $[\rho]$  is  $H^1(X, \text{Ad}_p)$ , where  $\text{Ad}_p$  is defined in Section 3.1.

Hubbard also proved in [Hub81] that in the case  $n = 0$ , the differential at  $p$  of the holonomy map, i.e.,  $d_p \text{Hol}_g(0)^* : T_p \mathcal{P}_g(0) \rightarrow H^1(X, \text{Ad}_p)$ , is given by the Kodaira-Spencer at  $p$  of the universal family  $\mathcal{U}_g^{\text{bps}}(0)$ . The situation is almost the same in the branched case ( $n > 1$ ), with two differences. Firstly the Kodaira-Spencer map  $\text{KS}^{\text{bps}}$  (see Section 3.4) is only defined on  $T_p \mathcal{P}_g(\kappa)$ , where  $\kappa$  is the partition of  $n$  associated to the branching divisor of  $p$ . Secondly  $\text{KS}^{\text{bps}}$  takes values in the first cohomology group of the subsheaf  $\Lambda_p$  of  $\text{Ad}_p$  (see Section 3.1). However there is a canonical map  $\phi : H^1(X, \Lambda_p) \rightarrow H^1(X, \text{Ad}_p)$ , coming from the inclusion  $\Lambda_p \subset \text{Ad}_p$ .

Thus the proof of Hubbard gives in the branched case:

$$d_p \text{Hol}_g(n)^*|_{T_p \mathcal{P}_g(\kappa)} = \phi \circ \text{KS}^{\text{bps}}.$$

Since  $\phi$  is surjective according to (9) and  $\text{KS}^{\text{bps}}$  is an isomorphism,  $d_p \text{Hol}_g(n)^*|_{T_p \mathcal{P}_g(\kappa)}$  is surjective, thus  $d_p \text{Hol}_g(n)^*$  is a fortiori surjective, which proves that  $\text{Hol}_g(n)^*$  is a submersion.  $\square$

## REFERENCES

- [AB20] D. G. L. ALLEGRETTI & T. BRIDGELAND — “The monodromy of meromorphic projective structures”, *Trans. Amer. Math. Soc.* **373** (2020), no. 9, p. 6321–6367.
- [BM14] D. BARLET & J. MAGNÚSSON — *Cycles analytiques complexes. I. Théorèmes de préparation des cycles*, Cours Spécialisés, vol. 22, Société Mathématique de France, Paris, 2014.
- [BD05] A. BEILINSON & V. DRINFELD — “Opers”, 2005, [arXiv:math/0501398](https://arxiv.org/abs/math/0501398).
- [BDG19] I. BISWAS, S. DUMITRESCU & S. GUPTA — “Branched projective structures on a Riemann surface and logarithmic connections”, *Doc. Math.* **24** (2019), p. 2299–2337.
- [BDH22] I. BISWAS, S. DUMITRESCU & S. HELLER — “Branched  $\text{SL}(r, \mathbb{C})$ -opers”, *Internat. Math. Res. Notices* **10** (2022), p. 8311–8355.
- [Blu79] R. A. BLUMENTHAL — “Transversely homogeneous foliations”, *Ann. Inst. Fourier (Grenoble)* **29** (1979), no. 4, p. 143–158.
- [CDF14] G. CALSAMIGLIA, B. DEROIN & S. FRANCAVIGLIA — “Branched projective structures with Fuchsian holonomy”, *Geom. Topol.* **18** (2014), no. 1, p. 379–446.
- [DG23] B. DEROIN & A. GUILLOT — “Foliated affine and projective structures”, *Compositio Math.* **159** (2023), no. 6, p. 1153–1187.
- [Dum09] D. DUMAS — “Complex projective structures”, in *Handbook of Teichmüller theory. Vol. II*, IRMA Lect. Math. Theor. Phys., vol. 13, European Mathematical Society, Zürich, 2009, p. 455–508.
- [EE69] C. J. EARLE & J. EELLS — “A fibre bundle description of Teichmüller theory”, *J. Differential Geometry* **3** (1969), p. 19–43.
- [FR21] S. FRANCAVIGLIA & L. RUFFONI — “Local deformations of branched projective structures: Schiffer variations and the Teichmüller map”, *Geom. Dedicata* **214** (2021), p. 21–48.
- [Fre07] E. FRENKEL — “Lectures on the Langlands program and conformal field theory”, in *Frontiers in number theory, physics, and geometry. II*, Springer, Berlin, 2007, p. 387–533.
- [FG10] E. FRENKEL & D. GAITSGORY — “Weyl modules and opers without monodromy”, in *Arithmetic and geometry around quantization*, Progress in Math., vol. 279, Birkhäuser Boston, Boston, MA, 2010, p. 101–121.

- [GKM00] D. GALLO, M. KAPOVICH & A. MARDEN – “The monodromy groups of Schwarzian equations on closed Riemann surfaces”, *Ann. of Math. (2)* **151** (2000), no. 2, p. 625–704.
- [God91] C. GODBILLON – *Feuilletages*, Progress in Math., vol. 98, Birkhäuser Verlag, Basel, 1991.
- [Gun66] R. C. GUNNING – *Lectures on Riemann surfaces*, Princeton Math. Notes, Princeton University Press, Princeton, NJ, 1966.
- [GM20] S. GUPTA & M. MJ – “Monodromy representations of meromorphic projective structures”, *Proc. Amer. Math. Soc.* **148** (2020), no. 5, p. 2069–2078.
- [GM21] ———, “Meromorphic projective structures, grafting and the monodromy map”, *Adv. Math.* **383** (2021), article no. 107673 (49 pages).
- [Hej75] D. A. HEJHAL – “Monodromy groups and linearly polymorphic functions”, *Acta Math.* **135** (1975), no. 1, p. 1–55.
- [Hub81] J. H. HUBBARD – “The monodromy of projective structures”, in *Riemann surfaces and related topics (Stony Brook, NY, 1978)*, Annals of Math. Studies, vol. 97, Princeton University Press, Princeton, NJ, 1981, p. 257–275.
- [LF23] T. LE FILS – “Holonomy of complex projective structures on surfaces with prescribed branch data”, *J. Topology* **16** (2023), no. 1, p. 430–487.
- [LMP09] F. LORAY & D. MARÍN PÉREZ – “Projective structures and projective bundles over compact Riemann surfaces”, in *Équations différentielles et singularités*, Astérisque, vol. 323, Société Mathématique de France, Paris, 2009, p. 223–252.
- [Man72] R. MANDELBAUM – “Branched structures on Riemann surfaces”, *Trans. Amer. Math. Soc.* **163** (1972), p. 261–275.
- [Man73] ———, “Branched structures and affine and projective bundles on Riemann surfaces”, *Trans. Amer. Math. Soc.* **183** (1973), p. 37–58.
- [Man75] ———, “Unstable bundles and branched structures on Riemann surfaces”, *Math. Ann.* **214** (1975), p. 49–59.
- [dSG16] H. P. DE SAINT-GERVAIS – *Uniformization of Riemann surfaces*, Heritage of European Math., European Mathematical Society, Zürich, 2016.
- [Scá97] B. A. SCÁRDUA – “Transversely affine and transversely projective holomorphic foliations”, *Ann. Sci. École Norm. Sup. (4)* **30** (1997), no. 2, p. 169–204.
- [Sér22] T. SÉRANDOUR – “Meromorphic projective structures, opers and monodromy”, PhD Thesis, Université de Rennes, 2022.
- [Ste43] N. E. STEENROD – “Homology with local coefficients”, *Ann. of Math. (2)* **44** (1943), p. 610–627.
- [Sun17] F. SUN – “An elementary proof for Poincaré duality with local coefficients”, 2017, [arXiv: 1709.00569](https://arxiv.org/abs/1709.00569).
- [Voi07] C. VOISIN – *Hodge theory and complex algebraic geometry. I*, Cambridge Studies in Advanced Math., vol. 76, Cambridge University Press, Cambridge, 2007.
- [Whi78] G. W. WHITEHEAD – *Elements of homotopy theory*, Graduate Texts in Math., vol. 61, Springer-Verlag, New York-Berlin, 1978.

Manuscript received 31st October 2023

accepted 17th October 2024

GUSTAVE BILLON, IRMA, CNRS, Université de Strasbourg,  
 7 rue René Descartes, 67084 Strasbourg Cedex, France  
*E-mail* : [g.billon@unistra.fr](mailto:g.billon@unistra.fr)  
*Url* : <https://irma.math.unistra.fr/~billon/>