

Journal de l'École polytechnique

Mathématiques

Hoel QUEFFELEC, Alistair SAVAGE, & Oded YACOBI

An equivalence between truncations of categorified quantum groups and Heisenberg categories

Tome 5 (2018), p. 197-238.

http://jep.cedram.org/item?id=JEP_2018__5__197_0

© Les auteurs, 2018.

Certains droits réservés.



Cet article est mis à disposition selon les termes de la licence
CREATIVE COMMONS ATTRIBUTION – PAS DE MODIFICATION 3.0 FRANCE.
<http://creativecommons.org/licenses/by-nd/3.0/fr/>

L'accès aux articles de la revue « Journal de l'École polytechnique — Mathématiques » (<http://jep.cedram.org/>), implique l'accord avec les conditions générales d'utilisation (<http://jep.cedram.org/legal/>).

Publié avec le soutien
du Centre National de la Recherche Scientifique

cedram

Article mis en ligne dans le cadre du
Centre de diffusion des revues académiques de mathématiques
<http://www.cedram.org/>

AN EQUIVALENCE BETWEEN TRUNCATIONS OF
CATEGORIFIED QUANTUM GROUPS AND
HEISENBERG CATEGORIES

BY HOEL QUEFFELEC, ALISTAIR SAVAGE & ODED YACOBI

ABSTRACT. — We introduce a simple diagrammatic 2-category \mathcal{A} that categorifies the image of the Fock space representation of the Heisenberg algebra and the basic representation of \mathfrak{sl}_∞ . We show that \mathcal{A} is equivalent to a truncation of the Khovanov–Lauda categorified quantum group \mathcal{U} of type A_∞ , and also to a truncation of Khovanov’s Heisenberg 2-category \mathcal{H} . This equivalence is a categorification of the principal realization of the basic representation of \mathfrak{sl}_∞ . As a result of the categorical equivalences described above, certain actions of \mathcal{H} induce actions of \mathcal{U} , and vice versa. In particular, we obtain an explicit action of \mathcal{U} on representations of symmetric groups. We also explicitly compute the Grothendieck group of the truncation of \mathcal{H} . The 2-category \mathcal{A} can be viewed as a graphical calculus describing the functors of i -induction and i -restriction for symmetric groups, together with the natural transformations between their compositions. The resulting computational tool is used to give simple diagrammatic proofs of (apparently new) representation theoretic identities.

RÉSUMÉ (Une équivalence entre des troncations de groupes quantiques catégorifiés et des catégories de Heisenberg)

Nous introduisons une 2-catégorie élémentaire \mathcal{A} qui catégorifie l’image de l’espace de Fock comme représentation de l’algèbre de Heisenberg, ainsi que la représentation basique de \mathfrak{sl}_∞ . Nous montrons que \mathcal{A} est équivalente à une troncation du groupe quantique catégorifié de Khovanov–Lauda \mathcal{U} en type A_∞ , ainsi qu’à une troncation de la 2-catégorie de Heisenberg \mathcal{H} introduite par Khovanov. Cette équivalence se comprend comme une catégorification de la réalisation principale de la représentation basique de \mathfrak{sl}_∞ . Il résulte des équivalences catégoriques précédentes que certaines actions de \mathcal{H} induisent des actions de \mathcal{U} , et vice versa. En particulier, nous obtenons une action explicite de \mathcal{U} sur les représentations des groupes symétriques. Nous calculons également explicitement le groupe de Grothendieck de la troncation de \mathcal{H} . La 2-catégorie \mathcal{A} s’interprète comme un calcul graphique décrivant les foncteurs de i -induction et i -restriction pour les groupes symétriques, ainsi que les transformations naturelles entre leurs compositions. Nous utilisons l’outil de calcul qui en découle pour donner des preuves diagrammatiques simples d’identités (apparemment nouvelles) en théorie des représentations.

2010 MATHEMATICS SUBJECT CLASSIFICATION. — 17B10, 17B65, 20C30, 16D90.

KEYWORDS. — Categorification, Heisenberg algebra, Fock space, basic representation, principal realization, symmetric group.

H.Q. was supported by a Discovery Project from the Australian Research Council and a grant from the Université de Montpellier. A.S. was supported by a Discovery Grant from the Natural Sciences and Engineering Research Council of Canada. O.Y. was supported by a Discovery Early Career Research Award from the Australian Research Council.

CONTENTS

1. Introduction.....	198
2. Algebraic preliminaries.....	201
3. Modules for symmetric groups.....	205
4. The 2-category \mathcal{A}	210
5. The 2-category \mathcal{H}^{tr}	216
6. Equivalence of \mathcal{H}_ϵ and \mathcal{A}	222
7. Actions on modules for symmetric groups.....	229
8. Applications and further directions.....	234
References.....	237

1. INTRODUCTION

Affine Lie algebras play a key role in many areas of representation theory and mathematical physics. One of their prominent features is that their highest-weight irreducible representations have explicit realizations. In particular, constructions of the so-called basic representation involve deep mathematics from areas as diverse as algebraic combinatorics (symmetric functions), number theory (modular forms), and geometry (Hilbert schemes).

Two of the most well-studied realizations of the basic representation are the *homogeneous* and *principal* realizations (see, for example [Kac90, Ch. 14]). The homogeneous realization in affine types ADE has been categorified in [CL11]. In the current paper we focus our attention on the principal realization in type A_∞ . The infinite-dimensional Lie algebra \mathfrak{sl}_∞ behaves in many ways like an affine Lie algebra, and in particular, it has a basic representation with a principal realization coming from a close connection to the infinite-rank Heisenberg algebra H .

The Heisenberg algebra H has a natural representation on the space Sym of symmetric functions (with rational coefficients), called the Fock space representation. The universal enveloping algebra $U = U(\mathfrak{sl}_\infty)$ also acts naturally on Sym , yielding the basic representation. So we have algebra homomorphisms

$$(1.1) \quad H \xrightarrow{r_H} \text{End}_{\mathbb{Q}} \text{Sym} \xleftarrow{r_U} U.$$

Consider the vector space decomposition

$$\text{Sym} = \bigoplus_{\lambda \in \mathcal{P}} \mathbb{Q}s_\lambda,$$

where the sum is over all partitions \mathcal{P} and s_λ denotes the Schur function corresponding to λ . Let $1_\lambda: \text{Sym} \rightarrow \mathbb{Q}s_\lambda$ denote the natural projection. While the images of the representations r_H and r_U are not equal, we have an equality of their idempotent modifications:

$$(1.2) \quad \bigoplus_{\lambda, \mu \in \mathcal{P}} 1_\mu r_H(H) 1_\lambda = \bigoplus_{\lambda, \mu \in \mathcal{P}} 1_\mu r_U(U) 1_\lambda.$$

This observation is an \mathfrak{sl}_∞ analogue of the fact that the basic representation of $\widehat{\mathfrak{sl}}_n$ remains irreducible when restricted to the principal Heisenberg subalgebra—a fact which is the crucial ingredient in the principal realization of the basic representation. We view (1.2) as an additive \mathbb{Q} -linear category \mathcal{A} whose set of objects is the free monoid $\mathbb{N}[\mathcal{P}]$ on \mathcal{P} and with

$$\mathrm{Mor}_{\mathcal{A}}(\lambda, \mu) = 1_\mu r_H(H)1_\lambda = 1_\mu r_U(U)1_\lambda = \mathrm{Hom}_{\mathbb{Q}}(\mathbb{Q}s_\lambda, \mathbb{Q}s_\mu).$$

In [Kho14], Khovanov introduced a monoidal category, defined in terms of planar diagrams, whose Grothendieck group contains (and is conjecturally isomorphic to) the Heisenberg algebra H . Khovanov’s category has a natural 2-category analogue \mathcal{H} . On the other hand, in [KL10], Khovanov and Lauda introduced a 2-category, which we denote \mathcal{U} , that categorifies quantum \mathfrak{sl}_n and can naturally be generalized to the \mathfrak{sl}_∞ case (see [CL15]). A related construction was also described by Rouquier in [Rou08]. These categorifications have led to an explosion of research activity, including generalizations, and applications to representation theory, geometry, and topology. It is thus natural to seek a connection between the 2-categories \mathcal{H} and \mathcal{U} that categorifies the principal embedding relationship between H and U discussed above. This is the goal of the current paper.

We define a 2-category \mathcal{A} whose 2-morphism spaces are given by planar diagrams modulo isotopy and local relations. The local relations of \mathcal{A} are exceedingly simple and we show that \mathcal{A} categorifies \mathcal{A} . We then describe precise relationships between \mathcal{A} and the 2-categories \mathcal{H} and \mathcal{U} . Our first main result is that \mathcal{A} is equivalent to a degree zero piece of a truncation of the categorified quantum group \mathcal{U} . More precisely, recalling that the objects of \mathcal{U} are elements of the weight lattice of \mathfrak{sl}_∞ , we consider the truncation $\mathcal{U}^{\mathrm{tr}}$ of \mathcal{U} where we kill weights not appearing in the basic representation. Specifically, we quotient the 2-morphism spaces by the identity 2-morphisms of the identity 1-morphisms of such weights. (This type of truncation has appeared before in the categorification literature, for example, in [MSV13, QR16].) The resulting 2-morphism spaces of $\mathcal{U}^{\mathrm{tr}}$ are nonnegatively graded, and we show that the degree zero part \mathcal{U}_0 of $\mathcal{U}^{\mathrm{tr}}$ is equivalent to the 2-category \mathcal{A} (Theorem 4.4).

Our next main result is that \mathcal{A} is also equivalent to a summand of an idempotent completion of a truncation of the Heisenberg 2-category \mathcal{H} . More precisely, recalling that the objects of \mathcal{H} are integers, we consider the truncation $\mathcal{H}^{\mathrm{tr}'}$ of \mathcal{H} obtained by killing objects corresponding to negative integers. We then take an idempotent completion $\mathcal{H}^{\mathrm{tr}}$ of $\mathcal{H}^{\mathrm{tr}'}$, show that we have a natural decomposition $\mathcal{H}^{\mathrm{tr}} \cong \mathcal{H}_\epsilon \oplus \mathcal{H}_\delta$, and that \mathcal{A} is equivalent to the summand \mathcal{H}_ϵ (Theorem 6.7). This summand can be obtained from $\mathcal{H}^{\mathrm{tr}}$ by imposing one extra local relation (namely, declaring a clockwise circle in a region labeled n to be equal to n). We note that the idempotent completion we consider in the above construction is larger than the one often appearing in the categorification literature since we complete with respect to both idempotent 1-morphisms and 2-morphisms (see Definition 5.1 and Remark 5.2). As a result, the idempotent completion has more objects, with the object n splitting into a direct sum of objects labeled by the partitions of n .

We thus have 2-functors

$$\mathcal{H} \xrightarrow{\text{truncate}} \mathcal{H}^{\text{tr}} \xrightarrow{\text{summand}} \mathcal{H}_\epsilon \cong \mathcal{A} \cong \mathcal{U}_0 \xleftarrow{\text{summand}} \mathcal{U}^{\text{tr}} \xleftarrow{\text{truncate}} \mathcal{U}.$$

that can be thought of as a categorification of (1.1). The equivalence $\mathcal{H}_\epsilon \cong \mathcal{U}_0$ is a categorification of the isomorphism (1.2) and yields a categorical analog of the principal realization of the basic representation of \mathfrak{sl}_∞ . In particular, any action of \mathcal{H} factoring through \mathcal{H}^{tr} (which is true of any action categorifying the Fock space representation) induces an explicit action of \mathcal{U} . Conversely, any action of \mathcal{U} factoring through \mathcal{U}^{tr} (which is true of any action categorifying the basic representation) induces an explicit action of \mathcal{H} . See Section 7.1.

In [Kho14], Khovanov described an action of his Heisenberg category on modules for symmetric groups. This naturally induces an action of the 2-category \mathcal{H} factoring through \mathcal{H}^{tr} . Applying the categorical principal realization to this action we obtain an explicit action of the Khovanov–Lauda categorified quantum group \mathcal{U} on modules for symmetric groups, relating our work to [BK09a, BK09b]. See Section 7.4.

By computations originally due to Chuang and Rouquier in [CR08, §7.1], one can easily deduce that there is a categorical action of \mathfrak{sl}_∞ on modules for symmetric groups. This action is constructed using i -induction and i -restriction functors, and thus is closely related to Khovanov’s categorical Heisenberg action. The equivalence $\mathcal{H}_\epsilon \cong \mathcal{U}_0$ gives the precise diagrammatic connection between these actions on the level of 2-categories. In particular, the 2-category \mathcal{A} yields a graphical calculus for describing i -induction and i -restriction functors, together with the natural transformations between them (see Proposition 7.3). This provides a computational tool for proving identities about the representation theory of the symmetric groups. See Section 8.1 for some examples of identities that, to the best of our knowledge, are new.

One of the most important open questions about Khovanov’s Heisenberg category is the conjecture that it categorifies the Heisenberg algebra (see [Kho14, Conj. 1]). In the framework of 2-categories, this conjecture is the statement that the Grothendieck group of \mathcal{H} is isomorphic to $\bigoplus_{m \in \mathbb{Z}} H$. (The presence of the infinite sum here arises from the fact that, in a certain sense, the 2-category \mathcal{H} contains countably many copies of the monoidal Heisenberg category defined in [Kho14].) We prove the analog of Khovanov’s conjecture for the truncated category \mathcal{H}^{tr} , namely that the Grothendieck group of \mathcal{H}^{tr} is isomorphic to $\bigoplus_{m \in \mathbb{N}} \mathcal{A}$. See Corollary 6.8.

We now give an overview of the contents of the paper. In Section 2 we recall some basic facts about the basic representation and define the category \mathcal{A} . We also set some category theoretic notation and conventions. In Section 3 we recall some facts about modules for symmetric groups, discuss eigenspace decompositions with respect to Jucys–Murphy elements, and prove some combinatorial identities that will be used elsewhere in the paper. Then, in Section 4, we introduce the 2-category \mathcal{A} and show that it is equivalent to \mathcal{U}_0 . We also prove some results about the structure of \mathcal{A} and prove that it categorifies \mathcal{A} . We turn our attention to the Heisenberg 2-category in Section 5. In particular, we introduce the truncated Heisenberg 2-category \mathcal{H}^{tr} ,

describe the decomposition $\mathcal{H}^{\text{tr}} \cong \mathcal{H}_\epsilon \oplus \mathcal{H}_\delta$, and prove that \mathcal{H}_ϵ is equivalent to \mathcal{A} . In Section 7 we discuss how our results yield categorical Heisenberg actions from categorified quantum group actions and vice versa. In particular, we describe an explicit action of the Khovanov–Lauda 2-category on modules for symmetric groups. Finally, in Section 8 we give an application of our results to diagrammatic computation and discuss some possible directions for further research.

Note on the arXiv version. — For the interested reader, the tex file of the arXiv version of this paper includes hidden details of some straightforward computations and arguments that are omitted in the present article. These details can be displayed by switching the `details` toggle to true in the tex file and recompiling.

Acknowledgements. — The authors would like to thank C. Bonnafé, M. Khovanov, A. Lauda, A. Licata, A. Molev, E. Wagner, and M. Zabrocki for helpful conversations.

2. ALGEBRAIC PRELIMINARIES

2.1. **BOSONIC FOCK SPACE AND THE CATEGORY \mathcal{A} .** — Let \mathcal{P} denote the set of partitions and write $\lambda \vdash n$ to denote that $\lambda = (\lambda_1, \lambda_2, \dots)$, $\lambda_1 \geq \lambda_2 \geq \dots$, is a partition of $n \in \mathbb{N}$. Let Sym be the algebra of symmetric functions with rational coefficients. Then we have

$$\text{Sym} = \bigoplus_{\lambda \in \mathcal{P}} \mathbb{Q}s_\lambda,$$

where s_λ denotes the Schur function corresponding to the partition λ . For $\lambda \in \mathcal{P}$, we let $1_\lambda: \text{Sym} \rightarrow \mathbb{Q}s_\lambda$ denote the corresponding projection.

Let $\mathbb{N}[\mathcal{P}]$ be the free monoid on the set of partitions. Define \mathcal{A} to be the additive \mathbb{Q} -linear category whose set of objects is $\mathbb{N}[\mathcal{P}]$, where we denote the zero object by $\mathbf{0}$. The morphisms between generating objects are

$$\text{Mor}_{\mathcal{A}}(\lambda, \mu) = 1_\mu(\text{End}_{\mathbb{Q}} \text{Sym})1_\lambda = \text{Hom}_{\mathbb{Q}}(\mathbb{Q}s_\lambda, \mathbb{Q}s_\mu), \quad \lambda, \mu \in \mathcal{P}.$$

If \mathcal{V} denotes the category of finite-dimensional \mathbb{Q} -vector spaces, then we have an equivalence of categories

$$(2.1) \quad \mathbf{r}: \mathcal{A} \longrightarrow \mathcal{V}, \quad \lambda \longmapsto \mathbb{Q}s_\lambda.$$

Let $\langle \cdot, \cdot \rangle$ be the inner product on Sym under which the Schur functions are orthonormal. For $f \in \text{Sym}$, let f^* denote the operator on Sym adjoint to multiplication by f :

$$\langle f^*(g), h \rangle = \langle g, fh \rangle \quad \text{for all } f, g, h \in \text{Sym}.$$

The *Heisenberg algebra* H is the subalgebra of $\text{End}_{\mathbb{Q}} \text{Sym}$ generated by the operators f and f^* , $f \in \text{Sym}$. The tautological action of H on Sym is called the (*bosonic*) *Fock space representation*.

For $\lambda, \mu \in \mathcal{P}$, we have

$$1_\mu H 1_\lambda = \text{Hom}_{\mathbb{Q}}(\mathbb{Q}s_\lambda, \mathbb{Q}s_\mu) = \text{Mor}_{\mathcal{A}}(\lambda, \mu),$$

where the first equality follows from the fact that $s_\mu s_\lambda^* 1_\lambda$ is the map $s_\nu \mapsto \delta_{\lambda, \nu} s_\mu$. Thus, \mathcal{A} may be viewed as an idempotent modification of H .

2.2. THE BASIC REPRESENTATION. — Let \mathfrak{sl}_∞ denote the Lie algebra of all trace zero infinite matrices $a = (a_{ij})_{i,j \in \mathbb{Z}}$ with rational entries such that the number of nonzero a_{ij} is finite, with the usual commutator bracket. Set

$$e_i = E_{i,i+1}, \quad f_i = E_{i+1,i}, \quad h_i = [e_i, f_i] = E_{i,i} - E_{i+1,i+1},$$

where $E_{i,j}$ is the matrix whose (i,j) -entry is equal to one and all other entries are zero. Let $U = U(\mathfrak{sl}_\infty)$ denote the universal enveloping algebra of \mathfrak{sl}_∞ .

To a partition $\lambda = (\lambda_1, \dots, \lambda_n)$, we associate the Young diagram with rows numbered from top to bottom, columns numbered left to right, and which has λ_1 boxes in the first row, λ_2 boxes in the second row, etc. A box in row k and column ℓ has content $\ell - k \in \mathbb{Z}$. A Young diagram will be said to have an *addable* i -box if one can add to it a box of content i and get a Young diagram. Similarly, a Young diagram has a *removable* i -box if there is a box of content i that can be removed yielding another Young diagram. If $\lambda \vdash n$ has an addable i -box we let $\lambda \boxplus i$ be the partition of $n + 1$ obtained from λ by adding the box of content i , and similarly define $\lambda \boxminus i$.

EXAMPLE 2.1. — Let $\lambda = (3, 2) \vdash 5$. Then we have

$$\lambda = \begin{array}{|c|c|c|} \hline \square & \square & \square \\ \hline \square & \square & \\ \hline \end{array}, \quad \lambda \boxplus 3 = \begin{array}{|c|c|c|c|} \hline \square & \square & \square & \square \\ \hline \square & \square & \square & \\ \hline \end{array}, \quad \lambda \boxminus 0 = \begin{array}{|c|c|c|} \hline \square & \square & \square \\ \hline \square & \square & \\ \hline \end{array}.$$

For $\lambda \in \mathcal{P}$, define

$$B^+(\lambda) = \{i \mid \lambda \text{ has an addable } i\text{-box}\} \quad \text{and} \quad B^-(\lambda) = \{i \mid \lambda \text{ has a removable } i\text{-box}\}.$$

Note that, for all $\lambda \in \mathcal{P}$, we have $B^+(\lambda) \cap B^-(\lambda) = \emptyset$. If $i \notin B^+(\lambda)$ (respectively $i \notin B^-(\lambda)$), then we consider $\lambda \boxplus i = \mathbf{0}$ (respectively $\lambda \boxminus i = \mathbf{0}$) when viewing partitions as objects in \mathcal{A} .

Consider the action of U on Sym given by

$$(2.2) \quad e_i \cdot s_\lambda = s_{\lambda \boxplus i}, \quad f_i \cdot s_\lambda = s_{\lambda \boxminus i},$$

where, by convention, $s_{\mathbf{0}} = 0$. This defines an irreducible representation of U on Sym known as the *basic representation*. In fact, one can write explicit expressions for the action of the generators e_i and f_i in terms of the action of the Heisenberg algebra H on Sym . This construction is known as the *principal realization*. We refer the reader to [Kac90, §§14.9–14.10] for details. The element s_λ spans the weight space of weight

$$(2.3) \quad \omega_\lambda := \Lambda_0 - \sum_{i \in C(\lambda)} \alpha_i,$$

where the sum is over the multiset $C(\lambda)$ of contents of the boxes of λ , Λ_0 is the zeroth fundamental weight, and α_i is the i -th simple root. In particular, the map

$$(2.4) \quad \lambda \longmapsto \omega_\lambda$$

is a bijection between \mathcal{P} and the set of weights of the basic representation.

2.3. A KAC–MOODY PRESENTATION OF \mathcal{A} . — Let \widehat{U} denote the image of U in $\text{End}_{\mathbb{Q}} \text{Sym}$ under the basic representation described in Section 2.2. Then, for $\lambda, \mu \in \mathcal{P}$, we have

$$1_{\mu} \widehat{U} 1_{\lambda} = \text{Hom}_{\mathbb{Q}}(\mathbb{Q}s_{\lambda}, \mathbb{Q}s_{\mu}) = \text{Mor}_{\mathcal{A}}(\lambda, \mu).$$

This observation allows us to deduce a Kac–Moody-type presentation of \mathcal{A} . Define morphisms

$$\begin{aligned} e_i 1_{\lambda} &\in \text{Mor}_{\mathcal{A}}(\lambda, \lambda \boxplus i), & s_{\lambda} &\longmapsto s_{\lambda \boxplus i}, \\ f_i 1_{\lambda} &\in \text{Mor}_{\mathcal{A}}(\lambda, \lambda \boxminus i), & s_{\lambda} &\longmapsto s_{\lambda \boxminus i}, \end{aligned}$$

for $i \in \mathbb{Z}$, $\lambda \in \mathcal{P}$. Since Young’s lattice is connected, these morphisms clearly generate all morphisms in \mathcal{A} .

PROPOSITION 2.2. — *The morphisms in \mathcal{A} are generated by $e_i 1_{\lambda}$, $f_i 1_{\lambda}$, for $i \in \mathbb{Z}$, $\lambda \in \mathcal{P}$, subject only to the relations*

$$(2.5) \quad e_i e_j 1_{\lambda} = e_j e_i 1_{\lambda}, \quad \text{if } |i - j| > 1,$$

$$(2.6) \quad f_i f_j 1_{\lambda} = f_j f_i 1_{\lambda}, \quad \text{if } |i - j| > 1,$$

$$(2.7) \quad e_i f_j 1_{\lambda} = f_j e_i 1_{\lambda}, \quad \text{if } i \neq j,$$

$$(2.8) \quad e_i f_i 1_{\lambda} = 1_{\lambda}, \quad \text{if } i \in B^+(\lambda),$$

$$(2.9) \quad f_i e_i 1_{\lambda} = 1_{\lambda}, \quad \text{if } i \in B^-(\lambda).$$

Proof. — Let \mathcal{C} be the category with objects $\mathbb{N}[\mathcal{P}]$ and morphisms given by the presentation in the statement of the proposition. Since the relations (2.5)–(2.9) are immediate in \mathcal{A} , we have a full and essentially surjective functor $\mathcal{C} \rightarrow \mathcal{A}$. Therefore it suffices to show that $\dim \text{Mor}_{\mathcal{C}}(\lambda, \mu) \leq 1$ for all $\lambda, \mu \in \mathcal{P}$.

In fact, we will prove that, for $\lambda, \mu \in \mathcal{P}$, $\text{Mor}_{\mathcal{C}}(\lambda, \mu)$ is spanned by a single morphism of the form

$$(2.10) \quad f_{i_1} f_{i_2} \cdots f_{i_k} e_{j_1} e_{j_2} \cdots e_{j_\ell} 1_{\lambda},$$

where $\{i_1, \dots, i_k\} \cap \{j_1, \dots, j_\ell\} = \emptyset$ and $\mu = \lambda \boxplus j_\ell \cdots \boxplus j_2 \boxplus j_1 \boxplus i_k \boxplus \cdots \boxplus i_2 \boxplus i_1$. This follows from the following three statements:

- (a) Morphisms of the form (2.10) span $\text{Mor}_{\mathcal{C}}(\lambda, \mu)$.
- (b) Suppose $\lambda \in \mathcal{P}$ and $j_1, \dots, j_\ell, i_1, \dots, i_\ell \in \mathbb{Z}$ such that

$$(2.11) \quad \lambda \boxplus i_\ell \boxplus \cdots \boxplus i_2 \boxplus i_1 = \lambda \boxplus j_\ell \boxplus \cdots \boxplus j_2 \boxplus j_1$$

are nonzero. Then

$$f_{i_1} f_{i_2} \cdots f_{i_\ell} 1_{\lambda} = f_{j_1} f_{j_2} \cdots f_{j_\ell} 1_{\lambda}.$$

- (c) Suppose $\lambda \in \mathcal{P}$ and $j_1, \dots, j_\ell, i_1, \dots, i_\ell \in \mathbb{Z}$ such that

$$\lambda \boxminus i_\ell \boxminus \cdots \boxminus i_2 \boxminus i_1 = \lambda \boxminus j_\ell \boxminus \cdots \boxminus j_2 \boxminus j_1$$

are nonzero. Then

$$e_{i_1} e_{i_2} \cdots e_{i_\ell} 1_{\lambda} = e_{j_1} e_{j_2} \cdots e_{j_\ell} 1_{\lambda}.$$

Proof of (a). — Given a morphism in \mathcal{C} that is a composition of $e_i 1_\lambda$ and $f_i 1_\lambda$, it follows from (2.7) and (2.8) that this composition is isomorphic to a 1-morphism of the form (2.10), possibly not satisfying the condition $\{i_1, \dots, i_k\} \cap \{j_1, \dots, j_\ell\} = \emptyset$. To see that we can also satisfy this condition, choose $a \in \{1, \dots, k\}$ and $b \in \{1, \dots, \ell\}$ such that $i_a = j_b$ and such that we cannot find $a' \in \{a, \dots, k\}$ and $b' \in \{1, \dots, b\}$ such that $i_{a'} = j_{b'}$ and either $a' > a$ or $b' < b$. (Intuitively speaking, we pick an “innermost” f_i, e_i pair.) We claim that none of the indices $a + 1, a + 2, \dots, k$ or $1, 2, \dots, b - 1$ is equal to $i_a - 1$ or $i_a + 1$. It will then follow from (2.5) and (2.6) that our morphism is equal to one in which f_{i_a} is immediately to the left of e_{j_b} , allowing us to use (2.9) to cancel this pair. Then statement (a) follows by induction.

To prove the claim, consider the morphism $1_\mu e_{j_b} e_{j_{b+1}} \cdots e_{j_\ell} 1_\lambda$. We then have $\mu = \lambda \boxplus j_\ell \boxminus \cdots \boxminus j_b$. In particular, μ has an addable j_b box. If we now remove a $j_b + 1$ box or a $j_b - 1$ box, the resulting Young diagram will no longer have an addable j_b box. Therefore, by our assumption that we have picked the innermost f_i, e_i pair, none of the indices $1, 2, \dots, b - 1$ is equal to $j_b + 1$ or $j_b - 1$. So $\mu \boxminus j_{b-1} \boxminus \cdots \boxminus j_1$ has an addable j_b box. But then it does not have an addable $j_b + 1$ box or an addable $j_b - 1$ box. Therefore, none of the indices i_{a+1}, \dots, i_k is equal to $i_a + 1$ or $i_a - 1$. This proves the claim.

Proof of (b). — Fix $\lambda \in \mathcal{P}$. We prove the statement by induction on ℓ . It is clear for $\ell = 0$ and $\ell = 1$. Suppose $\ell \geq 2$. The partition λ has a j_ℓ -addable box and an i_ℓ -addable box. If $j_\ell = i_\ell$, then the result follows by the inductive hypothesis applied to $\lambda \boxplus j_\ell$. So we assume $j_\ell \neq i_\ell$. By assumption, there must exist some $a \in \{1, \dots, \ell - 1\}$ such that $j_a = i_\ell$. Choose the maximal a with this property. By (2.11), λ has an addable i_ℓ -box. Thus, by an argument as in the proof of statement (a), none of the integers $j_\ell, j_{\ell-1}, \dots, j_{a+1}$ can be equal to $j_a \pm 1$. Then, by (2.6), we have

$$f_{j_1} f_{j_2} \cdots f_{j_\ell} 1_\lambda = f_{j_1} f_{j_2} \cdots f_{j_{a-1}} f_{j_{a+1}} \cdots f_{j_\ell} f_{j_a} 1_\lambda.$$

Then statement (b) follows by the inductive hypothesis applied to $\lambda \boxplus j_a = \lambda \boxplus i_\ell$.

Proof of (c). — The proof of statement (c) is analogous to that of statement (b). \square

2.4. NOTATION AND CONVENTIONS FOR 2-CATEGORIES. — We will use calligraphic font for 1-categories ($\mathcal{A}, \mathcal{C}, \mathcal{M}, \mathcal{V}$, etc.) and script font for 2-categories ($\mathcal{A}, \mathcal{C}, \mathcal{U}, \mathcal{H}$, etc.). We use bold lowercase for functors (\mathbf{a}, \mathbf{r} , etc.) and bold uppercase for 2-functors (\mathbf{F}, \mathbf{S} , etc.). The notation $\mathbf{0}$ will denote a zero object in a 1-category or 2-category. Other objects will be denoted with italics characters (x, y, e , etc.). We use sans serif font for 1-morphisms ($\mathbf{e}, \mathbf{x}, \mathbf{Q}$, etc.) and Greek letters for 2-morphisms.

If \mathcal{C} is a 2-category and x, y are objects of \mathcal{C} , we let $\mathcal{C}(x, y)$ denote the category of morphisms from x to y . We denote the class of objects of $\mathcal{C}(x, y)$, which are 1-morphisms in \mathcal{C} by $1\text{Mor}_{\mathcal{C}}(x, y)$. For \mathbf{P}, \mathbf{Q} objects in $\mathcal{C}(x, y)$, we denote the class of morphisms from \mathbf{P} to \mathbf{Q} , which are 2-morphisms in \mathcal{C} , by $2\text{Mor}_{\mathcal{C}}(\mathbf{P}, \mathbf{Q})$. For an object x of \mathcal{C} , we let 1_x denote the identity 1-morphism on x and let id_x denote the identity 2-morphism on 1_x . We denote vertical composition of 2-morphisms by \circ and

horizontal composition by juxtaposition. Whenever we speak of a linear category or 2-category, or a linear functor or 2-functor, we mean \mathbb{Q} -linear.

If \mathcal{C} is an additive linear 2-category, we define its Grothendieck group $K(\mathcal{C})$ to be the category with the same objects as \mathcal{C} and whose space of morphisms between objects x and y is $K(\mathcal{C}(x, y))$, the usual split Grothendieck group, over \mathbb{Q} , of the category $\mathcal{C}(x, y)$.

3. MODULES FOR SYMMETRIC GROUPS

In this section we recall some well-known facts about modules for symmetric groups and prove some combinatorial identities that we will need later on in our constructions.

3.1. MODULE CATEGORIES. — For an associative algebra A , we let $A\text{-mod}$ denote the category of finite-dimensional left A -modules. For $n \in \mathbb{N}$, we let $A_n = \mathbb{Q}S_n$ denote the group algebra of the symmetric group. By convention, we set $A_0 = A_1 = \mathbb{Q}$. We index the representations of A_n by partitions of n in the usual way, and for $\lambda \vdash n$, we let V_λ be the corresponding irreducible representation of A_n .

Let \mathcal{M}_λ denote the full subcategory of $A_n\text{-mod}$ whose objects are isomorphic to direct sums of V_λ (including the empty sum, which is the zero representation). We then have a decomposition

$$(3.1) \quad \mathcal{M}_n := A_n\text{-mod} = \bigoplus_{\lambda \vdash n} \mathcal{M}_\lambda.$$

We consider A_n to be a subalgebra of A_{n+1} in the natural way, where S_n is the subgroup of S_{n+1} fixing $n+1$. We use the notation (n) to denote A_n considered as an (A_n, A_n) -bimodule in the usual way. We use subscripts to denote restriction of the left and right actions. Thus, $(n+1)_n$ is A_{n+1} considered as an (A_{n+1}, A_n) -bimodule, ${}_n(n+1)$ is A_{n+1} considered as an (A_n, A_{n+1}) -bimodule, etc. Then

$$(n+1)_n \otimes_{A_n} -: A_n\text{-mod} \rightarrow A_{n+1}\text{-mod} \quad \text{and} \quad {}_n(n+1) \otimes_{A_{n+1}} -: A_{n+1}\text{-mod} \rightarrow A_n\text{-mod}$$

are the usual induction and restriction functors. Tensor products of such bimodules correspond in the same way to composition of induction and restriction functors, and bimodule homomorphisms correspond to natural transformations of the corresponding functors.

We define a 2-category \mathcal{M} as follows. The objects of \mathcal{M} are finite direct sums of \mathcal{M}_λ , $\lambda \in \mathcal{P}$, and a zero object $\mathbf{0}$. We adopt the conventions that $\mathcal{M}_n = \mathbf{0}$ when $n < 0$, $\mathcal{M}_{\lambda \boxplus i} = \mathbf{0}$ when $i \notin B^+(\lambda)$, and $\mathcal{M}_{\lambda \boxminus i} = \mathbf{0}$ when $i \notin B^-(\lambda)$. The 1-morphisms are generated, under composition and direct sum, by additive \mathbb{Q} -linear direct summands of the functors

$$(n+1)_n \otimes_{A_n} -: \mathcal{M}_n \rightarrow \mathcal{M}_{n+1}, \quad {}_{n-1}(n) \otimes_{A_n} -: \mathcal{M}_n \rightarrow \mathcal{M}_{n-1}.$$

The 2-morphisms of \mathcal{M} are natural transformations of functors.

REMARK 3.1. — In the above definition, it is important that we allow *direct summands* of the given functors. In Section 3.4 we will discuss the direct summands

$$(n+1)_n^i \otimes_{A_n} -: \mathcal{M}_\lambda \rightarrow \mathcal{M}_{\lambda \boxplus i}, \quad {}_{n-1}^i(n) \otimes_{A_n} -: \mathcal{M}_\lambda \rightarrow \mathcal{M}_{\lambda \boxminus i}, \quad \text{where } n = |\lambda|.$$

arising from decomposing induction and restriction according to eigenspaces for the action of Jucys–Murphy elements.

For $\lambda \vdash n$, consider the functors

$$(3.2) \quad \mathbf{i}_\lambda := \text{Hom}_{A_n}(V_\lambda, -) : \mathcal{M}_\lambda \longrightarrow \mathcal{V} \quad \text{and} \quad \mathbf{j}_\lambda := V_\lambda \otimes_{\mathbb{Q}} - : \mathcal{V} \longrightarrow \mathcal{M}_\lambda.$$

We have $\mathbf{i}_\lambda \circ \mathbf{j}_\lambda \cong 1_{\mathcal{V}}$ and $\mathbf{j}_\lambda \circ \mathbf{i}_\lambda \cong 1_{\mathcal{M}_\lambda}$, and hence an equivalence of categories $\mathcal{M}_\lambda \cong \mathcal{V}$.

3.2. DECATEGORYIFICATION. — Suppose $\lambda, \mu \in \mathcal{P}$ and consider an additive linear functor $\mathbf{a} : \mathcal{M}_\lambda \rightarrow \mathcal{M}_\mu$. Then the functor $\mathbf{i}_\mu \circ \mathbf{a} \circ \mathbf{j}_\lambda$ is naturally isomorphic to a direct sum of some finite number of copies of the identity functor. In other words, under the equivalences (3.2), every object in $\mathcal{M}(\mathcal{M}_\lambda, \mathcal{M}_\mu)$ is isomorphic to $1_{\mathcal{V}}^{\oplus n}$ for some $n \geq 0$, where $1_{\mathcal{V}} : \mathcal{V} \rightarrow \mathcal{V}$ is the identity functor.

It follows that $K(\mathcal{M})$ is the category given by

$$\text{Ob } K(\mathcal{M}) = \text{Ob } \mathcal{M} \quad \text{and} \quad \text{Mor}_{K(\mathcal{M})}(\mathcal{M}_\lambda, \mathcal{M}_\mu) = \mathbb{Q}, \quad \lambda, \mu \in \mathcal{P}.$$

Composition of morphisms is given by multiplication of the corresponding elements of \mathbb{Q} .

We have a natural functor $K(\mathcal{M}) \rightarrow \mathcal{V}$ given by

$$(3.3) \quad \mathcal{M}_\lambda \longmapsto \mathbb{Q}s_\lambda, \quad z \longmapsto (s_\lambda \mapsto zs_\mu),$$

for $\lambda, \mu \in \mathcal{P}$ and $z \in \mathbb{Q} = \text{Mor}_{K(\mathcal{M})}(\mathcal{M}_\lambda, \mathcal{M}_\mu)$. This functor is clearly an equivalence of categories.

3.3. BIADJUNCTION AND THE FUNDAMENTAL BIMODULE DECOMPOSITION

PROPOSITION 3.2. — *The maps*

$$\varepsilon_{\mathbb{R}} : (n+1)_n(n+1) \longrightarrow (n+1), \quad \varepsilon_{\mathbb{R}}(a \otimes b) = ab, \quad a \in (n+1)_n, \quad b \in {}_n(n+1),$$

$$\eta_{\mathbb{R}} : (n) \longleftarrow {}_n(n+1)_n, \quad \eta_{\mathbb{R}}(a) = a, \quad a \in (n),$$

$$\varepsilon_{\mathbb{L}} : {}_n(n+1)_n \longrightarrow (n), \quad \varepsilon_{\mathbb{L}}(g) = \begin{cases} g & \text{if } g \in S_n, \\ 0 & \text{if } g \in S_{n+1} \setminus S_n, \end{cases}$$

$$\eta_{\mathbb{L}} : (n+1) \longrightarrow (n+1)_n(n+1), \quad \eta_{\mathbb{L}}(a) = a \sum_{i \in \{1, \dots, n+1\}} s_i \cdots s_n \otimes s_n \cdots s_i, \quad a \in (n+1),$$

(where we interpret the expression $s_i \cdots s_n \otimes s_n \cdots s_i$ as $1 \otimes 1$ when $i = n+1$) are bimodule homomorphisms and satisfy the relations

$$(3.4) \quad (\varepsilon_{\mathbb{R}} \otimes \text{id}) \circ (\text{id} \otimes \eta_{\mathbb{R}}) = \text{id}, \quad (3.5) \quad (\text{id} \otimes \varepsilon_{\mathbb{R}}) \circ (\eta_{\mathbb{R}} \otimes \text{id}) = \text{id},$$

$$(3.6) \quad (\varepsilon_{\mathbb{L}} \otimes \text{id}) \circ (\text{id} \otimes \eta_{\mathbb{L}}) = \text{id}, \quad (3.7) \quad (\text{id} \otimes \varepsilon_{\mathbb{L}}) \circ (\eta_{\mathbb{L}} \otimes \text{id}) = \text{id}.$$

In particular, $(n+1)_n$ is both left and right adjoint to ${}_n(n+1)$ in the 2-category of bimodules over rings.

Proof. — The verification of these relations, which are a formulation of the well-known Frobenius reciprocity between induction and restriction for finite groups, is a straightforward computation. \square

It is well known (see, for example, [Kle05, Lem. 7.6.1]) that we have a decomposition

$$(3.8) \quad A_{n+1} = A_n \oplus (A_n s_n A_n),$$

and an isomorphism of (A_n, A_n) -bimodules

$$(3.9) \quad (n)_{n-1}(n) \xrightarrow{\cong} A_n s_n A_n \subseteq {}_n(n+1)_n, \quad a \otimes b \mapsto a s_n b.$$

This yields an isomorphism of (A_n, A_n) -bimodules

$$(3.10) \quad (n)_{n-1}(n) \oplus (n) \xrightarrow{\cong} {}_n(n+1)_n, \quad (a \otimes b, c) \mapsto a s_n b + c.$$

More precisely, the maps

$$(3.11) \quad (n)_{n-1}(n) \begin{array}{c} \xrightarrow{\rho} \\ \xleftarrow{\tau} \end{array} {}_n(n+1)_n \begin{array}{c} \xleftarrow{\eta_R} \\ \xrightarrow{\varepsilon_L} \end{array} (n)$$

where

$$(3.12) \quad \rho(a \otimes b) = a s_n b, \quad \text{for } a, b \in A_n,$$

$$(3.13) \quad \tau(a) = 0, \quad \tau(a s_n b) = a \otimes b, \quad \text{for } a, b \in A_n \subseteq A_{n+1},$$

satisfy

$$(3.14) \quad \varepsilon_L \circ \eta_R = \text{id}, \quad \tau \circ \rho = \text{id}, \quad \varepsilon_L \circ \rho = 0, \quad \tau \circ \eta_R = 0,$$

$$(3.15) \quad \rho \circ \tau + \eta_R \circ \varepsilon_L = \text{id}.$$

Note that

$$A_n s_n A_n = \text{Span}_{\mathbb{C}}(S_{n+1} \setminus S_n) \subseteq A_{n+1}.$$

3.4. THE JUCYS–MURPHY ELEMENTS AND THEIR EIGENSPACES. — Recall that the *Jucys–Murphy elements* of A_n are given by

$$(3.16) \quad J_1 = 0, \quad J_i = \sum_{k=1}^{i-1} (k, i), \quad i = 2, \dots, n,$$

where $(k, i) \in S_n$ denotes the transposition of k and i . The element J_i commutes with A_{i-1} . Thus, left multiplication by J_{n+1} is an endomorphism of the bimodule ${}_n(n+1)$. In fact, this action is semisimple and the set of eigenvalues is $\{-n, -n+1, \dots, n-1, n\}$.

We let ${}_n^i(n+1)$, $i \in \mathbb{Z}$, denote the i -eigenspace of ${}_n(n+1)$ under left multiplication by J_{n+1} . Similarly, we let $(n+1)_n^i$, $i \in \mathbb{Z}$, denote the i -eigenspace of $(n+1)_n$ under right multiplication by J_{n+1} . Since these two actions (right and left multiplication by J_{n+1}) commute, we can also consider the simultaneous eigenspaces ${}_n^i(n+1)_n^j$, $i, j \in \mathbb{Z}$. So we have

$$(3.17) \quad (n+1)_n = \bigoplus_{i \in \mathbb{Z}} (n+1)_n^i, \quad {}_n(n+1) = \bigoplus_{i \in \mathbb{Z}} {}_n^i(n+1), \quad {}_n(n+1)_n = \bigoplus_{i, j \in \mathbb{Z}} {}_n^i(n+1)_n^j.$$

Similarly, for $i, j \in \mathbb{Z}$, we let $(n+1)_{n-1}^{i, j}$ denote the simultaneous eigenspace of $(n+1)_{n-1}$ under right multiplication by J_{n+1} and J_n , with respective eigenvalues i and j . Similarly, we let ${}_{n-1}^{j, i}(n+1)$ denote the simultaneous eigenspace of ${}_{n-1}(n+1)$

under left multiplication by J_{n+1} and J_n , with respective eigenvalues i and j . We have

$$(3.18) \quad (n+1)_n^i \otimes_{A_n} V_\lambda \cong V_{\lambda \boxplus i} \quad \text{and} \quad {}_n^i(n+1) \otimes_{A_{n+1}} V_\mu \cong V_{\mu \boxminus i}$$

for $\lambda \vdash n$ and $\mu \vdash n+1$, where we define V_0 to be the zero module.

The primitive central idempotents in $\mathbb{Q}S_n$ are

$$(3.19) \quad e_\lambda = \frac{\dim V_\lambda}{n!} \sum_{w \in S_n} \text{tr}(w_\lambda^{-1})w, \quad \lambda \vdash n,$$

where w_λ^{-1} denotes the action of w^{-1} on the representation V_λ . (See, for example, [FH91, (2.13), p. 23].) Multiplication by e_λ is projection onto the V_λ -isotypic component. It follows that

$$(3.20) \quad (n+1)_n^i = \bigoplus_{\mu \vdash n} e_{\mu \boxplus i} (n+1)_n e_\mu,$$

$$(3.21) \quad {}_n^i(n+1) = \bigoplus_{\mu \vdash n} e_{\mu \boxminus i} (n+1)_n e_\mu.$$

3.5. COMBINATORIAL FORMULAS. — In this subsection we prove some combinatorial identities, used elsewhere in the paper, involving the dimensions $d_\lambda := \dim(V_\lambda)$ of irreducible representations of symmetric groups. By convention, $d_{\lambda \boxplus i} = 0$ if λ has no addable i -box, and $d_{\lambda \boxminus i} = 0$ if λ has no removable i -box.

It follows from (3.17) and (3.18) that

$$(3.22) \quad d_\lambda = \sum_{j \in B^-(\lambda)} d_{\lambda \boxplus j},$$

$$(3.23) \quad \sum_{i \in B^+(\lambda)} d_{\lambda \boxminus i} = (|\lambda| + 1)d_\lambda.$$

Recall the hook-length formula

$$(3.24) \quad d_\lambda = \frac{|\lambda|!}{\prod_{i,j} h_\lambda(i,j)}, \quad \lambda \in \mathcal{P}.$$

Here $h_\lambda(i,j)$ counts the number of boxes in the Young diagram of λ in the hook whose upper left corner is in position (i,j) and the product is over the positions (i,j) of all the boxes in the Young diagram of λ .

LEMMA 3.3. — *For a partition λ , we have*

$$(3.25) \quad \left(1 - \frac{1}{(i-j)^2}\right) \frac{|\lambda|+1}{|\lambda|+2} \frac{d_\lambda d_{\lambda \boxplus i \boxplus j}}{d_{\lambda \boxplus i} d_{\lambda \boxplus j}} = 1 \quad \forall i, j \in B^+(\lambda), |i-j| > 1,$$

$$(3.26) \quad \sum_{i \in B^+(\lambda)} \frac{d_{\lambda \boxplus i}}{(i-j)^2} = \frac{(|\lambda|+1)d_\lambda^2}{|\lambda|d_{\lambda \boxplus j}} \quad \forall j \in B^-(\lambda),$$

$$(3.27) \quad \sum_{j \in B^-(\lambda)} \frac{d_{\lambda \boxminus j}}{(i-j)^2} = \left(\frac{(|\lambda|+1)d_\lambda^2}{d_{\lambda \boxminus i}|\lambda|} - \frac{d_\lambda}{|\lambda|}\right) \quad \forall i \in B^+(\lambda).$$

Proof. — Relation (3.25) follows from a direct computation using (3.24). We omit the details.

To prove (3.26), assume $j \in B^-(\lambda)$. Then, for $|i - j| > 1$, (3.25) for $\lambda \boxplus j$ implies

$$(3.28) \quad \frac{1}{(i - j)^2} \frac{|\lambda| d_{\lambda \boxplus i} d_{\lambda \boxplus j}}{(|\lambda| + 1) d_\lambda} = \frac{|\lambda| d_{\lambda \boxplus i} d_{\lambda \boxplus j}}{(|\lambda| + 1) d_\lambda} - d_{\lambda \boxplus i \boxplus j}.$$

When $|i - j| = 1$, we have $d_{\lambda \boxplus i \boxplus j} = 0$, and so (3.28) in fact holds for all $i \neq j$. Therefore,

$$\begin{aligned} \sum_{i \in B^+(\lambda)} \frac{1}{(i - j)^2} \frac{|\lambda| d_{\lambda \boxplus i} d_{\lambda \boxplus j}}{(|\lambda| + 1) d_\lambda} &= \frac{|\lambda| d_{\lambda \boxplus j}}{(|\lambda| + 1) d_\lambda} \sum_{i \in B^+(\lambda)} d_{\lambda \boxplus i} - \sum_{i \in B^+(\lambda)} d_{\lambda \boxplus i \boxplus j} \\ &= \frac{|\lambda| d_{\lambda \boxplus j}}{(|\lambda| + 1) d_\lambda} \sum_{i \in B^+(\lambda)} d_{\lambda \boxplus i} - \sum_{i \in B^+(\lambda \boxplus j)} d_{\lambda \boxplus j \boxplus i} + d_\lambda \stackrel{(3.23)}{=} |\lambda| d_{\lambda \boxplus j} - |\lambda| d_{\lambda \boxplus j} + d_\lambda = d_\lambda, \end{aligned}$$

where the second equality follows from the fact that $\lambda \boxplus i \boxplus j = \mathbf{0}$ whenever $i \notin B^+(\lambda \boxplus j)$. Relation (3.26) follows.

To prove relation (3.27), suppose $i \in B^+(\lambda)$. Then

$$\begin{aligned} \sum_{j \in B^-(\lambda)} \frac{|\lambda| d_{\lambda \boxplus i} d_{\lambda \boxplus j}}{(i - j)^2 d_\lambda} &\stackrel{(3.28)}{=} \sum_{j \in B^-(\lambda)} \frac{|\lambda| d_{\lambda \boxplus i} d_{\lambda \boxplus j}}{d_\lambda} - \sum_{j \in B^-(\lambda)} (|\lambda| + 1) d_{\lambda \boxplus i \boxplus j} \\ &\stackrel{(3.22)}{=} |\lambda| d_{\lambda \boxplus i} - (|\lambda| + 1) \left(\sum_{j \in B^-(\lambda \boxplus i)} d_{\lambda \boxplus i \boxplus j} - d_\lambda \right) \\ &\stackrel{(3.22)}{=} |\lambda| d_{\lambda \boxplus i} - (|\lambda| + 1) (d_{\lambda \boxplus i} - d_\lambda) = (|\lambda| + 1) d_\lambda - d_{\lambda \boxplus i}, \end{aligned}$$

where, in the second equality, we have used the fact that $\lambda \boxplus i \boxplus j = \mathbf{0}$ whenever $j \notin B^-(\lambda \boxplus i)$. Relation (3.27) follows. \square

LEMMA 3.4

(a) For $\lambda \vdash n$, we have

$$(3.29) \quad \varepsilon_L(e_\lambda) = \sum_{i \in B^-(\lambda)} \frac{d_\lambda}{n d_{\lambda \boxplus i}} e_{\lambda \boxplus i}.$$

(b) For $\lambda \vdash n$ and $i \in \mathbb{Z}$, we have

$$(3.30) \quad J_{n+1} e_{\lambda \boxplus i} e_\lambda = i e_{\lambda \boxplus i} e_\lambda.$$

(c) For $\lambda \vdash n$ and $i \in \mathbb{Z}$, we have

$$(3.31) \quad \frac{1}{(n + 1)!} \sum_{w \in S_{n+1}} w e_\lambda w^{-1} = \sum_{i \in B^+(\lambda)} \frac{d_\lambda}{d_{\lambda \boxplus i}} e_{\lambda \boxplus i}.$$

Proof. — (a) Fix $\lambda \vdash n$. As $\mathbb{Q}S_{n-1}$ -modules, we have a decomposition $V_\lambda \cong \bigoplus_{i \in B^-(\lambda)} V_{\lambda \boxplus i}$. Then

$$\begin{aligned} \varepsilon_L(e_\lambda) &\stackrel{(3.19)}{=} \varepsilon_L \left(\frac{d_\lambda}{n!} \sum_{w \in S_n} \text{tr}(w_\lambda^{-1}) w \right) = \frac{d_\lambda}{n!} \sum_{w \in S_{n-1}} \text{tr}(w_\lambda^{-1}) w \\ &= \frac{d_\lambda}{n!} \sum_{i \in B^-(\lambda)} \sum_{w \in S_{n-1}} \text{tr}(w_{\lambda \boxplus i}^{-1}) w \stackrel{(3.19)}{=} \sum_{i \in B^-(\lambda)} \frac{d_\lambda}{n d_{\lambda \boxplus i}} e_{\lambda \boxplus i}. \end{aligned}$$

(b) Recall that if $\lambda \vdash n$ and V is an A_n -module, then $e_\lambda(V)$ is the λ -isotypic component of V . Now, to show that two elements of A_{n+1} are equal it suffices to show that they act identically on every irreducible representation of A_{n+1} . For $\mu \vdash (n+1)$ and $\lambda \vdash n$, let $V_{\mu,\lambda}$ be the λ -isotypic component of V_μ (this is either zero or isomorphic to V_λ as an A_n -module). Then

$$e_{\lambda \boxplus i} e_\lambda(V_\mu) = \delta_{\mu, \lambda \boxplus i} V_{\lambda \boxplus i, \lambda}$$

The result then follows from the fact that $V_{\lambda \boxplus i, \lambda}$ is the i -eigenspace of $V_{\lambda \boxplus i}$ under multiplication by J_{n+1} .

(c) Fix $\lambda \vdash n$. Then $\frac{1}{(n+1)!} \sum_{w \in S_{n+1}} w e_\lambda w^{-1}$ belongs to the center of $\mathbb{Q}S_{n+1}$, and so is a linear combination of the central idempotents e_ν , $\nu \vdash n+1$. Thus we have

$$\frac{1}{(n+1)!} \sum_{w \in S_{n+1}} w e_\lambda w^{-1} = \sum_{\nu \vdash n+1} c_\nu e_\nu$$

for some $c_\nu \in \mathbb{Q}$. Therefore, for $\mu \vdash n+1$, we have

$$c_\mu e_\mu = e_\mu \sum_{\nu \vdash n+1} c_\nu e_\nu = \frac{e_\mu}{(n+1)!} \sum_{w \in S_{n+1}} w e_\lambda w^{-1} = \frac{1}{(n+1)!} \sum_{w \in S_{n+1}} w e_\mu e_\lambda w^{-1}.$$

Then, considering the action of $\mathbb{Q}S_{n+1}$ on V_μ , we have

$$c_\mu d_\mu = \text{tr}(c_\mu e_\mu) = \frac{1}{(n+1)!} \sum_{w \in S_{n+1}} \text{tr}(e_\mu e_\lambda) = \begin{cases} d_\lambda & \text{if } \mu = \lambda \boxplus i \text{ for some } i \in \mathbb{Z}, \\ 0 & \text{otherwise.} \end{cases}$$

The result follows. □

4. THE 2-CATEGORY \mathcal{A}

In this section we define an additive linear 2-category \mathcal{A} and investigate some of its key properties. We will show in Section 4.2 that \mathcal{A} is equivalent to the degree zero part of a truncation of a categorified quantum group. Later, in Section 6, we will also show that \mathcal{A} is equivalent to a summand of a truncation of a Heisenberg 2-category. We will also see, in Proposition 7.3, that \mathcal{A} is equivalent to the category \mathcal{M} .

4.1. DEFINITION. — The set of objects of \mathcal{A} is the free monoid on the set of partitions:

$$\text{Ob } \mathcal{A} = \mathbb{N}[\mathcal{P}].$$

We denote the zero object by $\mathbf{0}$. The 1-morphisms of \mathcal{A} are generated by (i.e. direct sums of compositions of)

$$\begin{aligned} F_i 1_\lambda &= 1_{\lambda \boxplus i} F_i = 1_{\lambda \boxplus i} F_i 1_\lambda, \quad \text{and} \\ E_i 1_\lambda &= 1_{\lambda \boxminus i} E_i = 1_{\lambda \boxminus i} E_i 1_\lambda, \quad i \in \mathbb{Z}, \lambda \in \mathcal{P}. \end{aligned}$$

We adopt the convention that if λ does not have an addable (resp. removable) i -box, then $1_{\lambda \boxplus i} = 0$ (resp. $1_{\lambda \boxminus i} = 0$) and hence $\lambda \boxplus i \cong \mathbf{0}$ (resp. $\lambda \boxminus i \cong \mathbf{0}$) in \mathcal{A} . In particular,

$$(4.1) \quad E_i^2 1_\lambda = 0 \quad \text{and} \quad F_i^2 1_\lambda = 0 \quad \text{for all } \lambda \in \mathcal{P},$$

and similarly

$$(4.2) \quad E_i E_{i\pm 1} E_i 1_\lambda = 0 \quad \text{and} \quad F_i F_{i\pm 1} F_i 1_\lambda = 0 \quad \text{for all } \lambda \in \mathcal{P}.$$

The space of 2-morphisms between two 1-morphisms is the \mathbb{Q} -algebra generated by suitable planar diagrams modulo local relations. The diagrams consist of oriented compact one-manifolds immersed into the plane strip $\mathbb{R} \times [0, 1]$ modulo local relations, with strands labeled by integers and regions of the strip labeled/colored by elements of $\mathcal{P} \sqcup \{0\}$. In particular, the identity 2-morphism of $F_i 1_\lambda$ will be denoted by an upward strand labeled i , where the region to the right of the arrow is labeled λ , while the identity 2-morphism of $E_i 1_\lambda$ is denoted by a downward strand labeled i , where the region to the right of the strand is labeled λ :



Strands of distinct color may intersect transversely, but no triple intersections are allowed. The space of 2-morphisms is the space of such planar diagrams up to isotopy and modulo local relations. The domain and codomain are given by the orientations of the strands at the bottom and top of the diagram respectively.

The local relations are as follows, where i, j, k range over all integers satisfying

$$|i - j|, |i - k|, |j - k| > 1.$$

For relations when the regions are not labeled, we impose the relation for all labelings of the regions.

$$(4.3) \quad \begin{array}{c} \nearrow \\ \searrow \nearrow \\ i \quad j \quad k \end{array} = \begin{array}{c} \searrow \nearrow \\ \nearrow \\ i \quad j \quad k \end{array}$$

$$(4.4) \quad \begin{array}{c} \nearrow \\ \searrow \nearrow \\ i \quad j \end{array} = \begin{array}{c} \uparrow \\ \uparrow \\ i \quad j \end{array}$$

$$(4.5) \quad \begin{array}{c} \nearrow \\ \searrow \nearrow \\ i \quad j \end{array} = \begin{array}{c} \downarrow \\ \downarrow \\ i \quad j \end{array}$$

$$(4.6) \quad \begin{array}{c} \searrow \nearrow \\ \nearrow \\ i \quad j \end{array} = \begin{array}{c} \uparrow \\ \downarrow \\ i \quad j \end{array}$$

$$(4.7) \quad \begin{array}{c} \downarrow \\ \uparrow \\ i \quad i \end{array} = \begin{array}{c} \curvearrowright \\ \curvearrowleft \\ i \end{array}$$

$$(4.8) \quad \begin{array}{c} \uparrow \\ \downarrow \\ i \quad i \end{array} = \begin{array}{c} \curvearrowleft \\ \curvearrowright \\ i \end{array}$$

$$(4.9) \quad i \circlearrowleft^\lambda = \text{id}_\lambda \quad \text{for } i \in B^-(\lambda), \quad (4.10) \quad i \circlearrowright^\lambda = \text{id}_\lambda \quad \text{for } i \in B^+(\lambda).$$

REMARK 4.1. — Note that the crossings



are always zero for $|i - j| \leq 1$. In other words, nonzero diagrams can only have strands crossing if their colors differ by at least two. This is because $\lambda \boxplus i \boxplus j = \mathbf{0}$

when $|i - j| \leq 1$. Therefore, relations (4.4)–(4.6) allow us to resolve all nonzero double crossings.

4.2. TRUNCATED CATEGORIFIED QUANTUM GROUPS. — In [CL15, §2], Cautis and Lauda associated a graded additive linear 2-category to a Cartan datum and choice of scalars, generalizing the definition of [KL10]. We let \mathcal{U} be the 2-category defined in [CL15, §2] for the Cartan datum of type A_∞ and the choice of scalars

$$(4.11) \quad t_{ij} = 1, \quad s_{ij}^{pq} = 0, \quad r_i = 1, \quad \text{for all } i, j, p, q,$$

except that we enlarge the 2-category by allowing finite formal direct sums of objects. By [CL15, Rem. (3), p. 210], we in fact lose no generality in making the choices (4.11). The set of objects of \mathcal{U} is the free monoid generated by the weight lattice of \mathfrak{sl}_∞ . We define \mathcal{U}^{tr} to be the quotient of \mathcal{U} by the identity 2-morphisms of the identity 1-morphisms of all objects corresponding to weights that do not appear in the basic representation. Since the weights of the basic representation are in natural bijection with partitions (see Section 2.2), the objects of \mathcal{U}^{tr} can be identified with elements of the free monoid $\mathbb{N}[\mathcal{P}]$ on the set of partitions.

PROPOSITION 4.2. — *All the 2-morphism spaces of \mathcal{U}^{tr} are nonnegatively graded. The positive degree 2-morphism spaces are spanned by diagrams with dots.*

Proof. — The degrees of the 2-morphisms in \mathcal{U} are given by the degrees of the crossings, cups, caps, and dots in [CL15, Def. 1.1]. The crossing has degree

$$\deg \left(\begin{array}{c} \nearrow \lambda \\ \searrow \\ i \quad j \end{array} \right) = \begin{cases} -2, & \text{if } i = j, \\ 1, & \text{if } |i - j| = 1, \\ 0, & \text{otherwise.} \end{cases}$$

However, the leftmost region above is labeled $\lambda \boxplus j \boxminus i$, which is always isomorphic to $\mathbf{0}$ (i.e. is not a weight of the basic representation) when $|i - j| \leq 1$. (Note that the upward oriented strands in \mathcal{U} correspond to subtracting boxes under the bijection (2.4), as opposed to adding boxes, as in \mathcal{A} .) Therefore, the only crossings which are nonzero in \mathcal{U}^{tr} have degree zero. The situation for downwards oriented crossings is analogous.

Similarly, the right cup



has degree $1 + (\lambda, \alpha_i)$. However, this cup is zero in \mathcal{U}^{tr} unless λ has a removable i -box, in which case $(\lambda, \alpha_i) = -1$. Thus, the only right cups that are nonzero in \mathcal{U}^{tr} have degree zero. The situation for the other cups and caps is analogous.

The result now follows from the fact that a dot on an i -colored strand has degree $(\alpha_i, \alpha_i) = 2$. \square

Let \mathcal{U}_0 be the additive linear 2-category defined as follows. The objects of \mathcal{U}_0 are the same as the objects of \mathcal{U}^{tr} . The 1-morphisms in \mathcal{U}_0 are formal direct sums of compositions of the generating 1-morphisms given in [CL15, Def. 1.1] *without degree*

shifts. The 2-morphism spaces of \mathcal{U}_0 are the degree zero part of the corresponding 2-morphism spaces of \mathcal{U}^{tr} (equivalently, the quotient of the corresponding 2-morphism spaces of \mathcal{U}^{tr} by the ideal consisting of 2-morphisms of strictly positive degree).

REMARK 4.3. — It will follow from Theorem 4.4 and Corollary 4.10 that \mathcal{U}_0 is idempotent complete; hence there is no need to pass to the idempotent completion, as is done in [CL15] with the larger category \mathcal{U} .

THEOREM 4.4. — *The 2-categories \mathcal{A} and \mathcal{U}_0 are equivalent via a 2-functor that acts on 2-morphisms by reversing the orientation of strands.*

Proof. — The sets of objects and 1-morphisms of \mathcal{A} and \mathcal{U}_0 are clearly the same. Furthermore, the spaces of 2-morphisms both consist of string diagrams with strands labeled by integers. Therefore, it suffices to check that the local relations are the same. The relations of [CL15, §2.2] correspond in \mathcal{A} to the fact that we consider diagrams up to isotopy. Relations [CL15, (2.8)] and [CL15, (2.9)] become trivial since they involve strands of the same color crossing or parallel strands of the same color, which yield the zero 2-morphism in the truncation. Relation [CL15, (2.10)] corresponds to (4.4). Note that the $(\alpha_i, \alpha_j) \neq 0$ case of [CL15, (2.10)] is trivial in the truncation since strands of color i and $i + 1$ cannot cross. Relation [CL15, (2.12)] is not relevant since it involves dots. Relation [CL15, (2.13)] becomes (4.3). Note that [CL15, (2.14)] becomes trivial in the truncation since $(\alpha_i, \alpha_j) < 0$ implies $|i - j| = 1$, in which case we have two strands of the same color crossing.

The relations [CL15, (2.16)] correspond to (4.5) and (4.6). For $\lambda \in X$ and $i \in \mathbb{Z}$, we have

$$\langle i, \lambda \rangle = (\alpha_i, \lambda) = \begin{cases} 1 & \text{if } i \in B^+(\lambda), \\ -1 & \text{if } i \in B^-(\lambda), \\ 0 & \text{otherwise.} \end{cases}$$

Therefore, relations [CL15, (2.17)] become trivial for us since the conditions on m are never satisfied. On the other hand, the relations at the top of page 211 of [CL15] correspond to (4.9) and (4.10).

Now consider the extended \mathfrak{sl}_2 relations of [CL15, §2.6]. Here one considers relations involving strands of some fixed color i . In the truncation, the region labels in [CL15, §2.6] of nonzero diagrams are 0 (corresponding to a region label λ with no i -addable or i -removable boxes), 1 (corresponding to a λ with an i -removable box but no i -addable box), or -1 (corresponding to a λ with an i -addable box, but no i -removable box). The relations [CL15, (2.21)] become trivial in the truncation since a strand crossing itself is zero. It follows from [CL15, (2.19)] with $n = 1$ and $j = 0$, together with (4.9) (which tells us that a clockwise circle in outer region 1 is equal to one) that a counterclockwise fake bubble with dot label -2 and outer region 1 is equal to 1. Thus, the first relation of [CL15, (2.22)] becomes (4.7). The second relation in [CL15, (2.22)] becomes zero in the truncation, since $n > 0$ implies that n has no i -addable box. The relations [CL15, (2.23)] are trivial in the truncation since

they involve a strand crossing itself. The first relation in [CL15, (2.24)] is trivial in the truncation, while the second relation in [CL15, (2.24)] becomes (4.8). Finally, relations [CL15, (2.25), (2.26)] are trivial in the truncation. \square

4.3. 1-MORPHISM SPACES

LEMMA 4.5. — For $\lambda \in \mathcal{P}$ and $i, j \in \mathbb{Z}$, we have

$$(4.12) \quad E_i E_j 1_\lambda \cong E_j E_i 1_\lambda, \quad \text{if } |i - j| > 1,$$

$$(4.13) \quad F_i F_j 1_\lambda \cong F_j F_i 1_\lambda, \quad \text{if } |i - j| > 1,$$

$$(4.14) \quad E_i F_j 1_\lambda \cong F_j E_i 1_\lambda, \quad \text{if } i \neq j,$$

$$(4.15) \quad E_i F_i 1_\lambda \cong 1_\lambda, \quad \text{if } i \in B^+(\lambda),$$

$$(4.16) \quad F_i E_i 1_\lambda \cong 1_\lambda, \quad \text{if } i \in B^-(\lambda).$$

Proof. — The isomorphisms (4.12) and (4.13) follow immediately from (4.4), isomorphism (4.14) follows immediately from (4.5) and (4.6), isomorphism (4.15) follows from (4.7) and (4.10), and (4.16) follows from (4.8) and (4.9). \square

PROPOSITION 4.6. — We have

$$\dim \text{Mor}_{K(\mathcal{A})}(\lambda, \mu) = 1 \quad \text{for all } \lambda, \mu \in \mathcal{P}.$$

In particular, for $\lambda, \mu \in \mathcal{P}$, $\text{Mor}_{K(\mathcal{A})}(\lambda, \mu)$ is spanned by the class of a single 1-morphism in \mathcal{A} of the form

$$(4.17) \quad F_{i_1} F_{i_2} \cdots F_{i_k} E_{j_1} E_{j_2} \cdots E_{j_\ell} 1_\lambda,$$

where $\{i_1, \dots, i_k\} \cap \{j_1, \dots, j_\ell\} = \emptyset$ and $\mu = \lambda \boxplus j_\ell \cdots \boxplus j_2 \boxplus j_1 \boxplus i_k \boxplus \cdots \boxplus i_2 \boxplus i_1$.

Proof. — The proof of this statement is analogous to the proof of Proposition 2.2. \square

4.4. 2-MORPHISM SPACES

LEMMA 4.7 (Triple point moves). — Suppose $i, j, k \in \mathbb{Z}$. The relation

$$(4.18) \quad \begin{array}{c} \diagup \quad \diagdown \\ \diagdown \quad \diagup \\ i \quad j \quad k \end{array} = \begin{array}{c} \diagup \quad \diagdown \\ \diagdown \quad \diagup \\ i \quad j \quad k \end{array}$$

holds for all possible orientations of the strands.

Proof. — The case where all strands are pointed up is (4.3), and then the case where all strands are pointed down follows by isotopy invariance. We compute


$$\begin{array}{c} \diagup \quad \diagdown \\ \diagdown \quad \diagup \\ i \quad j \quad k \end{array} \stackrel{(4.4)}{=} \begin{array}{c} \diagup \quad \diagdown \\ \diagdown \quad \diagup \\ i \quad j \quad k \end{array} \stackrel{(4.3)}{=} \begin{array}{c} \diagup \quad \diagdown \\ \diagdown \quad \diagup \\ i \quad j \quad k \end{array} \stackrel{(4.4)}{=} \begin{array}{c} \diagup \quad \diagdown \\ \diagdown \quad \diagup \\ i \quad j \quad k \end{array}.$$

We omit proofs of the other cases, which are analogous. \square

LEMMA 4.8. — For all $\lambda \in \mathcal{P}$, we have $2\text{Mor}_{\mathcal{A}}(1_\lambda, 1_\lambda) \cong \mathbb{Q}\text{id}_\lambda$. In other words, all closed diagrams in a region labeled λ are isomorphic to some multiple of the empty diagram.

Proof. — Using the local relations, closed diagrams can be written as linear combinations of nested circles. By (4.9) and (4.10), these are equal to multiples of the empty diagram. \square

By definition, the 1-morphisms of \mathcal{A} are sequences of E_i 's and F_i 's, followed by 1_λ for some $\lambda \in \mathcal{P}$. If we think of such a sequence as a row of colored arrows, a down i -colored arrow for each E_i and an up i -colored arrow for each F_i , then the 2-morphisms between two 1-morphisms are strands connecting the arrows in such a way that the color and orientation of each strand agrees with its two endpoints. (We use Lemma 4.8 here to ignore closed diagrams.) By (4.4), (4.5), and (4.6), we may simplify such a diagram so that it contains no double crossings. Then, by isotopy invariance and Lemma 4.7, we see that the 2-morphism is uniquely determined by the matching of arrows induced by the strands. Furthermore, since strands of the same color cannot cross, the 2-morphism is determined by a crossingless matching of i -colored arrows for each $i \in \mathbb{Z}$. We call such a collection of crossingless matchings a *colored matching*. An example of such a colored matching is the following, where i, j, k , and ℓ are pairwise distinct.



Note that, since strands of colors i and j intersect, we must have $|i - j| > 1$ in order for the 2-morphism to be nonzero, by Remark 4.1. Similarly, we have $|i - k| > 1$ and $|j - k| > 1$. However, it is possible that $|k - \ell| = 1$.

PROPOSITION 4.9. — Suppose $\lambda, \mu \in \mathcal{P}$, and that $P, Q \in 1\text{Mor}_{\mathcal{A}}(\lambda, \mu)$ are two nonzero 1-morphisms that are sequences of E_i 's and F_i 's (followed by 1_λ). Then every nonzero 2-morphism from P to Q is an isomorphism. In particular, $1\text{Mor}_{\mathcal{A}}(P, Q)$ is one-dimensional.

Proof. — It is easy to see that there is always at least one colored matching. Repeated use of relations (4.7) and (4.8) allows us to see that any two crossingless matchings of i -colored arrows as described above are equal, up to scalar multiple.

Now let α be a nonzero 2-morphism from P to Q . By the above argument, α is a multiple of a colored matching from P to Q . Let β be a multiple of a colored matching from Q to P and consider the composition $\alpha \circ \beta$. We can resolve double crossings by (4.4), (4.5), and (4.6). Any circles may be slid into open regions (so that they do not intersect any other strands) using (4.4), (4.5), and (4.6) and then removed using (4.9)

and (4.10). Thus, $\alpha \circ \beta$ is a nonzero multiple of a colored matching. As above, it must be a multiple of the identity matching. Similarly $\beta \circ \alpha$ is a multiple of the identity matching. \square

COROLLARY 4.10. — *The 2-category \mathcal{A} is Krull–Schmidt. More precisely, for any two objects of \mathcal{A} , the morphism category between these two objects in Krull–Schmidt.*

Proof. — It follows from Proposition 4.6 that every 1-morphism in \mathcal{A} is a multiple of a 1-morphism of the form (4.17). Then the result follows from Proposition 4.9. \square

4.5. DECATEGORYIFICATION. — We now state one of our main results.

THEOREM 4.11. — *The functor $\mathcal{A} \rightarrow K(\mathcal{A})$ that is the identity on objects and, on 1-morphisms, is uniquely determined by*

$$e_i 1_\lambda \mapsto [E_i 1_\lambda], \quad f_i 1_\lambda \mapsto [F_i 1_\lambda],$$

is an isomorphism. In other words, \mathcal{A} categorifies \mathcal{A} .

Proof. — The fact that the functor is well-defined follows from Lemma 4.5. It is surjective since the images in the Grothendieck group of all the generating 1-morphisms $E_i 1_\lambda$ and $F_i 1_\lambda$ are in the image of the functor. Injectivity will be proven in Corollary 7.5. \square

5. THE 2-CATEGORY \mathcal{H}^{tr}

5.1. DEFINITION. — We introduce here a 2-category based on the diagrammatic monoidal category introduced by Khovanov in [Kho14]. We begin by defining an additive linear 2-category $\mathcal{H}^{\text{tr}'}$. The set of objects of $\mathcal{H}^{\text{tr}'}$ is the free monoid $\mathbb{N}[\mathbb{N}]$ on \mathbb{N} , where $\mathbf{0}$ is a zero object. The set of 1-morphisms of $\mathcal{H}^{\text{tr}'}$ is generated by

$$\mathbb{Q}_+ 1_k = 1_{k+1} \mathbb{Q}_+ = 1_{k+1} \mathbb{Q}_+ 1_k \quad \text{and} \quad \mathbb{Q}_- 1_{k+1} = 1_k \mathbb{Q}_- = 1_k \mathbb{Q}_- 1_{k+1}, \quad k \in \mathbb{N}.$$

In other words, if we let $\mathbb{Q}_c := \mathbb{Q}_{c_1} \otimes \cdots \otimes \mathbb{Q}_{c_\ell}$ for a finite sequence $c = c_1 \cdots c_\ell$ of $+$ and $-$ signs, then the 1-morphisms of $\mathcal{H}^{\text{tr}'}$ from n to m are finite direct sums of $\mathbb{Q}_c 1_n$ for $c = c_1 \cdots c_\ell$ satisfying

$$m - n = \#\{i \mid c_i = +\} - \#\{i \mid c_i = -\}.$$

If, for some $1 \leq k < \ell$, we have

$$n + \#\{1 \leq i \leq k \mid c_i = +\} - \#\{1 \leq i \leq k \mid c_i = -\} < 0,$$

then $\mathbb{Q}_c 1_n$ is the zero morphism. In other words, we view negative integers as the zero object $\mathbf{0}$.

The space of morphisms between two objects is the \mathbb{Q} -algebra generated by suitable planar diagrams. The diagrams consist of oriented compact one-manifolds immersed into the plane strip $\mathbb{R} \times [0, 1]$ modulo certain local relations, with regions of the strip labeled by nonnegative integers. The 2-morphism that is the identity on $\mathbb{Q}_+ 1_k$ will be denoted by an upward strand, where the region to the right of the arrow is labeled k (and the region to the left has label $k+1$), while the identity on $\mathbb{Q}_- 1_k$ will be denoted

by a downward strand, where the region to the right of the strand is labeled k (and the region to the left has label $k - 1$):



The endpoints of the strings are located at $\{1, \dots, m\} \times \{0\}$ and $\{1, \dots, k\} \times \{1\}$, where m and k are the lengths of the sequences c and c' respectively. The local relations are as follows (for any compatible labeling of the regions).

(5.1) ,

(5.2) ,

(5.3) ,

(5.4) ,

(5.5) = 1,

(5.6) = 0.

By convention, any string diagram containing a region labeled by a negative integer is the zero 2-morphism. This is compatible with our convention above for 1-morphisms.

DEFINITION 5.1 (Idempotent completion of a 2-category). — For a 2-category \mathcal{C} , we define an idempotent completion $\text{Kar}(\mathcal{C})$ as follows.

- The objects of $\text{Kar}(\mathcal{C})$ are triples (x, e, ϵ) , where $x \in \text{Ob } \mathcal{C}$, e is an idempotent 1-morphism of x in \mathcal{C} , and ϵ is an idempotent 2-morphism (under vertical composition) of e in \mathcal{C} .

- The 1-morphisms of $\text{Kar}(\mathcal{C})$ between objects (x, e, ϵ) and (x', e', ϵ') are pairs (g, β) , where $g: x \rightarrow x'$ is a 1-morphism in \mathcal{C} such that $e'ge = g$ and $\beta: g \rightarrow g$ is an idempotent 2-morphism in \mathcal{C} such that $\epsilon'\beta\epsilon = \beta$.

- The 2-morphisms between parallel 1-morphisms

$$(g, \beta), (h, \gamma): (x, e, \epsilon) \longrightarrow (x', e', \epsilon')$$

are 2-morphisms $\alpha: g \rightarrow h$ in \mathcal{C} such that $\gamma \circ \alpha \circ \beta = \alpha$.

Composition of 1-morphisms is pairwise composition of 1-morphisms as in \mathcal{C} and composition of 2-morphisms is as in \mathcal{C} .

If \mathcal{C} is a 2-category, we have a natural inclusion of \mathcal{C} into $\text{Kar}(\mathcal{C})$ sending the object x to $(x, 1_x, \text{id}_x)$ and the 1-morphism x to (x, id_x) . The idempotent completion of a 2-category \mathcal{C} is universal in the sense that any 2-functor $\mathcal{C} \rightarrow \mathcal{D}$ to a 2-category \mathcal{D} in which all idempotent 1-morphisms and idempotent 2-morphisms split factors through a 2-functor $\text{Kar}(\mathcal{C}) \rightarrow \mathcal{D}$.


We then define

$$\mathcal{H}^{\text{tr}} = \text{Kar}(\mathcal{H}^{\text{tr}'}) .$$


REMARK 5.2

(a) In the 2-category $\mathcal{H}^{\text{tr}'}$, the only idempotent 1-morphisms are the identity 1-morphisms. However, we state Definition 5.1 in full generality.

where A^{op} denotes the opposite algebra of an algebra A . For $z \in \mathbb{Q}S_n$, we will denote the corresponding elements of $2\text{End}_{\mathcal{H}^{\text{tr}}}(\mathbb{Q}_+^n 1_k)$ and $2\text{End}_{\mathcal{H}^{\text{tr}}}(\mathbb{Q}_-^n 1_k)$ by



respectively, where the dashed strand represents n strands (determined by z). It follows that



for all $z \in \mathbb{Q}S_n$.

LEMMA 5.4. — For all $n > 0$, we have

$$(5.8) \quad \begin{array}{c} \uparrow 0 \\ \text{---} \\ \downarrow 0 \end{array} = \sum_{z \in S_n} \begin{array}{c} \uparrow \\ \boxed{z} \\ \downarrow \end{array} \begin{array}{c} 0 \\ \text{---} \\ \downarrow \end{array}.$$

Proof. — This follows from repeated use of (5.3). □

Recall the definition of the central idempotent e_λ in (3.19).

LEMMA 5.5. — For $n \in \mathbb{N}$, in $2\text{End}_{\mathcal{H}^{\text{tr}}}(1_n)$ we have the following orthogonal idempotent decomposition of id_n :

$$(5.9) \quad \text{id}_n = \sum_{\lambda \vdash n} \epsilon_\lambda + \delta_n,$$

where, for $n > 0$, we define

$$(5.10) \quad \epsilon_\lambda = \frac{1}{n!} \begin{array}{c} \text{---} \\ \boxed{e_\lambda} \\ \uparrow \end{array} \begin{array}{c} 0 \\ \text{---} \\ \downarrow \end{array}, \quad \delta_n = 1 - \epsilon_n, \quad \epsilon_n = \sum_{\lambda \vdash n} \epsilon_\lambda = \frac{1}{n!} \begin{array}{c} \text{---} \\ \boxed{1_{S_n}} \\ \uparrow \end{array} \begin{array}{c} 0 \\ \text{---} \\ \downarrow \end{array},$$

where 1_{S_n} denotes the identity element of S_n . By convention, we define $\epsilon_0 = \epsilon_\emptyset = \text{id}_0$ and $\delta_0 = 0$.

Proof. — It is clear that (5.9) is satisfied. The fact that $\delta_n, \epsilon_\lambda, \lambda \vdash n$, are orthogonal idempotents follows from Lemma 5.4 and the fact that the e_λ are central. □

REMARK 5.6. — The fact that the ϵ_λ are idempotents relies on the fact that diagrams with negative regions are equal to zero. The analogous diagrams in the 2-category version \mathcal{H} of Khovanov’s category, where we allow regions labeled by any integer (see Remark 5.3), are not idempotents. In fact, it follows from [Kho14, Prop. 3] that the only idempotent 2-morphism of any 1_n , $n \in \mathbb{Z}$, is the trivial diagram. So we

see that passing to the truncation, where we kill negative regions, introduces a large number of idempotents.

LEMMA 5.7. — For all $n \in \mathbb{N}$, we have

$$\epsilon_{n+1}Q_+ = Q_+\epsilon_n, \quad \epsilon_nQ_- = Q_-\epsilon_{n+1}, \quad \delta_{n+1}Q_+ = Q_+\delta_n, \quad \delta_nQ_- = Q_-\delta_{n+1},$$

in $2\text{End}_{\mathcal{H}^{\text{tr}}}(Q_+1_n)$.

Proof. — This follows from successive applications of (5.3), followed by (5.2). \square

Recall Definition 5.1 of the idempotent completion of a 2-category. We define \mathcal{H}_ϵ to be the full sub-2-category of \mathcal{H}^{tr} whose objects are direct sums of triples $(n, 1_n, \epsilon)$ where $n \in \mathbb{N}$, and ϵ is an idempotent 2-morphism of 1_n such that $\epsilon\epsilon_n = \epsilon$. Similarly, we define \mathcal{H}_δ to be the full sub-2-category of \mathcal{H}^{tr} whose objects are direct sums of triples $(n, 1_n, \epsilon)$ where $n \in \mathbb{N}$, and ϵ is an idempotent 2-morphism of 1_n such that $\epsilon\delta_n = \epsilon$.

Recall that a 2-functor is an equivalence of 2-categories if and only if it is essentially surjective on objects, essentially full on 1-morphisms, and fully faithful on 2-morphisms.

PROPOSITION 5.8. — We have an equivalence of 2-categories

$$\mathcal{H}^{\text{tr}} \cong \mathcal{H}_\epsilon \oplus \mathcal{H}_\delta.$$

Proof. — It follows from Lemma 5.5 that, for all objects $(n, 1_n, \epsilon)$ of \mathcal{H}^{tr} , we have the decomposition

$$(n, 1_n, \epsilon) \cong (n, 1_n, \epsilon\epsilon_n) \oplus (n, 1_n, \epsilon\delta_n).$$

By Lemma 5.7, together with the fact that $\epsilon_n\delta_n = 0$ for all $n \in \mathbb{N}$, we see that any 1-morphism between an object of the form $(n, 1_n, \epsilon\epsilon_n)$ and an object of the form $(n', 1_{n'}, \epsilon'\delta_{n'})$ is isomorphic to zero. \square

PROPOSITION 5.9. — The 2-category \mathcal{H}_ϵ is isomorphic to the 2-category whose objects and 1-morphisms are the same as those of \mathcal{H}^{tr} , and whose 2-morphisms spaces are quotients of the 2-morphism spaces of \mathcal{H}^{tr} by the local relation that a clockwise circle in a region labeled n is equal to n :

$$(5.11) \quad \overset{n}{\circlearrowright} = n, \quad n \in \mathbb{N}.$$

Proof. — This follows from the fact that, in any quotient of the 2-morphism spaces,

$$\delta_n = 0 \quad \forall n \in \mathbb{N} \iff \epsilon_n = 1 \quad \forall n \in \mathbb{N} \iff (5.11) \text{ holds.}$$

Indeed, the reverse implication in the final \Leftrightarrow is clear. For the forward implication, note that $\epsilon_1 = 1$ gives (5.11) for $n = 1$. Then, supposing (5.11) holds for $n \leq k$, the equation $\epsilon_{k+1} = 1$ gives (5.11) for $n = k + 1$. \square

PROPOSITION 5.10. — *The objects*

$$(n, 1_n, \epsilon_\lambda), \quad n \in \mathbb{N}, \lambda \vdash n,$$

form a complete list of pairwise-nonisomorphic indecomposable objects of \mathcal{H}_ϵ . Furthermore, any object of \mathcal{H}_ϵ can be written uniquely (up to permutation) as a direct sum of indecomposable objects.

Proof. — Fix $n \in \mathbb{N}$. By [LRS, Prop. 4.12], the 2-endomorphism space of 1_n is spanned by

$$(5.12) \quad \begin{array}{c} n \\ \text{---} \\ \boxed{z} \\ \text{---} \\ \uparrow \end{array}, \quad z \in \mathbb{Q}S_k, \quad k \in \mathbb{N}.$$

Now, if $k > n$, then the innermost region of the above diagram is negative, and so the diagram is zero. On the other hand, if $k < n$, then by Proposition 5.9, we can insert additional clockwise circles in the center region, up to a scalar multiple. Therefore, the 2-endomorphism space of 1_n is spanned by the diagrams (5.12) for $z \in \mathbb{Q}S_n$. Now

$$\begin{array}{c} \text{---} \\ \boxed{z} \\ \text{---} \\ \uparrow \end{array} \circlearrowleft = \frac{1}{n!} \sum_{w \in S_n} \begin{array}{c} \text{---} \\ \boxed{z} \\ \text{---} \\ \uparrow \end{array} \begin{array}{c} \text{---} \\ \boxed{w} \\ \text{---} \\ \downarrow \\ \boxed{w^{-1}} \\ \text{---} \\ \uparrow \end{array} \circlearrowleft = \begin{array}{c} \text{---} \\ \boxed{z'} \\ \text{---} \\ \uparrow \end{array} \circlearrowleft,$$

where $z' = \frac{1}{n!} \sum_{w \in S_n} w^{-1}zw \in Z(\mathbb{Q}S_n)$. Therefore, the 2-endomorphism space of 1_n is spanned by the diagrams (5.12) for z in the center $Z(\mathbb{Q}S_n)$ of $\mathbb{Q}S_n$. But then this 2-endomorphism space is spanned by the diagrams (5.12) as z ranges over the central idempotents e_λ , $\lambda \vdash n$. In other words, this space is spanned by the ϵ_λ , $\lambda \vdash n$. It follows from Lemma 7.1 below that these elements are also linearly independent. \square

5.3. REGION SHIFTING. — We define a shift 2-functor

$$\partial: \mathcal{H}^{\text{tr}'} \longrightarrow \mathcal{H}^{\text{tr}'},$$

given by lowering region labels by one. More precisely, on objects we define

$$\partial(0) = \partial(\mathbf{0}) = \mathbf{0}, \quad \partial(n) = n - 1, \quad n > 1.$$

On 1-morphisms, we define

$$\partial(\mathbb{Q}_c 1_n) = \mathbb{Q}_c 1_{n-1},$$

for $n \in \mathbb{N}$ and c a (possibly empty) sequence of $+$ and $-$. On 2-morphisms, ∂ is given by lowering the region labels of diagrams by one. The 2-functor ∂ induces a 2-functor

$$\partial: \mathcal{H}^{\text{tr}} \longrightarrow \mathcal{H}^{\text{tr}}$$

on the idempotent completion $\mathcal{H}^{\text{tr}} = \text{Kar}(\mathcal{H}^{\text{tr}'})$.

PROPOSITION 5.11. — *The 2-functor ∂ sends \mathcal{H}_ϵ to zero. Furthermore, the restriction of ∂ to \mathcal{H}_δ induces an equivalence of 2-categories $\mathcal{H}_\delta \cong \mathcal{H}^{\text{tr}}$.*

Proof. — Since ∂ maps ϵ_λ to zero for all partitions λ , the first statement follows. Now let $\partial_\delta: \mathcal{H}_\delta \rightarrow \mathcal{H}^{\text{tr}}$ be the restriction of the 2-functor ∂ . We define a 2-functor $s': \mathcal{H}^{\text{tr}'} \rightarrow \mathcal{H}_\delta$. On objects, we define

$$s'(\mathbf{0}) = \mathbf{0}, \quad s'(n) = (n + 1, 1_{n+1}, \delta_{n+1}), \quad n \in \mathbb{N}.$$

On 1-morphisms, we define

$$s'(\mathbb{Q}_c 1_n) = (\mathbb{Q}_c 1_{n+1}, \mathbb{Q}_c \delta_{n+1}).$$

On 2-morphisms, we define s' by increasing the region labels in diagrams by one. Note that the definition of s' is compatible with our convention that diagrams with negative region labels are zero since Lemma 5.7, together with the fact that $\delta_0 = 0$, ensures that s' maps any diagram with a negative region label to zero. The 2-functor s' induces a 2-functor $s: \mathcal{H}^{\text{tr}} \rightarrow \mathcal{H}_\delta$.

It is clear that $\partial_\delta \circ s$ is isomorphic to the identity 2-functor since $\partial(\delta_{n+1}) = \text{id}_n$ for all $n \in \mathbb{N}$. Similarly, $s \circ \partial_\delta$ is isomorphic to the identity 2-functor since the object $(0, 1_0, \delta_0)$ is already isomorphic to zero in \mathcal{H}_δ . \square

COROLLARY 5.12. — *We have an equivalence of 2-categories $\mathcal{H}^{\text{tr}} \cong \bigoplus_{m=0}^\infty \mathcal{H}_\epsilon$.*

Proof. — It is clear that, for any 2-morphism θ in \mathcal{H}^{tr} , we have $\partial^m(\theta) = 0$ for sufficiently large m . Thus, the result follows from Propositions 5.8 and 5.11. \square

6. EQUIVALENCE OF \mathcal{H}_ϵ AND \mathcal{A}

6.1. A 2-FUNCTOR FROM \mathcal{A} TO \mathcal{H}_ϵ

LEMMA 6.1. — *For $\mu \vdash n + 1$, we have the following equality of 2-morphisms in \mathcal{H}^{tr} :*

$$(6.1) \quad \epsilon_\mu \left| \begin{array}{c} \uparrow \\ n \end{array} \right. = \frac{1}{n!} \left| \begin{array}{c} \uparrow \\ n \\ \boxed{e_\mu} \\ \uparrow \uparrow \\ 0 \end{array} \right. .$$

Proof. — We have

$$\epsilon_\mu \left| \begin{array}{c} \uparrow \\ n \end{array} \right. = \frac{1}{(n+1)!} \left| \begin{array}{c} \uparrow \\ n \\ \boxed{e_\mu} \\ \uparrow \\ 0 \end{array} \right. = \frac{1}{(n+1)!} \sum_{\ell=0}^n \left| \begin{array}{c} \uparrow \\ n \\ \boxed{e_\mu} \\ \uparrow \uparrow \\ 0 \end{array} \right. = \frac{1}{n!} \left| \begin{array}{c} \uparrow \\ n \\ \boxed{e_\mu} \\ \uparrow \uparrow \\ 0 \end{array} \right. ,$$

where the second equality follows from repeated use of (5.3) together with the fact that diagrams with negative region labels are equal to zero, and the last equality follows from the fact that the idempotent e_μ is central (so we can slide crossings through the box labeled e_μ). \square

COROLLARY 6.2. — *For $\lambda \vdash n$ and $\mu \vdash n + 1$, we have*

$$\epsilon_\mu \mathbb{Q}_+ \epsilon_\lambda = 0 \quad \text{unless } \lambda \subseteq \mu.$$

Proof. — We have

$$(6.2) \quad \epsilon_\mu \uparrow \epsilon_\lambda \stackrel{(6.1)}{=} \frac{1}{n!(n-1)!} \left[\begin{array}{c} \uparrow^n \\ \boxed{e_\mu} \quad \boxed{e_\lambda} \\ \uparrow \end{array} \right] = \frac{1}{n!} \left[\begin{array}{c} \uparrow^n \\ \boxed{e_\mu} \quad \boxed{e_\lambda} \\ \uparrow \end{array} \right] = \frac{1}{n!} \left[\begin{array}{c} \uparrow^n \\ \boxed{e_\mu e_\lambda} \\ \uparrow \end{array} \right],$$

where the second equality follows from repeated use of (5.3), the fact that diagrams with negative region labels are equal to zero, and the fact that the idempotent e_λ is central (as in the proof of Lemma 6.1). The result then follows from the fact that $e_\mu e_\lambda = 0$ unless $\lambda \subseteq \mu$. \square

LEMMA 6.3. — For $\lambda \vdash n$, we have

$$(6.3) \quad \left(\begin{array}{c} \epsilon_\lambda \\ \circlearrowright \end{array} \right) = \sum_{i \in \mathbb{Z}} \frac{(n+1)d_\lambda}{d_{\lambda \boxplus i}} \epsilon_{\lambda \boxplus i}, \quad (6.4) \quad \left(\begin{array}{c} \epsilon_\lambda \\ \circlearrowleft \end{array} \right) = \sum_{i \in \mathbb{Z}} \frac{d_\lambda}{nd_{\lambda \boxminus i}} \epsilon_{\lambda \boxminus i}.$$

Proof. — Suppose $\lambda \vdash n$. To prove (6.3), we compute

$$\begin{aligned} \left(\begin{array}{c} \epsilon_\lambda \\ \circlearrowright \end{array} \right) &= \sum_{i \in \mathbb{Z}} \epsilon_{\lambda \boxplus i} \left(\begin{array}{c} \epsilon_\lambda \\ \circlearrowright \end{array} \right) \stackrel{(6.2)}{=} \sum_{i \in \mathbb{Z}} \frac{1}{n!} \left[\begin{array}{c} \uparrow^n \\ \boxed{e_{\lambda \boxplus i} e_\lambda} \\ \uparrow \end{array} \right] = \frac{1}{n!(n+1)!} \sum_{w \in S_{n+1}} \left[\begin{array}{c} \uparrow^n \\ \boxed{e_{\lambda \boxplus i} e_\lambda} \quad \begin{array}{c} \boxed{w} \\ \downarrow \\ \boxed{w^{-1}} \end{array} \\ \uparrow \end{array} \right] \\ &\stackrel{(3.31)}{=} \sum_{i \in \mathbb{Z}} \frac{d_\lambda}{n!d_{\lambda \boxplus i}} \left[\begin{array}{c} \uparrow^n \\ \boxed{e_{\lambda \boxplus i}} \\ \uparrow \end{array} \right] = \sum_{i \in \mathbb{Z}} \frac{(n+1)d_\lambda}{d_{\lambda \boxplus i}} \epsilon_{\lambda \boxplus i}. \end{aligned}$$

To prove (6.4), we compute

$$\left(\begin{array}{c} \epsilon_\lambda \\ \circlearrowleft \end{array} \right) \stackrel{(6.1)}{=} \frac{1}{(n-1)!} \left[\begin{array}{c} \uparrow^n \\ \boxed{e_\lambda} \quad 0 \\ \uparrow \end{array} \right] \stackrel{(5.6)}{=} \frac{d_\lambda}{n!d_{\lambda \boxminus i}} \left[\begin{array}{c} \uparrow^n \\ \boxed{e_{\lambda \boxminus i}} \quad 0 \\ \uparrow \end{array} \right] = \frac{d_\lambda}{nd_{\lambda \boxminus i}} \epsilon_{\lambda \boxminus i}. \quad \square$$

We will use an open circle to denote a right curl in \mathcal{H}^{tr} :

$$(6.5) \quad \left(\begin{array}{c} \uparrow \\ \circlearrowright \end{array} \right) := \left(\begin{array}{c} \uparrow \\ \circlearrowright \end{array} \right).$$

LEMMA 6.4. — For a partition λ and $i \in \mathbb{Z}$, we have

$$(6.6) \quad \epsilon_{\lambda \boxplus i} \uparrow \left(\begin{array}{c} \uparrow \\ \circlearrowright \end{array} \right) \epsilon_\lambda = i \left(\begin{array}{c} \epsilon_{\lambda \boxplus i} \uparrow \epsilon_\lambda \end{array} \right).$$

Proof. — For $\lambda \vdash n + 1$, we have

$$\begin{array}{c} \epsilon_{\lambda \boxplus i} \uparrow \\ \epsilon_{\lambda} \uparrow \\ \circ \end{array} \stackrel{(6.2)}{=} \frac{1}{n!} \begin{array}{c} \epsilon_{\lambda \boxplus i} \uparrow \\ \circ \\ \boxed{e_{\lambda \boxplus i} e_{\lambda}} \\ \uparrow \end{array} \stackrel{(3.30)}{=} \frac{1}{n!} \begin{array}{c} \uparrow^n \\ \boxed{J_{n+1}} \\ \uparrow \\ \boxed{e_{\lambda \boxplus i} e_{\lambda}} \\ \uparrow \end{array} \stackrel{(6.2)}{=} \frac{i}{n!} \begin{array}{c} \uparrow \\ \boxed{e_{\lambda \boxplus i} e_{\lambda}} \\ \uparrow \end{array} \stackrel{(6.2)}{=} i \left(\begin{array}{c} \epsilon_{\lambda \boxplus i} \uparrow \\ \epsilon_{\lambda} \uparrow \end{array} \right),$$

where the second equality follows from repeated use of (5.3) together with the fact that diagrams with negative region labels are equal to zero. \square

LEMMA 6.5. — For a partition λ and $i, j \in \mathbb{Z}$, $i \neq j$, we have

$$(6.7) \quad \begin{array}{c} \epsilon_{\lambda \boxplus i} \nearrow \\ \epsilon_{\lambda \boxplus i \boxplus j} \times \epsilon_{\lambda} \\ \epsilon_{\lambda \boxplus i} \searrow \end{array} = \frac{1}{j - i} \left(\begin{array}{c} \epsilon_{\lambda \boxplus i \boxplus j} \uparrow \\ \epsilon_{\lambda \boxplus i} \uparrow \\ \epsilon_{\lambda} \uparrow \end{array} \right).$$

Proof. — We have

$$(j - i) \left(\begin{array}{c} \epsilon_{\lambda \boxplus i} \nearrow \\ \epsilon_{\lambda \boxplus i \boxplus j} \times \epsilon_{\lambda} \\ \epsilon_{\lambda \boxplus i} \searrow \end{array} \right) \stackrel{(6.6)}{=} \begin{array}{c} \epsilon_{\lambda \boxplus i} \nearrow \\ \circ \\ \epsilon_{\lambda \boxplus i \boxplus j} \times \epsilon_{\lambda} \\ \epsilon_{\lambda \boxplus i} \searrow \end{array} - \begin{array}{c} \epsilon_{\lambda \boxplus i} \nearrow \\ \epsilon_{\lambda \boxplus i \boxplus j} \times \epsilon_{\lambda} \\ \circ \\ \epsilon_{\lambda \boxplus i} \searrow \end{array} = \begin{array}{c} \epsilon_{\lambda \boxplus i \boxplus j} \uparrow \\ \epsilon_{\lambda \boxplus i} \uparrow \\ \epsilon_{\lambda} \uparrow \end{array},$$

where the last equality follows from the relation in [Kho14, §2.1] concerning sliding right curls through crossings. \square

We now define an additive linear 2-functor

$$\mathbf{S}: \mathcal{A} \rightarrow \mathcal{H}_{\epsilon}$$

as follows. On objects, we define

$$\mathbf{S}(\mathbf{0}) = \mathbf{0}, \quad \mathbf{S}(\lambda) = (n, 1_n, \epsilon_{\lambda}) \quad \text{for } \lambda \vdash n.$$

On 1-morphisms, \mathbf{S} is determined by

$$\mathbf{S}(E_i 1_{\lambda}) = (\mathbf{Q} - 1_n, \epsilon_{\lambda \boxplus i} \mathbf{Q} - \epsilon_{\lambda}), \quad \mathbf{S}(F_i 1_{\lambda}) = (\mathbf{Q} + 1_n, \epsilon_{\lambda \boxplus i} \mathbf{Q} + \epsilon_{\lambda}), \quad \text{for } \lambda \vdash n,$$

where, by convention, we set $\epsilon_{\mathbf{0}} = 0$. On 2-morphisms, \mathbf{S} is determined as follows. Suppose θ is a diagram representing a 2-morphism in \mathcal{A} . Then $\mathbf{S}(\theta)$ is the diagram obtained from θ by placing a ϵ_{λ} in each region labeled λ and then acting as follows on crossings, cups, and caps:

$$(6.8) \quad \mathbf{S} \left(\begin{array}{c} \nearrow^{\lambda} \\ i \times j \\ \searrow \end{array} \right) = \xi_{i,j} \begin{array}{c} \epsilon_{\lambda \boxplus i} \nearrow \\ \epsilon_{\lambda \boxplus i \boxplus j} \times \epsilon_{\lambda} \\ \epsilon_{\lambda \boxplus j} \searrow \end{array}, \quad (6.9) \quad \mathbf{S} \left(\begin{array}{c} \searrow^{\lambda} \\ i \times j \\ \nearrow \end{array} \right) = \xi_{i,j} \begin{array}{c} \epsilon_{\lambda \boxplus i} \searrow \\ \epsilon_{\lambda \boxplus i \boxplus j} \times \epsilon_{\lambda} \\ \epsilon_{\lambda \boxplus j} \nearrow \end{array},$$

$$(6.10) \quad \mathbf{S} \left(\begin{array}{c} i \times j \\ \searrow^{\lambda} \end{array} \right) = \xi_{i,j} \frac{(|\lambda| + 1) d_{\lambda} d_{\lambda \boxplus i \boxplus j}}{|\lambda| d_{\lambda \boxplus i} d_{\lambda \boxplus j}} \begin{array}{c} \epsilon_{\lambda \boxplus j} \searrow \\ \epsilon_{\lambda \boxplus i \boxplus j} \times \epsilon_{\lambda} \\ \epsilon_{\lambda \boxplus i} \searrow \end{array},$$

$$(6.11) \quad \mathbf{S} \left(\begin{array}{c} i \\ \times \\ j \end{array} \lambda \right) = \xi_{i,j} \begin{array}{c} \epsilon_{\lambda \boxplus j} \\ \epsilon_{\lambda \boxplus i \boxplus j} \\ \epsilon_{\lambda} \\ \epsilon_{\lambda \boxplus i} \end{array} ,$$

$$(6.12) \quad \mathbf{S} \left(\begin{array}{c} \curvearrowright \\ i \end{array} \lambda \right) = \frac{d_\lambda}{|\lambda| d_{\lambda \boxplus i}} \begin{array}{c} \epsilon_{\lambda \boxplus i} \\ \epsilon_{\lambda} \end{array} , \quad (6.13) \quad \mathbf{S} \left(\begin{array}{c} i \\ \curvearrowleft \\ \lambda \end{array} \right) = \frac{(|\lambda| + 1) d_\lambda}{d_{\lambda \boxplus i}} \begin{array}{c} \epsilon_{\lambda \boxplus i} \\ \epsilon_{\lambda} \end{array} ,$$

$$(6.14) \quad \mathbf{S} \left(\begin{array}{c} \curvearrowright \\ i \end{array} \lambda \right) = \begin{array}{c} \epsilon_{\lambda \boxplus i} \\ \epsilon_{\lambda} \end{array} , \quad (6.15) \quad \mathbf{S} \left(\begin{array}{c} i \\ \curvearrowleft \\ \lambda \end{array} \right) = \begin{array}{c} \epsilon_{\lambda \boxplus i} \\ \epsilon_{\lambda} \end{array} ,$$

where

$$(6.16) \quad \xi_{i,j} = \frac{i - j}{i - j - 1}.$$

Note that

$$(6.17) \quad \frac{1}{\xi_{i,j} \xi_{j,i}} = 1 - \frac{1}{(i - j)^2}.$$

PROPOSITION 6.6. — *The map \mathbf{S} described above is a well-defined 2-functor.*

Proof. — We first verify that \mathbf{S} respects isotopy invariance. It is straightforward to verify that

$$\mathbf{S}(\cap) = \mathbf{S}(\cup) \quad \text{and} \quad \mathbf{S}(\cup) = \mathbf{S}(\cap)$$

with all possible orientations of the strands and all possible labelings of the regions and strands. We also have

$$\begin{aligned} \mathbf{S} \left(\begin{array}{c} j \ i \\ \curvearrowright \\ \lambda \end{array} \right) &= \xi_{i,j} \frac{d_\lambda}{n d_{\lambda \boxplus j}} \frac{d_{\lambda \boxplus j}}{(n - 1) d_{\lambda \boxplus j \boxplus i}} \frac{n d_{\lambda \boxplus i} (n - 1) d_{\lambda \boxplus i \boxplus j}}{d_\lambda d_{\lambda \boxplus i}} \begin{array}{c} \epsilon_{\lambda \boxplus j} \\ \epsilon_{\lambda \boxplus i \boxplus j} \\ \epsilon_{\lambda} \\ \epsilon_{\lambda \boxplus i} \end{array} \\ &= \mathbf{S} \left(\begin{array}{c} \times \\ i \ j \end{array} \lambda \right). \end{aligned}$$

Similarly, one easily verifies that

$$\begin{aligned} \mathbf{S} \left(\begin{array}{c} j \ i \\ \curvearrowleft \\ \lambda \end{array} \right) &= \mathbf{S} \left(\begin{array}{c} \times \\ i \ j \end{array} \lambda \right), \\ \mathbf{S} \left(\begin{array}{c} \curvearrowright \\ j \ i \end{array} \lambda \right) &= \mathbf{S} \left(\begin{array}{c} \curvearrowright \\ j \ i \end{array} \lambda \right) = \mathbf{S} \left(\begin{array}{c} \times \\ j \ i \end{array} \lambda \right), \\ \mathbf{S} \left(\begin{array}{c} \curvearrowleft \\ j \ i \end{array} \lambda \right) &= \mathbf{S} \left(\begin{array}{c} \curvearrowleft \\ j \ i \end{array} \lambda \right) = \mathbf{S} \left(\begin{array}{c} \times \\ j \ i \end{array} \lambda \right). \end{aligned}$$

It follows that \mathbf{S} respects isotopy invariance. It remains to check that \mathbf{S} respects the local relations (4.3)–(4.10).

Relation (4.3). — We have

$$\begin{aligned}
 \frac{1}{\xi_{i,j}\xi_{j,k}\xi_{i,k}} \mathbf{S} \left(\begin{array}{c} \text{Diagram with crossings } \lambda \\ i \quad j \quad k \end{array} \right) &= \begin{array}{c} \text{Diagram with crossings } \lambda \text{ and } \epsilon_\lambda \\ \epsilon_{\lambda \boxplus i \boxplus j \boxplus k} \quad \epsilon_{\lambda \boxplus i} \\ \epsilon_{\lambda \boxplus k \boxplus j} \quad \epsilon_{\lambda \boxplus k} \\ \epsilon_{\lambda \boxplus k \boxplus j} \end{array} = \begin{array}{c} \text{Diagram with crossings } \lambda \text{ and } \epsilon_\lambda \\ \epsilon_{\lambda \boxplus i \boxplus j \boxplus k} \quad \epsilon_{\lambda \boxplus i} \\ \epsilon_{\lambda \boxplus k \boxplus j} \quad \epsilon_{\lambda \boxplus k} \\ \epsilon_{\lambda \boxplus k \boxplus j} \end{array} \\
 \stackrel{(5.1)}{=} \begin{array}{c} \text{Diagram with crossings } \lambda \text{ and } \epsilon_\lambda \\ \epsilon_{\lambda \boxplus i \boxplus j \boxplus k} \quad \epsilon_{\lambda \boxplus i} \\ \epsilon_{\lambda \boxplus k \boxplus j} \quad \epsilon_{\lambda \boxplus k} \\ \epsilon_{\lambda \boxplus k \boxplus j} \end{array} &= \begin{array}{c} \text{Diagram with crossings } \lambda \text{ and } \epsilon_\lambda \\ \epsilon_{\lambda \boxplus i \boxplus j \boxplus k} \quad \epsilon_{\lambda \boxplus i} \\ \epsilon_{\lambda \boxplus k \boxplus j} \quad \epsilon_{\lambda \boxplus k} \\ \epsilon_{\lambda \boxplus k \boxplus j} \end{array} = \frac{1}{\xi_{i,j}\xi_{j,k}\xi_{i,k}} \mathbf{S} \left(\begin{array}{c} \text{Diagram with crossings } \lambda \\ i \quad j \quad k \end{array} \right),
 \end{aligned}$$

where the second and fourth equalities follow from (5.9) and Corollary 6.2.

Relation (4.4). — For $\lambda \vdash n$, we have

$$\begin{aligned}
 \frac{1}{\xi_{i,j}\xi_{j,i}} \mathbf{S} \left(\begin{array}{c} \text{Diagram with crossings } \lambda \\ i \quad j \end{array} \right) &= \begin{array}{c} \text{Diagram with crossings } \lambda \text{ and } \epsilon_\lambda \\ \epsilon_{\lambda \boxplus i \boxplus j} \quad \epsilon_\lambda \\ \epsilon_{\lambda \boxplus i} \quad \epsilon_{\lambda \boxplus j} \end{array} = \begin{array}{c} \text{Diagram with crossings } \lambda \text{ and } \epsilon_\lambda \\ \epsilon_{\lambda \boxplus i \boxplus j} \quad \epsilon_\lambda \\ \epsilon_{\lambda \boxplus i} \quad \epsilon_{\lambda \boxplus j} \end{array} - \begin{array}{c} \text{Diagram with crossings } \lambda \text{ and } \epsilon_\lambda \\ \epsilon_{\lambda \boxplus i \boxplus j} \quad \epsilon_\lambda \\ \epsilon_{\lambda \boxplus i} \quad \epsilon_{\lambda \boxplus j} \end{array} \\
 \stackrel{(5.2)}{=} \left(1 - \frac{1}{(i-j)^2} \right) \begin{array}{c} \uparrow \quad \uparrow \\ \epsilon_{\lambda \boxplus i \boxplus j} \quad \epsilon_{\lambda \boxplus j} \\ \downarrow \quad \downarrow \\ \epsilon_\lambda \end{array} &\stackrel{(6.17)}{=} \frac{1}{\xi_{i,j}\xi_{j,i}} \mathbf{S} \left(\begin{array}{c} \text{Diagram with crossings } \lambda \\ i \quad j \end{array} \right),
 \end{aligned}$$

where the second equality follows from (5.9) and Corollary 6.2.

Relation (4.5). — Let $\lambda \vdash n$ and $i, j \in \mathbb{Z}, i \neq j$. By (3.25) and (6.17), we have

$$\frac{1}{\xi_{i,j}\xi_{j,i}} \frac{nd_{\lambda \boxplus i} d_{\lambda \boxplus j}}{(n+1)d_\lambda d_{\lambda \boxplus i \boxplus j}} = 1.$$

Thus

$$\begin{aligned}
 \mathbf{S} \left(\begin{array}{c} \text{Diagram with crossings } \lambda \\ i \quad j \end{array} \right) &= \begin{array}{c} \text{Diagram with crossings } \lambda \text{ and } \epsilon_\lambda \\ \epsilon_{\lambda \boxplus i \boxplus j} \quad \epsilon_\lambda \\ \epsilon_{\lambda \boxplus i} \quad \epsilon_{\lambda \boxplus j} \end{array} = \begin{array}{c} \text{Diagram with crossings } \lambda \text{ and } \epsilon_\lambda \\ \epsilon_{\lambda \boxplus i \boxplus j} \quad \epsilon_\lambda \\ \epsilon_{\lambda \boxplus i} \quad \epsilon_{\lambda \boxplus j} \end{array} \\
 \stackrel{(5.3)}{=} \begin{array}{c} \uparrow \quad \uparrow \\ \epsilon_{\lambda \boxplus i \boxplus j} \quad \epsilon_{\lambda \boxplus j} \\ \downarrow \quad \downarrow \\ \epsilon_\lambda \end{array} - \begin{array}{c} \text{Diagram with crossings } \lambda \text{ and } \epsilon_\lambda \\ \epsilon_{\lambda \boxplus i \boxplus j} \quad \epsilon_\lambda \\ \epsilon_{\lambda \boxplus i} \quad \epsilon_{\lambda \boxplus j} \end{array} &= \begin{array}{c} \uparrow \quad \uparrow \\ \epsilon_{\lambda \boxplus i \boxplus j} \quad \epsilon_{\lambda \boxplus j} \\ \downarrow \quad \downarrow \\ \epsilon_\lambda \end{array} = \mathbf{S} \left(\begin{array}{c} \text{Diagram with crossings } \lambda \\ i \quad j \end{array} \right),
 \end{aligned}$$

where the second equality follows from (5.9) and Corollary 6.2, since the only partition μ of size $n - 1$ satisfying $\mu \subseteq \lambda \boxplus i \boxplus j$ and $\mu \subseteq \lambda$ is $\mu = \lambda \boxplus i$.

Relation (4.6). — Let $\lambda \vdash n$ and $i, j \in \mathbb{Z}$, $i \neq j$. Again, using (3.25) and (6.17), we have

$$\mathbf{S} \left(\begin{array}{c} \lambda \\ \swarrow \quad \searrow \\ i \quad j \end{array} \right) = \begin{array}{c} \epsilon_{\lambda \boxplus i \boxplus j} \\ \swarrow \quad \searrow \\ \epsilon_{\lambda \boxplus i} \\ \swarrow \quad \searrow \\ \epsilon_{\lambda \boxplus j} \end{array} = \begin{array}{c} \epsilon_{\lambda \boxplus i \boxplus j} \\ \swarrow \quad \searrow \\ \epsilon_{\lambda} \\ \swarrow \quad \searrow \\ \epsilon_{\lambda \boxplus j} \end{array} \stackrel{(5.4)}{=} \begin{array}{c} \uparrow \\ \epsilon_{\lambda \boxplus i \boxplus j} \\ \downarrow \\ \epsilon_{\lambda \boxplus j} \\ \epsilon_{\lambda} \end{array} = \mathbf{S} \left(\begin{array}{c} \uparrow \\ \downarrow \\ i \quad j \end{array} \right),$$

where the second equality follows from (5.9) and Corollary 6.2, since the only partition μ of size $n + 1$ satisfying $\lambda \boxplus i \boxplus j \subseteq \mu$ and $\lambda \subseteq \mu$ is $\mu = \lambda \boxplus i$.

Relation (4.7). — For $\lambda \vdash n$ and $i \in \mathbb{Z}$, we have

$$\begin{aligned} \mathbf{S} \left(\begin{array}{c} \uparrow \\ \downarrow \\ i \quad i \end{array} \right) &= \begin{array}{c} \epsilon_{\lambda} \\ \uparrow \\ \epsilon_{\lambda \boxplus i} \\ \downarrow \\ \epsilon_{\lambda} \end{array} \stackrel{(5.3)}{=} \begin{array}{c} \epsilon_{\lambda \boxplus i} \\ \uparrow \\ \epsilon_{\lambda} \\ \downarrow \\ \epsilon_{\lambda \boxplus i} \end{array} + \begin{array}{c} \epsilon_{\lambda \boxplus i} \\ \swarrow \quad \searrow \\ \epsilon_{\lambda} \\ \swarrow \quad \searrow \\ \epsilon_{\lambda \boxplus i} \end{array} = \begin{array}{c} \epsilon_{\lambda \boxplus i} \\ \uparrow \\ \epsilon_{\lambda} \\ \downarrow \\ \epsilon_{\lambda \boxplus i} \end{array} + \sum_{j \in \mathbb{Z}} \begin{array}{c} \epsilon_{\lambda \boxplus i} \\ \swarrow \quad \searrow \\ \epsilon_{\lambda \boxplus j} \\ \swarrow \quad \searrow \\ \epsilon_{\lambda \boxplus i} \end{array} \\ &\stackrel{(6.7)}{=} \begin{array}{c} \epsilon_{\lambda \boxplus i} \\ \uparrow \\ \epsilon_{\lambda} \\ \downarrow \\ \epsilon_{\lambda \boxplus i} \end{array} \left(1 + \sum_{j \in \mathbb{Z}} \frac{1}{(i-j)^2} \begin{array}{c} \epsilon_{\lambda \boxplus j} \end{array} \right) \stackrel{(6.3)}{=} \begin{array}{c} \epsilon_{\lambda \boxplus i} \\ \uparrow \\ \epsilon_{\lambda} \\ \downarrow \\ \epsilon_{\lambda \boxplus i} \end{array} \left(1 + \sum_{j \in \mathbb{Z}} \frac{nd_{\lambda \boxplus j}}{d_{\lambda}(i-j)^2} \right) \\ &\stackrel{(3.27)}{=} \frac{(n+1)d_{\lambda}}{d_{\lambda \boxplus i}} \begin{array}{c} \epsilon_{\lambda \boxplus i} \\ \uparrow \\ \epsilon_{\lambda} \\ \downarrow \\ \epsilon_{\lambda \boxplus i} \end{array} = \mathbf{S} \left(\begin{array}{c} i \\ \uparrow \quad \downarrow \\ \lambda \end{array} \right). \end{aligned}$$

Relation (4.8). — For $\lambda \vdash n$ and $i \in \mathbb{Z}$, we have

$$\begin{aligned} \mathbf{S} \left(\begin{array}{c} \uparrow \\ \downarrow \\ i \quad i \end{array} \right) &= \begin{array}{c} \epsilon_{\lambda} \\ \uparrow \\ \epsilon_{\lambda \boxplus i} \\ \downarrow \\ \epsilon_{\lambda} \end{array} \stackrel{(5.4)}{=} \begin{array}{c} \epsilon_{\lambda \boxplus i} \\ \swarrow \quad \searrow \\ \epsilon_{\lambda} \\ \swarrow \quad \searrow \\ \epsilon_{\lambda \boxplus i} \end{array} = \sum_{j \in \mathbb{Z}} \begin{array}{c} \epsilon_{\lambda \boxplus i} \\ \swarrow \quad \searrow \\ \epsilon_{\lambda \boxplus j} \\ \swarrow \quad \searrow \\ \epsilon_{\lambda \boxplus i} \end{array} \stackrel{(6.7)}{=} \sum_{j \in \mathbb{Z}} \frac{1}{(j-i)^2} \begin{array}{c} \epsilon_{\lambda \boxplus i} \\ \uparrow \\ \epsilon_{\lambda} \\ \downarrow \\ \epsilon_{\lambda \boxplus i} \end{array} \begin{array}{c} \epsilon_{\lambda \boxplus j} \end{array} \\ &\stackrel{(6.4)}{=} \sum_{j \in \mathbb{Z}} \frac{d_{\lambda \boxplus j}}{(n+1)d_{\lambda}(j-i)^2} \begin{array}{c} \epsilon_{\lambda \boxplus i} \\ \uparrow \\ \epsilon_{\lambda} \\ \downarrow \\ \epsilon_{\lambda \boxplus i} \end{array} \stackrel{(3.26)}{=} \frac{d_{\lambda}}{nd_{\lambda \boxplus i}} \begin{array}{c} \epsilon_{\lambda \boxplus i} \\ \uparrow \\ \epsilon_{\lambda} \\ \downarrow \\ \epsilon_{\lambda \boxplus i} \end{array} = \mathbf{S} \left(\begin{array}{c} i \\ \uparrow \quad \downarrow \\ \lambda \end{array} \right). \end{aligned}$$

Relation (4.9). — Suppose $\lambda \vdash n$. If $i \notin B^-(\lambda)$, then it clear that \mathbf{S} maps the left-hand side of (4.9) to zero since $\epsilon_{\lambda \boxplus i} = \epsilon_{\mathbf{0}} = 0$. If $i \in B^-(\lambda)$, we have

$$\mathbf{S} \left(i \circlearrowleft \lambda \right) = \frac{d_\lambda}{nd_{\lambda \boxplus i}} \left(\epsilon_{\lambda \boxplus i} \right)^{\epsilon_\lambda} \stackrel{(6.3)}{=} \epsilon_\lambda.$$

Relation (4.10). — Suppose $\lambda \vdash n$. If $i \notin B^+(\lambda)$, then it clear that \mathbf{S} maps the left-hand side of (4.10) to zero since $\epsilon_{\lambda \boxplus i} = \epsilon_{\mathbf{0}} = 0$. If $i \in B^+(\lambda)$, we have

$$\mathbf{S} \left(i \circlearrowright \lambda \right) = \frac{(n+1)d_\lambda}{d_{\lambda \boxplus i}} \left(\epsilon_{\lambda \boxplus i} \right)^{\epsilon_\lambda} \stackrel{(6.4)}{=} \epsilon_\lambda. \quad \square$$

The following theorem is one of the main results of the current paper.

THEOREM 6.7. — *The 2-functor $\mathbf{S}: \mathcal{A} \rightarrow \mathcal{H}_\epsilon$ is an equivalence of 2-categories.*

Proof. — The 2-functor \mathbf{S} is essentially surjective on objects by Proposition 5.10. By Corollary 6.2, it is also essentially full on 1-morphisms by and full on 2-morphisms. We will show in Corollary 7.4 that it is also faithful on 2-morphisms. \square

COROLLARY 6.8. — *We have an equivalence of categories $K(\mathcal{H}^{\text{tr}}) \cong \bigoplus_{m=0}^\infty \mathcal{A}$.*

Proof. — This follows by combining Theorem 6.7 with Corollary 5.12 and Theorem 4.11. \square

REMARK 6.9. — Corollary 6.8 can be viewed as an analogue of Khovanov’s Heisenberg categorification conjecture [Kho14, Conj. 1]. In the framework of 2-categories, Khovanov’s conjecture is that Grothendieck group of \mathcal{H} is isomorphic to $\bigoplus_{m \in \mathbb{Z}} H$.

6.2. A 2-FUNCTOR FROM \mathcal{H}^{tr} TO \mathcal{A} . — It follows from Theorem 6.7 that we have an equivalence $\mathbf{T}: \mathcal{H}_\epsilon \rightarrow \mathcal{A}$ of 2-categories, obtained by inverting \mathbf{S} . The domain of this 2-functor can be extended by zero to $\mathcal{H}^{\text{tr}} \cong \mathcal{H}_\epsilon \oplus \mathcal{H}_\delta$. Since \mathcal{A} is idempotent complete, it follows from the universal property of the idempotent completion that this 2-functor is uniquely determined by its restriction $\mathbf{T}: \mathcal{H}^{\text{tr}'} \rightarrow \mathcal{A}$ (which we continue to denote by \mathbf{T}). For future reference, we describe this 2-functor explicitly.

The additive linear 2-functor $\mathbf{T}: \mathcal{H}^{\text{tr}'} \rightarrow \mathcal{A}$ is determined on objects by

$$\mathbf{0} \mapsto \mathbf{0}, \quad n \mapsto \bigoplus_{\lambda \vdash n} \lambda, \quad n \in \mathbb{N},$$

and on 1-morphisms by

$$Q_+ 1_n \mapsto \sum_{\lambda \vdash n} \sum_{i \in \mathbb{Z}} F_i 1_\lambda = \sum_{\lambda \vdash n} \sum_{i \in B^+(\lambda)} F_i 1_\lambda, \quad Q_- 1_n \mapsto \sum_{\lambda \vdash n} \sum_{i \in \mathbb{Z}} E_i 1_\lambda = \sum_{\lambda \vdash n} \sum_{i \in B^-(\lambda)} E_i 1_\lambda.$$

On 2-morphisms, \mathbf{T} is given as follows:

$$(6.18) \quad \mathbf{T} \left(\begin{array}{c} \times \\ \nearrow \quad \searrow \\ i \quad j \\ n \end{array} \right) = \sum_{\substack{\lambda \vdash n \\ i, j \in \mathbb{Z}}} \left(\frac{i-j-1}{i-j} \begin{array}{c} \times \\ \nearrow \quad \searrow \\ i \quad j \\ \lambda \end{array} + \frac{1}{i-j} \begin{array}{c} \uparrow \quad \uparrow \\ i \quad j \\ \lambda \end{array} \right),$$

diagram of 2-functors:

$$(7.1) \quad \begin{array}{ccccccc} & & & & \mathcal{U} & & \\ & & & & \downarrow & \searrow \mathbf{A} & \\ \mathcal{H} & \xrightarrow{(5.7)} & \mathcal{H}^{\text{tr}} & \xrightarrow{\mathbf{T}} & \mathcal{A} & \longrightarrow & \mathcal{U}^{\text{tr}} \longrightarrow \mathcal{C} \end{array}$$

Thus, we obtain a 2-functor $\mathcal{H} \rightarrow \mathcal{C}$. That is, we have an action of the 2-category version of Khovanov’s monoidal category in \mathcal{C} . This is a categorical restriction of the basic representation to the principal Heisenberg subalgebra.

Conversely, suppose $\mathbf{B}: \mathcal{H} \rightarrow \mathcal{C}$ is an additive linear 2-functor for some additive linear 2-category \mathcal{C} mapping all objects $n \in \mathbb{Z}, n < 0$, to zero. Then \mathbf{B} factors through \mathcal{H}^{tr} . Since \mathcal{A} is equivalent to a quotient of \mathcal{U}^{tr} , we have the following commutative diagram of 2-functors:

$$(7.2) \quad \begin{array}{ccccccc} & & & & \mathcal{H} & & \\ & & & & \downarrow & \searrow \mathbf{B} & \\ \mathcal{U} & \longrightarrow & \mathcal{U}^{\text{tr}} & \longrightarrow & \mathcal{A} & \xrightarrow{\mathbf{S}} & \mathcal{H}^{\text{tr}} \longrightarrow \mathcal{C} \end{array}$$

where we identify the 2-functor $\mathbf{S}: \mathcal{A} \rightarrow \mathcal{H}_\epsilon$ with its composition with the inclusion $\mathcal{H}_\epsilon \rightarrow \mathcal{H}^{\text{tr}}$. Therefore we obtain a representation $\mathcal{U} \rightarrow \mathcal{C}$, which can be viewed as a categorical induced action. An example of this will be developed in detail in Section 7.2.

Passing to Grothendieck groups, the 2-functor \mathbf{B} induces a representation of the Heisenberg algebra H on $K(\mathcal{C})$. If this representation is irreducible then, by the Stone–von Neumann Theorem, it is isomorphic to the Fock space representation. Then the induced action of \mathcal{U} on \mathcal{C} categorifies the basic representation. It follows from the results of Section 4.3 that the categories $\mathcal{A}(\lambda, \mu)$ are semisimple (with at most one simple object) for all $\lambda, \mu \in \mathcal{P}$. Therefore, equalities of 1-morphisms in the Grothendieck group \mathcal{A} imply isomorphisms in \mathcal{A} of the corresponding 1-morphisms. It thus follows from Theorems 4.4, 4.11, and 6.7 that we have isomorphisms of 1-morphisms in \mathcal{A} corresponding to the expressions of generators of \mathfrak{sl}_∞ in terms of generators of H appearing in the principal realization (see Section 2.2). Therefore, the results of the current paper yield a categorification of the principal realization.

We note that these techniques are very different from those used in the categorification of the homogeneous realization of the basic representation in affine types ADE described in [CL11]. The main tools of [CL11] are categorical vertex operators, which are certain complexes in Heisenberg 2-representations. By contrast, our construction does not involve complexes and thus does not require passing to the homotopy category.

7.2. ACTION OF \mathcal{H}^{tr} . — We now describe an action of the 2-category \mathcal{H}^{tr} that arises from the action of Khovanov’s Heisenberg category on modules for symmetric groups described in [Kho14, §3.3].

Recall the 2-category \mathcal{M} of Section 3.1. We define an additive linear 2-functor $\mathbf{F}_{\mathcal{H}}: \mathcal{H}^{\text{tr}'} \rightarrow \mathcal{M}$ as follows. On objects,

$$(7.3) \quad \mathbf{F}_{\mathcal{H}}(\mathbf{0}) = \mathbf{0}, \quad \mathbf{F}_{\mathcal{H}}(n) = \mathcal{M}_n, \quad n \in \mathbb{N}.$$

On 1-morphisms, for $n \in \mathbb{N}$, we define,

$$(7.4) \quad \mathbf{F}_{\mathcal{H}}(\mathbf{Q}_+ 1_n) = (n+1)_n \otimes_{A_n} -: \mathcal{M}_n \rightarrow \mathcal{M}_{n+1},$$

$$(7.5) \quad \mathbf{F}_{\mathcal{H}}(\mathbf{Q}_- 1_{n+1}) = {}_n(n+1) \otimes_{A_{n+1}} -: \mathcal{M}_{n+1} \rightarrow \mathcal{M}_n.$$

We now define $\mathbf{F}_{\mathcal{H}}$ on 2-morphisms. The 2-functor $\mathbf{F}_{\mathcal{H}}$ will map 2-morphisms of $\mathcal{H}^{\text{tr}'}$ to natural transformations of functors given by tensoring with bimodules. These natural transformations are given by homomorphisms of the corresponding bimodules. We define

$$(7.6) \quad \mathbf{F}_{\mathcal{H}} \left(\begin{array}{c} \nearrow \\ \times \\ \searrow \end{array} \right)_{n-1} = R_n, \quad (7.7) \quad \mathbf{F}_{\mathcal{H}} \left(\begin{array}{c} \nwarrow \\ \times \\ \swarrow \end{array} \right)_{n-1} = L_n,$$

$$(7.8) \quad \mathbf{F}_{\mathcal{H}} \left(\begin{array}{c} \nearrow \\ \times \\ \nearrow \end{array} \right)_n = \rho, \quad (7.9) \quad \mathbf{F}_{\mathcal{H}} \left(\begin{array}{c} \nwarrow \\ \times \\ \nwarrow \end{array} \right)_n = \tau,$$

$$(7.10) \quad \mathbf{F}_{\mathcal{H}} \left(\begin{array}{c} \curvearrowright \\ \downarrow \end{array} \right)_{n+1} = \varepsilon_R, \quad (7.11) \quad \mathbf{F}_{\mathcal{H}} \left(\begin{array}{c} \cup \\ \uparrow \end{array} \right)_n = \eta_R,$$

$$(7.12) \quad \mathbf{F}_{\mathcal{H}} \left(\begin{array}{c} \curvearrowright \\ \uparrow \end{array} \right)_n = \varepsilon_L, \quad (7.13) \quad \mathbf{F}_{\mathcal{H}} \left(\begin{array}{c} \cup \\ \downarrow \end{array} \right)_{n+1} = \eta_L,$$

where

$$(7.14) \quad R_n: (n+1)_{n-1} \rightarrow (n+1)_{n-1}, \quad a \mapsto as_n,$$

$$(7.15) \quad L_n: {}_{n-1}(n+1) \rightarrow {}_{n-1}(n+1), \quad a \mapsto s_n a,$$

ρ and τ are defined in (3.12) and (3.13), respectively, and $\varepsilon_R, \eta_R, \varepsilon_L,$ and η_L are the adjunction maps defined in Proposition 3.2. It follows from the results of [Kho14, §3.3] that $\mathbf{F}_{\mathcal{H}}$ respects the local relations and topological invariance in the definition of $\mathcal{H}^{\text{tr}'}$.

Since the only idempotent 1-morphisms in \mathcal{M} are the identity 1-morphisms and all idempotent 2-morphisms in \mathcal{M} split, the 2-functor $\mathbf{F}_{\mathcal{H}}$ induces a 2-functor (which we denote by the same symbol)

$$\mathbf{F}_{\mathcal{H}}: \mathcal{H}^{\text{tr}} \rightarrow \mathcal{M}.$$

LEMMA 7.1. — *The 2-functor $\mathbf{F}_{\mathcal{H}}$ maps the 2-morphism ϵ_λ to the bimodule map $(n) \rightarrow (n)$ given by multiplication by the central idempotent e_λ .*

Proof. — We compute that $\mathbf{F}_{\mathcal{H}}(\epsilon_\lambda)$ is the bimodule map $(n) \rightarrow (n)$ given by

$$a \mapsto a \frac{1}{n!} \sum_{w \in S_n} we_\lambda w^{-1} = ae_\lambda. \quad \square$$

COROLLARY 7.2. — *The 2-functor $\mathbf{F}_{\mathcal{H}}$ maps the object $(n, 1_n, \epsilon_\lambda)$ of \mathcal{H}^{tr} to \mathcal{M}_λ and maps \mathcal{H}_δ to zero.*

7.3. ACTION OF \mathcal{A} . — The composition

$$\mathbf{F}_{\mathcal{A}} := \mathbf{F}_{\mathcal{H}} \circ \mathbf{S}$$

is an additive linear 2-functor that defines an action of \mathcal{A} on modules for symmetric groups. For future reference, we describe this 2-functor explicitly here.

On objects, we have

$$(7.16) \quad \mathbf{F}_{\mathcal{A}}(\mathbf{0}) = \mathbf{0}, \quad \mathbf{F}_{\mathcal{A}}(\lambda) = \mathcal{M}_{\lambda}, \quad \lambda \in \mathcal{P}.$$

On 1-morphisms, for $\lambda \in \mathcal{P}$ with an addable i -box, it follows from (3.20) that

$$(7.17) \quad \mathbf{F}_{\mathcal{A}}(\mathbf{F}_i 1_{\lambda}) = (n+1)_n^i \otimes_{A_n} -: \mathcal{M}_{\lambda} \longrightarrow \mathcal{M}_{\lambda \boxplus i}, \quad \text{where } n = |\lambda|.$$

Similarly, for $\lambda \in \mathcal{P}$ with a removable i -box, it follows from (3.21) that

$$(7.18) \quad \mathbf{F}_{\mathcal{A}}(\mathbf{E}_i 1_{\lambda}) = {}_{n-1}^i(n) \otimes_{A_n} -: \mathcal{M}_{\lambda} \longrightarrow \mathcal{M}_{\lambda \boxminus i}, \quad \text{where } n = |\lambda|.$$

We now describe $\mathbf{F}_{\mathcal{A}}$ on 2-morphisms. The 2-functor $\mathbf{F}_{\mathcal{A}}$ maps 2-morphisms of \mathcal{A} to natural transformations of functors given by tensoring with bimodules. These natural transformations are given by homomorphisms of the corresponding bimodules. For $i, j \in \mathbb{Z}$ with $i \neq j$, and $\lambda \vdash n$, we have

$$(7.19) \quad \mathbf{F}_{\mathcal{A}} \left(\begin{array}{c} \nearrow \lambda \\ i \quad j \end{array} \right) = R_{n+1}|_{i,j}^{j,i}, \quad (7.20) \quad \mathbf{F}_{\mathcal{A}} \left(\begin{array}{c} \searrow \lambda \\ i \quad j \end{array} \right) = L_{n-1}|_{i,j}^{j,i},$$

$$(7.21) \quad \mathbf{F}_{\mathcal{A}} \left(\begin{array}{c} i \quad \nearrow \lambda \\ j \quad \searrow \end{array} \right) = \frac{i-j}{i-j-1} \frac{(n+1)d_{\lambda}d_{\lambda \boxplus i \boxplus j}}{|\lambda|d_{\lambda \boxplus i}d_{\lambda \boxplus j}} \rho|_{j,i}^{i,j},$$

$$(7.22) \quad \mathbf{F}_{\mathcal{A}} \left(\begin{array}{c} i \quad \searrow \lambda \\ j \quad \nearrow \end{array} \right) = \frac{i-j}{i-j-1} \tau|_{j,i}^{i,j},$$

$$(7.23) \quad \mathbf{F}_{\mathcal{A}} \left(\begin{array}{c} \curvearrowright \lambda \\ i \end{array} \right) = \frac{d_{\lambda}}{nd_{\lambda \boxplus i}} \varepsilon_{\mathbb{R}}^{i,i}, \quad (7.24) \quad \mathbf{F}_{\mathcal{A}} \left(\begin{array}{c} \curvearrowleft \lambda \\ i \end{array} \right) = \frac{(n+1)d_{\lambda}}{d_{\lambda \boxplus i}} \eta_{\mathbb{R}}^{i,i},$$

$$(7.25) \quad \mathbf{F}_{\mathcal{A}} \left(\begin{array}{c} \curvearrowright \lambda \\ i \end{array} \right) = \varepsilon_{\mathbb{L}}^{i,i}, \quad (7.26) \quad \mathbf{F}_{\mathcal{A}} \left(\begin{array}{c} \curvearrowleft \lambda \\ i \end{array} \right) = \eta_{\mathbb{L}}^{i,i}.$$

where, for $i, j, k, \ell \in \mathbb{Z}$, we define the components

$$(7.27) \quad R_{n+1}|_{i,j}^{k,\ell} : (n+2)_n^{i,j} \hookrightarrow (n+2)_n \xrightarrow{R_{n+1}} (n+2)_n \twoheadrightarrow (n+2)_n^{k,\ell},$$

$$(7.28) \quad L_{n-1}|_{i,j}^{k,\ell} : {}_{n-2}^{i,j}(n) \hookrightarrow {}_{n-2}(n) \xrightarrow{L_{n-1}} {}_{n-2}(n) \twoheadrightarrow {}_{n-2}^{k,\ell}(n),$$

$$(7.29) \quad \rho|_{j,i}^{k,\ell} : (n)_{n-1}^j \hookrightarrow (n)_{n-1}(n) \xrightarrow{\rho} {}_n(n+1)_n \twoheadrightarrow {}_n^{k,\ell}(n+1)_n^{\ell},$$

$$(7.30) \quad \tau_{j,i}^{k,\ell} : {}_n^j(n+1)_n^i \hookrightarrow {}_n(n+1)_n \xrightarrow{\tau} (n)_{n-1}(n) \twoheadrightarrow (n)_{n-1}^k \ell(n),$$

$$(7.31) \quad \varepsilon_{\mathbf{R}}^{i,j} : (n+1)_n^j(n+1) \hookrightarrow (n+1)_n(n+1) \xrightarrow{\varepsilon_{\mathbf{R}}} (n+1),$$

$$(7.32) \quad \eta_{\mathbf{R}}^{i,j} : (n) \xrightarrow{\eta_{\mathbf{R}}} {}_n(n+1)_n \twoheadrightarrow {}_n^i(n+1)_n^j,$$

$$(7.33) \quad \varepsilon_{\mathbf{L}}^{i,j} : {}_n^i(n+1)_n^j \hookrightarrow {}_n(n+1)_n \xrightarrow{\varepsilon_{\mathbf{L}}} (n),$$

$$(7.34) \quad \eta_{\mathbf{L}}^{i,j} : (n+1) \xrightarrow{\eta_{\mathbf{L}}} (n+1)_n(n+1) \twoheadrightarrow (n+1)_n^i j(n+1).$$

Note that any time the denominator in a coefficient in (7.21)–(7.24) is zero, the corresponding diagram is zero and so we can ignore such expressions. For instance, in (7.21), when $i = j + 1$, the crossing of strands colored i and j is zero (see Remark 4.1).

It is possible to write the equations (7.19)–(7.22) in a manner that avoids one of the eigenspace projections. For example, we have

$$(7.35) \quad \mathbf{F}_{\mathcal{A}} \left(\begin{array}{c} \nearrow \lambda \\ \searrow i \quad \nearrow j \end{array} \right) = \mathbf{F}_{\mathcal{H}} \left(\begin{array}{c} \nearrow \epsilon_{\lambda \boxplus i} \\ \searrow \epsilon_{\lambda \boxplus i \boxplus j} \quad \nearrow \epsilon_{\lambda} \\ \searrow \epsilon_{\lambda \boxplus j} \end{array} \right) = \mathbf{F}_{\mathcal{H}} \left(\begin{array}{c} \nearrow \epsilon_{\lambda \boxplus i \boxplus j} \quad \nearrow \epsilon_{\lambda} \\ \searrow \epsilon_{\lambda \boxplus j} \end{array} - \begin{array}{c} \nearrow \epsilon_{\lambda \boxplus j} \\ \searrow \epsilon_{\lambda \boxplus j} \quad \nearrow \epsilon_{\lambda} \end{array} \right) \\ = \frac{i-j}{i-j-1} R_{n+1} {}^r(e_{\lambda \boxplus i \boxplus j} e_{\lambda \boxplus j} e_{\lambda}) - \frac{1}{i-j-1},$$

where ${}^r z$ denotes right multiplication by an element z , so that ${}^r(e_{\lambda \boxplus i \boxplus j} e_{\lambda \boxplus j} e_{\lambda})$ is projection onto $(n+2)_n^{i,j} \subseteq (n+2)_n$.

PROPOSITION 7.3. — *The 2-functor $\mathbf{F}_{\mathcal{A}} = \mathbf{F}_{\mathcal{H}} \circ \mathbf{S} : \mathcal{A} \rightarrow \mathcal{M}$ is an equivalence of 2-categories.*

Proof. — By definition, $\mathbf{F}_{\mathcal{A}}$ is essentially surjective on objects. Consider $\lambda, \mu \in \mathcal{P}$. By Proposition 4.6 any 1-morphism in $1\text{Mor}_{\mathcal{A}}(\lambda, \mu)$ is isomorphic to a multiple of one of the form $P = F_{i_1} F_{i_2} \cdots F_{i_k} E_{j_1} E_{j_2} \cdots E_{j_\ell} 1_\lambda$.

Recall that we have canonical equivalences $\mathcal{M}_\lambda \cong \mathcal{V}$ and $\mathcal{M}_\mu \cong \mathcal{V}$ (see Section 3.1). Under these equivalences, $1\text{Mor}_{\mathcal{M}}(\mathcal{M}_\lambda, \mathcal{M}_\mu) = \{1_{\mathcal{V}}^{\oplus n} : n \geq 0\}$, and $\mathbf{F}_{\mathcal{A}}(P) = 1_{\mathcal{V}}$. It follows that $\mathbf{F}_{\mathcal{A}}$ is essentially full on 1-morphisms.

Given $P, Q \in 1\text{Mor}_{\mathcal{A}}(\lambda, \mu)$ two nonzero 1-morphisms as above, by Proposition 4.9 we have that $2\text{Mor}_{\mathcal{A}}(P, Q)$ is one-dimensional. Since $2\text{Mor}(1_{\mathcal{V}}, 1_{\mathcal{V}})$ is also one-dimensional and $\mathbf{F}_{\mathcal{A}}$ preserves 2-isomorphisms, it follows that $\mathbf{F}_{\mathcal{A}}$ induces an isomorphism

$$2\text{Mor}_{\mathcal{A}}(P, Q) \cong 2\text{Mor}_{\mathcal{M}}(\mathbf{F}_{\mathcal{A}}(P), \mathbf{F}_{\mathcal{A}}(Q)).$$

By linearity we conclude that $\mathbf{F}_{\mathcal{A}}$ is fully faithful on 2-morphisms. Thus $\mathbf{F}_{\mathcal{A}}$ is an equivalence. □

We can now complete the proofs of Theorems 6.7 and 4.11.

COROLLARY 7.4. — *The 2-functor \mathbf{S} is faithful on 2-morphisms.*

COROLLARY 7.5. — *The functor $\mathcal{A} \rightarrow K(\mathcal{A})$ of Theorem 4.11 is faithful.*

Proof. — We have commutative diagram

$$(7.36) \quad \begin{array}{ccc} \mathcal{A} & \xrightarrow{\mathbf{r}} & \mathcal{V} \\ \downarrow & & \uparrow \\ K(\mathcal{A}) & \xrightarrow{[\mathbf{F}_{\mathcal{A}}]} & K(\mathcal{M}) \end{array}$$

Thus, the corollary follows from the fact that the functor \mathbf{r} of (2.1) is faithful. \square

7.4. ACTION OF CATEGORIFIED QUANTUM GROUPS. — From the results of Sections 4.2 and 7.3, we immediately obtain an explicit action of the 2-category \mathcal{U} of [CL15] (the categorified quantum group of type A_∞) on modules for symmetric groups. This categorifies the fundamental representation $L(\Lambda_0)$ of \mathfrak{sl}_∞ .

For ease of reference, we describe this action here, which is an additive linear 2-functor

$$\mathbf{F}_{\mathcal{U}} : \mathcal{U} \longrightarrow \mathcal{M}.$$

Recall that the set of objects of \mathcal{U} is the free monoid on the weight lattice of \mathfrak{sl}_∞ and recall the definition of ω_λ in (2.3). On objects, we have

$$\mathbf{F}_{\mathcal{U}}(x) = \begin{cases} \mathcal{M}_\lambda & \text{if } x = \omega_\lambda, \lambda \in \mathcal{P}, \\ \mathbf{0} & \text{if } x \text{ is not of the form } \omega_\lambda \text{ for any } \lambda \in \mathcal{P}. \end{cases}$$

On 1-morphisms, $\mathbf{F}_{\mathcal{U}}$ acts just as $\mathbf{F}_{\mathcal{A}}$ does in (7.17) and (7.18). On 2-morphisms, $\mathbf{F}_{\mathcal{U}}$ maps any diagram with dots to zero. On diagrams without dots, $\mathbf{F}_{\mathcal{U}}$ acts just as $\mathbf{F}_{\mathcal{A}}$ does in (7.19)–(7.26), but with orientations of strands reversed (see Theorem 4.4).

REMARK 7.6. — In [BK09a], Brundan and Kleshchev constructed an explicit isomorphism between blocks of cyclotomic Hecke algebras and sign-modified cyclotomic Khovanov–Lauda algebras in type A . They then used this isomorphism to describe actions on categories of modules for cyclotomic Hecke algebras in [BK09b]. This is related to the action described above, using (7.19)–(7.26), since level one cyclotomic Hecke algebras are isomorphic to group algebras of symmetric groups.

8. APPLICATIONS AND FURTHER DIRECTIONS

8.1. DIAGRAMMATIC COMPUTATION. — As an application of the constructions of the current paper, we give some examples of how one can prove combinatorial identities related to the dimensions of modules for symmetric groups using the diagrammatics of the categories introduced above.

PROPOSITION 8.1. — *If λ is a partition, then*

$$(8.1) \quad \sum_{j \in B^+(\lambda)} \frac{d_{\lambda \boxplus j}}{j - i} = 0 \quad \forall i \in B^-(\lambda), \quad \text{and}$$

$$(8.2) \quad \sum_{j \in B^-(\lambda)} \frac{d_{\lambda \boxplus j}}{i - j} = i \frac{d_\lambda}{|\lambda|} \quad \forall i \in B^+(\lambda).$$

Proof. — For $i \in B^-(\lambda)$, we have

$$\begin{aligned}
 0 \stackrel{(5.6)}{=} \begin{array}{c} \uparrow \\ \textcircled{\epsilon_\lambda} \\ \epsilon_\lambda \end{array} \epsilon_{\lambda \boxplus i} &= \sum_{j \in B^+(\lambda)} \begin{array}{c} \epsilon_{\lambda \boxplus j} \\ \textcircled{\epsilon_\lambda} \\ \epsilon_\lambda \end{array} \epsilon_{\lambda \boxplus i} \stackrel{(6.7)}{=} \sum_{j \in B^+(\lambda)} \frac{1}{j-i} \begin{array}{c} \epsilon_{\lambda \boxplus j} \\ \textcircled{\epsilon_\lambda} \\ \epsilon_\lambda \end{array} \epsilon_{\lambda \boxplus i} \\
 &\stackrel{(6.4)}{=} \frac{1}{(n+1)d_\lambda} \sum_{j \in B^+(\lambda)} \frac{d_{\lambda \boxplus j}}{j-i} \begin{array}{c} \epsilon_\lambda \\ \uparrow \\ \epsilon_{\lambda \boxplus i} \end{array} .
 \end{aligned}$$

Since the final diagram above is nonzero (it is sent to i -induction from $\mathcal{M}_{\lambda \boxplus i}$ to \mathcal{M}_λ under the 2-functor $\mathbf{F}_{\mathcal{H}}$), relation (8.1) follows.

Now suppose $i \in B^+(\lambda)$. Then

$$\begin{aligned}
 i \left(\begin{array}{c} \uparrow \\ \epsilon_{\lambda \boxplus i} \\ \epsilon_\lambda \end{array} \right) &\stackrel{(6.6)}{=} \begin{array}{c} \epsilon_{\lambda \boxplus i} \\ \textcircled{\epsilon_\lambda} \\ \epsilon_\lambda \end{array} = \sum_{j \in B^-(\lambda)} \begin{array}{c} \epsilon_{\lambda \boxplus i} \\ \textcircled{\epsilon_{\lambda \boxplus j}} \\ \epsilon_\lambda \end{array} \stackrel{(6.7)}{=} \sum_{j \in B^-(\lambda)} \frac{1}{i-j} \begin{array}{c} \epsilon_{\lambda \boxplus i} \\ \textcircled{\epsilon_\lambda} \\ \epsilon_\lambda \end{array} \\
 &\stackrel{(6.3)}{=} \sum_{j \in B^-(\lambda)} \frac{|\lambda| d_{\lambda \boxplus j}}{d_\lambda (i-j)} \begin{array}{c} \epsilon_{\lambda \boxplus i} \\ \uparrow \\ \epsilon_\lambda \end{array} .
 \end{aligned}$$

Since the final diagram above is nonzero (as above), relation (8.2) follows. □

It is possible to prove the identities (8.1) and (8.2) algebraically, using a careful analysis of the representation theory of the symmetric group. However, such a proof is considerably longer than the above diagrammatic one. To the best of the authors' knowledge, these identities have not appeared previously in the literature. It would be interesting to find purely combinatorial proofs.

8.2. FURTHER DIRECTIONS. — The results of the current paper suggest a number of future research directions. We briefly describe of few of these here.

8.2.1. *Symmetric groups in positive characteristic.* — In light of Proposition 7.3, the 2-category \mathcal{A} can be viewed as a graphical calculus describing the functors of i -induction and i -restriction, together with the natural transformations between them. Throughout this paper, we have worked over the field \mathbb{Q} . It would be natural to instead consider the representation theory of the symmetric group in characteristic $p > 0$. We believe that most of the results presented here have positive characteristic analogues that would yield a relationship between categorified quantum $\widehat{\mathfrak{sl}}_p$ (instead of \mathfrak{sl}_∞) and Heisenberg categories that categorifies the principal embedding. We refer

the reader to the survey [Kle14] for an overview of the modular representation theory of the symmetric group in the context of categorification.

8.2.2. Cyclotomic Hecke algebras. — Group algebras of symmetric groups are isomorphic to level one degenerate cyclotomic Hecke algebras. It is natural to expect that the results of the current paper can be extended to higher level degenerate cyclotomic Hecke algebras. On the Heisenberg side, this corresponds to the higher level Heisenberg categories defined in [MS17], which involve planar diagrams decorated by dots corresponding to the polynomial generators of the degenerate cyclotomic Hecke algebras. On the categorified quantum group side, this should correspond to modifying the definition of \mathcal{U}_0 (see Section 4.2) so as not to kill all dots. This would be related to the results of [BK09a, BK09b].

8.2.3. More general Heisenberg categories. — The Heisenberg category considered here is a special case of a much more general construction, described in [RS17], that associates a Heisenberg category (or 2-category) to any graded Frobenius superalgebra. (The Heisenberg 2-category considered in the current paper corresponds to the case where this Frobenius algebra is simply the base field.) It would be interesting to generalize the results of the current paper to these more general Heisenberg categories. Representation theoretically, this amounts to replacing the group algebra of the symmetric group by wreath product algebras associated to the Frobenius algebra in question. Of special interest would be the case where the Frobenius algebra is the zigzag algebra associated to a finite-type Dynkin diagram, in which case the corresponding Heisenberg categories are the ones considered in [CL12].

A q -deformation of Khovanov’s category was also defined in [LS13]. This deformation corresponds to replacing group algebras of symmetric groups by Hecke algebras of type A . One could form a truncated q -deformed Heisenberg 2-category and attempt to relate such a truncation to q -deformations of categorified quantum groups.

8.2.4. Trace decategorification. — In contrast to passing to the Grothendieck group, there is another natural method of decategorification: taking the trace or zeroth Hochschild homology. The trace of Khovanov’s Heisenberg category has been related to W -algebras in [CLLS16]. On the other hand, traces of categorified quantum groups have been related to current algebras in [BHLW17, SVV17]. It would be interesting to investigate the relationship between these two trace decategorifications implied by the results of the current paper and their generalizations mentioned above.

8.2.5. Geometry. — Heisenberg categories are closely related to the geometry of the Hilbert scheme (see [CL12]). Similarly, the geometry of quiver varieties [Nak98] can be used to build categorifications of quantum group representations (see for example [VV11, Zhe14, CKL13, Web12]). It is thus natural to expect that the results of the current paper are related to geometric constructions relating these spaces, such as [Sav06, Nag09, LS10, Lem16].

REFERENCES

- [BHLW17] A. BELIAKOVA, K. HABIRO, A. D. LAUDA & B. WEBSTER – “Current algebras and categorified quantum groups”, *J. London Math. Soc. (2)* **95** (2017), no. 1, p. 248–276.
- [BK09a] J. BRUNDAN & A. KLESHCHEV – “Blocks of cyclotomic Hecke algebras and Khovanov-Lauda algebras”, *Invent. Math.* **178** (2009), no. 3, p. 451–484.
- [BK09b] ———, “Graded decomposition numbers for cyclotomic Hecke algebras”, *Adv. Math.* **222** (2009), no. 6, p. 1883–1942.
- [CKL13] S. CAUTIS, J. KAMNITZER & A. LICATA – “Coherent sheaves on quiver varieties and categorification”, *Math. Ann.* **357** (2013), no. 3, p. 805–854.
- [CL15] S. CAUTIS & A. D. LAUDA – “Implicit structure in 2-representations of quantum groups”, *Selecta Math. (N.S.)* **21** (2015), no. 1, p. 201–244.
- [CLLS16] S. CAUTIS, A. D. LAUDA, A. LICATA & J. SUSSAN – “W-algebras from Heisenberg categories”, *J. Inst. Math. Jussieu* (2016), online, doi:10.1017/S1474748016000189.
- [CL11] S. CAUTIS & A. LICATA – “Vertex operators and 2-representations of quantum affine algebras”, arXiv:1112.6189, 2011.
- [CL12] ———, “Heisenberg categorification and Hilbert schemes”, *Duke Math. J.* **161** (2012), no. 13, p. 2469–2547.
- [CR08] J. CHUANG & R. ROUQUIER – “Derived equivalences for symmetric groups and \mathfrak{sl}_2 -categorification”, *Ann. of Math. (2)* **167** (2008), no. 1, p. 245–298.
- [FH91] W. FULTON & J. HARRIS – *Representation theory*, Graduate Texts in Math., vol. 129, Springer-Verlag, New York, 1991.
- [Kac90] V. G. KAC – *Infinite-dimensional Lie algebras*, third ed., Cambridge University Press, Cambridge, 1990.
- [Kho14] M. KHOVANOV – “Heisenberg algebra and a graphical calculus”, *Fund. Math.* **225** (2014), no. 1, p. 169–210.
- [KL10] M. KHOVANOV & A. D. LAUDA – “A categorification of quantum $\mathfrak{sl}(n)$ ”, *Quantum Topol.* **1** (2010), no. 1, p. 1–92.
- [Kle05] A. KLESHCHEV – *Linear and projective representations of symmetric groups*, Cambridge Tracts in Mathematics, vol. 163, Cambridge University Press, Cambridge, 2005.
- [Kle14] ———, “Modular representation theory of symmetric groups”, arXiv:1405.3326, 2014.
- [Lem16] J. LEMAY – “Geometric realizations of the basic representation of $\widehat{\mathfrak{gl}}_r$ ”, *Selecta Math. (N.S.)* **22** (2016), no. 1, p. 341–387.
- [LRS] A. LICATA, D. ROSSO & A. SAVAGE – “Categorification and Heisenberg doubles arising from towers of algebras”, *J. Combinatorial Theory Ser. A*, to appear, arXiv:1610.01862.
- [LS10] A. LICATA & A. SAVAGE – “Vertex operators and the geometry of moduli spaces of framed torsion-free sheaves”, *Selecta Math. (N.S.)* **16** (2010), no. 2, p. 201–240.
- [LS13] ———, “Hecke algebras, finite general linear groups, and Heisenberg categorification”, *Quantum Topol.* **4** (2013), no. 2, p. 125–185.
- [MS17] M. MACKAAY & A. SAVAGE – “Degenerate cyclotomic Hecke algebras and higher level Heisenberg categorification”, arXiv:1705.03066, 2017.
- [MSV13] M. MACKAAY, M. STOŠIĆ & P. VAZ – “A diagrammatic categorification of the q -Schur algebra”, *Quantum Topol.* **4** (2013), no. 1, p. 1–75.
- [Nag09] K. NAGAO – “Quiver varieties and Frenkel-Kac construction”, *J. Algebra* **321** (2009), no. 12, p. 3764–3789.
- [Nak98] H. NAKAJIMA – “Quiver varieties and Kac-Moody algebras”, *Duke Math. J.* **91** (1998), no. 3, p. 515–560.
- [QR16] H. QUEFFelec & D. E. V. ROSE – “The \mathfrak{sl}_n foam 2-category: A combinatorial formulation of Khovanov–Rozansky homology via categorical skew Howe duality”, *Adv. Math.* **302** (2016), p. 1251–1339.
- [RS17] D. ROSSO & A. SAVAGE – “A general approach to Heisenberg categorification via wreath product algebras”, *Math. Z.* **286** (2017), no. 1–2, p. 603–655.
- [Rou08] R. ROUQUIER – “2-Kac-Moody algebras”, arXiv:0812.5023v1, 2008.
- [Sav06] A. SAVAGE – “A geometric boson-fermion correspondence”, *C. R. Math. Rep. Acad. Sci. Canada* **28** (2006), no. 3, p. 65–84.

- [SVV17] P. SHAN, M. VARAGNOLO & E. VASSEROT – “On the center of quiver Hecke algebras”, *Duke Math. J.* **166** (2017), no. 6, p. 1005–1101.
- [VV11] M. VARAGNOLO & E. VASSEROT – “Canonical bases and KLR-algebras”, *J. reine angew. Math.* **659** (2011), p. 67–100.
- [Web12] B. WEBSTER – “A categorical action on quantized quiver varieties”, [arXiv:1208.5957](https://arxiv.org/abs/1208.5957), 2012.
- [Zhe14] H. ZHENG – “Categorification of integrable representations of quantum groups”, *Acta Mech. Sinica (English Ed.)* **30** (2014), no. 6, p. 899–932.

Manuscript received April 11, 2017
accepted December 4, 2017

HOEL QUEFFELEC, Institut Montpellierain Alexander Grothendieck, Université de Montpellier, CNRS
Case courrier 051, Place Eugène Bataillon, 34095 Montpellier Cedex 5, France
E-mail : hoel.queffelec@umontpellier.fr
Url : <http://www.math.univ-montp2.fr/~queffelec/>

ALISTAIR SAVAGE, Department of Mathematics and Statistics, University of Ottawa
585 King Edward Ave, Ottawa, Ontario, Canada K1N 6N5
E-mail : alistair.savage@uottawa.ca
Url : <http://alistairsavage.ca>

ODED YACOBI, School of Mathematics and Statistics
University of Sydney NSW 2006, Australia
E-mail : oded.yacobi@sydney.edu.au
Url : <http://www.maths.usyd.edu.au/u/oyacobi/>